

ASSOCIATE EDITOR: MARTIN C. MICHEL

Brothers in Arms: ABCA1- and ABCG1-Mediated Cholesterol Efflux as Promising Targets in Cardiovascular Disease Treatment[§]

Sanne J. C. M. Frambach, Ria de Haas, Jan A. M. Smeitink, Gerard A. Rongen, Frans G. M. Russel, and Tom J. J. Schirris

Department of Pharmacology and Toxicology, Radboud Institute for Molecular Life Sciences (S.J.C.M.F., G.A.R., F.G.M.R., T.J.J.S.), Radboud Center for Mitochondrial Medicine (S.J.C.M.F., R.d.H., J.A.M.S., F.G.M.R., T.J.J.S.), Department of Pediatrics (R.d.H., J.A.M.S.), and Department of Internal Medicine, Radboud Institute for Health Sciences (G.A.R.), Radboud University Medical Center, Nijmegen, The Netherlands

Abstract	153
Significance Statement	153
I. Introduction	153
A. The Importance of Plasma Cholesterol for Cardiovascular Disease	153
B. Statins as the Cornerstone of Cardiovascular Disease Treatment.....	155
C. High-Density Lipoprotein and Cardiovascular Disease: High-Density Lipoprotein Levels Are a Poor Reflection of Reverse Cholesterol Transport Capacity	155
D. Systematic Review: Scope and Methodology.....	156
II. Reverse Cholesterol Transport Pathway	156
A. Initiation and Propagation of Reverse Cholesterol Transport	156
B. ATP-Binding Cassette A1 and ATP-Binding Cassette G1 as Master Effectors of Cholesterol Efflux.....	158
C. Regulation of ATP-Binding Cassette A1- and ATP-Binding Cassette G1-Mediated Cholesterol Efflux.....	161
III. Apolipoprotein A-I and Apolipoprotein E Mimetics.....	161
IV. Regulation and Pharmacological Manipulation of Nuclear Receptor-Mediated ATP-Binding Cassette A1 and ATP-Binding Cassette G1 Expression	163
A. Nuclear Receptors Are Important Regulators of ATP-Binding Cassette A1 and ATP-Binding Cassette G1 Expression.....	163
B. Liver X Receptor Activation to Induce ATP-Binding Cassette A1 and ATP-Binding Cassette G1 Expression	164
C. Peroxisome Proliferator-Activated Receptor Activation to Enhance ATP-Binding Cassette A1 and ATP-Binding Cassette G1 Expression.....	169
D. Enhancement of ATP-Binding Cassette A1 and ATP-Binding Cassette G1 Expression by Retinoid X Receptor Agonists.....	173
V. ATP-Binding Cassette A1 and ATP-Binding Cassette G1 mRNA Stability.....	173
A. mRNA Degradation as a Post-Translational Mechanism to Regulate ATP-Binding Cassette A1 and ATP-Binding Cassette G1 Expression.....	173
B. Targeting Post-Transcriptional Regulation of ATP-Binding Cassette A1 mRNA.....	174
VI. ATP-Binding Cassette A1 and ATP-Binding Cassette G1 Protein Degradation as a Target to Increase Cholesterol Efflux	175

Address correspondence to: Dr. Frans G.M. Russel, Department of Pharmacology and Toxicology, Radboud University Medical Center, PO Box 9101, 6500 HB Nijmegen, The Netherlands. E-mail: frans.russel@radboudumc.nl or Dr. Tom J.J. Schirris, Department of Pharmacology and Toxicology, Radboud University Medical Center, PO Box 9101, 6500 HB Nijmegen, The Netherlands. E-mail: tom.schirris@radboudumc.nl

F.G.M.R. and T.J.J.S. contributed equally as last authors.

This work was supported by the Princes Beatrix Muscle Foundation (Prinses Beatrix Spierfonds) [Grant WO.R16-19].

J.A.M.S. holds a partial position at Khondrion, a Radboud University Medical Center spin-out company founded by J.A.M.S.

<https://doi.org/10.1124/pr.119.017897>.

[§]This article has supplemental material available at pharmrev.aspetjournals.org.

A.	The Role of ATP-Binding Cassette A1 and ATP-Binding Cassette G1 Reuptake and Degradation in the Regulation of Their Plasma Membrane Abundance	175
B.	Pharmacological Inhibition of ATP-Binding Cassette A1 and ATP-Binding Cassette G1 Protein Degradation	176
VII.	ATP-Binding Cassette A1 Function and Cyclic Adenosine Monophosphate.....	179
A.	cAMP Is a Potent Regulator of ATP-Binding Cassette A1 Function and Expression.....	179
B.	Stimulation of cAMP Levels Enhances ATP-Binding Cassette A1-Mediated Cholesterol Efflux	180
VIII.	Increasing Cellular Cholesterol Efflux via Unknown Mechanisms	180
IX.	Conclusions and Future Directions	181
	References.....	183

Abstract—Atherosclerosis is a leading cause of cardiovascular disease worldwide, and hypercholesterolemia is a major risk factor. Preventive treatments mainly focus on the effective reduction of low-density lipoprotein cholesterol, but their therapeutic value is limited by the inability to completely normalize atherosclerotic risk, probably due to the disease complexity and multifactorial pathogenesis. Consequently, high-density lipoprotein cholesterol gained much interest, as it appeared to be cardioprotective due to its major role in reverse cholesterol transport (RCT). RCT facilitates removal of cholesterol from peripheral tissues, including atherosclerotic plaques, and its subsequent hepatic clearance into bile. Therefore, RCT is expected to limit plaque formation and progression. Cellular cholesterol efflux is initiated and propagated by the ATP-binding cassette (ABC) transporters ABCA1 and ABCG1. Their expression and function are expected to be rate-limiting for cholesterol efflux, which makes them interesting targets to stimulate RCT and lower atherosclerotic risk. This systematic review discusses the molecular mechanisms relevant for RCT and ABCA1 and ABCG1 function, followed by a critical overview of potential pharmacological

strategies with small molecules to enhance cellular cholesterol efflux and RCT. These strategies include regulation of ABCA1 and ABCG1 expression, degradation, and mRNA stability. Various small molecules have been demonstrated to increase RCT, but the underlying mechanisms are often not completely understood and are rather unspecific, potentially causing adverse effects. Better understanding of these mechanisms could enable the development of safer drugs to increase RCT and provide more insight into its relation with atherosclerotic risk.

Significance Statement—Hypercholesterolemia is an important risk factor of atherosclerosis, which is a leading pathological mechanism underlying cardiovascular disease. Cholesterol is removed from atherosclerotic plaques and subsequently cleared by the liver into bile. This transport is mediated by high-density lipoprotein particles, to which cholesterol is transferred via ATP-binding cassette transporters ABCA1 and ABCG1. Small-molecule pharmacological strategies stimulating these transporters may provide promising options for cardiovascular disease treatment.

I. Introduction

A. The Importance of Plasma Cholesterol for Cardiovascular Disease

Cardiovascular disease (CVD) comprises a wide range of disorders, including myocardial infarction and stroke, for which atherosclerosis is the major pathologic mechanism. CVD is associated with severe morbidity and one of the leading causes of mortality worldwide, with 17.5 million annual deaths accounting for almost

one-third of all deaths (WHO, 2008, 2014; Taylor et al., 2011). The etiology of atherosclerosis is complex and multifactorial, but hypercholesterolemia and particularly increased low-density lipoprotein (LDL) cholesterol (LDL-C) levels are acknowledged as major risk factors (Levine et al., 1995; De Backer et al., 2003; Taylor et al., 2011; Ridker, 2014; Piepoli et al., 2016). Together with very LDL (VLDL) and intermediate-density lipoprotein particles, cholesterol- and triglyceride-loaded LDL particles facilitate lipid and cholesterol transport from the

ABBREVIATIONS: ABC, ATP-binding cassette; AC, adenylate cyclase; AICAR, 5-aminoimidazole-4-carboxyamide ribonucleoside; AMPK, AMP-activated protein kinase; apo, apolipoprotein; CETP, cholesteryl ester transfer protein; CVD, cardiovascular disease; DMHCA, *N,N*-dimethyl-3 β -hydroxycholeamide; ERK, extracellular signal-regulated kinase; GPR, G protein-coupled receptor; HDL, high-density lipoprotein; HDL-C, HDL cholesterol; HMG-CoA, 3-hydroxy-3-methylglutaryl-coenzyme A; HO-1, heme oxygenase-1; IDL, intermediate density lipoprotein; IMM-H007, triacetyl-3 hydroxyphenyl-adenosine; LDL, low-density lipoprotein; LDL-C, LDL cholesterol; LDLR, LDL receptor; LXR, liver X receptor; LXRE, LXR response element; MAPK, mitogen-activated protein kinase; MC1-R, melanocortin 1 receptor; miRNA, microRNA; NBD, nucleotide binding domain; Nrf, nuclear factor-like; ox-LDL, oxidized LDL; PDE, phosphodiesterase; PEST, Pro-Glu-Ser-Thr; PKA, protein kinase A; PKC, protein kinase C; PKD, protein kinase D; PPAR, peroxisome proliferator-activated receptor; RA, retinoic acid; RAR, retinoid-activated receptor; RCT, reverse cholesterol transport; RXR, retinoid X receptor; SR-B1, scavenger receptor B1; SREBP-1c, sterol regulatory-binding element protein 1c; TMD, transmembrane domain; TNF, tumor necrosis factor; TNPB, 4-[(E)-2-(5,6,7,8-Tetrahydro-5,5,8,8-tetramethyl-2-naphthalenyl)-1-propenyl]benzoic acid; VLDL, very LDL.

liver to peripheral tissues (Hegele, 2009; Kingwell et al., 2014) (Fig. 1). Hepatic lipases and lipoprotein lipases mediate the formation of LDL particles from VLDL particles, the latter containing high triglycerides, but

moderate cholesterol and phospholipid concentrations (Fig. 1, right panels), whereas LDL particles contain high cholesterol and moderate phospholipid concentrations and almost no triglycerides (Kwiterovich, 2000;




Fig. 1. Cholesterol uptake, distribution, and peripheral utilization. (I) Schematic overview illustrating intestinal cholesterol (light yellow spheres) uptake by enterocytes, including cholesterol esterification and apoB-48 binding, which facilitates cholesterol efflux to the lymphatic system. (II) The resulting nascent chylomicrons are subsequently transported to the blood, where they collide with HDL particles that transfer apoE and apoC-II to chylomicrons, leading to their maturation. (III) Mature chylomicrons can be converted to chylomicron remnants by endothelial lipoprotein lipase (LPL) activated by apoC-II, which releases free fatty acids (yellow spheres) for uptake in peripheral tissues (e.g., muscles and fat) and apoC-II for translocation to HDL. (IV) Next, chylomicron remnants are imported into hepatocytes via the chylomicron remnant receptor (CRR), which releases the remaining cholesterol and apoE by lysosomal degradation. The resulting hepatic free cholesterol, which may also originate from de novo synthesis, can be exported as VLDL particles upon binding to apoB-100. (III) Removal of triglycerides by endothelial LPL converts VLDL particles into intermediate density lipoprotein (IDL) particles. They can collide with HDL to acquire apoE. (IV) Hepatic lipase (HL) subsequently hydrolyzes the remaining triglycerides in IDL, which forms LDL particles (III) that can also be formed by LPL (III) out of IDL particles, and that are able to release cholesterol to peripheral tissues via LDLR.

Guyton and Hall, 2011). Once formed, circulating LDL particles are removed from the plasma via the interaction of apolipoprotein (apo)B-100 (i.e., one of the LDL particle lipoproteins) with LDL receptors (LDLRs) mainly expressed on liver, but also various other tissues (e.g., lungs, kidneys, urinary bladder, gastrointestinal tract, and adipose tissue). However, if the plasma levels of LDL particles increase, they may accumulate in the arterial wall, where they become proinflammatory by enzymatic oxidation. Subsequently, these proinflammatory LDL particles stimulate endothelial cells and smooth muscle cells in the *tunica intima* (i.e., the innermost arterial layer) to express adhesion molecules and chemoattractants for the recruitment of monocytes, lymphocytes, and neutrophils. In addition, oxidized LDL (ox-LDL) particles can also directly induce a proinflammatory state in monocytes (i.e., inflammatory priming) and accelerate the formation of foam cells (Bekkering et al., 2014). This is preceded by macrophage formation from monocytes, after they have entered the arterial wall guided by chemoattractants. These macrophages recognize and engulf cholesterol, finally yielding foam cells (Insull, 2009; Libby et al., 2011; Maiolino et al., 2013; Bentzon et al., 2014). Consequently, these lipid-loaded macrophages can result in atherosclerotic plaque formation, contributing to an increased risk of CVD (Levine et al., 1995; Kwiterovich, 2000; De Backer et al., 2003; Guyton and Hall, 2011; Taylor et al., 2011; Ridker, 2014; Piepoli et al., 2016).

B. Statins as the Cornerstone of Cardiovascular Disease Treatment

The strong association between high LDL-C levels, atherosclerotic plaque formation, and CVD led to the development of multiple LDL-lowering therapies. To date, statins are among the most widely used LDL-C-lowering therapies (Downs et al., 1998; Baigent et al., 2005; Karalis, 2009; Piepoli et al., 2016), leading to a decrease of 22% in CVD risk with every millimole per liter LDL-C reduction, as quantified in a meta-analysis (Baigent et al., 2010). Statins inhibit 3-hydroxy-3-methylglutaryl-coenzyme A (HMG-CoA) reductase, which is the rate-limiting enzyme in cholesterol synthesis that mediates the conversion of HMG-CoA into mevalonate (Sacks et al., 1996; Taylor et al., 2011; Cruz et al., 2013). Because statins effectively reduce LDL-C, they comprise the cornerstone of atherosclerosis prevention strategies. Although generally well-tolerated, statin-induced muscle complaints are observed in 7%–29% of all users (Bitzur et al., 2013; Wilkinson et al., 2014; Stroes et al., 2015), which contribute significantly to the high discontinuation rate observed with statin therapy. Myopathic symptoms range from very rare cases of rhabdomyolysis to muscle complaints with normal or minimally elevated creatine kinase levels (Bitzur et al., 2013; Wilkinson et al., 2014; Stroes et al., 2015).

These complaints are expected to result from a combination of drug-related (e.g., relative potency, statin metabolism, drug–drug interactions, dose, and lipophilicity) and patient-related (e.g., age, comorbidities, ethnicity, and gender) factors. At the molecular level, many mechanisms have been proposed, with a pivotal role for mitochondrial dysfunction (Schirris et al., 2015a; Stroes et al., 2015). This is supported by the recent discovery of the first statin off-target associated with these muscle complaints, demonstrating that statins in their pharmacologically inactive lactone form inhibit the third complex of the mitochondrial respiratory chain (Schirris et al., 2015a). The cholesterol-lowering statin acid is converted into the lactone form by uridine 5'-diphospho-glycoronyltransferases in the liver (Schirris et al., 2015b). Besides a low adherence due to muscle complaints, statin treatment does not completely normalize the risk of LDL-associated CVD, despite effective LDL reduction (Kuhnast et al., 2015), as demonstrated by several multiple large controlled clinical trials, and follow-up trials illustrated that CVD events persist after treatment in two-thirds of all patients (Sacks et al., 1996; Long-Term Intervention with Pravastatin in Ischaemic Disease (LIPID) Study Group, 1998; Pedersen et al., 2004; Libby, 2005). Similar effects were observed in a subset of patients with coronary heart disease, coronary heart disease risk equivalents, and diabetes mellitus, in which there was a failure to sufficiently lower plasma LDL-C levels (Davidson et al., 2005). More recently, monoclonal antibody proprotein convertase subtilisin/kexin type 9 inhibitors have been introduced (i.e., evolocumab and alirocumab) that drastically reduce LDL-C up to 59% (Sabatine et al., 2017). Contradictory to the large LDL-C reduction, treatment of patients suffering from atherosclerotic CVD with the proprotein convertase subtilisin/kexin type 9 inhibitor evolocumab still resulted in a residual cardiovascular event incidence of 9.8% (Sabatine et al., 2017). This gives rise to the idea that lowering LDL-C cholesterol alone is not sufficient to fully prevent CVD and warrants the exploration of novel therapies to interfere with the pathologic mechanism underlying atherosclerosis.

C. High-Density Lipoprotein and Cardiovascular Disease: High-Density Lipoprotein Levels Are a Poor Reflection of Reverse Cholesterol Transport Capacity

In contrast to blood LDL-C levels, high-density lipoprotein (HDL) cholesterol (HDL-C) levels have been inversely correlated to atherosclerotic events (Tang and Oram, 2009; Uehara and Saku, 2014; Westerterp et al., 2014; Kuhnast et al., 2015). HDL particles have a high protein and low cholesterol and phospholipid content and are involved in the reverse cholesterol transport (RCT) pathway, explaining the beneficial effects of these particles in atherosclerosis (Murphy et al., 2013; Kingwell et al., 2014; Kuhnast et al., 2015).

However, the concept that increased HDL cholesterol levels do uniformly translate into reduced myocardial infarction risk is challenged by a Mendelian randomization study (Voight et al., 2012). The RCT pathway consists of cholesterol transport by HDL particles from peripheral tissue to the liver, where cholesterol is subsequently excreted into the bile by the ATP-binding cassette (ABC) transporters ABCG5 and ABCG8 (Zanlungo et al., 2004; Kingwell et al., 2014) (Fig. 2). As HDL particles are the key players of RCT, they have the potential to degrade atherosclerotic plaques and prevent the formation of new plaques. At the site of the atherosclerotic plaque, cholesterol is transferred to HDL particles by macrophages, which after engulfment transfer cholesterol to HDL via ABC transporter-mediated efflux across the plasma membrane by ABCA1 and ABCG1 (Fig. 2). In addition, the inverse correlation between HDL-C levels and atherosclerotic events has been associated with anti-inflammatory, antioxidant, antiplatelet, and vasodilatory effects (Murphy et al., 2013; Kingwell et al., 2014; Kuhnast et al., 2015; Ramasamy, 2016). Consequently, stimulation of HDL particle levels as well as RCT provides interesting treatment strategies for atherosclerosis. In contrast to previous studies (Fazio and Linton, 2003), recent clinical trials evaluating the infusion of apo mimetics did not show atherosclerotic protection (Karalis and Jukema, 2018). This indicates that increasing HDL levels alone may not be sufficient to stimulate RCT and lower atherosclerotic risk. This also initiated the development of cholesteryl ester transfer protein (CETP) inhibitors, which increase HDL-C levels by reducing the transfer of cholesterol from HDL particles to LDL particles and triglyceride-loaded lipoprotein particles (Barter et al., 2015). However, next to the involvement of CETP in the heterotypic cholesterol transfer pathway (i.e., cholesterol and triglyceride (TG) movement between VLDL or LDL and HDL), it also contributes to the homotypic cholesterol transfer pathway (Lagrost et al., 1990; Rye et al., 1999; Niesor et al., 2010; Barter and Rye, 2012; Mohammadpour and Akhlaghi, 2013; Lauer et al., 2016). In the homotypic pathway, CETP induces the formation of pre- β HDL and cholesterol efflux between subparticles of HDL, including HDL3 and HDL2 (Lagrost et al., 1990; Rye et al., 1999; Niesor et al., 2010). The effect of CETP inhibitors on the heterotypic and homotypic cholesterol transfer pathway depends on the type of inhibitor, which most likely explains their difference in efficacy to reduce cardiovascular risk. For instance, both the homotypic and heterotypic transfer are inhibited by torcetrapib, evacetrapib, and anacetrapib, whereas dalcetrapib more selectively inhibited the heterotypic transfer without affecting the homotypic transfer (Hewing and Fisher, 2012; Mohammadpour and Akhlaghi, 2013). To date, development of CETP inhibitors has been discontinued due to adverse events or lack of efficacy

(Tall and Rader, 2018). The latter also substantiates the notion that functionality of HDL, rather than its absolute HDL-C level, determines effectivity of reducing atherosclerotic risk. Consequently, stimulation of ABCA1- and ABCG1-mediated cellular cholesterol efflux to HDL particles could, due to its pivotal role in RCT, provide a more effective strategy to stimulate RCT and decrease atherosclerotic risk.

D. Systematic Review: Scope and Methodology

In this review, we discuss the major players of RCT as well as therapeutic strategies explored to stimulate them. A systematic literature search was conducted using both Medline and Embase, which resulted in 2928 abstracts (Fig. 3; Supplemental Tables 1 and 2). Removal of duplicates resulted in 2809 unique publications that were independently evaluated for their relevance by two reviewers. First, all publications not involving cholesterol efflux stimulation were excluded based on the title (i.e., title screen, Fig. 3). Following strict exclusion criteria, 175 publications remained, based on their abstract for a more in-depth study and inclusion in this review (Fig. 3), including publications describing the effects on cholesterol efflux of endogenous compounds, natural compounds, apolipoprotein mimetics, microRNAs (miRNAs), as well as nonresearch publications and publications of which only the abstract or no English version was available.

We first provide an overview of the main players involved in RCT, focusing on the crucial role of ABCA1 and ABCG1 cholesterol efflux transporters in the initiation and propagation of this process. We conclude with a critical discussion of small-molecule pharmaceutical interventions that could affect RCT beneficially.

II. Reverse Cholesterol Transport Pathway

A. Initiation and Propagation of Reverse Cholesterol Transport

Each RCT cycle is initiated by the removal of cellular cholesterol from peripheral tissues, of which the determinants (i.e., HDL size and composition, gender, body mass index, and age) have been reviewed recently by Talbot et al. (2018). Active transport is involved in 70% of all cellular cholesterol efflux pathways (Ono, 2012; Phillips, 2014). Passive bidirectional cholesterol flux between HDL particles and the plasma membrane is mediated by aqueous diffusion and via interaction of HDL particles with scavenger receptor B1 (SR-B1). SR-B1 mediates cholesterol uptake in the cell without endocytic uptake and degradation of HDL particles by ABCG1 (Phillips, 2014). However, the contribution of this mechanism to peripheral cholesterol efflux is expected to be minimal, as SR-B1 is most abundantly expressed in the adrenal gland, and at lower levels in peripheral tissues. Aqueous diffusion includes passive transport through the intervening aqueous phase until collision occurs between cholesterol and an extra- or




Fig. 2. Reverse cholesterol transport. Schematic overview of reverse cholesterol transport, (I) which is initiated by the efflux of cholesterol (yellow spheres) by ABCA1 and ABCG1 transporters. They are expressed on a variety of peripheral tissues, including macrophages that engulf cholesterol from atherosclerotic plaques (i.e., foam cells). ABCA1-mediated cholesterol efflux transfers cholesterol to lipid-poor apoA-I, leading to the formation of pre-βHDL. These particles can be converted by circulating lecithin:cholesterol acyltransferase (LCAT) into HDL particles, which function as cholesterol acceptor for ABCG1. Cholesterol efflux by ABCG1 is mediated via reorganization of cholesterol in the plasma membrane, which increases plasma membrane cholesterol concentrations. Subsequently, aqueous diffusion could increase the cholesterol efflux out of the cell to HDL without necessity of HDL to bind to the plasma membrane. (II) SR-BI mediates cholesterol influx into hepatocytes without whole HDL particle uptake, allowing unbound apoA-I to circulate and enter a new RCT cycle. Finally, hepatic free cholesterol can be directly removed to bile canaliculi by ABCG5 and ABCG8 transporters, or indirect via conversion by cytochrome P450 (CYP)7A1 into bile acids (green spheres) that can subsequently be transported into bile via the multidrug resistance protein (MRP)2/ABCC2 and bile salt export pump (BSEP/ABCB11) transporters, located on the canalicular membrane.

intracellular cholesterol acceptor (Yancey et al., 2003; Rosenson et al., 2012; Phillips, 2014). ATP-dependent cholesterol efflux is mainly facilitated by ABCA1 and ABCG1 (Fig. 2) (Oram, 2003; Yancey et al., 2003;

Yvan-Charvet et al., 2010b; Phillips, 2014; Westerterp et al., 2014). ABCA1-mediated cholesterol and phospholipid efflux is initiated by the interaction of lipid- and cholesterol-free apoA-I particles with ABCA1




Fig. 3. Flow chart of systematic literature search strategy. Overview of the number of publications retrieved from all MEDLINE and EMBASE searches and exclusion criteria applied to these publications, including removal of duplicates and title- and abstract-based screenings conducted by two independent reviewers. For a detailed overview of the search strategies, see Supplemental Tables 1 and 2.

transporters on peripheral tissue (i.e., highest expression in macrophages and foam cells) and hepatocytes (Fig. 2). This results in the formation of lipid-poor pre- β HDL particles, which upon rapid esterification by lecithin:cholesterol acyltransferase are converted into lipid-rich HDL particles (Morgado et al., 2005). These particles mediate cellular cholesterol efflux via interaction with ABCG1, thereby further facilitating transport of cholesterol from peripheral tissues toward the liver, followed by uptake into hepatocytes via SR-B1 (Tall and Yvan-Charvet, 2015). As the final step of RCT, cholesterol is secreted into the bile via an ABCG5/8 heterodimer complex present at the canalicular membrane (Thomas et al., 2003; Roglans et al., 2004; Zanlungo et al., 2004; Valasek et al., 2007; Lee et al., 2016). These two ABCG transporters are only expressed in hepatocytes,

gallbladder epithelium, and enterocytes (Tauscher and Kuver, 2003; Wang et al., 2015a; Patel et al., 2018). Formation of a heterodimer of ABCG5 and ABCG8 in the endoplasmic reticulum is required before the complex can be translocated to the apical membrane, where it facilitates sterol transport into the bile or gut lumen (Graf et al., 2003; Yu et al., 2014). The biliary excretion rate of cholesterol is associated with the level of hepatic ABCG5 and ABCG8 expression (Yu et al., 2005). Consequently, the ABCG5/G8 complex is important in the correction of high plasma and hepatic cholesterol and sterol levels (Yu et al., 2002, 2014).

B. ATP-Binding Cassette A1 and ATP-Binding Cassette G1 as Master Effectors of Cholesterol Efflux

ABCA1 plays a crucial role in the cellular cholesterol efflux, and its importance is illustrated by Tangier

disease, which is characterized by a severe deficiency in plasma HDL and cholesterol caused by autosomal recessive mutations in the *ABCA1* gene (Brooks-Wilson et al., 1999; Langmann et al., 1999; Rust et al., 1999; Oram, 2000; Uehara et al., 2011; Phillips, 2018). The human *ABCA1* gene has a total length of 149 kb, including a 1,453-bp promoter, 146,581 bp of introns and exons, and a 1-kb 3' flanking region. Of the exons, 58 are comprised in the *ABCA1* gene, and multiple bindings sites for transcription factors are detected in the promoter region (Gene, 1982–2019a; Santamarina-Fojo et al., 2000). Like many other transporters of the ABCA subfamily, ABCA1 is a full transporter, which contains 2,261 amino acids and is integrated into the membrane via two transmembrane domains (TMDs) that both comprise six transmembrane helices. In addition, ABCA1 has two nucleotide binding domains (NBDs), which contain the conserved Walker-A and Walker-B peptide motifs (Langmann et al., 1999; Santamarina-Fojo et al., 2000; Uehara et al., 2011). The tissue distribution of ABCA1 is ubiquitous, and expression is especially high in placenta, liver, small intestine, macrophages, adrenal glands, lung, and fetal tissues (Langmann et al., 1999; Uehara et al., 2011). ABCA1 initiates cellular cholesterol efflux and RCT to the lymph and bloodstream via a specific interaction with apoA-I, allowing cellular phospholipids and cholesterol to bind to these apolipoproteins, which almost exclusively mediates HDL biosynthesis and is therefore seen as its rate-limiting step (Lee et al., 2002; Murthy et al., 2002; Ohama et al., 2002; Mulligan et al., 2003; Singaraja et al., 2003). It remains, however, unknown whether phospholipid and cholesterol binding occurs at the plasma membrane surface or whether apoA-I is internalized and targeted to late endosomes after binding to ABCA1 at the plasma membrane, where it forms complexes with lipids that are subsequently released by exocytosis (Takahashi and Smith, 1999; Neufeld et al., 2001; Vaughan and Oram, 2003; Denis et al., 2008; Lorenzi et al., 2008; Azuma et al., 2009; Tang and Oram, 2009; von Eckardstein and Rohrer, 2009; Yvan-Charvet et al., 2010b; Westerterp et al., 2014; Du et al., 2015a) (Fig. 4). This is also supported by the localization of ABCA1 on early and late endosome and lysosome membranes, next to its plasma membrane localization, and in line with its reuptake to regulate cholesterol efflux rates, as described in the next section (Neufeld et al., 2001; Tang and Oram, 2009; von Eckardstein and Rohrer, 2009; Uehara and Saku, 2014; Westerterp et al., 2014; Du et al., 2015a). However, the majority of apoA-I lipidation is expected to occur at the plasma membrane (Denis et al., 2008; Faulkner et al., 2008). The exact modes of interaction between apoA-I and ABCA1 still need to be elucidated. At least six mechanisms have been proposed, as follows: 1) direct apoA-I binding to phosphatidylserine, which is translocated outward by ABCA1 floppase activity (Chambenoit et al., 2001; Alder-Baerens et al., 2005); 2) direct binding of

apoA-I to extracellular ABCA1 loop domains (Wang et al., 2001; Fitzgerald et al., 2004); 3) apoA-I binding to protrusions as a result of ABCA1 floppase activity (Vedhachalam et al., 2007a); 4) outward translocation of phosphatidylinositol 4,5-bisphosphate mediated by ABCA1 floppase activity, which allows apoA-I to bind and unfold, followed by microsolvubilization of the membrane (Gulshan et al., 2016); 5) low-affinity apoA-I binding to ABCA1 and high-affinity binding to cholesterol (Hassan et al., 2007; Vedhachalam et al., 2007b); 6) apoA-I binding to extracellular domains of ABCA1 dimers that have been formed from two ABCA1 monomers. These monomers have undergone a conformational change as a result of the translocation of phosphatidylcholine and cholesterol leading to dimer formation (Ishigami et al., 2018). Although ABCA1 mainly interacts with apoA-I, it can also associate with lipid-free apoE, which is most efficient when these apolipoproteins originate from small and dense HDL subfractions like HDL3b and HDL3c (Neufeld et al., 2001; Tang and Oram, 2009; von Eckardstein and Rohrer, 2009; Uehara and Saku, 2014; Westerterp et al., 2014; Du et al., 2015a).

In contrast to ABCA1, ABCG1 has a larger ambiguity regarding its lipid acceptors, which include HDL, LDL, and phospholipid vesicles (Wang et al., 2004; Vaughan and Oram, 2005; Kobayashi et al., 2006; Sankaranarayanan et al., 2009; Yvan-Charvet et al., 2010b; Phillips, 2014; Westerterp et al., 2014). Unlike ABCA1, ABCG1 is a half transporter, which only contains a single TMD comprising of six transmembrane helices and a single NBD at the C terminus of the TMD that is responsible for ATP binding and hydrolysis (Kerr et al., 2011; Uehara et al., 2011). Therefore, the ABCG1 protein needs to either homo- or heterodimerize with other ABCG proteins to become functional. The human *ABCG1* gene comprises 23 exons spanning 98 kb (Gene, 1982–2019b; Kennedy et al., 2001). Eight different isoforms of ABCG1 are produced by alternative splicing, with a length varying between 644 and 785 amino acids. The ABCG1 protein is expressed in many cell types, including macrophages, neurons, astrocytes, endothelial, and epithelial cells, and in many tissues, such as the liver, intestine, kidney, spleen, lung, and brain (Nakamura et al., 2004; Kennedy et al., 2005; Wang et al., 2008; Bojanic et al., 2010). Although the precise cellular localization of ABCG1 needs to be elucidated, the transporter was detected in the plasma membrane and membranes of the Golgi apparatus and endosomes of ABCG1-overexpressing HEK293 cells and macrophages (Kobayashi et al., 2006; Wang et al., 2006; Tarling and Edwards, 2011; Neufeld et al., 2014; Phillips, 2014; Westerterp et al., 2014). It has been demonstrated that cholesterol efflux facilitated by ABCG1 does not increase lipoprotein binding to the cell surface (Wang et al., 2004; Kobayashi et al., 2006; Sankaranarayanan et al., 2009; Yvan-Charvet et al., 2010b; Phillips, 2014), which makes a mechanism similar to




Fig. 4. Potential ABCA1- and ABCG1-driven cholesterol efflux modes. ABCA1-driven cholesterol efflux to lipid-poor apoA-I is hypothesized to occur either at the plasma membrane (upper left panel), or via endocytosis of the apoA-I-bound ABCA1 transporter, followed by intracellular lipid loading and exocytosis (upper right panel). ABCA1 is ubiquitously expressed, particularly at high levels in liver, small intestine, macrophages, adrenal glands, lungs, placenta, and fetal tissue. The transporter initiates cellular cholesterol efflux to the lymph and bloodstream via a specific interaction with apoA-I. Six different mechanisms have been proposed for the interaction between apoA-I and ABCA1 (small boxes), including the following: outward translocation of phosphatidylserine (PS) by ABCA1 floppase activity allowing apoA-I binding; direct binding of apoA-I to extracellular ABCA1 loop domains; ABCA1 floppase activity leading to the formation of protrusions facilitating apoA-I binding; ABCA1 floppase activity leading to the outward translocation of phosphatidylinositol 4,5-bisphosphate (PIP2), allowing apoA-I to bind and unfold, followed by microsolubilization of the membrane; low-affinity binding to ABCA1 and high-affinity binding to cholesterol; dimerization of ABCA1 transporter proteins is initiated by loading of the extracellular loop domains with cholesterol, followed by apoA-I binding to the dimerized cholesterol-loaded extracellular loop domains (ED). ABCG1-driven cholesterol efflux to HDL particles in the bloodstream or lymph is expected to be the result of a collision of these particles with cholesterol molecules that protrude from the plasma membrane (lower left panel), mediated either by a direct effect of the ABCG1 dimer on the membrane structure or by outward translocation via ABCG1 floppase activity (lower right panel). ABCG1 is expressed in many cell types, including macrophages, neurons, astrocytes, endothelial and epithelial cells, and many tissues (e.g., liver, intestine, kidney, spleen, lung, and brain), where it mediates basolateral cholesterol efflux.

ABCA1 unlikely. Several pathways have been suggested to explain the ABCG1-mediated cellular cholesterol efflux mechanism (Yvan-Charvet et al., 2010b; Neufeld et al., 2014; Phillips, 2014). One of the proposed mechanisms suggests that ABCG1 facilitates protrusion of cholesterol from the membrane pool into the hydrophilic water layer lining the plasma membrane (Fig. 4). Subsequently, cholesterol uptake by an acceptor occurs after transient collision (Small, 2003; Yvan-Charvet et al., 2010b; Neufeld et al., 2014; Phillips, 2014). A second model suggests that ABCG1 promotes changes in the organization of plasma membrane phospholipids functioning as a phospholipid floppase, which leads to a redistribution of sterols to the plasma membrane (Yvan-Charvet et al., 2010b; Phillips, 2014). This could result in an increased efflux of cholesterol out of the cell by aqueous diffusion because of increased cholesterol concentrations at the plasma membrane (Sankaranarayanan et al., 2009; Yvan-Charvet et al., 2010b; Phillips, 2014).

In summary, ABCA1- and ABCG1-mediated cholesterol efflux is indispensable in cholesterol and phospholipid loading of apolipoproteins and thereby the maturation of HDL particles, which makes both ABC transporters essential for the initiation of RCT (Tang and Oram, 2009; Phillips, 2014; Uehara and Saku, 2014; Westerterp et al., 2014). Consequently, stimulation of cellular cholesterol removal via ABCA1 and ABCG1 could provide a promising target to enhance RCT.

C. Regulation of ATP-Binding Cassette A1- and ATP-Binding Cassette G1-Mediated Cholesterol Efflux

Cholesterol efflux by ABCA1 and ABCG1 is regulated by a variety of different mechanisms. Serum HDL and its key apoA-I are key regulators of the cholesterol efflux rate, in which high apolipoprotein levels are associated with high cellular cholesterol efflux rates. Cellular mechanisms mainly regulate ABCA1- and ABCG1-mediated cholesterol efflux by controlling the plasma membrane expression of both transporters. This is accomplished by various mechanisms. First, the nuclear receptors retinoid X receptor (RXR), peroxisome proliferator-activated receptors (PPARs), and liver X receptor (LXR) regulate the transcription of the genes encoding *ABCA1* and *ABCG1* (Fig. 5). Second, the stability of *ABCA1* and *ABCG1* mRNA can be influenced by their protein expression. Third, the expression of the transporter on the plasma membrane is regulated via modulation of its internalization, degradation, and recycling. Finally, activity and localization are enhanced by cAMP-mediated transporter phosphorylation and the stimulatory role of extracellular ATP thereon (Lee et al., 2011). The different cellular mechanisms influencing ABCA1- and ABCG1-mediated cholesterol efflux are discussed in more detail below, along with an overview of small-molecule treatment strategies known to influence these mechanisms.

III. Apolipoprotein A-I and Apolipoprotein E Mimetics

Although we focus on small-molecule therapies that enhance ABCA1- and ABCG1-mediated cholesterol efflux, we will briefly address recent promising progress in the development of recombinant apolipoproteins to stimulate cholesterol efflux by ABCA1 and ABCG1 (Zhang et al., 2003; Tall et al., 2008; Bielicki et al., 2010; Khera et al., 2011), which was extensively reviewed by others (Sherman et al., 2010; White et al., 2014; Stoekenbroek et al., 2015; Cao et al., 2017). Many efforts have been directed to mimic or overexpress apoA-I as potential atherosclerosis treatment strategy, which is a promising therapy for several reasons. First, apoA-I is the major functional and most abundant structural lipoprotein in HDL, which accounts for approximately two-thirds of the total HDL protein content (Zhang et al., 2003; Tall et al., 2008; Bielicki et al., 2010; Getz and Reardon, 2011; Khera et al., 2011; Uehara and Saku, 2014; Kontush et al., 2015). Moreover, formation of cholesterol-loaded apoA-I by ABCA1 is considered to be the rate-limiting step of HDL particle biogenesis. Consequently, increasing apoA-I levels using apoA-I mimetics (i.e., short synthetic peptides that share structural and biologic features of native apolipoproteins) is expected to stimulate cholesterol transport. The potential of this strategy is demonstrated in several in vitro and in vivo studies, in which treatment with apoA-I mimetics increased plasma HDL levels and reduced atherosclerotic lesions (Zhang et al., 2003; Tall et al., 2008; Bielicki et al., 2010; Khera et al., 2011; Osei-Hwedie et al., 2011; Uehara and Saku, 2014; Kuhnast et al., 2015). More recently, clinical trials evaluating the infusion of these apoA-I mimetics, including the potent mimetic apoA-I Milano, did not show atherosclerotic protection (Karalis and Jukema, 2018). This also substantiates the notion that not the HDL levels, but other parts of RCT, like ABCA1- and ABCG1-mediated cholesterol efflux, contribute to the inverse correlation between HDL-C levels and atherosclerotic events.

Besides apoA-I mimetics, compounds that act like apoE are of great interest for the treatment of atherosclerosis. ApoE possesses antiatherogenic properties by its cholesterol-reducing potential. These effects are mediated via the clearance of plasma apoB-containing lipoprotein remnants (e.g., VLDL and chylomicrons) by apoE LDLR-binding domain (Sharifov et al., 2011; Xu et al., 2016). Moreover, they are involved in the initiation of RCT from peripheral tissue and macrophages upon lipid binding (Sharifov et al., 2011; Xu et al., 2016) and binding to ABCG1 to facilitate cholesterol efflux by this transporter. Several in vivo studies revealed that apoE mimetics have cholesterol-lowering, anti-inflammatory, and atheroprotective properties (Nayyar et al., 2012; Handattu et al., 2013; Xu et al., 2016). Moreover,




Fig. 5. Cellular regulation of ABCA1 and ABCG1 transporter expression. Expression of ABCA1 and ABCG1 transporters is increased by stimulation of the nuclear PPAR, which upon dimerization with RXR (i.e., another nuclear factor) bind to the PPAR-responsive element (PPRE), leading to the transcription of the nuclear LXR and RAR, respectively (right upper panel). Next, dimerization of LXR or RAR with RXR enables binding to the LXRE, inducing the transcription of *ABCA1* and *ABCG1*, respectively. Once transcribed, the stability of *ABCA1* and *ABCG1* mRNA can either be enhanced or decreased upon binding of miRNAs, of which an overview is provided in Table 4 (right middle panel). ABCA1 and ABCG1 plasma membrane expression is also mediated via modulation of its lysosomal (blue vesicle) or endosomal (orange vesicle) degradation. The latter is stimulated by phosphorylation of the PEST sequence by apelin-13 (AP-13) via PKC α (left lower panel). Upon phosphorylation, calpain (CALP) can bind and initiate proteolysis, a process that is stimulated by calpastatin (CPSTAT), and negatively affected by calmodulin (CM) or by the HO-1 axis. Finally, the ability of ABCA1 and ABCG1 to bind lipid poor apoA-I is stimulated upon transporter phosphorylation by PKA at Ser-1042 and Ser-2054, located in the nucleotide binding domain of ABCA1. PKA is positively regulated by cAMP levels that depend on the balance of cAMP breakdown by PDE, formation by adenosine receptor-stimulated AC, and efflux by multidrug resistance protein (MRP) 4 and 5 (right lower panel).

a hybrid peptide containing properties of both apoA-I and apoE improved arterial endothelial function and protected against atherogenesis in vivo (Gupta et al., 2005).

Although these apolipoprotein mimetics showed promising results for the treatment of atherosclerosis, their peptide nature may make them less favorable to administer as compared with orally available small-molecule drugs. ABCA1 and ABCG1 transporters provide an interesting target for small molecules to stimulate RCT, as their expression and function are expected to be rate-limiting for cellular cholesterol efflux (Oram, 2003; Tall et al., 2008; Yvan-Charvet et al., 2010b; Phillips, 2014; Westerterp et al., 2014).

IV. Regulation and Pharmacological Manipulation of Nuclear Receptor-Mediated ATP-Binding Cassette A1 and ATP-Binding Cassette G1 Expression

A. Nuclear Receptors Are Important Regulators of ATP-Binding Cassette A1 and ATP-Binding Cassette G1 Expression

Multiple nuclear receptors are involved in the regulation of *ABCA1* and *ABCG1* mRNA expression, of which the two LXR isoforms, LXR α and LXR β , have demonstrated to play an important role in the cholesterol homeostasis, including RCT (Zhao and Dahlman-Wright, 2010; Hong and Tontonoz, 2014). LXR α is predominantly expressed in metabolically active tissue (e.g., liver, kidney, macrophages, adipose tissue, and small intestine), whereas LXR β has a more ubiquitous distribution and is in particular highly expressed in the developing brain (Zhao and Dahlman-Wright, 2010; Li et al., 2016). LXRs act as cellular cholesterol sensors, as they are activated by accumulation of oxysterols, oxidized derivatives of cholesterol, which subsequently induce transcription of genes involved in the protection of cells against cholesterol overload (Zhao and Dahlman-Wright, 2010; Di et al., 2012). After activation by oxysterols, LXR α and LXR β form heterodimers with isoforms of RXRs, including RXR α , RXR β , or RXR γ (Di et al., 2012; Hong and Tontonoz, 2014) (Fig. 5). After heterodimerization, LXR/RXR initiate transcription of target genes (e.g., genes involved in lipid synthesis and metabolism, including, but not limited to, the following: *ABCA1*, *ABCG1*, *ABCG5*, *ABCG8*, *SREBP-1C*, and *FAS*) by binding to the LXR response element (LXRE), which consists of two direct repeats (i.e., a AGGTCA sequence) separated by four nucleotides (Edwards et al., 2002; Zhao and Dahlman-Wright, 2010; Jakobsson et al., 2012; Hong and Tontonoz, 2014; Li et al., 2016). These LXREs are found in the proximal promoters of genes involved in fatty acid, bile acid, cholesterol, and glucose regulation, but also of genes with high relevance to RCT, including *ABCA1* and *ABCG1* (Zhao and Dahlman-Wright, 2010; Di et al., 2012; Hong and Tontonoz, 2014). Combined

with its high expression in macrophages, LXR α is an important regulator in the initiation of RCT via modulation of *ABCA1* and *ABCG1* expression.

Another class of nuclear receptors involved in the regulation of *ABCA1* and *ABCG1* expression is PPAR, which comprises the following three isoforms: PPAR α , PPAR β/δ , and PPAR γ (Berger et al., 2005; Ogata et al., 2009). PPAR α is mainly expressed in liver, kidney, heart, muscle, and other metabolically active tissues that rely predominantly on fatty acid β -oxidation. PPAR β/δ is ubiquitously expressed (e.g., cardiovascular, urinary, respiratory, digestive, endocrine, nervous, and hematopoietic organ system), whereas PPAR γ is mainly found in adipose tissue, skeletal and cardiac muscles, and human monocytes (Escher et al., 2001; Berger et al., 2005; Higashiyama et al., 2007; Ogata et al., 2009). Like LXR, PPARs form obligate heterodimers with RXR upon activation by their ligands (e.g., fatty acid metabolites), which bind to isotype-specific peroxisome proliferator response elements in target genes (Berger et al., 2005; Grygiel-Górniak, 2014) (Fig. 5). Targets of PPAR α and PPAR γ include genes involved in lipid metabolism (e.g., *SLC25A20*, *APOA1*, *LXR α* , and *SCD-1*) and glucose metabolism (e.g., *G6PC*, *PCK*, and *PDK4*) (Muio et al., 2002; Li and Glass, 2004; Rakhshandehroo et al., 2010). PPARs have been associated with an increased *ABCA1* expression and consequently enhanced HDL biogenesis in vitro and in vivo (Chinetti et al., 2001; Ogata et al., 2009). All three PPAR isoforms mediate these effects via LXR α . However, only PPAR γ is known to directly affect LXR α expression via interaction with a peroxisome proliferator response element proximal to the LXR α promoter (Chawla et al., 2001), whereas the exact mechanism remains unknown for the other isoforms and is expected to be indirect (e.g., via effects on its endogenous ligands) (Ogata et al., 2009).

In addition to RXR, retinoid-activated receptors (RARs; i.e., another group of retinoid nuclear receptors) play a role in cholesterol homeostasis. RARs are expressed in a wide variety of tissues and, like other retinoid nuclear receptors, also need to form a heterodimer with RXR (Matsumoto et al., 2007; Jung et al., 2010; Kuntz et al., 2015; Manna et al., 2015; Zhou et al., 2015). For all retinoid nuclear receptors, this heterodimerization allows them to bind to a response element in the promoter region of their target genes, including *ABCA1*, *ABCG1*, *APOA1*, *GCK*, *UCP1* and *UCP3*, and *FGF21* (Puigserver et al., 1996; Solanes et al., 2000; Balmer and Blomhoff, 2002; Cadoudal et al., 2008; Nishimaki-Mogami et al., 2008; Cui et al., 2011; Ayaori et al., 2012; Li et al., 2013; Zhang et al., 2013a, 2015b; Kuntz et al., 2015). PPAR/RXR, LXR/RXR, and RAR/RXR can be activated by agonists for either RXR and any other nuclear receptor (Nishimaki-Mogami et al., 2008; Cui et al., 2011; Kuntz et al., 2015) (Fig. 5). Consequently, RXR exerts pleiotropic effects due to crosstalk of RXR with other nuclear receptors, resulting in the simultaneous activation of multiple converging

signaling pathways (Matsumoto et al., 2007; Nishimaki-Mogami et al., 2008). Both retinoic nuclear receptors are activated by their endogenous ligand retinoic acid (RA; i.e., all-*trans*-RA and 9-*cis*-RA) (Costet et al., 2003; Koldamova et al., 2003; Nishimaki-Mogami et al., 2008; Zhou et al., 2015). Recently, the relevance of this pathway for RCT was demonstrated in a mouse model of atherosclerosis (*apoE*^{-/-} mice on a high-fat diet) by the LXRE-dependent stimulatory effects of 9-*cis*-RA on macrophage ABCA1 and ABCG1 expression, cholesterol efflux, and HDL-C levels (Zhou et al., 2015). Consequently, RXR as well as the other nuclear factors are important effectors of the cellular cholesterol homeostasis, which makes them valuable targets to induce ABCA1 and ABCG1 expression and to eventually enhance cellular cholesterol efflux and RCT.

B. Liver X Receptor Activation to Induce ATP-Binding Cassette A1 and ATP-Binding Cassette G1 Expression

The increased understanding about the pivotal role of LXR target genes in cholesterol metabolism, including ABCA1 and ABCG1, has enhanced the interest to pharmacologically modulate LXR signaling. One of the first discovered synthetic steroidal LXR agonists, T0901317, a full and potent agonist of both LXR α and LXR β , appeared to protect against atherosclerotic development in vitro and in animal studies (Terasaka et al., 2003; Thomas et al., 2003; Beyer et al., 2004; Quinet et al., 2004; Wang et al., 2006; Dai et al., 2008; Sato et al., 2008; Larrede et al., 2009; Verschuren et al., 2009; Yan et al., 2010; Honzumi et al., 2011; Chen et al., 2012; Kirchgessner et al., 2015; Manna et al., 2015; Jiang and Li, 2017) (Table 1). T0901317 enhanced the in vitro and in vivo expression of LXR target genes, including ABCA1 and ABCG1, leading to increased cholesterol efflux to apoA-I and HDL and decreased foam cell and atherosclerotic plaque formation (Fukumoto et al., 2002; Murthy et al., 2002; Terasaka et al., 2003; Thomas et al., 2003; Beyer et al., 2004; Miao et al., 2004; Quinet et al., 2004, 2006; Wu et al., 2004; Panzenboeck et al., 2006; Wang et al., 2006; Delvecchio et al., 2007, 2008; Fujiyoshi et al., 2007; Dai et al., 2008; DiBlasio-Smith et al., 2008; Sato et al., 2008; Zanotti et al., 2008; Larrede et al., 2009; Verschuren et al., 2009; Mogilenko et al., 2010; Morrow et al., 2010; Yan et al., 2010; Honzumi et al., 2011; Maejima et al., 2011; Chen et al., 2012; Di et al., 2012; Elali and Hermann, 2012; Jiang et al., 2012; Ma et al., 2014; Kaneko et al., 2015; Kirchgessner et al., 2015; Manna et al., 2015; Tamehiro et al., 2015; Carter et al., 2017; Jiang and Li, 2017; Marinozzi et al., 2017; Monzel et al., 2017; Kaseda et al., 2018). Unfortunately, T0901317 was associated with enhanced lipogenesis, resulting in elevated serum triglyceride levels and hepatic steatosis, which is most likely explained by LXR α -induced activation of the sterol regulatory-binding element protein 1c (SREBP-1c) pathway (Terasaka et al.,

2003; Thomas et al., 2003; Beyer et al., 2004; Miao et al., 2004; Quinet et al., 2004, 2006; Delvecchio et al., 2008; DiBlasio-Smith et al., 2008; Sato et al., 2008; Verschuren et al., 2009; Yan et al., 2010; Honzumi et al., 2011; Chen et al., 2012; Kaneko et al., 2015; Kirchgessner et al., 2015; Manna et al., 2015; Carter et al., 2017; Marinozzi et al., 2017). A nonsteroidal LXR agonist, GW3965, also protected against atherosclerosis development (Ruan et al., 2003), but less effectively as T0901317 (Sparrow et al., 2002; Bennett et al., 2006; Brunham et al., 2006; Naik et al., 2006; Quinet et al., 2006; Delvecchio et al., 2007; DiBlasio-Smith et al., 2008; Di et al., 2012; Kannisto et al., 2014). Moreover, it amplified *SREBP-1c* expression leading to increased hepatic and plasma triglyceride levels (Sparrow et al., 2002; Quinet et al., 2004). Various other LXR agonists, including acetyl-podocarpic dimer, LXR-623 (i.e., also known as WAY-252623), ritonavir (i.e., an antiretroviral drug), side-chain modified sterol, ergosterol derivatives, and C24-hydroxylated stigmastane derivatives, also effectively increased ABCA1 expression (Sparrow et al., 2002; DiBlasio-Smith et al., 2008; Pou et al., 2008; Quinet et al., 2009; Marinozzi et al., 2017; Castro Navas et al., 2018). However, they all produced unwanted effects on plasma triglyceride levels by stimulation of *SREBP-1c* gene transcription and LXR-623 enhanced plasma triglyceride levels, yet all to a lower extent than T0901317 (Sparrow et al., 2002; DiBlasio-Smith et al., 2008; Pou et al., 2008; Quinet et al., 2009; Marinozzi et al., 2017; Castro Navas et al., 2018).

Although the adverse effects limit the clinical use of these LXR agonists, they raised great interest in LXR stimulation as a potential target to stimulate RCT. Novel LXR agonists have been developed, like *N,N*-dimethyl-3 β -hydroxycholelamide (DMHCA), a synthetic oxysterol that has the potential to enhance cholesterol transport in an LXR-dependent manner without increasing plasma triglyceride levels. Moreover, hepatic *SREBP-1c* mRNA expression was only slightly increased by DMHCA in rat aortic endothelial cells, whereas in macrophages *SREBP-1c* was reduced (Quinet et al., 2004; Kratzer et al., 2009; Hammer et al., 2017). The potential of this type of LXR agonist was emphasized by methyl-3 β -hydroxy-5 α ,6 α -epoxy-cholanate, a compound with similar structure and effects as DMHCA (Yan et al., 2010), and a novel analog of *N,N*-disubstituted 2,8-diazaspiro [4.5]decane, also known as IMB-151 (Li et al., 2014). This compound also upregulated ABCA1 and ABCG1 expression in RAW264.7 macrophages in an LXR α -dependent manner, whereas SREBP-1c protein expression levels in HepG2 cells were only slightly increased (Li et al., 2014). Other LXR-agonists, ibrolipim (also known as NO-1886) and BMS-779788, increased ABCA1 expression and cellular cholesterol efflux and reduced *SREBP-1c* expression in vitro and markedly lowered plasma triglyceride levels in vivo (Zhang et al., 2006; Ma et al., 2009; Chen et al., 2010; Tsou et al., 2014; Kick et al., 2015; Kirchgessner et al., 2015). The LXR β -specific

TABLE 1
LXR-activating compounds

Overview of compounds that activate LXR, including their primary pharmacological action, effects on ABCA1 and ABCG1 expression (arrows: *mRNA*, protein), and effects on cellular cholesterol efflux. Effects on ABCA1, ABCG1, and cholesterol efflux are presented as follows: (↓) decreased; (=) no effect; (↑) increased. Italicized symbols indicate changes in *ABCA1* and *ABCG1* mRNA levels, whereas nonitalicized symbols indicate protein levels.

Compound	Primary Action	ABCA1	ABCG1	Cholesterol Efflux	Reference
Acetyl-podocarpic dimer	Selective LXR agonist	↑ THP-1 ↑ Caco-2 ↑ Primary hepatocytes ^a	↑ Caco-2 Primary hepatocytes ^a	↑ THP-1 apoA-I ↑ Caco-2 ↑ Monocytes ^a Primary fibroblasts ^a : ↑ apoA-I	Sparrow et al., 2002
Baclofen	GABA _B receptor agonist	—	—	MDM = apoA-I = HDL	Yang et al., 2014
Digoxin	Na ⁺ -K ⁺ -ATPase inhibitor	↑ H9c2	—	↑ H9c2	Campia et al., 2012
Disodium ascorbyl phytostanol phosphate (FM-VP4)	MDR1 antagonist	↑ Liver ^b = Small intestine ^b	—	—	Méndez-González et al., 2010
Daunorubicin	Anthracycline antibiotic	= HL-1	= HL-1	—	Monzel et al., 2017
DMHCA	LXR α agonist	↑ THP-1 ↑ J774 ↑ HepG2 ↑ BREC ↑ PM ^{b,c} ↑ Liver ^a = Liver ^d ↑ Ileum ^d ↑ Aorta ^d	↑ BREC ↑ PM ^c ↑ Liver ^{a,d} ↑ Ileum ^d ↑ Aorta ^d	↑ THP-1	Quinet et al., 2004; Kratzer et al., 2009; Hammer et al., 2017
Doxorubicin	Anthracycline antibiotic	↑ HL-1	↑ HL-1	HL-1 ↑ apoA-I ↑ HDL	Monzel et al., 2017
E17110	Benzofuran-2-carboxylate analog	↑↑ RAW264.7	↑↑ RAW264.7	↑↑ RAW264.7 ↑ apoA-I ↑ HDL	Li et al., 2016
(E)-1-(e,4-diisopropoxyphenyl)-3-(4-isopropoxy-3-methoxyphenyl)-2-en-1-one	Chalcone derivative	↑↑ THP-1	↑ THP-1	—	Teng et al., 2018
Ergosterol derivatives	Ergosterol analog	↑ U937	—	—	Marinozzi et al., 2017
Etoposide	DNA topoisomerase II inhibitor	↑ RAW264.7 ↑ PM ^b ↑ THP-1 Blood monocyte ^a	↑ PM ^b	↑ PM ↑ RCT ^b	Zhang et al., 2013a
EXEL-04286651/BMS-779788	LXR partial agonist	↑ Murine blood cells	↑ Murine blood cells ↑ Blood cells ^e	—	Kick et al., 2015; Kirchgessner et al., 2015
FTY720-P	Sphingosine-1-phosphate analog	↑ Monocytes ^a	—	↑ Monocytes ^a	Blom et al., 2010
G004	Unknown (synthetic sulfonylurea compound)	↑↑ RAW264.7 ↑ Liver ^c	↑↑ RAW264.7 ↑ Liver ^c	↑ RAW264.7 ↑ RCT ^c	Qian et al., 2017
GW3965	LXR α agonist	↑ THP-1 ↑ J774 ↑ HepG2 ↑ haSMC ^a ↑ BMM ↑ PM ^{b,h,i} ↑↑ Liver ^{a,h} ↑ Liver ^{a,h} ↑ Peripheral blood ^f ↑ Spleen ^f ↑↑ Prox small intestine ^b = Prox small intestine ^g	↑ haSMC ^a ↑ Liver ^b ↑ Peripheral blood ^f ↑ Spleen ^f ↑ Prox small intestine ^{b,g}	↑ THP-1 ↑ RCT ^b	Miao et al., 2004; Quinet et al., 2004, 2006; Brunham et al., 2006; Naik et al., 2006; Delvecchio et al., 2007; DiBlasio-Smith et al., 2008; Kannisto et al., 2014

(continued)

TABLE 1—Continued

Compound	Primary Action	ABCA1	ABCG1	Cholesterol Efflux	Reference
Ibandronate	Osteoclast inhibitor	↑ Kidney ^{b,h} ↑ Duodenum ^{b,h} ↑ MM6	—	↑ MM6	Strobach and Lorenz, 2003
Ibrolipim (NO-1886)	Lipoprotein lipase activator	↑ PBMCS ^a ↑↑ THP-1 ↑↑ Liver ^j ↑↑ Adipose tissue ^j ↑↑ Aorta ^j	↑↑ THP-1	↑ THP-1	Zhang et al., 2006; Ma et al., 2009; Chen et al., 2010
Idarubicin	Anthracycline antibiotic	↑ HL-1	↑ HL-1	—	Monzel et al., 2017
IMB-151	Unknown	↑↑ RAW264.7	↑↑ RAW264.7	↑ RAW264.7	Li et al., 2014
Infliximab	Anti-TNF- α mAb	↑↑ THP-1	= THP-1	↑ THP-1 (restores effect of TNF- α)	Voloshyna et al., 2014
Lansoprazole	Proton pump inhibitor	↑ H4 neuroglioma cells ^a ↑ U-87 astrocytoma ↑↑ CCF astrocytoma ↑ U-118 astrocytoma ↑ Primary astrocytes ^b	—	—	Cronican et al., 2010
LXR-623	Synthetic LXR agonist	↑ Duodenum ^g = Liver ^g ↑ Spleen ^f ↑ Peripheral blood ^{a,b,f} ↑↑ PBMCS ^a ↑ Whole blood ^{a,e} ↑ THP-1 ↑ Aorta ^c	↑ Duodenum ^g = Liver ^g ↑ Spleen ^f ↑ Peripheral blood ^{a,b,f} ↑↑ PBMCS ^a ↑ Whole blood ^{a,e}	—	DiBlasio-Smith et al., 2008; Katz et al., 2009; Quinet et al., 2009
Methyl-3 β -hydroxy-5 α ,6 α -epoxycholesterol	LXR α agonist	↑ THP-1	—	—	Yan et al., 2010
Omeprazole	Proton pump inhibitor	↑ H4 neuroglioma cells ^a	—	—	Cronican et al., 2010
Pantoprazole	Proton pump inhibitor	↑ H4 neuroglioma cell ^a	—	—	Cronican et al., 2010
Ouabain	Na ⁺ -K ⁺ -ATPase inhibitor	↑ H9c2	—	↑ H9c2	Campia et al., 2012
Ritonavir	Viral proteinase inhibitor	↑↑ THP-1	= THP-1	—	Pou et al., 2008
Sirolimus	FK-binding protein-12 inhibitor	↑ hVSMC	= hVSMC	= hVSMC	Ma et al., 2007
(24S)-stigmasta-5,28-diene-3 β ,24-ol	LXR agonist	↑ U937	—	—	Castro Navas et al., 2018
(24S)-stigmasta-5-ene-3 β ,24-ol	LXR agonist	↑ U937	—	—	Castro Navas et al., 2018
Stigmasterol derivatives T0901317	LXR agonist LXR agonist	↑ U937 ↑↑ THP-1 ↑↑ RAW264.7 ↑ J774 ↑ U937 ↑↑ HepG2 ↑↑ Caco-1 ↑ HL-1 ↑↑ pBCECs	↑ THP-1 ↑↑ RAW264.7 ↑ MCF-7 ↑ Caco-1 ↑ HL-1 ↑ TR-CSFB3 ↑ haSMC ^a ↑ Cerebral endothelial cells	↑ THP-1 ↑ HDL ↑ J774 ↑ Caco-1 ↑ HL-1 ↑ apoA-I ↑ HDL ↑ SAS	Marinozzi et al., 2017 Fukumoto et al., 2002; Murthy et al., 2002; Terasaka et al., 2003; Thomas et al., 2003; Beyer et al., 2004; Miao et al., 2004; Quinet et al., 2004, 2006; Wu et al., 2004; Panzenboeck et al., 2006; Wang et al., 2006; Delvecchio et al., 2007, 2008; Fujiyoshi et al., 2007; Sprecher et al., 2007; Dai et al., 2008; DiBlasio-Sato et al., 2008; Smith et al., 2008; Zanotti et al., 2008; Larrede et al., 2009; Verschuren et al., 2009; Mogilenko et al., 2010; Morrow et al., 2010; Yan et al., 2010; Maejima et al., 2011; Honzumi et al., 2011; Chen et al., 2012; Di et al., 2012; El Roz et al., 2012; Elali and Hermann, 2012;
		↑ TR-CSFB3 ↑↑ SAS	↑↑ Blood-derived macrophages ^a ↑ Aorta endothelial cells ^c	↑ Jurkat ↑ Fu5AH	
		↑↑ haSMC ↑↑ McARH7777 ↑↑ CD4 ⁺ T cells ^a	↑ Liver ^{b,j} ↑ Aorta ^{i,o} ↑ Peripheral blood ^{e,f}	↑ COS-7 ↑ pBCECs ↑ Monocyte-derived macrophages ^a ↑ CD4 ⁺ T cells ^a ↑ HSKMc ^a	
		↑ Jurkat ↑ Cerebral endothelial cells			

(continued)

TABLE 1—Continued

Compound	Primary Action	ABCA1	ABCG1	Cholesterol Efflux	Reference
		↑ Murine immortal macrop.		haSMC ^a	Jiang et al., 2012; Ma et al., 2014; Kaneko et al., 2015; Kirchgesner et al., 2015; Manna et al., 2015; Tamehiro et al., 2015; Li et al., 2016; Carter et al., 2017; Jiang and Li, 2017; Marinozzi et al., 2017; Monzel et al., 2017; Kaseda et al., 2018
		↑ Murine neuro2A		= apoA-I	
		↑ Murine BV-2		↑ HDL	
		↑ Rat C6		MCF-7	
		↑↑ Blood-derived macrop ^g		= apoA-I	
		↑ Aorta endothelial cells ^c		↑ HDL	
		↑ Renal glomerular mesangial cells ⁱ		↑ Murine primary macrop.	
		↑↑ PM ^{b,d,g,h}		↑ PM ^{b,d,g,h}	
		↑↑ Liver ^{b,c,h,i,l,k}		↑ Murine immortal macrop.	
		↑↑ Aorta ^{b,c,i,k,o}		↑ Renal glomerular mesangial cells ⁱ	
		↑ Small intestine ^c			
		↑ Kidney ^{b,h}			
		↑ Duodenum ^{b,h}			
		↑ Peripheral blood ^f			
		↑ Proximal intestine ^k			
		↑ Distal intestine ^k			
		↑↑ Brain ^{m,n}			
Tacrolimus	FK-binding protein-12 inhibitor	↑= THP-1	—	—	Jin et al., 2004
Teniposide	DNA topoisomerase II inhibitor	↑ RAW264.7 ↑↑ PM ^b ↑ THP-1 ↑ Blood monocyte	↑ PM ^b	↑ PM ^b ↑ RCT	Zhang et al., 2013a
Topiramate	GABA-A receptor agonist	↑↑ MDM	↑↑ MDM	MDM ↑ apoA-I ↑ HDL	Yang et al., 2014
YC-1	Soluble guanylyl cyclase activator	↑↑ J774A.1 ↑ Aorta ^c	= J774A.1 = Aorta ^c	↑ J774A.1	Tsou et al., 2014

BMM, bone marrow– derived macrophage; BREC, bovine retinal endothelial cells; haSMC, human airway smooth muscle cells; HSKM, human skeletal muscle cells; hVSMC, human vascular smooth muscle cells; macrop, macrophage; MCF-7, Michigan Cancer Foundation-7; MDM, monocyte-derived macrophage; pBCECs, porcine brain capillary endothelial cells; PBMC, peripheral blood mononuclear cell; PM, peritoneal macrophages; prox, proximal; SAS, human squamous cell carcinoma cell line.

^aHuman.

^bC75BL/6 mice.

^c*apoE*^{−/−} C57BL/6 mice.

^d*LXRβ*^{−/−} C75BL/6 mice.

^eCynomolgus monkeys.

^fMale long Evans rats.

^g*LDLR*^{−/−} C75BL/6 mice.

^h*LXRα*^{−/−} C75BL/6 mice.

ⁱNew Zealand White rabbits.

^jMale chine Bama minipigs.

^kMale SD rats.

^l129Sv mice.

^m*APP/PS1Δ9/APOE4*^{+/+}/ABCA1^{+/−} C57 BL/6 mice.

ⁿ*APP/PS1Δ9/APOE3*^{+/+}/ABCA1^{+/−} C57 BL/6 mice.

^oE3L mice.

agonist E17110 (i.e., a benzofuran-2-carboxylate analog) and LXRα-specific synthetic chalcone derivative, (E)-1-(3,4-di-isopropoxyphenyl)-3-(4-isopropoxy-3-methoxyphenyl)prop-2-en-1-one, both induced ABCA1 expression in macrophages, whereas an enhanced cholesterol efflux was only described after treatment with E17110 (Li et al., 2016; Teng et al., 2018). Moreover, commonly used proton pump inhibitors

(i.e., lansoprazole, omeprazole, pantoprazole) demonstrated to act as LXR agonists, of which lansoprazole was most potent (Cronican et al., 2010). By inducing LXR, proton pump inhibitors enhanced ABCA1 expression in human and mouse cells (Cronican et al., 2010).

Type II DNA topoisomerase inhibitors, which are used as chemotherapeutic medication, also enhanced

ABCA1 expression in an LXR α -dependent manner (Zhang et al., 2013a; Tsou et al., 2014; Shen et al., 2015). Inhibition of type II DNA topoisomerase by etoposide and teniposide induced in vitro ABCA1 expression, free cholesterol efflux, and in vivo RCT. Teniposide exerted its effect by inhibiting expression of receptor-interacting protein 140, a corepressor of LXR, and consequently mediated its favorable effects at lower concentrations compared with etoposide (Zhang et al., 2013a). Like other LXR agonists, etoposide and teniposide both enhanced FAS expression, indicating that these inhibitors also activate genes involved in lipogenesis, which could limit their use in the treatment of atherosclerosis (Zhang et al., 2013a) in addition to their unfavorable cytostatic effects. In contrast, 3-(5'-hydroxymethyl-2'-furyl)-1-benzyl indazole (YC-1), which in addition to LXR also activates soluble guanylyl cyclase, upregulated ABCA1 expression, enhanced cholesterol efflux, and decreased ox-LDL particle accumulation in macrophages in an LXR α -dependent manner, whereas ABCG1, SR-B1, SR-1, and CD36 expression levels were not altered (Tsou et al., 2014).

Indirect activation of LXR by a synthetic sphingosine analog (i.e., 2-amino-2-[2-(4-*n*-octylphenyl)ethyl]-1,3-propanediol hydrochloride, FTY720-P) and a synthetic sulfonylurea compound (i.e., G004) also enhanced in vitro expression of *ABCA1* and *ABCG1* and cholesterol efflux (Blom et al., 2010). FTY720-P upregulated *ABCA1* expression in human macrophages via increased production of the endogenous LXR ligand 27-hydroxycholesterol, which subsequently enhanced LXR activity independent of sphingosine 1-phosphate receptor activation (Blom et al., 2010; Qian et al., 2017). In contrast, G004 increased *ABCA1* and *ABCG1* expression by targeting sirtuin 1 and thereby acting as an atheroprotective agent in vivo (Qian et al., 2017). Ibandronate, a bisphosphonate, also stimulated LXR indirectly, which was reversed upon addition of geranylgeranyl pyrophosphate, suggesting that low levels of this mevalonate pathway intermediate increased LXR expression via a negative feedback mechanism (Strobach and Lorenz, 2003). Anthracyclines, which are used as chemotherapeutic agents, also demonstrated to indirectly activate LXR-ABCA1/ABCG1 pathway by enhancing several oxysterol and cholesterol precursor levels, although a direct binding between anthracyclines and LXR cannot be excluded. Their clinical use is generally strongly limited due to cardiotoxic effects (Monzel et al., 2017), and their cytostatic effect omits the use of these drugs to stimulate RCT clinically. An indirect stimulatory effect on LXR expression by a yet unknown mechanism has also been observed with disodium ascorbyl phytostanol phosphate (FM-VP4), a potential cholesterol-lowering drug (Méndez-González et al., 2010).

The cardiac glycosides, digoxin and ouabain, stimulated cholesterol efflux via LXR-mediated upregulation of *ABCA1* expression in cardiomyocytes, but only slightly increased *ABCG1* expression (Campia et al., 2012). They also increased cholesterol and ubiquinone

synthesis via upregulation of HMG-CoA reductase expression without affecting the intracellular cholesterol concentration. This could be explained by the stimulation of cholesterol efflux in cardiomyocytes, which might also contribute to the beneficial effect of cardiac glycosides in CVD (i.e., next to their antiarrhythmic effects). However, further research is warranted to demonstrate their antiatherosclerotic potential. Finally, the anti-tumor necrosis factor (TNF)- α monoclonal antibody infliximab was linked to a reduced foam cell formation in THP-1 macrophages induced by TNF- α , which was dependent on the reversal of TNF- α -induced inhibition of cholesterol efflux and LXR- α , *ABCA1*, and *ABCG1* mRNA and protein expression (Voloshyna et al., 2014). This could also provide an explanation for the improved vascular function, as recently observed in patients with rheumatoid arthritis with TNF inhibitor therapy (Rongen et al., 2018). Although Voloshyna et al. (2014) demonstrated a lowering effect of TNF- α on LXR- α and *ABCA1* levels in THP-1 monocytes, no differences in the mRNA levels of the LXR-target gene *ABCA1* and apoA-I-dependent cholesterol efflux were reported in TNF- α -treated C57BL/6 mice peritoneal macrophages (Castrillo et al., 2003). The uncertainty about the exact role of TNF- α on LXR- α and *ABCA1* expression is further emphasized by different findings in several cell types. TNF- α reduced LXR- α expression and LXRE activity level in HK-2 proximal tubular cells and Hep3B liver cells, whereas in Caco-2 cells *ABCA1* was decreased without attenuation of LXR- α upon stimulation with TNF- α (Wang et al., 2005; Kim et al., 2007; Field et al., 2010). A stimulatory effect on cellular cholesterol efflux was found after lowering TNF- α production in human macrophages by GABA and the GABA agonist topiramate (Yang et al., 2014). These effects could not be observed with baclofen, another GABA agonist (Yang et al., 2014), which questions the direct involvement of GABA in the previously observed effects with GABA agonists. Interestingly, a reversal of TNF- α -induced cholesterol efflux inhibition could also be observed with the immunosuppressant sirolimus, which increased cholesterol efflux in human vascular smooth muscle cells accompanied by an increased *ABCA1* expression (Ma et al., 2007). Although the structurally related drug tacrolimus did also increase *ABCA1* expression, its effect seemed to be mediated via PPAR γ and not via a TNF- α -dependent mechanism (Jin et al., 2004). Because both immunosuppressants have been also associated with elevated plasma cholesterol and triglyceride levels (Włodarczyk et al., 2005; Kido et al., 2018), their clinical applicability to reduce atherosclerotic risk is limited.

Of all LXR agonists, only two (i.e., BMS-779788, also known as EXEL-04286652 and LXR-623) were studied in a clinical trial (Hong and Tontonoz, 2014). Unfortunately, no published report is available on the completed study of BMS-779788 (Hong and Tontonoz, 2014).

LXR-623 stimulated *ABCA1* and *ABCG1* expression in peripheral blood cells of healthy individuals, which is expected to result in an enhanced RCT. However, the phase I clinical trial was terminated due to unfortunate neurologic adverse effects at the two highest concentrations tested. It remains unknown whether this effect was LXR-mediated or via other unknown off-target mechanisms of LXR-623 (Katz et al., 2009).

C. Peroxisome Proliferator-Activated Receptor Activation to Enhance ATP-Binding Cassette A1 and ATP-Binding Cassette G1 Expression

Fibrates have been used for decades as cholesterol-lowering drugs, but currently they are mainly administered as comedication with statins to treat severe hypertriglyceridemia ($>15 \text{ mmol} \times \text{l}^{-1}$). Although fibrates are mainly agonists of PPAR α , they stimulate all three PPAR isoforms with different potencies, except for bezafibrate, which is an equipotent agonist of all isoforms. Gemfibrozil, fenofibrate, bezafibrate, and clofibrate all increased *ABCA1* mRNA and protein expression upon PPAR activation in vitro (Oliver et al., 2001; Forcheron et al., 2002; Lin and Bornfeldt, 2002; Guan et al., 2003; Ruan et al., 2003; Thomas et al., 2003; Arakawa et al., 2005; Kooistra et al., 2006; Hossain et al., 2008; Tanabe et al., 2008; Ogata et al., 2009; Kobayashi et al., 2011; Jiang and Li, 2017), which was associated with enhanced apoA-I-mediated cholesterol efflux (Oliver et al., 2001; Ruan et al., 2003; Arakawa et al., 2005; Kooistra et al., 2006; Ogata et al., 2009) (Table 2). Next to an effect on peripheral *ABCA1* expression, fenofibrate and gemfibrozil also induced hepatic *ABCA1*, *ABCG1*, and *ABCG5/8* mRNA expression in a PPAR α -dependent manner (Hossain et al., 2008; Rotllan et al., 2011). This positive effect on *ABCA1* expression could also in part explain the antiatherosclerotic effects observed with fibrates in clinical trials (Steiner, 2005). The coronary risk reduction found in these trials could also be related to their LDL-C-lowering capacity. However, the therapeutic value of fibrates remains equivocal due to absence of a reduction in total cardiovascular mortality in several trials. Two large-scale intervention studies with fenofibrate (FIELD trial and ACCORD-LIPID trial) in patients with type 2 diabetes mellitus found no differences in coronary events, nonfatal myocardial infarction, and stroke (Keech et al., 2005; Ginsberg et al., 2010). Similar results were reported in two intervention trials with bezafibrate, in which no significant effect was found on both fatal and nonfatal myocardial infarction in the Bezafibrate Infarction Prevention (BIP) study, and no reduction in the incidence of coronary heart disease and of strokes in the LEADER trial (Bezafibrate Infarction Prevention (BIP) Study, 2000; Meade et al., 2002). In contrast, two trials using gemfibrozil (Helsinki Heart Study (HHS) trial and Veterans Affairs-HDL Intervention Trial (VA-HIT)

study] showed a significant reduction in the incidence of coronary heart disease and myocardial infarction or cardiovascular death, demonstrating an overall benefit of gemfibrozil treatment (Frick et al., 1987; Rubins et al., 1999).

Like gemfibrozil and fenofibrate, the specific PPAR α agonist, Wy14643, enhanced *ABCA1* protein expression and apoA-I-mediated cholesterol release in vitro, primarily mediated via LXR α activation (Chinetti et al., 2001; Ruan et al., 2003; Thomas et al., 2003; Beyer et al., 2004; Arakawa et al., 2005; Lee et al., 2008; Maejima et al., 2011). Although beneficial effects on *ABCA1* expression and cholesterol efflux were observed after treatment with gemfibrozil, fenofibrate, and Wy14643, all three compounds stimulated PPAR α with a relatively low affinity in the micromolar range (Ferri et al., 2017). This initiated the exploration of high-affinity PPAR α agonists like GW7647, which effectively increased *ABCA1* mRNA. However, no effect of GW7647 on apoA-I-driven cholesterol efflux was observed in THP-1 macrophages (Oliver et al., 2001; Li et al., 2004; Wang et al., 2010; Nakaya et al., 2011). Another high-affinity PPAR α agonist, LY518674, did increase both *ABCA1* expression and apoA-I-mediated cholesterol efflux via LXR α activation, resulting in increased HDL biogenesis, whereas Wy14563 only increased cholesterol efflux via PPAR α stimulation (Chawla et al., 2001; Hossain et al., 2008; Ogata et al., 2009; Ferri et al., 2017). Surprisingly, high aspirin concentrations ($\geq 250 \mu\text{M}$) slightly enhanced *ABCA1* expression and cholesterol efflux in macrophages in a PPAR α -dependent manner (Viñals et al., 2005; Wang et al., 2010).

Statins most likely also affect *ABCA1*-dependent cholesterol efflux beneficially via indirect stimulation of PPAR α (Zanotti et al., 2004, 2006; Argmann et al., 2005; Kobayashi et al., 2011; Maejima et al., 2011; Song et al., 2011; Shimizu et al., 2014; Nicholls et al., 2015). However, the results of a variety of studies are equivocal and report large differences with different statins in vitro and in vivo (e.g., in various cell types, animal models, and patients). In hepatocytes, only pitavastatin seemed to enhance *ABCA1* mRNA expression at low micromolar concentrations in all studies (Zanotti et al., 2004, 2006; Kobayashi et al., 2011; Maejima et al., 2011; Song et al., 2011). Mechanistically, PPAR α activation was seen in all studies, but LXR was activated as well as suppressed in hepatocytes after statin treatment. Moreover, this effect seems to depend on downstream products of the cholesterol synthesis pathway (e.g., mevalonate, geranylgeranyl pyrophosphate), with a major role for Ras homolog gene family member A, which fully reversed statin-mediated *ABCA1* upregulation and *ABCA1*-mediated cholesterol efflux (Zanotti et al., 2004; Argmann et al., 2005). Not all statins did increase macrophage *ABCA1* expression in vitro (Zanotti et al., 2006), but rosuvastatin exposure in vivo increased total and *ABCA1*-dependent cholesterol efflux (Shimizu et al.,

TABLE 2
PPAR-activating compounds

Overview of compounds that activate PPARs, including their primary pharmacological action, effects on ABCA1 and ABCG1 expression (arrows: *mRNA*, protein), and effects on cellular cholesterol efflux. Effects on ABCA1, ABCG1, and cholesterol efflux are presented as follows: (↓) decreased; (=) no effect; (↑) increased. Italicized symbols indicate changes in *ABCA1* and *ABCG1* mRNA levels, whereas nonitalicized symbols indicate protein levels.

Compound	Primary Action	ABCA1	ABCG1	Cholesterol Efflux	Reference
PPAR γ -activating compounds					
15d-PGJ2	Prostanoid-specific receptor inhibitor	↑ THP-1 ↑ HMC ↓ PM ^a	↑ PM ^a	Lipid-loaded HMC ↑ apoA-I PM ^a ↓ apoA-I ↑ HDL	Ruan et al., 2003; Jiang and Li, 2017
4010B-30	Benzamide analog	↑ RAW264.7 ↑ HepG2	—	RAW264.7 ↑ apoA-I	Du et al., 2015b
Ciglitazone	PPAR γ agonist	↑ THP-1	↑ THP-1	↑ THP-1	Argmann et al., 2003, 2005
E3317	PPAR γ agonist	↑ RAW264.7 ↑ LO2	—	↑ RAW264.7 ↑ apoA-I	Wang et al., 2018
GQ-11	PPAR γ/α agonist	↑ Liver ^b	—	—	Silva et al., 2018
GW1929	PPAR γ agonist	↑ HepG2	—	—	Mogilenko et al., 2010
GW7845	PPAR γ agonist	↑ THP-1	—	↑ THP-1	Chawla et al., 2001; Oliver et al., 2001
Propofol	GABA _B receptor agonist	↑ THP-1 ↑ RAECs	↑ THP-1	↑ THP-1 ↑ RAECs	Ma et al., 2015; Hsu et al., 2018
Lysophosphatidylcholine	Unknown	↑ PM ^a	—	↑ HDL	Hou et al., 2007
Mycophenolic acid	Inosine-5'-monophosphate dehydrogenase inhibitor	↑ HepG2	—	↑ PM ^a	Xu et al., 2011
Pioglitazone	PPAR γ agonist	↑ THP-1 ↑ HepG2	↑ THP-1 ↑ Monocyte-derived macrop	↑ THP-1 ↑ apoA-I	Panzenboeck et al., 2006; Nakaya et al., 2007; Tanabe et al., 2008; Ogata et al., 2009; Cocks et al., 2010; Ozasa et al., 2011; Wang et al., 2014b, 2015b; Jiang and Li, 2017; Silva et al., 2018
		↑ pBCECs ↑ gBECs	↑ PM ^a ↑ Diabetic patients ^c	↑ HDL ↓ pBCECs	
		↑ WI38 fibroblasts	↑ Rat cortical neurons	↑ gBECs	
		↑ Monocyte-derived macrop ^c		↑ WI38 fibroblasts	
		↓ PM ^a = Liver ^b		PM ^a	
		↑ Diabetic patients ^c		↓ apoA-I	
		↑ Rat cortical neurons		↑ HDL	
				↑ Diabetic patients ^c	
Rosiglitazone	PPAR γ agonist	↑ THP-1 ↑ RAW264.7 ↑ HepG2 == macrop ^d ↑ PM ^e ↑ Hepatocytes ^e = Aorta ^d Aorta lesion ^e	↑ THP-1 ↑ macrop ^d ↑ Aorta ^d	↑ THP ↑ RAW264.7 ↑ macrop ^d ↑ PM ^e ↑ Hepatocytes ^e	Chawla et al., 2001; Chinetti et al., 2001; Claudel et al., 2001; Li et al., 2004, 2015; Llaverias et al., 2006
Telmisartan	Angiotensin receptor 1 antagonist	↑ THP-1 ↑ Monocyte-derived macrop ^c	↑ THP-1 ↑ Monocyte-derived macrop ^c	↑ THP-1	Nakaya et al., 2007
Troglitazone	PPAR γ agonist	↑ THP-1 ↑ pBCECs ↑ gBECs = Monocyte-derived macrop ^c ↓ PM ^a	↑ PM ^a	↑ pBCECs PM ^a ↓ apoA-I ↑ HDL	Cabrero et al., 2003; Panzenboeck et al., 2006; Lee et al., 2008; Jiang and Li, 2017
PPAR α -activating compounds					
Aspirin	COX-1/2 inhibitor	↑ THP-1 ↑ RAW264.7	—	RAW264.7 ↑ apoA-I	Viñals et al., 2005; Wang et al., 2010
Atorvastatin	HMG-CoA reductase inhibitor	↑ THP-1 ↑ McARH7777	↑ THP-1	↑ THP-1 ↑ apoA-I ↑ HDL	Argmann et al., 2005; Maejima et al., 2011; Nicholls et al., 2015, 2017
Bezafibrate	Pan-PPAR agonist	↑ THP-1 ↑ HepG2 == pBCECs ↑ W138 fibroblast ↑ HMC	—	↑ THP-1 ↑ HepG2 = pBCECs ↑ W138 fibroblast	Cabrero et al., 2003; Ruan et al., 2003; Panzenboeck et al., 2006; Hossain et al., 2008; Inaba et al., 2008; Ogata et al., 2009

(continued)

TABLE 2—Continued

Compound	Primary Action	ABCA1	ABCG1	Cholesterol Efflux	Reference
				Lipid-loaded HMC apoA-I	
Clofibrate	PPAR α agonist	↑↑ Primary hepatocytes = Monocyte-derived macrop ^c = aorta ^b ↑ HepG2 = Primary hepatocytes ^f	—	↑ Primary hepatocytes	Guan et al., 2003; Kobayashi et al., 2011
Fenofibrate	PPAR α agonist	↑ Liver ^f ↑↑ THP-1 ↑↑ RAW264.7 ↑↑ HepG2 =↑ pBCECs ↑ Balb/3T3 ↑ W138 fibroblast	↑ Liver ^g	↑ THP-1 ↑ RAW264.7 ↑ HepG2 = pBCECs = Balb/3T3 ↑ W138 fibroblast ↑ Primary hepatocytes	Forcheron et al., 2002; Lin and Bornfeldt, 2002; Cabrero et al., 2003; Thomas et al., 2003; Arakawa et al., 2005; Kooistra et al., 2006; Hossain et al., 2008; Tanabe et al., 2008; Ogata et al., 2009; Jiang and Li, 2017
Gemfibrozil	PPAR α agonist	↑ PM ^a ↑ Liver ^h ↑ Diabetic patients ^c ↑ Aorta ⁱ ↑↑ THP-1 ↑↑ HepG2 ↑↑ W138 fibroblast	—	↑ THP-1 ↑ HepG2 ↑ W138 fibroblast	Hossain et al., 2008; Ogata et al., 2009
GW7647	PPAR α agonist	↑↑ Primary hepatocytes ↑ THP-1 ↑↑ RAW264.7 == macrop ^d == Atherosclerotic lesion ^d	= macrop ^d = Aorta ^d ↑↑ BMM ^a	= THP-1 = RAW264.7 ↑ apoA-I = macrop ^d	Oliver et al., 2001; Li et al., 2004; Wang et al., 2010; Nakaya et al., 2011
LY518674	PPAR α agonist	↑↑ BMM ^a ↑↑ THP-1 ↑↑ HepG2 ↑↑ W138 fibroblast	—	↑ BMM ^a ↑ THP-1 ↑ HepG2 ↑ W138 fibroblast	Hossain et al., 2008; Ogata et al., 2009
Pitavastatin	HMG-CoA reductase inhibitor	↑↑ Primary hepatocytes ↓ J774 ↑ HepG2 ↑↑ McARH7777 ↑ Liver ^f	—	↑ Primary hepatocytes ↓ J774 ↑ Fu5AH ↓ PM ^{a,b,d} = LXR ^{-/-} mice	Zanotti et al., 2004, 2006; Kobayashi et al., 2011; Maejima et al., 2011
Pravastatin	HMG-CoA reductase inhibitor	↑↑ 3T3-L1 = McARH7777	↓↓ 3T3-L1	↓ 3T3-L1	Maejima et al., 2011; Mostafa et al., 2016
Rosuvastatin	HMG-CoA reductase inhibitor	= Hepatocytes ^a		↑ J774 ↑ BMM ^a ↑ RCT ^a	Shimizu et al., 2014; Mostafa et al., 2016
Simvastatin	HMG-CoA reductase inhibitor	↑ McARH7777 ↑ Diabetic patients with hyperlipidemia == PM ^d ↑↑ Liver ⁱ ↑↑ THP-1 ↑↑ RAW264.7 ↑↑ gBECs ↑↑ Balb/3T3 = McARH7777 ↑ Liver ^a	== PM ⁱ == Liver ⁱ	↑ THP-1 ↑ apoA-I	Argmann et al., 2005; Guan et al., 2008; Maejima et al., 2011; Song et al., 2011; Ying et al., 2013; Gong et al., 2014
WY14643	PPAR α agonist	↑ Liver ⁱ ↑↑ THP-1 ↑↑ RAW264.7 ↑↑ gBECs ↑↑ Balb/3T3	↑ THP-1	↑ HDL ↑ RAW264.7 ↑ THP-1 ↑ RAW264.7 = Balb/3T3 = Lipid-loaded HMC ↑ apoA-I	Chinetti et al., 2001; Ruan et al., 2003; Beyer et al., 2004; Arakawa et al., 2005; Lee et al., 2008; Maejima et al., 2011
Wy,14,563	PPAR δ/β -activating compounds	—	—	↑ THP-1	Chawla et al., 2001
Carbaprostacyclin (cPGI)	PPAR δ agonist	—	—	↑ THP-1	Chawla et al., 2001
GW0742	PPAR δ agonist	= Liver ^a = Small intestine ^a	= Liver ^a = Small intestine ^a	↑ BMM ^a = macrop ^a	Li et al., 2004; Briand et al., 2009

(continued)

TABLE 2—Continued

Compound	Primary Action	ABCA1	ABCG1	Cholesterol Efflux	Reference
		= = macrop ^d = Atherosclerotic lesion ^d	= macrop ^d = Aorta ^d		
GW501515	PPARδ agonist	↑↑ THP-1	= HSKM	↑ THP-1	Oliver et al., 2001; Sprecher et al., 2007; Ogata et al., 2009
		↑↑ W138 fibroblast		↑ W138 fibroblast	
		↑ 1BR3N fibroblast		↑ 1BR3N fibroblast	
		↑ Intestinal FHS74		↑ Intestinal	
		↑ HSKM		↑ HSKM	

BMM, bone marrow–derived macrophage; gBECS, gallbladder epithelial cells; HMC, human mast cell; HSKM, human skeletal muscle cell; macrop, macrophage; pBCECs, porcine brain capillary endothelial cells; PM, peritoneal macrophages; RAECs, rat aortic endothelial cells.
^aC75BL/6 mice.
^b*LDLR*^{−/−} C75BL/6 mice.
^cHuman.
^d*LDLR*^{−/−} C75BL/6 hypercholesterolemic mice.
^eNew Zealand White rabbits.
^fMale Wistar rats.
^gMale Zucker diabetic fatty rats.
^h129SV mice.
ⁱFemale E3L transgenic mice.

2014). Although cotreatment with atorvastatin and evacetrapib revealed an enhanced cholesterol efflux, whereas single treatment with atorvastatin resulted in a decreased cholesterol efflux, the clinical relevance is limited, because evacetrapib also increased overall atherogenic risk (Nicholls et al., 2017). Therefore, the absence of an effect or even negative effect of statins on ABCA1-mediated cholesterol efflux in some clinical studies (Nicholls et al., 2015, 2017), as well as the high incidence of dose-dependent muscle complaints associated with these drugs, limits their ability to increase RCT.

Next to compounds that beneficially affect PPARα, a vast number of drugs can stimulate PPARγ. Thiazolidinediones are probably the most well-known and widely used PPARγ agonists, due to their antidiabetic properties. Thiazolidinediones, including pioglitazone, ciglitazone, and rosiglitazone, enhanced LXRα expression, which subsequently induced ABCA1 protein expression and cholesterol efflux toward apoA-I in vitro (Chawla et al., 2001; Chinetti et al., 2001; Claudel et al., 2001; Cabrero et al., 2003; Li et al., 2004, 2015; Llaverias et al., 2006; Panzenboeck et al., 2006; Nakaya et al., 2007; Lee et al., 2008; Tanabe et al., 2008; Ogata et al., 2009; Cocks et al., 2010; Ozasa et al., 2011; Wang et al., 2014b, 2015b; Jiang and Li, 2017) (Table 2). Contradictory to these results with pioglitazone in THP-1 macrophages and J774 macrophages, an attenuated or unaffected ABCA1 expression was found in peritoneal macrophages isolated from 15PGJ2-, troglitazone-, and pioglitazone-treated C57BL/6 mice, and in the liver of pioglitazone-treated *LDLR*^{−/−} C57BL/6 mice (Ruan et al., 2003; Ogata et al., 2009; Ozasa et al., 2011; Zhao et al., 2015; Jiang and Li, 2017; Silva et al., 2018). Clinically, the cardioprotective antiatherosclerotic effects of thiazolidinediones have been heavily debated (Liebson, 2010). Although increased HDL-C levels were observed after pioglitazone and rosiglitazone treatment, increased cardiovascular morbidity and heart failure have been associated with rosiglitazone (Gerstein et al., 2006;

Home et al., 2009; Liebson, 2010; Chandra et al., 2017). Although in various clinical trials cardioprotective effects have been observed with pioglitazone (Chandra et al., 2017), its chronic use is limited by the observed association with an increased risk of developing bladder cancer, bone fractures, and congestive heart failure (Tang et al., 2018). Another PPARγ agonist, GW7845, which is an analog of the PPARα agonist GW7647, did not affect apoA-I-mediated cholesterol efflux, whereas *ABCA1* mRNA expression was enhanced (Chawla et al., 2001; Oliver et al., 2001). The PPARγ agonist, GW1929, reduced ABCA1 protein expression in HepG2 cells, whereas *ABCG1* gene expression was increased probably due to an interplay between PPARγ and LXRβ, resulting in dissociation of LXRβ from the ABCA1/LXRβ complex (Mogilenko et al., 2010). In contrast, mycophenolic acid, 4010B-30, telmisartan, propofol, and lysophosphatidylcholine exposure did result in an increased in vitro ABCA1 expression and apoA-I-mediated cholesterol efflux, mediated via the PPARγ–LXRα–ABCA1 axis (Hou et al., 2007; Nakaya et al., 2007; Xu et al., 2011; Du et al., 2015b; Ma et al., 2015; Hsu et al., 2018). Lysophosphatidylcholine treatment enhanced apoE secretion from peritoneal macrophages (Hou et al., 2007) and 4010B-30 enhanced apoA-I production, whereas telmisartan and propofol also enhanced ABCG1 expression in macrophages (Nakaya et al., 2007; Du et al., 2015b; Ma et al., 2015). The clinical antiatherosclerotic applicability of propofol seems limited, because of its sedative effect and risk of hypertriglyceridemia induced by the lipid emulsion formulation (Eddleston and Shelly, 1991). Additionally, E3317 enhanced in vitro ABCA1 expression and apoA-I-dependent cholesterol efflux via PPARγ, and a novel thiazolidine, GQ-11, which is a partial PPARγ and PPARα agonist, increased hepatic *ABCA1*, *APOA-I*, and *HDL* mRNA expression in *LDLR*^{−/−} C57BL/6 mice (Silva et al., 2018; Wang et al., 2018). Thus, PPARγ agonists show controversial results in the stimulation of ABCA1 expression, which could be due to their

relatively low potency for PPAR α and high PPAR γ potency.

Although explored to a lesser extent, activation of PPAR β/δ may be promising, as activation with GW501515 promoted RCT by enhancing in vitro apoA-I and HDL levels, *ABCA1* expression, and apoA-I-mediated cholesterol efflux (Oliver et al., 2001; Sprecher et al., 2007; Ogata et al., 2009) (Table 2). Another PPAR β/δ agonist, GW0742, did not affect *ABCA1* or *ABCG1* mRNA expression and only stimulated cholesterol efflux in bone marrow-derived macrophages, but not in peritoneal macrophages from *LDLR*^{-/-} mice (Li et al., 2004; Briand et al., 2009). Carbaprostacyclin, a PPAR β/δ agonist, only increased cholesterol efflux in macrophages (Chawla et al., 2001).

Combinations of PPAR and LXR agonists were investigated to overcome disadvantages observed with either PPAR or LXR agonists (e.g., increased plasma triglyceride concentrations), which demonstrated to be a promising strategy. Fenofibrate could abolish the LXR-mediated induction of *SREBP1* by T0901317 in THP-1 macrophages without affecting the endogenous *ABCA1* expression (Thomas et al., 2003). Similar results were observed after coadministration of T0901317 and Wy14643, as shown by the attenuation of T0901317-induced increases in serum and plasma triglycerides, without posing an effect on *ABCA1* mRNA levels in C57BL/6 mice (Beyer et al., 2004).

In summary, both PPAR α and PPAR γ agonists increase *ABCA1* and *ABCG1* expression in an LXR α -dependent manner, whereas PPAR α agonists were more effective inducers of apoA-I levels and apoA-I-mediated cholesterol efflux. Furthermore, coadministration of LXR and PPAR α agonists may overcome the adverse effects on plasma triglyceride levels of LXR agonists. However, more studies are needed to elucidate the mechanism and effectivity of such combinatorial strategies for treatment of atherosclerosis.

D. Enhancement of ATP-Binding Cassette A1 and ATP-Binding Cassette G1 Expression by Retinoid X Receptor Agonists

The notion that RXR activates and orchestrates the signaling of other nuclear receptors, including LXR and PPAR, led to the idea that RXR stimulation could be of special interest to stimulate RCT (Costet et al., 2003; Nishimaki-Mogami et al., 2008; Cui et al., 2011; Zhang et al., 2013a; Zhou et al., 2015). Such crosstalk with other nuclear receptors is illustrated by RXR/LXR α heterodimer formation after stimulation of RXR by endogenously synthesized 9-*cis*-RA (Manna et al., 2015; Zhou et al., 2015) and with synthetic RXR agonists, including PA024 and HX630 (Nishimaki-Mogami et al., 2008; Zhou et al., 2015), which all increased *ABCA1* expression (Table 3). In RAW264 cells, PA024, unlike HX630, directly influenced LXRE and positively modified the promoter activity to enhance *ABCA1* mRNA expression, resulting in generation of HDL particles and stimulation

of cholesterol efflux (Nishimaki-Mogami et al., 2008). In contrast, HX630 is expected to stimulate *ABCA1* expression by activating PPAR γ /RXR heterodimer, leading to an enhanced LXR expression in RAW264 cells (Nishimaki-Mogami et al., 2008). Similar mechanisms were observed with two other RXR agonists, tributyltin chloride and LG268, which activated PPAR γ /RXR and LXR α /RXR signaling. This treatment modulated cellular lipid homeostasis and cholesterol efflux in RAW264 and THP-1 macrophages via increased expression of *ABCA1* and *ABCG1* (Repa et al., 2000; Chawla et al., 2001; Cui et al., 2011; Sun et al., 2015). Additionally, LG101305 enhanced *ABCA1* mRNA expression and cholesterol efflux in macrophages (Claudel et al., 2001). The RXR agonists, bexarotene and methoprene, enhanced *ABCA1* expression and cholesterol efflux in astrocytes and increased the overall cerebral cholesterol efflux into the circulation (LaClair et al., 2013; Kuntz et al., 2015; Tachibana et al., 2016).

However, as RXR interacts with many other nuclear receptors, RXR agonists are associated with a wide spectrum of adverse events (Costet et al., 2003), including enhanced lipogenesis due to LXR/RXR heterodimerization (Manna et al., 2015), which severely limits their therapeutic potential to stimulate RCT.

Compounds that stimulate RAR do not suffer from these adverse mechanisms, as they are less ambiguous in the targets they activate. The RAR agonists, 4-[(E)-2-(5,6,7,8-Tetrahydro-5,5,8,8-tetramethyl-2-naphthalenyl)-1-propenyl]benzoic acid (TTNPB) and Am580, enhanced *ABCG1* expression by modulating *ABCG1* promoter activity, especially by interacting with LXRE-B in macrophages (Ayaori et al., 2012). Furthermore, TTNPB stimulated *ABCA1* expression in mouse and human macrophages mediated via a stimulatory effect of RAR γ /RXR on the *ABCA1* promoter (Costet et al., 2003). Chen et al. (2011b) found opposite effects of TTNPB in astrocytes, indicating that the effects may be cell-type dependent. The beneficial effects in macrophages are, however, most relevant for a potential antiatherosclerotic effect of RAR agonists, which makes them an interesting class of compounds to target RCT.

V. ATP-Binding Cassette A1 and ATP-Binding Cassette G1 mRNA Stability

A. mRNA Degradation as a Post-Translational Mechanism to Regulate ATP-Binding Cassette A1 and ATP-Binding Cassette G1 Expression

Another mechanism regulating plasma membrane abundance of *ABCG1* and *ABCA1* transporters is the rapid post-translational degradation of their mRNA transcripts. For *ABCG1* this depends on the interaction of the 3' untranslated region of its transcript with miRNAs, resulting in repression of translation or mRNA degradation (Li et al., 2010; Rayner et al., 2011; Rotllan and Fernandez-Hernando, 2012;

TABLE 3
Synthetic retinoid nuclear receptor agonists

Overview of retinoid nuclear receptor agonists, including their primary pharmacological action, effects on ABCA1 and ABCG1 expression (arrows: *mRNA*, protein), and effects on cellular cholesterol efflux. Effects on ABCA1, ABCG1, and cholesterol efflux are presented as follows: (↓) decreased; (=) no effect; (↑) increased. Italicized symbols indicate changes in *ABCA1* and *ABCG1* mRNA levels, whereas nonitalicized symbols indicate protein levels.

Compound	Primary Action	ABCA1	ABCG1	Cholesterol Efflux	Reference
<i>RXR agonists</i>					
Bexarotene	pan-RXR agonist	↑↑ BLECs ↑ Cortex ^a ↑ Cortex ^b ↑ Cortex ^c	—	↑ BLECs	LaClair et al., 2013; Kuntz et al., 2015; Tachibana et al., 2016
HX630	RXR agonist	↑ THP-1 ↑ RAW264.7	↑ THP-1	↑ THP-1	Nishimaki-Mogami et al., 2008
LG101305	RXR agonist	↑ RAW264.7	—	↑ RAW264.7	Claudel et al., 2001
LG268	RXR agonist	↑ Small intestine ^d ↑ PM	—	↑ THP-1	Chawla et al., 2001
Methoprene	RXR agonist	↑ Astrocyte	↑ Astrocyte	Astrocyte ↑ apoA-I ↑ HDL	Repa et al., 2000; Chen et al., 2011b
PA024	RXR agonist	↑ THP-1 ↑ RAW264.7	↑ THP-1	↑ THP-1	Nishimaki-Mogami et al., 2008
Tri-butyltin chloride	RXRα agonist	↑↑ RAW264.7 ↑↑ Primary mouse astrocyte cells ↑↑ Cortex ^c	↑↑ Primary mouse astrocyte cells ↑↑ Cortex ^c	↑ RAW264.7	Cui et al., 2011; Sun et al., 2015
<i>RAR agonists</i>					
AM580	RARα agonist	—	↑ THP-1	—	Ayaori et al., 2012
TTNPB	Synthetic RAR agonist	↑ THP-1 ↑ HEK293 ↓ Astrocytes = Liver ^d ↑ PM ^d	↑ THP-1 ↓ Astrocytes = Liver ^d	↑ RAW264.7 ↓ Astrocytes	Costet et al., 2003; Chen et al., 2011b; Ayaori et al., 2012

BLEC, bovine lens epithelial cells; PM, peritoneal macrophages.
^aLrp1^{fllox/flox}; αCamKII-Cre^{-/-} mice.
^bLrp1^{fllox/flox}; αCamKII-Cre^{+/-} mice.
^cAPPSWE/PSE1^{ΔE} mice.
^dC75BL/6/A129Sv mice.

Lv et al., 2014). Consequently, these small noncoding RNAs appear to be important modulators of gene expression (Rotllan and Fernandez-Hernando, 2012), which upon binding with other mRNA regions can lead to degradation as well as increased mRNA expression and translation (Rotllan and Fernandez-Hernando, 2012) (Fig. 5). Several miRNAs, which are summarized in Table 4, are known to affect *ABCA1* mRNA expression. Although less is known about *ABCG1* regulation, some miRNAs directly decreased *ABCG1* mRNA expression levels (Table 4) Moore et al., 2010; 2011; Fernández-Hernando et al., 2011; Fernández-Hernando and Moore, 2011; Rayner et al., 2011, 2012; Hazen and Smith, 2012; Iatan et al., 2012; Rotllan and Fernandez-Hernando, 2012; Sun et al., 2012; Wang et al., 2012, 2014a; Adlakha et al., 2013; Dávalos and Fernandez-Hernando, 2013; de Aguiar Vallim et al., 2013; Kang et al., 2013; Ramírez et al., 2013; Canfrán-Duque et al., 2014; Goedeke et al., 2014; Mao et al., 2014; DiMarco and Fernandez, 2015; He et al., 2015; Mandolini et al., 2015; Yang et al., 2015; Feinberg and Moore, 2016; Ono, 2016; Rotllan et al., 2016; Aryal et al., 2017).

B. Targeting Post-Transcriptional Regulation of ATP-Binding Cassette A1 mRNA

The therapeutic potential of miRNA in the regulation of cholesterol metabolism will not be further addressed

in this review, as this was extensively reviewed by others (Moore et al., 2010, 2011; Fernández-Hernando and Moore, 2011; Rotllan and Fernandez-Hernando, 2012; Dávalos and Fernandez-Hernando, 2013; Canfrán-Duque et al., 2014; Goedeke et al., 2014; DiMarco and Fernandez, 2015; Rotllan et al., 2016; Aryal et al., 2017). In this study, other drugable mechanisms involved in the post-transcriptional regulation of *ABCA1* and *ABCG1* mRNA expression will be discussed (Moore et al., 2011; Rotllan and Fernandez-Hernando, 2012; Dangwal and Thum, 2014; van Rooij and Kauppinen, 2014; Rotllan et al., 2016). One of these mechanisms is mediated by the cellular energy sensor AMP-activated protein kinase (AMPK), through its regulation of cholesterol metabolism. AMPK activation by 5-aminoimidazole-4-carboxyamide ribonucleoside (AICAR; an AMP mimetic) has led to enhanced *ABCG1* mRNA and protein expression, reduced ox-LDL uptake, and enhanced HDL-mediated cholesterol efflux (Table 5). These effects were found to be independent of LXRα, but mediated through *ABCG1* mRNA 3' untranslated region without affecting *ABCA1*, SR-A, CD36, and SR-B1 protein expression (Li et al., 2010). In contrast, Kemmerer et al. (2016) demonstrated predominant LXRα-mediated upregulation of *ABCA1* mRNA expression in macrophages after exposure to AICAR and the allosteric AMPK activators A769662 and salicylate (Li et al., 2010). The increased

TABLE 4
Overview of microRNAs enhancing or reducing ABCA1 and ABCG1 mRNA

Overview of microRNAs, including their effects on ABCA1 and ABCG1 expression. Effects on ABCA1, ABCG1, and cholesterol efflux are presented as follows: (↓) decreased; (=) no effect; (↑) increased.

MicroRNA	ABCA1	ABCG1	Reference
miR-10b	↓	↓	Hazen and Smith, 2012; Wang et al., 2012; Dávalos and Fernandez-Hernando, 2013; Goedeke et al., 2014; Rayner and Moore, 2014; Rotllan et al., 2016; Aryal et al., 2017
miR-17	↓		He et al., 2015
miR-19b	↓		Lv et al., 2014, 2015; DiMarco and Fernandez, 2015
miR-20a/b	↓		Liang et al., 2017
miR-21	↓	↓	Canfrán-Duque et al., 2017
miR-26	↓	↓ (Indirect via LXR)	Sun et al., 2012; Dávalos and Fernandez-Hernando, 2013; Canfrán-Duque et al., 2014; Goedeke et al., 2014; Rayner and Moore, 2014; DiMarco and Fernandez, 2015; Yang et al., 2015; Feinberg and Moore, 2016; Rotllan et al., 2016
miR-27a/b	↓	= (↓ Indirect)	Kang et al., 2013; Canfrán-Duque et al., 2014; Goedeke et al., 2014, 2015b; Zhang et al., 2014; DiMarco and Fernandez, 2015; Yang et al., 2015; Rotllan et al., 2016
miR-33a/33b	↓	↓	Moore et al., 2010, 2011; Fernandez-Hernando et al., 2011; Fernández-Hernando and Moore, 2011; Rayner et al., 2011, 2012; Iatan et al., 2012; Rotllan and Fernandez-Hernando, 2012; Dávalos and Fernandez-Hernando, 2013; Kang et al., 2013; Canfrán-Duque et al., 2014; Goedeke et al., 2014; Mao et al., 2014; Rayner and Moore, 2014; DiMarco and Fernandez, 2015; He et al., 2015; Mandolini et al., 2015; Yang et al., 2015; Feinberg and Moore, 2016; Ono, 2016; Rotllan et al., 2016; Aryal et al., 2017
miR-93	↓		He et al., 2015
miR-96	↓		Moazzeni et al., 2017
miR-101	↓		Zhang et al., 2015a; Aryal et al., 2017
miR-106b	↓		Kim et al., 2012; Rotllan and Fernandez-Hernando, 2012; Dávalos and Fernandez-Hernando, 2013; Goedeke et al., 2014; Rayner and Moore, 2014; Feinberg and Moore, 2016
miR-128-1	↓		Wagschal et al., 2015; Feinberg and Moore, 2016; Rotllan et al., 2016; Aryal et al., 2017
miR-128-2	↓	↓	Adlakha et al., 2013; DiMarco and Fernandez, 2015
miR-130b	↓		Wagschal et al., 2015; Feinberg and Moore, 2016
miR-144	↓	↓ (Indirect via RXR)	de Aguiar Vallim et al., 2013; Kang et al., 2013; Ramírez et al., 2013; Canfrán-Duque et al., 2014; Goedeke et al., 2014; Rayner and Moore, 2014; DiMarco and Fernandez, 2015; Feinberg and Moore, 2016; Rotllan et al., 2016; Aryal et al., 2017
miR-145	↓		Kang et al., 2013; Canfrán-Duque et al., 2014; Goedeke et al., 2014; Sala et al., 2014; DiMarco and Fernandez, 2015
miR-148a	↓		Kang et al., 2013; Goedeke et al., 2015a; Wagschal et al., 2015; Feinberg and Moore, 2016; Rotllan et al., 2016; Aryal et al., 2017
miR-223	↑		DiMarco and Fernandez, 2015; Rotllan et al., 2016
miR-301b	↓		Wagschal et al., 2015; Feinberg and Moore, 2016
miR-302a	↓		DiMarco and Fernandez, 2015; Meiler et al., 2015; Rotllan et al., 2016; Aryal et al., 2017
miR-378		↓ (Indirect)	Wang et al., 2014a; DiMarco and Fernandez, 2015; Yang et al., 2015
miR-613	↓		Zhao et al., 2014; DiMarco and Fernandez, 2015
miR-758	↓		Ramirez et al., 2011; Kim et al., 2012; Rayner et al., 2012; Rotllan and Fernandez-Hernando, 2012; Dávalos and Fernandez-Hernando, 2013; Canfrán-Duque et al., 2014; Goedeke et al., 2014; Rayner and Moore, 2014; DiMarco and Fernandez, 2015; Mandolini et al., 2015; Yang et al., 2015; Feinberg and Moore, 2016

miR, microRNA.

ABCA1 expression by A769662 and AICAR was proposed to be mediated via inhibition of extracellular signal-regulated kinase (ERK) or the mammalian target of rapamycin pathways (Kemmerer et al., 2016). Indeed, ERK signaling is described as a negative regulator of ABCA1 protein stability (Mogilenko et al., 2010). Compounds that affect the ERK pathway will be discussed in Section VI.B (Pharmacological Inhibition of ATP-Binding Cassette A1 and ATP-Binding Cassette G1 Protein Degradation). Thus, AMPK agonists could increase cholesterol efflux via a dual mechanism, either by enhancing *ABCG1* mRNA stability or via increased mRNA expression of *ABCG1* and *ABCA1*.

VI. ATP-Binding Cassette A1 and ATP-Binding Cassette G1 Protein Degradation as a Target to Increase Cholesterol Efflux

A. The Role of ATP-Binding Cassette A1 and ATP-Binding Cassette G1 Reuptake and Degradation in the Regulation of Their Plasma Membrane Abundance

Like other plasma membrane transporters and receptors, ABCA1 and ABCG1 protein abundance is dependent

on their cellular reuptake and degradation (Yokoyama et al., 2012; Wang et al., 2017). After internalization, ABCA1 is mainly degraded via lysosomal and ubiquitin- or calpain-mediated proteolysis (Yokoyama et al., 2012; Huang et al., 2015; Wang et al., 2017) (Fig. 5). Although the function of lysosomal and ubiquitin-mediated proteolytic degradation in ABCA1 turnover and activity needs to be elucidated, the importance of calpain-mediated ABCA1 breakdown has been established, especially in THP-1 macrophages (Yokoyama et al., 2012; Wang et al., 2017).

ABCA1 proteolysis is initiated by calpain binding to a calpain-specific cleavage sequence [Pro-Glu-Ser-Thr (PEST)] (Iwamoto et al., 2010; Huang et al., 2015; Wang et al., 2017), which is located within the large cytosolic loop of ABCA1 (Reiss and Cronstein, 2012) (Fig. 5). The relevance of this region was illustrated by deletion of the PEST sequence, which prevented the breakdown of ABCA1 (Yokoyama et al., 2012). This region is also important for ABCA1 half-life, which is 1 to 2 hours in the absence of helical apolipoproteins, such as apoA-I, apoA-II, and apoE (Iwamoto et al., 2010; Yokoyama et al., 2012).

TABLE 5
Compounds enhancing ABCA1 and ABCG1 mRNA stability

Overview of compounds that enhance *ABCA1* and *ABCG1* mRNA stability, including their primary pharmacological action, effects on *ABCA1* and *ABCG1* expression (arrows: *mRNA*, protein), and effects on cellular cholesterol efflux. Effects on *ABCA1*, *ABCG1*, and cholesterol efflux are presented as follows: (↓) decreased; (=) no effect; (↑) increased. Italicized symbols indicate changes in *ABCA1* and *ABCG1* mRNA levels, whereas nonitalicized symbols indicate protein levels.

Compound	Primary Action	ABCA1		ABCG1		Cholesterol Efflux	Reference
AICAR	AMPK agonist	=	J774.A1	↑	J774.A1	J774.A1	Li et al., 2010; Kemmerer et al., 2016
		↑	THP-1	↑	PM ^a	HDL	
A769662	Allosteric AMPK agonist	↑	THP-1	↑	J774.A1	↑	Li et al., 2010; Kemmerer et al., 2016
Salicylate	Allosteric AMPK agonist	↑	THP-1	—	—	—	

^a*apoE*^{−/−} C75BL/6 mice.
PM, peritoneal macrophages.

Phosphorylation of the PEST sequence is suggested to reduce ABCA1 half-life (Iwamoto et al., 2010; Yokoyama et al., 2012), whereas apoA-I binding or pre-exposure (i.e., apoA-I presence before internalization of prebiotinylated surface ABCA1 expressed on THP-1) prolongs the half-life by increasing the resistance of ABCA1 calpain-mediated proteolysis (Lu et al., 2008; Iwamoto et al., 2010; Yokoyama et al., 2012). This did not affect ABCA1 internalization in endosomes, but resulted in a larger portion of ABCA1 that is recycled from the endosomes to the plasma membrane. Besides apoA-I, calpain-mediated ABCA1 degradation is regulated by various endogenous ligands, including calmodulin, calpeptin, calpastatin, protein kinase C (PKC)α, and heme oxygenase-1 (HO-1). Calmodulin interacts with a region in the large cytoplasmic loop near the PEST sequence, which protects ABCA1 against calpain-mediated proteolysis (Iwamoto et al., 2010). In contrast to calmodulin, calpastatin and the nuclear factor-like (Nrf)2–HO-1 axis directly inhibit calpain activity and decrease ABCA1 degradation (Tsai et al., 2010; Wissel et al., 2015). Finally, phosphorylation of the PEST region by activation of apelin-13–mediated PKCα also protects ABCA1 against calpain-mediated proteolysis (Liu et al., 2013). In short, inhibition of calpain activity and ABCA1 internalization, or stimulation of its turnover rate, may result in an increased ABCA1-mediated cellular cholesterol efflux and increased HDL biosynthesis.

B. Pharmacological Inhibition of ATP-Binding Cassette A1 and ATP-Binding Cassette G1 Protein Degradation

As proteolytic breakdown of ABCA1 and ABCG1 plays a significant role in the regulation of their plasma membrane abundance, different pharmacological treatments have been investigated and aimed at increasing RCT via this mechanism (Table 6). One is the use of inhibitors of thiol proteases (e.g., calpain), like leupeptin and N-acetyl-leu-leu-norleucinal (ALLN), which both prevented ABCA1 degradation (Arakawa and Yokoyama, 2002; Yokoyama, 2004). In contrast, other thiol protease inhibitors (i.e., pepstatin A, aprotinin, and phosphoramidon) did not affect ABCA1 degradation (Arakawa and Yokoyama, 2002), questioning the specificity of the effect of leupeptin and ALLN. In addition, ABCA1

protein expression levels were not affected by non-specific protease inhibitors like the proteasome inhibitor lactacystin (Arakawa and Yokoyama, 2002).
The stabilizing role of PKC on ABCA1 also seems to hold for ABCG1, as the ABCG1-dependent cholesterol efflux is decreased by the PKC inhibitors calphostin C and PKC19-36 (Gelissen et al., 2012). As mentioned above, apelin-13, a cleaved peptide from the adipocytokine apelin, phosphorylated ABCA1 in THP-1 macrophages, leading to ABCA1 stabilization and enhanced cholesterol efflux (Liu et al., 2013). Remarkably, several other PKC inhibitors (i.e., Gö6983, Gö6976, rottlerin, and doxazosin) enhanced ABCA1 mRNA and protein expression as well as cholesterol efflux in macrophages via an inhibitory effect on protein kinase D (PKD), with the strongest effect by Gö6983 (Iwamoto et al., 2007, 2008; Tsunemi et al., 2014). This effect is probably mediated by a decreased phosphorylation of activator protein 2α, which represses transcription of *ABCA1* as phosphorylated activator protein 2α binds to the promoter region of *ABCA1* (Iwamoto et al., 2007, 2008; Remaley, 2007). The potential of this promoter region as a target to increase cellular cholesterol efflux is emphasized by pyrrole-imidazole polyamides that bind to this region and enhanced ABCA1 mRNA and protein levels as well as cholesterol efflux in 3T3-L1 adipocytes and RCT in C57BL/6 mice (Tsunemi et al., 2014). In this study, ABCA1 levels were increased via PKD inhibition, which illustrates the importance of the balance between PKC and PKD activity.
A more specific calpain inhibitor, triacetyl-3 hydroxyphenyl-adenosine (IMM-H007), increased ABCA1 plasma membrane expression in THP-1 cells and enhanced ABCA1-mediated cholesterol efflux to apoA-I (Huang et al., 2015). In *apoE*^{−/−} mice, IMM-H007 delayed ABCA1 protein degradation, promoted ABCA1 cell surface localization, enhanced RCT, and suppressed atherosclerotic lesion development (Huang et al., 2015). Interestingly, IMM-H007 activated AMPK, which may have also contributed to these effects via ABCA1 stabilization, as described above. In summary, the adenosine analog IMM-H007 may be a promising candidate to upregulate ABCA1 expression with proven capacity to increase RCT in vivo.
More indirect strategies to inhibit calpain, including the induction of HO-1 production, have also been shown

TABLE 6
Inhibitors of ABCA1 and ABCG1 protein breakdown

Overview of compounds that inhibit ABCA1 and ABCG1 protein breakdown, including their primary pharmacological action, effects on ABCA1 and ABCG1 expression (arrows: *mRNA*, protein), and effects on cellular cholesterol efflux. Effects on ABCA1, ABCG1, and cholesterol efflux are presented as follows: (↓) decreased; (=) no effect; (↑) increased. Italicized symbols indicate changes in *ABCA1* and *ABCG1* mRNA levels, whereas nonitalicized symbols indicate protein levels.

Compound	Primary Action	ABCA1	ABCG1	Cholesterol Efflux	Reference
Acifran	GPR109A agonist	—	—	—	Gaidarov et al., 2013
Acipimox	GPR109A agonist	—	—	—	Gaidarov et al., 2013
ALLN	Thiol-protease inhibitor	=↑ THP-1	—	↑ THP-1	Arakawa and Yokoyama, 2002; Yokoyama, 2004
Calphostin C	PKC inhibitor	—	—	↓ CHO-K1 ABCG1(+12) ↓ CHO-K1 ABCG1(−12)	Gelissen et al., 2012
Diphenoquinone	Unknown, probucol metabolite	↑ THP-1 ↑ RAW264.7 ↑ HEK293 ↑ Balb/3T3 ↑ MEFs =↑ Liver ^a	↑ RAW264.7	↑ THP-1 ↑ RAW264.7 ↑ apoA-I ↑ HDL ↑ Hek293 ↑ MEFs	Arakawa et al., 2009; Lu et al., 2016; Yakushiji et al., 2016
Exendin-4	GLP-1R agonist	↑↑ 3T3-L1 adipocytes	↑↑ 3T3-L1 adipocytes	↑ 3T3-L1 adipocytes	Mostafa et al., 2015; Yin et al., 2016
Ezetimibe	NPC1L1 inhibitor	↑↑ glomeruli ^b ↑ VSCMs ↓↓ Liver ^c ↓↓ Proximal small intestine ^c	—	↑ glomeruli ^b —	Gong et al., 2014; Kannisto et al., 2014
Gö6976	PKC inhibitor	↑↑ THP-1	—	↑ THP-1	Iwamoto et al., 2008
Gö6983	PKC inhibitor	↑↑ THP-1 ↑↑ Liver ^c	—	↑ THP-1	Iwamoto et al., 2008
IMM-H007	AMPK agonist	↑ THP-1 ↑ Liver ^d	—	↑ J774 ↑ THP-1 ↑ RCT ^e	Huang et al., 2015
LD211	MC1-R agonist	↑ BMDM ^d	↑ BMDM ^d	↑ BMDM ^d ↑ apoA-I ↑ HDL	Rinne et al., 2017
Leupeptin	Thiol-protease inhibitor	=↑ THP-1	—	—	Arakawa and Yokoyama, 2002
MK-0354	GPR109A agonist	= MDM ^d	= MDM ^d	= MDM ^d	Gaidarov et al., 2013
MK-1903	GPR109A agonist	↑ MDM ^d	↑ MDM ^d	↑ MDM ^d	Gaidarov et al., 2013
MSG606	MC1-R agonist	↑ BMDM ^d = Aorta ^d ↑ Liver ^d	↑ BMDM ^d = Aorta ^d ↑ Liver ^d	BMDM ^d ↑ apoA-I ↑ HDL	Rinne et al., 2017
<i>N</i> -acetyl cysteine	Glutathione synthase stimulator	↓ J774	↑ J774	J774 ↓ apoA-I ↑ HDL	Machado et al., 2014
Niacin	GPR109A agonist	↑↑ HepG2 ↑↑ 3T3-L1 adipocytes ↑ MM6sr ↑ MDM ^d ↑ Monocyte ^d ↑↑ Liver ^e	↑ MDM ^d	↑ THP-1 ↑ HepG2 ↑ 3T3-L1 adipocytes ↑ MM6sr ↑ MDM ^d	Rubic et al., 2004; Siripurkpong and Na-Bangchang, 2009; Wu and Zhao, 2009; Yvan-Charvet et al., 2010a; Zhang et al., 2012; Gaidarov et al., 2013
Nicardipine	Calcium channel blocker	—	—	↑ THP-1 ↑ RAW264.7	Suzuki et al., 2004
Nifedipine	Calcium channel blocker	↑↑ RAW264.7 ↑↑ Aorta sinus ^f ↑↑ PM ^a	↑↑ RAW264.7	↑ THP-1 ↑ RAW264.7 ↑ apoA-I ↑ HDL	Suzuki et al., 2004; Ishii et al., 2010; Zhang et al., 2013b
PD98059	MEK1/2 inhibitor	↑ THP-1 ↑↑ RAW264.7 ↓ HuH7 ↓ CHO	↑ THP-1 ↑ RAW264.7 ↓ CHO ↓ HEK293 ↑↑ Mouse primary macrop	↓ CHO PM ^d ↑ HDL	Zhou et al., 2010; Mulay et al., 2013; Zhang et al., 2016
PKC19-36	PKC inhibitor	—	—	↓ CHO-K1 ABCG1(+12)	Gelissen et al., 2012

(continued)

TABLE 6—Continued

Compound	Primary Action	ABCA1	ABCG1	Cholesterol Efflux	Reference
Rottlerin Spiroquinone	PKC inhibitor Unknown, probucol metabolite	↑↑ THP-1	—	↓ CHO-K1 ABCG1(–12)	Iwamoto et al., 2008 Yokoyama, 2004; Arakawa et al., 2009; Lu et al., 2016; Yakushiji et al., 2016
		↑ THP-1	↑ RAW264.7	= THP-1	
		↑ RAW264.7		↑ THP-1	
		↑ HEK293		↑ apoA-I	
		↑ Balb/3T3		= HDL	
Tert- butylhydroquinone	Synthetic phenolic antioxidant	↑ MEFs		↑ Hek293	Lu et al., 2013
		=↑ Liver ^a		↑ MEFs	
		↑↑ THP-1	—	↑ THP-1	
U0126	MEK1/2 inhibitor	↑↑ RAW264.7	↑ THP-1	RAW264.7	Mogilenko et al., 2010; Zhou et al., 2010; Mulay et al., 2013; Xue et al., 2016; Zhang et al., 2016; Liu et al., 2019
		↑↑ HepG2	↑↑ RAW264.7	↑ apoA-I	
		↑↑ PM ^b	↑↑ Mouse	↑ HDL	
			primary macrop		
		↑↑ Jurkat	↑ PM ^d	PM	
Verapamil	Calcium channel blocker		↑↑ PM ^g	↑ apoA-I	Suzuki et al., 2004
			↑↑ Jurkat	↑ HDL	
				↑ Jurkat	
				↑ apoA-I	
				↑ THP-1	
Vildagliptin	GLP-1R agonist	↑↑ 3T3-L1	↑↑ 3T3-L1	↑ RAW264.7	Mostafa et al., 2015, 2016
		adipocytes	adipocytes	↑ 3T3-L1	

ALLN, *N*-acetyl-leu-leu-norleucinal; GLP-1R, glucagon-like peptide-1 receptor; MDM, monocyte-derived macrophage; MEF, murine embryonic fibroblast; MEK, mitogen-activated protein kinase kinase; NPC1L1, Niemann-Pick C1-like 1; PM, peritoneal macrophages.

^aNew Zealand White rabbits.

^b*apoE*^{−/−} C75BL/6 diabetic mice.

^c*apoE*^{−/−} C75BL/6 mice.

^dC75BL/6 mice.

^eGolden Syrian Hamster.

^fC3H/He mice.

^gSprague Dawley rats.

to suppress ABCA1 degradation. Tertiary-butylhydroquinone, a synthetic phenolic antioxidant, mediates these effects by activation of Nrf2 via its dissociation from kelch-like ECH-associated protein (Keap) 1, translocation of Nrf2 toward the nucleus, and activation of antioxidant responsive element (ARE)-dependent transcription of *HO-1* gene. The increased production of HO-1, which is an endogenous inhibitor of calpain, resulted in reduced ABCA1 degradation (Lu et al., 2013).

In addition, two oxidative products of probucol, spiroquinone and diphenoquinone, suppressed ABCA1 and ABCG1 degradation both in vitro and in vivo, and thereby increased RCT and reduced lipid deposition in atherosclerotic plaques in vivo (Yokoyama, 2004; Arakawa et al., 2009; Lu et al., 2016; Yakushiji et al., 2016). These effects are expected to be due to disruption of the caveolin-1 interaction with ABCA1 by spiroquinone and diphenoquinone, resulting in the stabilization of ABCA1 protein against degradation (Arakawa et al., 2009). This mechanism is also expected to mediate the increase in ABCA1 abundance by ezetimibe, which lowers caveolin-1 expression through suppression of SREBP-1 expression (Lu et al., 2016). Another mechanism directly affecting ABCA1 stability is the reduction of its disulfide bonds (Bungert et al., 2001), as described with *N*-acetyl-cysteine (Machado et al., 2014). *N*-acetyl-cysteine treatment is, however, associated with an increased cellular cholesterol efflux, which was mediated by increased ABCG1 expression and HDL-dependent

cholesterol efflux, whereas ABCA1 expression and apoA-I-dependent cholesterol efflux were decreased in J774 cells (Machado et al., 2014).

Finally, calpain-mediated ABCA1 proteolysis can also be inhibited by alteration of the phosphorylation status of the ABCA1 PEST sequence, as described above (Reiss et al., 2008; Chen et al., 2011a). This could explain the beneficial effects of the ERK1/2 inhibitors, PD98059 and U0126, on ABCA1 and ABCG1 mRNA and protein expression, cholesterol efflux in macrophages, and RCT in *apoE*^{−/−} mice (Mulay et al., 2013; Kemmerer et al., 2016; Xue et al., 2016; Zhang et al., 2016). Besides a direct effect, ERK1/2 inhibition could also increase *ABCA1* and *ABCG1* mRNA stability and expression via an LXRα-dependent mechanism, possibly, by enhancing the LXRα to LXRE-A binding, without affecting PPARγ and LXRα expression (Xue et al., 2016; Liu et al., 2019). In addition to an increased cholesterol efflux, lipid deposition and CD36 expression were suppressed upon U0126 treatment in ox-LDL-stimulated macrophages (Xue et al., 2016). The effect of ERK1/2 inhibition is cell-type dependent, as ABCA1 and ABCG1 protein stability was reduced in CHO and HuH7 cells after ERK1/2 inhibition (Gelissen et al., 2012; Mulay et al., 2013). In addition, the beneficial effect on ABCA1 and ABCG1 expression of zerumbone (i.e., a wild ginger-derived natural compound) and tanshinone IIA was abolished in THP-1 macrophages by ERK inhibitor PD98059 (Liu et al., 2014; Mostafa et al., 2015, 2016; Zhu and Liu, 2015;

Yin et al., 2016). Moreover, LXR α , PPAR α , ABCA1, and ABCG1 expression were not altered after inhibition of c-Jun N-terminal kinase and P38 mitogen-activated protein kinase (MAPK) phosphorylation by SP600125 and SB203580, respectively (Liu et al., 2019). The uncertainty about the precise role of ERK inhibition in cellular cholesterol efflux is emphasized by the observation that stimulation of MAPK/ERK increased LXR-mediated ABCA1 expression, as found for the glucagon-like peptide 1 receptor agonist exendin-4 and the dipeptidyl peptidase-4 inhibitor vildagliptin (Mostafa et al., 2015, 2016; Yin et al., 2016). Similar effects upon MAPK/ERK stimulation were observed using the melanocortin 1 receptor (MC1-R) agonist MSG606, which upregulates cholesterol efflux and ABCA1 and ABCG1 protein levels in bone marrow-derived macrophages, probably via ERK stimulation without increasing cAMP levels (Rinne et al., 2017). The MC1-R agonist, LD211, which stimulated ERK1/2 and p38 MAPK phosphorylation along with a strong increase in cAMP levels, also positively affected cholesterol efflux (Rinne et al., 2017). However, the exact mechanism behind the effect of MC1-R agonists on ERK1/2 is yet unidentified.

Stimulation of the ERK1/2–PPAR γ –LXL α axis has been associated with the favorable effects of the G protein-coupled receptor (GPR)109 agonist, niacin, on triglyceride and total, LDL, and HDL cholesterol levels in plasma. Moreover, it has been shown that this first clinically available cholesterol-lowering drug may upregulate ABCA1 expression, stabilize newly produced HDL, and promote cholesterol efflux possibly via cAMP/protein kinase A (PKA) and PPAR γ –LXR α pathways (Rubic et al., 2004; Siripurkpong and Na-Bangchang, 2009; Wu and Zhao, 2009; Yvan-Charvet et al., 2010a; Zhang et al., 2012; Connolly et al., 2013; Gaidarov et al., 2013). The GPR109A agonists, acifran and acipimox, but not isoniaicin, also activated ERK1/2 and Ca²⁺ flux (Gaidarov et al., 2013). Although niacin exerted beneficial effects, it is also associated with vasodilation and flushing side effects, possibly mediated via the secretion of prostaglandins as a result of activation of GPR109A (Rubic et al., 2004; Gaidarov et al., 2013). Besides niacin, the full GPR109A agonist, MK-1903, gave similar effects on ABCA1- and apoA-I-mediated cholesterol efflux, whereas the effect on ABCG1 expression was lower compared with niacin (Gaidarov et al., 2013). Another GPR109A agonist, MK-0354, induced neither flushing nor activation of GPR109A signaling in macrophages and HDL modulation (Gaidarov et al., 2013). This suggests that GPR109A agonists that do not cause flushing are probably also unable to exert an antiatherogenic effect.

ERK1/2 inhibition by the commonly used calcium channel blocker nifedipine leads to in vitro antiatherogenic effects at clinically relevant low nanomolar concentrations via inhibition of monocyte chemoattractant protein-1 and stimulation of ABCA1 expression

(Ishii et al., 2010). Two other calcium channel blockers, verapamil and nifedipine, also increased ABCA1- but not ABCG1-mediated cholesterol and phospholipid efflux at suprapharmacological concentrations in the low micromolar range (Suzuki et al., 2004). Verapamil was able to enhance ABCA1 promoter activity in an LXR-independent manner (Suzuki et al., 2004). In summary, ERK1/2 has the potential to beneficially affect cellular cholesterol efflux, but some of the controversies of this relation will need to be clarified to assess its true therapeutic relevance.

To conclude, inhibition of ABCA1 and ABCG1 degradation has demonstrated to be a promising strategy to enhance their plasma membrane expression, in combination with beneficial in vivo effects on RCT and the progression of atherosclerotic plaque formation. However, many compounds are rather unspecific, which may limit their therapeutic applicability due to adverse effects as a consequence of off-target activities.

VII. ATP-Binding Cassette A1 Function and Cyclic Adenosine Monophosphate

A. cAMP Is a Potent Regulator of ATP-Binding Cassette A1 Function and Expression

ABCA1 expression and function can also be enhanced by increasing cellular cAMP levels (Sakr et al., 1999). The transporter is phosphorylated upon cAMP-mediated activation of PKA, which leads to an increased capacity to interact with apoA-I and subsequent apoA-I-dependent cellular cholesterol efflux (Haidar et al., 2002, 2004) (Fig. 5). PKA has two major phosphorylation sites, Ser-1042 and Ser-2054, which are located at the NBDs of ABCA1 (See et al., 2002). These effects are most likely specific to macrophages (Bortnick et al., 2000; Suzuki et al., 2004). Interestingly, phosphorylation can also be autoregulatory, as apoA-I binding to ABCA1 has been demonstrated to stimulate adenylate cyclase (AC) activity by an unknown mechanism and boost intracellular cAMP levels (Haidar et al., 2002). Besides influencing cholesterol efflux, cAMP is also involved in the regulation of glucose, lipid, and cholesterol metabolism (Haidar et al., 2002, 2004; Tresguerres et al., 2011), and its cellular concentrations are a result of a delicate balance between its production from ATP by AC and its hydrolysis into AMP by phosphodiesterases (PDEs) and cellular efflux by the ABC transporters multidrug resistance protein 4/ABCC4 and multidrug resistance protein 5/ABCC5 (Lin and Bornfeldt, 2002; Wielinga et al., 2003; Copsel et al., 2011; Tresguerres et al., 2011) (Fig. 5). The activity of AC is regulated by different adenosine receptor subtypes, of which A₁ and A₃ receptors have an inhibitory and A_{2a} and A_{2b} a stimulatory effect (Reiss and Cronstein, 2012). Consequently, A_{2a} receptor activation enhanced cAMP levels and promoted RCT via ABCA1-mediated cholesterol efflux (Bingham et al., 2010).

B. Stimulation of cAMP Levels Enhances ATP-Binding Cassette A1-Mediated Cholesterol Efflux

Treatment with cAMP analogs, 8-bromo-cAMP, 8-(4-chlorophenylthio)adenosine-cAMP, (Bu)₂cAMP, and dibutyryl cAMP, stimulated apoA-I-dependent cholesterol efflux in different cell types (Sakr et al., 1999; Abe-Dohmae et al., 2000; Haidar et al., 2002; Lin and Bornfeldt, 2002; Kellner-Weibel et al., 2003; Bingham et al., 2010; Gaidarov et al., 2013; Manna et al., 2015) (Table 7). The 8-bromo-cAMP enhanced cholesterol efflux toward apoA-I, most likely via increased PKA-mediated ABCA1 phosphorylation (Iwamoto et al., 2008; Katz et al., 2009), as PKA inhibition by H89 reversed the beneficial effects of 8-bromo-cAMP on D4-F-mediated cholesterol efflux in RAW264.7 macrophages (Bingham et al., 2010; Xie et al., 2010). Similar results were observed with 8-(4-chlorophenylthio)adenosine-cAMP, which elevated ABCA1 protein expression (Sakr et al., 1999; Haidar et al., 2002; Lin and Bornfeldt, 2002; Kellner-Weibel et al., 2003). A direct PKA agonist, 6-Benz-cAMP, also stimulated ABCA1 protein expression and cellular cholesterol expression, which emphasizes the role of PKA-dependent phosphorylation in enhancing ABCA1-mediated cholesterol efflux capacity (Bingham et al., 2010). Surprisingly, PKA seemed to have contradictory effects on ABCG1 as inhibition of its activity by H89 or KT5720 in CHO-K1 cells enhanced ABCG1-mediated cholesterol efflux and stabilized ABCG1. However, this effect was only observed in cells overexpressing a mutant ABCG1 containing a 12-amino-acid internal segment (Gelissen et al., 2012). Interestingly, treatment with the platelet inhibitor dipyridimole, which increases intracellular cAMP levels, demonstrated antiatherosclerotic effects in the ESPRIT trial (Halkes et al., 2006)).

In line with a central role for cAMP, modulation of AC and PDEs has demonstrated to be another useful strategy to enhance RCT. The AC activator forskolin enhanced RCT via elevated ABCA1 protein levels and phosphorylation (Haidar et al., 2002; Lin and Bornfeldt, 2002). AC stimulation via A_{2a} receptor activation with CGS-21680 and ATL313 enhanced cholesterol efflux and ABCA1 protein expression (Bingham et al., 2010; Voloshyna et al., 2013). Moreover, stimulation with 8-pcPT-2'-O-Me-cAMP of Epac, a signaling molecule downstream of A_{2a} receptor activation, showed similar results. A_{2a} receptor activation decreased foam cell formation in THP-1-derived macrophages by regulation of cholesterol influx and efflux (Bingham et al., 2010). Activation of this adenosine receptor is also associated with an increased ABCA1 expression in peripheral blood mononuclear cells by methotrexate, which is used to treat cancer and autoimmune diseases (Reiss et al., 2008; Chen et al., 2011a).

Increased cAMP levels as a result of reduced breakdown with PDE4 inhibitors (3-isobutyl-1-methylxanthine (IBMX), rolipram, and cilomast) also resulted in elevated ABCA1

expression levels and apoA-I-dependent cholesterol efflux in macrophages (Haidar et al., 2002; Lin and Bornfeldt, 2002). Similar effects were observed with the PDE3 inhibitor cilostazol through increased ABCA1 and ABCG1 expression and cholesterol efflux in vitro and in vivo (Nakaya et al., 2010).

In summary, cAMP analogs, A_{2a} receptor agonists, and other compounds that increase intracellular cAMP levels have the potential to induce ABCA1 expression, which could increase apoA-I-mediated cholesterol efflux. The involvement of cAMP in many different metabolic pathways (i.e., glucose, lipid, and cholesterol metabolism) may though render these strategies vulnerable to adverse effects.

VIII. Increasing Cellular Cholesterol Efflux via Unknown Mechanisms

Stimulation of cellular cholesterol efflux has also been observed with various other compounds by yet unidentified mechanisms, including four antimicrobial drugs (aclerubicin, daidzein, pratensein, pyrromycin) that upregulated ABCA1 expression in ABCA1-overexpressing HepG2 cells (Table 8). Similar effects were observed with the sphingolipid synthesis inhibitor 1-phenyl-2-decanoylamino-3-morpholino-1-propanol (a potential therapeutic for Gaucher's disease), which was accompanied by an increased ABCA1-dependent cholesterol efflux (Glaros et al., 2005). Although the mechanism remains unknown, a structurally similar sphingolipid synthesis inhibitor, *N*-butyldeoxy-nojirimycin, did not increase cellular cholesterol efflux. Enhanced cholesterol efflux was also observed in macrophages of patients treated with the CETP inhibitors anacetrapib, dalcetrapib and torcetrapib, whereas torcetrapib also increased ABCA1 expression (Yvan-Charvet et al., 2007, 2010a; Brodeur et al., 2017). This is, however, expected to be a result of the increased HDL-C levels due to CETP inhibition. The clinical application of these drugs seems to be limited, as the development of all three CETP inhibitors has been discontinued because of an unfavorable benefit-risk ratio (Tall and Rader, 2018). Upregulation of ABCA1 has also been observed with ibrutinib and MCC950, which decreased ox-LDL uptake in THP-1 macrophages mediated via inhibition of nucleotide binding oligomerization domain receptor family pyrin domain-containing protein (NLRP) 3 inflammasome (Chen et al., 2018). Inhibition of the NLRP3 inflammasome resulted in reduced foam cell formation in macrophages and promoted cholesterol efflux (Chen et al., 2018). Moreover, inhibition of soluble epoxide hydrolase using *trans*-4-[4-(3-Adamanthan-1-yl-uneido)-cyclohexyloxy]-benzoic acid (t-AUCB) increased hepatic ABCA1 expression and cholesterol efflux in *LDLR*^{-/-} mice via a yet unknown mechanism, whereas hepatic ABCG1 expression was not affected (Shen et al., 2014; Shen et al., 2015). The blood glucose-lowering drug, metformin, which

TABLE 7
Compounds increasing cAMP levels

Overview of compounds that increase intracellular cAMP levels, including their primary pharmacological action, effects on ABCA1 and ABCG1 expression (arrows: *mRNA*, protein), and effects on cellular cholesterol efflux. Effects on ABCA1, ABCG1, and cholesterol efflux are presented as follows: (↓) decreased; (=) no effect; (↑) increased. Italicized symbols indicate changes in *ABCA1* and *ABCG1* mRNA levels, whereas nonitalicized symbols indicate protein levels.

Compound	Primary Action	ABCA1	ABCG1	Cholesterol Efflux	Reference
6-Benz-cAMP	PKA agonist	↑ THP-1	—	↑ THP-1	Bingham et al., 2010
8-Bromo-cAMP	cAMP analog	=↑ Skin fibroblasts ^a	—	↑ THP-1	Haidar et al., 2002; Lin and Bornfeldt, 2002
8-(4-Chlorophenylthio) adenosine-cAMP	cAMP analog	↑ PM ↑ J774.A1	—	↑ Skin fibroblasts ^a = THP-1 ↑ J774.A1 ↑ L cells = CHO = Fu5AH = Skin fibroblasts ^a ↑ Mouse PM	Sakr et al., 1999; Kellner-Weibel et al., 2003
8-pcPT-2'-O-Me-cAMP	Epac agonist	↑ THP-1	—	↑ THP-1	Bingham et al., 2010
ATL313	A2AR agonist	↑↑ THP-1	↑↑ THP-1	—	Voloshyna et al., 2013
(Bu) ₂ cAMP	cAMP analog	↑ RAW264.7	—	↑ RAW264.7	Manna et al., 2015
CGS-21680	A2AR agonist	↑↑ THP-1	↑↑ THP-1	↑ THP-1	Bingham et al., 2010; Voloshyna et al., 2013
Cilomast	PDE4 inhibitor	↑ THP-1 ↑ J774.A1	—	↑ THP-1 ↑ J774.A1	Lin and Bornfeldt, 2002
Cilostazol	PDE3 inhibitor	↑↑ THP-1 ↑↑ RAW264.7 ↑ MDM ^a ↑ PM ^b = Liver ^b = Small intestine ^b	↑↑ THP-1 ↑↑ RAW264.7 ↑ MDM ^a	↑ THP-1 ↑ RAW264.7 ↑ MDM ^a ↑ RCT ^b	Nakaya et al., 2010
Dibutylr cyclic AMP	cAMP analog	↑ RAW264 ↑ Monocyte-derived macroph ^a	↑ Monocyte-derived macroph ^a	↑ RAW264.7 ↑ Monocyte-derived macroph ^a	Abe-Dohmae et al., 2000; Gaidarov et al., 2013
Doxazosin	α-A1 adrenergic receptor antagonist	↑↑ THP-1 ↑↑ RAW264.7 ↑ NCTC clone 1469 ↑ CHO-K1 ↑ Liver ^b	—	↑ THP-1 ↑ RAW264.7	Iwamoto et al., 2007; Tsunemi et al., 2014
Forskolin	Adenylyl cyclase activator	↑ Skin fibroblasts ^a	—	↑ THP-1 ↑ Skin fibroblasts ^a	Haidar et al., 2002; Lin and Bornfeldt, 2002
H89	PKA inhibitor	—	↑ CHO-K1 = ABCG1(+12) = CHO-K1 = ABCG1(−12)	↑ CHO-K1 = ABCG1(+12) = CHO-K1 = ABCG1(−12)	Gelissen et al., 2012
IBMX	Nonselective PDE inhibitor	—	—	↑ THP-1 ↑ Skin fibroblasts ^a	Haidar et al., 2002; Lin and Bornfeldt, 2002
KT5720	PKA inhibitor	—	—	↑ CHO-K1 = ABCG1(+12) = CHO-K1 = ABCG1(−12)	Gelissen et al., 2012
Methotrexate	Dihydrofolate reductase inhibitor	↑ PBMCs ^a	—	—	Chen et al., 2011a
Rolipram	PDE4-selective inhibitor	↑↑ THP-1 ↑ J774.A1	—	↑ THP-1 ↑ J774.A1	Lin and Bornfeldt, 2002

IBMX, 3-isobutyl-1-methylxanthine; macroph, macrophage; MDM, monocyte-derived macrophage; PBMC, peripheral blood mononuclear cell; PM, peritoneal macrophages.

^aHuman.

^bC57BL/6 mice.

among other effects is an AMPK activator, did not alter ABCA1 expression and apoA-I-dependent cholesterol efflux, but enhanced ABCG1 expression and HDL-dependent cholesterol efflux in RAW264.7 macrophages (He et al., 2019).

Finally, an experimental xanthone compound, IMB2026791, enhanced the binding between apoA-I and ABCA1 and consequently increased apoA-I-mediated cholesterol in vitro. Although the exact mechanism underlying the effect of this xanthone still needs to be elucidated, it is the only compound to date that directly affects the mechanism of ABCA1 cholesterol efflux (Liu et al., 2012).

IX. Conclusions and Future Directions

The discovery of the inverse association between HDL and risk of atherosclerosis and the consequent potency of ABCA1- and ABCG1-mediated cholesterol transport as cardioprotective therapeutic targets (Tang and Oram, 2009; Kuhnast et al., 2015) started the search for strategies to enhance this transport step and thereby enhance RCT. A decade of research resulted in several lipoprotein mimetics, many natural compounds, and miRNA-based strategies with the ability to increase the cellular cholesterol efflux to HDL and over 100 small-molecule-based approaches discussed in this review.

TABLE 8
Compounds increasing ABCA1/G1 expression or function by an unknown mechanism

Overview of compounds that increase ABCA1 or ABCG1 expression or function by a yet unknown mechanism, including their primary pharmacological action, effects on ABCA1 and ABCG1 expression (arrows: *mRNA*, protein), and effects on cellular cholesterol efflux. Effects on ABCA1, ABCG1, and cholesterol efflux are presented as follows: (↓) decreased; (=) no effect; (↑) increased. Italicized symbols indicate changes in *ABCA1* and *ABCG1* mRNA levels, whereas nonitalicized symbols indicate protein levels.

Compound	Primary Action	ABCA1	ABCG1	Cholesterol Efflux	Reference
1-Phenyl-2-Decanoylamino-3-morpholino-1-propanol	Glycosylceramide transferase inhibitor	↑↑ Skin fibroblasts ^a	—	↑ Skin fibroblasts ^a ↑ MDM ^a	Glaros et al., 2005
Aclarubicin	Topoisomerase I and II inhibitor	↑↑ HepG2	—	—	Gao et al., 2008
Anacetrapib	CETP inhibitor	—	—	↑ THP-1 ↑ BHK ↑ ABCA1-expressing BHK	Yvan-Charvet et al., 2010a; Brodeur et al., 2017
Daidzein	Mitochondrial aldehyde dehydrogenase inhibitor	↑↑ HepG2	—	—	Gao et al., 2008
Dalcetrapib	CETP inhibitor	—	—	↑ BHK ↑ ABCA1-expressing BHK	Brodeur et al., 2017
Ibrutinib	NLRP3 inflammasome inhibitor	↑ THP-1	= THP-1	THP-1 = apoA-I ↑ HDL	Chen et al., 2018
IMB2026791	Unknown	—	—	↑ THP-1 ↑ CHO-ABCA1 ↑ CHO	Liu et al., 2012
MCC950	NLRP3 inflammasome inhibitor	↑ THP-1	= THP-1	THP-1 = apoA-I ↑ HDL	Chen et al., 2018
Metformin	Antihyperglycemic agent	= RAW264.7	↑↑ RAW264.7	↑ RAW264.7 = apoA-I ↑ HDL	He et al., 2019
<i>N</i> -butyldeoxynojirimycin	Glycosylceramide transferase inhibitor	—	—	= Skin fibroblasts ^a	Glaros et al., 2005
Pratensein	Unknown	↑↑ HepG2	—	—	Gao et al., 2008
Pyromycin	Microbial protein synthesis inhibitor	↑↑ HepG2	—	—	Gao et al., 2008
t-AUCB	Soluble epoxide hydrolase inhibitor	↑↑ Liver ^b	= Liver ^b	↑ 3T3-L1 adipocytes ↑ Epididymal fatc ↑ RCT ^b	Shen et al., 2015
Torcetrapib	CEPT inhibitor	↑ THP-1	—	↑ THP-1	Yvan-Charvet et al., 2007

BHK, baby hamster kidney cell; fatc., fat cell; MDM, monocyte-derived macrophage; NLRP3, nucleotide binding oligomerization domain receptor family, pyrin domain-containing protein 3; t-AUCB, *trans*-4-[4-(3-Adamantan-1-yl-uneido)-cyclononyloxy]-benzoic acid.

^aHuman.

^b*LXRα*^{-/-} C75BL/6 mice.

Besides the discovery of a large variety of different experimental compounds and existing drugs that enhance ABCA1- and ABCG1-mediated cholesterol efflux, this quest has also provided us with enhanced insights into the mechanisms that regulate the expression and function of these two key players in the initiation and propagation of RCT. It also demonstrated one of the important challenges in this field, namely to develop specific and potent therapies, which is currently a major limiting factor in the clinical applicability of many compounds described in this review. Most of them are also involved in other mechanisms, including stimulation of lipogenesis by LXR and RXR agonists, and regulation of other metabolic pathways (e.g., lipogenesis, fatty acid β -oxidation, glycolysis) by AMPK and cAMP agonists. In addition, several other drugs that increased ABCA1- or ABCG1-dependent cholesterol efflux can probably not be directly used as

cardioprotectants due to their toxic effects. Recent developments though seem to be promising in overcoming these adverse mechanisms, like combining LXR and PPAR α agonists (Thomas et al., 2003; Beyer et al., 2004), or the use of novel RAR agonists instead of promiscuous RXR agonists (Costet et al., 2003; Ayaori et al., 2012; Chen et al., 2012).

Interestingly, a high-throughput screening approach searching for compounds that directly affect apoA-I and ABCA1 binding identified the xanthone compound IMB2026791 as a first direct activator of ABCA1-mediated cellular cholesterol efflux (Liu et al., 2012). Along with many other novel experimental as well as existing drugs, such developments indicate that we are heading toward promising compounds for clinical evaluation, as these steps have previously been limited by unforeseen adverse effects due to lack of specificity.

To conclude, increasing ABCA1- and ABCG1-mediated cellular efflux seems to be a promising strategy to lower cardiovascular risk and, combined with cholesterol-lowering therapies, to reduce mortality and morbidity associated with atherosclerosis.

Authorship Contributions

Participated in research design: Frambach, Schirris.

Performed data analysis: Frambach, Schirris.

Wrote or contributed to the writing of the manuscript: Frambach, de Haas, Smeitink, Rongen, Russel, Schirris.

References

- Abe-Dohmae S, Suzuki S, Wada Y, Aburatani H, Vance DE, and Yokoyama S (2000) Characterization of apolipoprotein-mediated HDL generation induced by cAMP in a murine macrophage cell line. *Biochemistry* **39**:11092–11099.
- Adlakha YK, Khanna S, Singh R, Singh VP, Agrawal A, and Saini N (2013) Pro-apoptotic miRNA-128-2 modulates ABCA1, ABCG1 and RXR α expression and cholesterol homeostasis. *Cell Death Dis* **4**:e780.
- Alder-Baerens N, Müller P, Pohl A, Korte T, Hamon Y, Chimini G, Pomorski T, and Herrmann A (2005) Headgroup-specific exposure of phospholipids in ABCA1-expressing cells. *J Biol Chem* **280**:26321–26329.
- Arakawa R, Tamehiro N, Nishimaki-Mogami T, Ueda K, and Yokoyama S (2005) Fenofibric acid, an active form of fenofibrate, increases apolipoprotein A-I-mediated high-density lipoprotein biogenesis by enhancing transcription of ATP-binding cassette transporter A1 gene in a liver X receptor-dependent manner. *Arterioscler Thromb Vasc Biol* **25**:1193–1197.
- Arakawa R, Tsujita M, Iwamoto N, Ito-Ohsumi C, Lu R, Wu CA, Shimizu K, Aotsuka T, Kanazawa H, Abe-Dohmae S, et al. (2009) Pharmacological inhibition of ABCA1 degradation increases HDL biogenesis and exhibits antiatherogenesis. *J Lipid Res* **50**:2299–2305.
- Arakawa R and Yokoyama S (2002) Helical apolipoproteins stabilize ATP-binding cassette transporter A1 by protecting it from thiol protease-mediated degradation. *J Biol Chem* **277**:22426–22429.
- Argmann CA, Edwards JY, Sawyez CG, O'Neil CH, Hegele RA, Pickering JG, and Huff MW (2005) Regulation of macrophage cholesterol efflux through hydroxymethylglutaryl-CoA reductase inhibition: a role for RhoA in ABCA1-mediated cholesterol efflux. *J Biol Chem* **280**:22212–22221.
- Argmann CA, Sawyez CG, McNeil CJ, Hegele RA, and Huff MW (2003) Activation of peroxisome proliferator-activated receptor gamma and retinoid X receptor results in net depletion of cellular cholesteryl esters in macrophages exposed to oxidized lipoproteins. *Arterioscler Thromb Vasc Biol* **23**:475–482.
- Aryal B, Singh AK, Rotllan N, Price N, and Fernández-Hernando C (2017) MicroRNAs and lipid metabolism. *Curr Opin Lipidol* **28**:273–280.
- Ayari M, Yakushiji E, Ogura M, Nakaya K, Hisada T, Uto-Kondo H, Takiguchi S, Terao Y, Sasaki M, Komatsu T, et al. (2012) Retinoic acid receptor agonists regulate expression of ATP-binding cassette transporter G1 in macrophages. *Biochim Biophys Acta* **1821**:561–572.
- Azuma Y, Takada M, Shin HW, Kioka N, Nakayama K, and Ueda K (2009) Retroendocytosis pathway of ABCA1/apoA-I contributes to HDL formation. *Genes Cells* **14**:191–204.
- Baigent C, Blackwell L, Emberson J, Holland LE, Reith C, Bhalra N, Peto R, Barnes EH, Keech A, Simes J, et al.; Cholesterol Treatment Trialists' (CTT) Collaboration (2010) Efficacy and safety of more intensive lowering of LDL cholesterol: a meta-analysis of data from 170,000 participants in 26 randomised trials. *Lancet* **376**:1670–1681.
- Baigent C, Keech A, Kearney PM, Blackwell L, Buck G, Pollicino C, Kirby A, Sourjina T, Peto R, Collins R, et al.; Cholesterol Treatment Trialists' (CTT) Collaborators (2005) Efficacy and safety of cholesterol-lowering treatment: prospective meta-analysis of data from 90,056 participants in 14 randomised trials of statins [published correction appears in *Lancet* (2005) 366:1358; *Lancet* (2008) 371:2084]. *Lancet* **366**:1267–1278.
- Balmer JE and Blomhoff R (2002) Gene expression regulation by retinoic acid. *J Lipid Res* **43**:1773–1808.
- Barter PJ, Nicholls SJ, Kastelein JJ, and Rye KA (2015) Is cholesteryl ester transfer protein inhibition an effective strategy to reduce cardiovascular risk? CETP inhibition as a strategy to reduce cardiovascular risk: the pro case. *Circulation* **132**:423–432.
- Barter PJ and Rye KA (2012) Cholesteryl ester transfer protein inhibition as a strategy to reduce cardiovascular risk. *J Lipid Res* **53**:1755–1766.
- Bekkering S, Quintin J, Joosten LA, van der Meer JW, Netea MG, and Riksen NP (2014) Oxidized low-density lipoprotein induces long-term proinflammatory cytokine production and foam cell formation via epigenetic reprogramming of monocytes. *Arterioscler Thromb Vasc Biol* **34**:1731–1738.
- Bennett DJ, Cooke AJ, and Edwards AS (2006) Non-steroidal LXR agonists; an emerging therapeutic strategy for the treatment of atherosclerosis. *Recent Pat Cardiovasc Drug Discov* **1**:21–46.
- Bentzon JF, Otsuka F, Virmani R, and Falk E (2014) Mechanisms of plaque formation and rupture. *Circ Res* **114**:1852–1866.
- Berger JP, Akiyama TE, and Meinke PT (2005) PPARs: therapeutic targets for metabolic disease. *Trends Pharmacol Sci* **26**:244–251.
- Beyer TP, Schmidt RJ, Foxworthy P, Zhang Y, Dai J, Bensch WR, Kauffman RF, Gao H, Ryan TP, Jiang XC, et al. (2004) Coadministration of a liver X receptor agonist and a peroxisome proliferator activator receptor-alpha agonist in mice: effects of nuclear receptor interplay on high-density lipoprotein and triglyceride metabolism in vivo. *J Pharmacol Exp Ther* **309**:861–868.
- Bezafibrate Infarction Prevention (BIP) Study (2000) Secondary prevention by raising HDL cholesterol and reducing triglycerides in patients with coronary artery disease. *Circulation* **102**:21–27.
- Bielicki JK, Zhang H, Cortez Y, Zheng Y, Narayanaswami V, Patel A, Johansson J, and Azhar S (2010) A new HDL mimetic peptide that stimulates cellular cholesterol efflux with high efficiency greatly reduces atherosclerosis in mice. *J Lipid Res* **51**:1496–1503.
- Bingham TC, Fisher EA, Parathath S, Reiss AB, Chan ES, and Cronstein BN (2010) A2A adenosine receptor stimulation decreases foam cell formation by enhancing ABCA1-dependent cholesterol efflux. *J Leukoc Biol* **87**:683–690.
- Bitzur R, Cohen H, Kamari Y, and Harats D (2013) Intolerance to statins: mechanisms and management. *Diabetes Care* **36** (Suppl 2):S325–S330.
- Blom T, Bäck N, Mutka A-L, Bittman R, Li Z, de Lera A, Kovanen PT, Diczfalussy U, and Ikonen E (2010) FTY720 stimulates 27-hydroxycholesterol production and confers atheroprotective effects in human primary macrophages. *Circ Res* **106**:720–729.
- Bojanic DD, Tarr PT, Gale GD, Smith DJ, Bok D, Chen B, Nusinowitz S, Lövgren-Sandblom A, Björkhem I, and Edwards PA (2010) Differential expression and function of ABCG1 and ABCG4 during development and aging. *J Lipid Res* **51**:169–181.
- Bortnick AE, Rothblat GH, Stoudt G, Hoppe KL, Royer LJ, McNeish J, and Francone OL (2000) The correlation of ATP-binding cassette 1 mRNA levels with cholesterol efflux from various cell lines. *J Biol Chem* **275**:28634–28640.
- Briand F, Naik SU, Fukui I, Millar JS, Macphree C, Walker M, Billheimer J, Rothblat G, and Rader DJ (2009) Both the peroxisome proliferator-activated receptor delta agonist, GW0742, and ezetimibe promote reverse cholesterol transport in mice by reducing intestinal reabsorption of HDL-derived cholesterol. *Clin Transl Sci* **2**:127–133.
- Broderick MR, Rhoads D, Charpentier D, Mihalache-Avram T, Mecteau M, Brand G, Chapat E, Perez A, Niesor EJ, Rhéaume E, et al. (2017) Dalcetrapib and anacetrapib differently impact HDL structure and function in rabbits and monkeys. *J Lipid Res* **58**:1282–1291.
- Brooks-Wilson A, Marcil M, Clee SM, Zhang LH, Roomp K, van Dam M, Yu L, Brewer C, Collins JA, Molhuizen HO, et al. (1999) Mutations in ABC1 in Tangier disease and familial high-density lipoprotein deficiency. *Nat Genet* **22**:336–345.
- Brunham LR, Kruit JK, Pape TD, Parks JS, Kuipers F, and Hayden MR (2006) Tissue-specific induction of intestinal ABCA1 expression with a liver X receptor agonist raises plasma HDL cholesterol levels. *Circ Res* **99**:672–674.
- Bungert S, Molday LL, and Molday RS (2001) Membrane topology of the ATP binding cassette transporter ABCR and its relationship to ABC1 and related ABCA transporters: identification of N-linked glycosylation sites. *J Biol Chem* **276**:23539–23546.
- Cabrero A, Cubero M, Llaverías G, Jové M, Planavila A, Alegret M, Sánchez R, Laguna JC, and Carrera MV (2003) Differential effects of peroxisome proliferator-activated receptor activators on the mRNA levels of genes involved in lipid metabolism in primary human monocyte-derived macrophages. *Metabolism* **52**:652–657.
- Cadoudal T, Glorian M, Massias A, Fouque F, Forest C, and Benelli C (2008) Retinoids upregulate phosphoenolpyruvate carboxykinase and glycerooneogenesis in human and rodent adipocytes. *J Nutr* **138**:1004–1009.
- Campia I, Sala V, Kopecka J, Leo C, Mitro N, Costamagna C, Caruso D, Pescarmona G, Crepaldi T, Ghigo D, et al. (2012) Digoxin and ouabain induce the efflux of cholesterol via liver X receptor signalling and the synthesis of ATP in cardiomyocytes. *Biochem J* **447**:301–311.
- Canfrán-Duque A, Ramírez CM, Goedeke L, Lin CS, and Fernández-Hernando C (2014) microRNAs and HDL life cycle. *Cardiovasc Res* **103**:414–422.
- Canfrán-Duque A, Rotllan N, Zhang X, Fernández-Fuertes M, Ramírez-Hidalgo C, Araldi E, Daimiel L, Busto R, Fernández-Hernando C, and Suárez Y (2017) Macrophage deficiency of miR-21 promotes apoptosis, plaque necrosis, and vascular inflammation during atherogenesis. *EMBO Mol Med* **9**:1244–1262.
- Cao YN, Xu L, Han YC, Wang YN, Liu G, and Qi R (2017) Recombinant high-density lipoproteins and their use in cardiovascular diseases. *Drug Discov Today* **22**:180–185.
- Carter AY, Letronne F, Fitz NF, Mounier A, Wolfe CM, Nam KN, Reeves VL, Kamboh H, Lefterov I, and Koldamova R (2017) Liver X receptor agonist treatment significantly affects phenotype and transcriptome of APOE3 and APOE4 Abca1 haplo-deficient mice. *PLoS One* **12**:e0172161.
- Castrillo A, Joseph SB, Vaidya SA, Haberland M, Fogelman AM, Cheng G, and Tontonoz P (2003) Crosstalk between LXR and Toll-like receptor signaling mediates bacterial and viral antagonism of cholesterol metabolism. *Mol Cell* **12**:805–816.
- Castro Navas FF, Giorgi G, Maggioni D, Pacciarini M, Russo V, and Marinuzzi M (2018) C24-hydroxylated stigmastane derivatives as liver X receptor agonists. *Chem Phys Lipids* **212**:44–50.
- Chambenoit O, Hamon Y, Marguet D, Rigneault H, Rosseneu M, and Chimini G (2001) Specific docking of apolipoprotein A-I at the cell surface requires a functional ABCA1 transporter. *J Biol Chem* **276**:9955–9960.
- Chandra M, Miriyala S, and Panchatcharam M (2017) PPAR γ and its role in cardiovascular diseases. *PPAR Res* **2017**:6404638.
- Chawla A, Boisvert WA, Lee CH, Laffitte BA, Barak Y, Joseph SB, Liao D, Nagy L, Edwards PA, Curtiss LK, et al. (2001) A PPAR gamma-LXR-ABCA1 pathway in macrophages is involved in cholesterol efflux and atherogenesis. *Mol Cell* **7**:161–171.
- Chen DY, Chih HM, Lan JL, Chang HY, Chen WW, and Chiang EP (2011a) Blood lipid profiles and peripheral blood mononuclear cell cholesterol metabolism gene expression in patients with and without methotrexate treatment. *BMC Med* **9**:4.
- Chen J, Costa LG, and Guizzetti M (2011b) Retinoic acid isomers up-regulate ATP binding cassette A1 and G1 and cholesterol efflux in rat astrocytes: implications for their therapeutic and teratogenic effects. *J Pharmacol Exp Ther* **338**:870–878.

- Chen J, Zhao L, Sun D, Narsinh K, Li C, Zhang Z, Qi S, Wei G, Li W, Guo W, et al. (2012) Liver X receptor activation attenuates plaque formation and improves vasomotor function of the aortic artery in atherosclerotic ApoE(-/-) mice. *Inflamm Res* **61**:1299–1307.
- Chen L, Yao Q, Xu S, Wang H, and Qu P (2018) Inhibition of the NLRP3 inflammasome attenuates foam cell formation of THP-1 macrophages by suppressing ox-LDL uptake and promoting cholesterol efflux. *Biochem Biophys Res Commun* **495**:382–387.
- Chen SG, Xiao J, Liu XH, Liu MM, Mo ZC, Yin K, Zhao GJ, Jiang J, Cui LB, Tan CZ, et al. (2010) Ibrulipim increases ABCA1/G1 expression by the LXRA signaling pathway in THP-1 macrophage-derived foam cells. *Acta Pharmacol Sin* **31**:1343–1349.
- Chinetti G, Lestavel S, Bocher V, Remaley AT, Neve B, Torra IP, Teissier E, Minnich A, Jaye M, Duverger N, et al. (2001) PPAR-alpha and PPAR-gamma activators induce cholesterol removal from human macrophage foam cells through stimulation of the ABCA1 pathway. *Nat Med* **7**:53–58.
- Claudel T, Leibowitz MD, Fiévet C, Tailleux A, Wagner B, Repa JJ, Torpier G, Lobaccaro JM, Paterniti JR, Mangelsdorf DJ, et al. (2001) Reduction of atherosclerosis in apolipoprotein E knockout mice by activation of the retinoid X receptor. *Proc Natl Acad Sci USA* **98**:2610–2615.
- Cocks G, Wilde JI, Graham SJ, Bousgouni V, Virley D, Lovestone S, and Richardson J (2010) The thiazolidinedione pioglitazone increases cholesterol biosynthetic gene expression in primary cortical neurons by a PPARgamma-independent mechanism. *J Alzheimers Dis* **19**:631–646.
- Connolly BA, O'Connell DP, Lamont-Fava S, LeBlanc DF, Kuang Y-L, Schaefer EJ, Coggale AL, Benedict CP, and Bachovchin WW (2013) The high-fat high-fructose hamster as an animal model for niacin's biological activities in humans. *Metabolism* **62**:1840–1849.
- Copsey S, Garcia C, Diez F, Vermeulen M, Baldi A, Bianciotti LG, Russel FG, Shayo C, and Davio C (2011) Multidrug resistance protein 4 (MRP4/ABCC4) regulates cAMP cellular levels and controls human leukemia cell proliferation and differentiation. *J Biol Chem* **286**:6979–6988.
- Costet P, Lallanne F, Gerbod-Giannone MC, Molina JR, Fu X, Lund EG, Gudas LJ, and Tall AR (2003) Retinoic acid receptor-mediated induction of ABCA1 in macrophages. *Mol Cell Biol* **23**:7756–7766.
- Cronican AA, Fitz NF, Pham T, Fogg A, Kifer B, Koldamova R, and Lefterov I (2010) Proton pump inhibitor lansoprazole is a nuclear liver X receptor agonist. *Biochem Pharmacol* **79**:1310–1316.
- Cruz PM, Mo H, McConathy WJ, Sabnis N, and Lacko AG (2013) The role of cholesterol metabolism and cholesterol transport in carcinogenesis: a review of scientific findings, relevant to future cancer therapeutics. *Front Pharmacol* **4**:119.
- Cui H, Okuhira K, Ohoka N, Naito M, Kagechika H, Hirose A, and Nishimaki-Mogami T (2011) Tributyltin chloride induces ABCA1 expression and apolipoprotein A1-mediated cellular cholesterol efflux by activating LXRA/RXR. *Biochem Pharmacol* **81**:819–824.
- Dai X-Y, Ou X, Hao X-R, Cao D-L, Tang Y-L, Hu Y-W, Li X-X, and Tang C-K (2008) The effect of T0901317 on ATP-binding cassette transporter A1 and Niemann-Pick type C1 in apoE(-/-) mice. *J Cardiovasc Pharmacol* **51**:467–475.
- Dangwal S and Thum T (2014) microRNA therapeutics in cardiovascular disease models. *Annu Rev Pharmacol Toxicol* **54**:185–203.
- Dávalos A and Fernández-Hernando C (2013) From evolution to revolution: miRNAs as pharmacological targets for modulating cholesterol efflux and reverse cholesterol transport. *Pharmacol Ther* **75**:60–72.
- Davidson MH, Maki KC, Pearson TA, Pasternak RC, Deedwania PC, McKenney JM, Fonarow GC, Maron DJ, Ansell BJ, Clark LT, et al. (2005) Results of the National Cholesterol Education (NCEP) Program Evaluation Project T Utilizing Novel E-Technology (NEPTUNE) II survey and implications for treatment under the recent NCEP Writing Group recommendations. *Am J Cardiol* **96**:556–563.
- de Aguiar Vallim TQ, Tarling EJ, Kim T, Civelek M, Baldán A, Esau C, and Edwards PA (2013) MicroRNA-144 regulates hepatic ATP binding cassette transporter A1 and plasma high-density lipoprotein after activation of the nuclear receptor farnesoid X receptor. *Circ Res* **112**:1602–1612.
- De Backer G, Ambrosioni E, Borch-Johnsen K, Brotons C, Cifkova R, Dallongeville J, Ebrahim S, Faergeman O, Graham I, Mancía G, et al.; European Society of Cardiology Committee for Practice Guidelines (2003) European guidelines on cardiovascular disease prevention in clinical practice: third joint task force of European and other societies on cardiovascular disease prevention in clinical practice (constituted by representatives of eight societies and by invited experts). *Eur J Cardiovasc Prev Rehabil* **10**:S1–S10.
- Delvecchio CJ, Bilan P, Nair P, and Capone JP (2008) LXR-induced reverse cholesterol transport in human airway smooth muscle is mediated exclusively by ABCA1. *Am J Physiol Lung Cell Mol Physiol* **295**:L949–L957.
- Delvecchio CJ, Bilan P, Radford K, Stephen J, Trigatti BL, Cox G, Parameswaran K, and Capone JP (2007) Liver X receptor stimulates cholesterol efflux and inhibits expression of proinflammatory mediators in human airway smooth muscle cells. *Mol Endocrinol* **21**:1324–1334.
- Denis M, Landry YD, and Zha X (2008) ATP-binding cassette A1-mediated lipidation of apolipoprotein A-I occurs at the plasma membrane and not in the endocytic compartments. *J Biol Chem* **283**:16178–16186.
- Di D, Wang Z, Liu Y, Luo G, Shi Y, Berggren-Söderlund M, Nilsson-Ehle P, Zhang X, and Xu N (2012) ABCA1 upregulating apolipoprotein M expression mediates via the RXR/LXR pathway in HepG2 cells. *Biochem Biophys Res Commun* **421**:152–156.
- DiBlasio-Smith EA, Arai M, Quinet EM, Evans MJ, Kornaga T, Basso MD, Chen L, Feingold I, Halpern AR, Liu Q-Y, et al. (2008) Discovery and implementation of transcriptional biomarkers of synthetic LXR agonists in peripheral blood cells. *J Transl Med* **6**:59.
- DiMarco DM and Fernandez ML (2015) The regulation of reverse cholesterol transport and cellular cholesterol homeostasis by microRNAs. *Biology (Basel)* **4**:494–511.
- Downs JR, Clearfield M, Weis S, Whitney E, Shapiro DR, Beere PA, Langendorfer A, Stein EA, Kruyer W, and Gotto AM Jr (1998) Primary prevention of acute coronary events with lovastatin in men and women with average cholesterol levels: results of AFCAPS/TexCAPS. Air Force/Texas Coronary Atherosclerosis Prevention Study. *JAMA* **279**:1615–1622.
- Du XM, Kim MJ, Hou L, Le Goff W, Chapman MJ, Van Eck M, Curtiss LK, Burnett JR, Cartland SP, Quinn CM, et al. (2015a) HDL particle size is a critical determinant of ABCA1-mediated macrophage cellular cholesterol export. *Circ Res* **116**:1133–1142.
- Du Y, Wang L, Si S, Yang Y, and Hong B (2015b) A novel compound 4010B-30 upregulates apolipoprotein A-I gene expression through activation of PPARγ in HepG2 cells. *Atherosclerosis* **239**:589–598.
- Eddleston JM and Shelly MP (1991) The effect on serum lipid concentrations of a prolonged infusion of propofol-hypertriglyceridaemia associated with propofol administration. *Intensive Care Med* **17**:424–426.
- Edwards PA, Kennedy MA, and Mak PA (2002) LXRs; oxysterol-activated nuclear receptors that regulate genes controlling lipid homeostasis. *Vascul Pharmacol* **38**:249–256.
- ElAli A and Hermann DM (2012) Liver X receptor activation enhances blood-brain barrier integrity in the ischemic brain and increases the abundance of ATP-binding cassette transporters ABCB1 and ABCG1 on brain capillary cells. *Brain Pathol* **22**:175–187.
- El Roz A, Bard JM, Huvelin JM, and Nazih H (2012) LXR agonists and ABCG1-dependent cholesterol efflux in MCF-7 breast cancer cells: relation to proliferation and apoptosis. *Anticancer Res* **32**:3007–3013.
- Escher P, Braissant O, Basu-Modak S, Michalik L, Wahli W, and Desvergne B (2001) Rat PPARs: quantitative analysis in adult rat tissues and regulation in fasting and refeeding. *Endocrinology* **142**:4195–4202.
- Faulkner LE, Panagiotopoulos SE, Johnson JD, Woollett LA, Hui DY, Witting SR, Maiorano JN, and Davidson WS (2008) An analysis of the role of a retroendocytosis pathway in ABCA1-mediated cholesterol efflux from macrophages. *J Lipid Res* **49**:1322–1332.
- Fazio S and Linton MF (2003) Apolipoprotein AI as therapy for atherosclerosis: does the future of preventive cardiology include weekly injections of the HDL protein? *Mol Interv* **3**:436–440.
- Feinberg MW and Moore KJ (2016) MicroRNA regulation of atherosclerosis. *Circ Res* **118**:703–720.
- Fernández-Hernando C and Moore KJ (2011) MicroRNA modulation of cholesterol homeostasis. *Arterioscler Thromb Vasc Biol* **31**:2378–2382.
- Fernández-Hernando C, Suárez Y, Rayner KJ, and Moore KJ (2011) MicroRNAs in lipid metabolism. *Curr Opin Lipidol* **22**:86–92.
- Ferri N, Corsini A, Sirtori C, and Ruscica M (2017) PPAR-α agonists are still on the rise: an update on clinical and experimental findings. *Expert Opin Investig Drugs* **26**:593–602.
- Field FJ, Watt K, and Mathur SN (2010) TNF-alpha decreases ABCA1 expression and attenuates HDL cholesterol efflux in the human intestinal cell line Caco-2. *J Lipid Res* **51**:1407–1415.
- Fitzgerald ML, Morris AL, Chroni A, Mendez AJ, Zannis VI, and Freeman MW (2004) ABCA1 and amphipathic apolipoproteins form high-affinity molecular complexes required for cholesterol efflux. *J Lipid Res* **45**:287–294.
- Forcheron F, Cachefo A, Thevenon S, Pinteaur C, and Beylot M (2002) Mechanisms of the triglyceride- and cholesterol-lowering effect of fenofibrate in hyperlipidemic type 2 diabetic patients. *Diabetes* **51**:3486–3491.
- Frick MH, Elo O, Haapa K, Heinonen OP, Heinsalmi P, Helo P, Huttunen JK, Kaitaniemi P, Koskenp B, Manninen V, et al. (1987) Helsinki Heart Study: primary-prevention trial with gemfibrozil in middle-aged men with dyslipidemia. Safety of treatment, changes in risk factors, and incidence of coronary heart disease. *N Engl J Med* **317**:1237–1245.
- Fujiyoshi M, Ohtsuki S, Hori S, Tachikawa M, and Terasaki T (2007) 24S-hydroxycholesterol induces cholesterol release from choroid plexus epithelial cells in an apical- and apoE isoform-dependent manner concomitantly with the induction of ABCA1 and ABCG1 expression. *J Neurochem* **100**:968–978.
- Fukumoto H, Deng A, Irizarry MC, Fitzgerald ML, and Rebeck GW (2002) Induction of the cholesterol transporter ABCA1 in central nervous system cells by liver X receptor agonists increases secreted Abeta levels. *J Biol Chem* **277**:48508–48513.
- Gaidarov I, Chen X, Anthony T, Maciejewski-Lenoir D, Liaw C, and Unett DJ (2013) Differential tissue and ligand-dependent signaling of GPR109A receptor: implications for anti-atherosclerotic therapeutic potential. *Cell Signal* **25**:2003–2016.
- Gao J, Xu Y, Yang Y, Yang Y, Zheng Z, Jiang W, Hong B, Yan X, and Si S (2008) Identification of upregulators of human ATP-binding cassette transporter A1 via high-throughput screening of a synthetic and natural compound library. *J Biomol Screen* **13**:648–656.
- Gelissen IC, Sharpe LJ, Sandoval C, Rao G, Kockx M, Kritharides L, Jessup W, and Brown AJ (2012) Protein kinase A modulates the activity of a major human isoform of ABCG1. *J Lipid Res* **53**:2133–2140.
- Gene A (1982–2019a) *Gene ABCA1 [Homo sapiens (human)]* [Gene ID: 19], National Library of Medicine (US), National Center for Biotechnology Information, Bethesda, MD.
- Gene A (1982–2019b) *Gene ABCG1 [Homo sapiens (human)]* [Gene ID: 9619], National Library of Medicine (US), National Center for Biotechnology Information, Bethesda, MD.
- Gerstein HC, Yusuf S, Bosch J, Pogue J, Sheridan P, Dinccag N, Hanefeld M, Hoogwerf B, Laakso M, Mohan V, et al.; DREAM (Diabetes REduction Assessment with ramipril and rosiglitazone Medication) Trial Investigators (2006) Effect of rosiglitazone on the frequency of diabetes in patients with impaired glucose tolerance or impaired fasting glucose: a randomised controlled trial. *Lancet* **368**:1096–1105.
- Getz GS and Reardon CA (2011) Apolipoprotein A-I and A-I mimetic peptides: a role in atherosclerosis. *J Inflamm Res* **4**:83–92.
- Ginsberg HN, Elam MB, Lovato LC, Crouse JR III, Leiter LA, Linz P, Friedewald WT, Buse JB, Gerstein HC, Probstfield J, et al. ACCORD Study Group (2010) Effects of combination lipid therapy in type 2 diabetes mellitus. *N Engl J Med* **362**:1563–1574.
- Glaros EN, Kim WS, Quinn CM, Wong J, Gelissen I, Jessup W, and Garner B (2005) Glycosphingolipid accumulation inhibits cholesterol efflux via the ABCA1/apolipoprotein A-I pathway: 1-phenyl-2-decanoylamino-3-morpholino-1-propanol is a novel cholesterol efflux accelerator. *J Biol Chem* **280**:24515–24523.

- Goedeke L, Aranda JF, and Fernández-Hernando C (2014) microRNA regulation of lipoprotein metabolism. *Curr Opin Lipidol* **25**:282–288.
- Goedeke L, Rotllan N, Canfrán-Duque A, Aranda JF, Ramírez CM, Araldi E, Lin CS, Anderson NN, Wagschal A, de Cabo R, et al. (2015a) MicroRNA-148a regulates LDL receptor and ABCA1 expression to control circulating lipoprotein levels. *Nat Med* **21**:1280–1289.
- Goedeke L, Rotllan N, Ramírez CM, Aranda JF, Canfrán-Duque A, Araldi E, Fernández-Hernando A, Langhi C, de Cabo R, Baldán Á, et al. (2015b) miR-27b inhibits LDLR and ABCA1 expression but does not influence plasma and hepatic lipid levels in mice. *Atherosclerosis* **243**:499–509.
- Gong Y-Z, Sun S-W, Yuan H-Y, Xie X-J, Tuo Q-H, Yang X-F, Tan X, Zheng X-L, Xiong Y, and Liao D-F (2014) Ezetimibe-mediated protection of vascular smooth muscle cells from cholesterol accumulation through the regulation of lipid metabolism-related gene expression. *Pharmacology* **94**:214–222.
- Graf GA, Yu L, Li WP, Gerard R, Tuma PL, Cohen JC, and Hobbs HH (2003) ABCG5 and ABCG8 are obligate heterodimers for protein trafficking and biliary cholesterol excretion. *J Biol Chem* **278**:48275–48282.
- Grygiel-Górniak B (2014) Peroxisome proliferator-activated receptors and their ligands: nutritional and clinical implications—a review. *Nutr J* **13**:17.
- Guan J-Z, Tamasawa N, Murakami H, Matsui J, Tanabe J, Matsuki K, Yamashita M, and Suda T (2008) HMG-CoA reductase inhibitor, simvastatin improves reverse cholesterol transport in type 2 diabetic patients with hyperlipidemia. *J Atheroscler Thromb* **15**:20–25.
- Guan J-Z, Tamasawa N, Murakami H, Matsui J, Yamato K, and Suda T (2003) Clofibrate, a peroxisome-proliferator, enhances reverse cholesterol transport through cytochrome P450 activation and oxysterol generation. *Tohoku J Exp Med* **201**:251–259.
- Gulshan K, Brubaker G, Conger H, Wang S, Zhang R, Hazen SL, and Smith JD (2016) PI(4,5)P2 is translocated by ABCA1 to the cell surface where it mediates apolipoprotein A1 binding and nascent HDL assembly. *Circ Res* **119**:827–838.
- Gupta H, White CR, Handattu S, Garber DW, Datta G, Chaddha M, Dai L, Gianturco SH, Bradley WA, and Anantharamaiah GM (2005) Apolipoprotein E mimetic peptide dramatically lowers plasma cholesterol and restores endothelial function in watanabe heritable hyperlipidemic rabbits. *Circulation* **111**:3112–3118.
- Guyton A and Hall J (2011) *Textbook of Medical Physiology*, Saunders Elsevier, Philadelphia.
- Haidar B, Denis M, Krimbou L, Marcil M, and Genest J Jr (2002) cAMP induces ABCA1 phosphorylation activity and promotes cholesterol efflux from fibroblasts. *J Lipid Res* **43**:2087–2094.
- Haidar B, Denis M, Marcil M, Krimbou L, and Genest J Jr (2004) Apolipoprotein A-I activates cellular cAMP signaling through the ABCA1 transporter. *J Biol Chem* **279**:9963–9969.
- Halkes PH, van Gijn J, Kappelle LJ, Koudstaal PJ, and Algra A; ESPRIT Study Group (2006) Aspirin plus dipyridamole versus aspirin alone after cerebral ischaemia of arterial origin (ESPRIT): randomised controlled trial [published correction appears in *Lancet* (2007) 369:274]. *Lancet* **367**:1665–1673.
- Hammer SS, Beli E, Kady N, Wang Q, Wood K, Lydic TA, Malek G, Saban DR, Wang XX, Hazra S, et al. (2017) The mechanism of diabetic retinopathy pathogenesis unifying key lipid regulators, sirtuin 1 and liver X receptor. *EBioMedicine* **22**:181–190.
- Handattu SP, Nayyar G, Garber DW, Palgunachari MN, Monroe CE, Keenum TD, Mishra VK, Datta G, and Anantharamaiah GM (2013) Two apolipoprotein E mimetic peptides with similar cholesterol reducing properties exhibit differential atheroprotective effects in LDL-R null mice. *Atherosclerosis* **227**:58–64.
- Hassan HH, Denis M, Lee DY, Iatan I, Nyholt D, Ruel I, Krimbou L, and Genest J (2007) Identification of an ABCA1-dependent phospholipid-rich plasma membrane apolipoprotein A-I binding site for nascent HDL formation: implications for current models of HDL biogenesis. *J Lipid Res* **48**:2428–2442.
- Hazen SL and Smith JD (2012) An antiatherosclerotic signaling cascade involving intestinal microbiota, microRNA-10b, and ABCA1/ABCG1-mediated reverse cholesterol transport. *Circ Res* **111**:948–950.
- He X, Chen X, Wang L, Wang W, Liang Q, Yi L, Wang Y, and Gao Q (2019) Metformin ameliorates Ox-LDL-induced foam cell formation in raw264.7 cells by promoting ABCG1 mediated cholesterol efflux. *Life Sci* **216**:67–74.
- He Y, Lin L, Cao J, Mao X, Qu Y, and Xi B (2015) Up-regulated miR-93 contributes to coronary atherosclerosis pathogenesis through targeting ABCA1. *Int J Clin Exp Med* **8**:674–681.
- Hegele RA (2009) Plasma lipoproteins: genetic influences and clinical implications. *Nat Rev Genet* **10**:109–121.
- Hewing B and Fisher EA (2012) Rationale for cholesteryl ester transfer protein inhibition. *Curr Opin Lipidol* **23**:372–376.
- Higashiyama H, Billin AN, Okamoto Y, Kinoshita M, and Asano S (2007) Expression profiling of peroxisome proliferator-activated receptor-delta (PPAR-delta) in mouse tissues using tissue microarray. *Histochem Cell Biol* **127**:485–494.
- Home PD, Pocock SJ, Beck-Nielsen H, Curtis PS, Gomis R, Hanefeld M, Jones NP, Komajda M, and McMurray JJ; RECORD Study Team (2009) Rosiglitazone evaluated for cardiovascular outcomes in oral agent combination therapy for type 2 diabetes (RECORD): a multicentre, randomised, open-label trial. *Lancet* **373**:2125–2135.
- Hong C and Tontonoz P (2014) Liver X receptors in lipid metabolism: opportunities for drug discovery. *Nat Rev Drug Discov* **13**:433–444.
- Honzumi S, Shima A, Hiroshima A, Koieyama T, and Terasaka N (2011) Synthetic LXR agonist inhibits the development of atherosclerosis in New Zealand White rabbits. *Biochim Biophys Acta* **1811**:1136–1145.
- Hossain MA, Tsujita M, Gonzalez FJ, and Yokoyama S (2008) Effects of fibrate drugs on expression of ABCA1 and HDL biogenesis in hepatocytes. *J Cardiovasc Pharmacol* **51**:258–266.
- Hou M, Xia M, Zhu H, Wang Q, Li Y, Xiao Y, Zhao T, Tang Z, Ma J, and Ling W (2007) Lysophosphatidylcholine promotes cholesterol efflux from mouse macrophage foam cells via PPARgamma-LXRalpha-ABCA1-dependent pathway associated with apoE. *Cell Biochem Funct* **25**:33–44.
- Hsu CP, Lin CH, and Kuo CY (2018) Endothelial-cell inflammation and damage by reactive oxygen species are prevented by propofol via ABCA1-mediated cholesterol efflux. *Int J Med Sci* **15**:978–985.
- Huang L, Fan B, Ma A, Shaul PW, and Zhu H (2015) Inhibition of ABCA1 protein degradation promotes HDL cholesterol efflux capacity and RCT and reduces atherosclerosis in mice. *J Lipid Res* **56**:986–997.
- Iatan I, Palmyre A, Alrasheed S, Ruel I, and Genest J (2012) Genetics of cholesterol efflux. *Curr Atheroscler Rep* **14**:235–246.
- Inaba T, Yagyu H, Itabashi N, Tazoe F, Fujita N, Nagashima S, Okada K, Okazaki M, Furukawa Y, and Ishibashi S (2008) Cholesterol reduction and atherosclerosis inhibition by bezafibrate in low-density lipoprotein receptor knockout mice. *Hypertens Res* **31**:999–1005.
- Insull W Jr (2009) The pathology of atherosclerosis: plaque development and plaque responses to medical treatment. *Am J Med* **122** (Suppl):S3–S14.
- Ishigami M, Ogasawara F, Nagao K, Hashimoto H, Kimura Y, Kioka N, and Ueda K (2018) Temporal sequestration of cholesterol and phosphatidylcholine within extracellular domains of ABCA1 during nascent HDL generation. *Sci Rep* **8**:6170.
- Ishii N, Matsumura T, Kinoshita H, Fukuda K, Motoshima H, Senokuchi T, Nakao S, Tsutsumi A, Kim-Mitsuyama S, Kawada T, et al. (2010) Nifedipine induces peroxisome proliferator-activated receptor-gamma activation in macrophages and suppresses the progression of atherosclerosis in apolipoprotein E-deficient mice. *Arterioscler Thromb Vasc Biol* **30**:1598–1605.
- Iwamoto N, Abe-Dohmae S, Ayaori M, Tanaka N, Kusuha M, Ohsuzu F, and Yokoyama S (2007) ATP-binding cassette transporter A1 gene transcription is downregulated by activator protein 2alpha: doxazosin inhibits activator protein 2alpha and increases high-density lipoprotein biogenesis independent of alpha1-adrenoceptor blockade. *Circ Res* **101**:156–165.
- Iwamoto N, Abe-Dohmae S, Lu R, and Yokoyama S (2008) Involvement of protein kinase D in phosphorylation and increase of DNA binding of activator protein 2 alpha to downregulate ATP-binding cassette transporter A1. *Arterioscler Thromb Vasc Biol* **28**:2282–2287.
- Iwamoto N, Lu R, Tanaka N, Abe-Dohmae S, and Yokoyama S (2010) Calmodulin interacts with ATP binding cassette transporter A1 to protect from calpain-mediated degradation and upregulates high-density lipoprotein generation. *Arterioscler Thromb Vasc Biol* **30**:1446–1452.
- Jakobsson T, Treuter E, Gustafsson JA, and Steffensen KR (2012) Liver X receptor biology and pharmacology: new pathways, challenges and opportunities. *Trends Pharmacol Sci* **33**:394–404.
- Jiang H, Badralmaa Y, Yang J, Lempicki R, Hazen A, and Natarajan V (2012) Retinoic acid and liver X receptor agonist synergistically inhibit HIV infection in CD4+ T cells by up-regulating ABCA1-mediated cholesterol efflux. *Lipids Health Dis* **11**:69.
- Jiang M and Li X (2017) Activation of PPARγ does not contribute to macrophage ABCA1 expression and ABCA1-mediated cholesterol efflux to apoA1. *Biochem Biophys Res Commun* **482**:849–856.
- Jin S, Mathis AS, Gioia K, Minko T, Friedman GS, Rosenblatt J, Peng F, Serur DS, and Knipp GT (2004) Effect of tacrolimus on the expression of macrophage scavenger and nuclear hormone receptors in THP-1-derived human macrophages. *Transplantation* **77**:1281–1287.
- Jung CG, Horike H, Cha BY, Uhm KO, Yamauchi R, Yamaguchi T, Hosono T, Iida K, Woo JT, and Michikawa M (2010) Honokiol increases ABCA1 expression level by activating retinoid X receptor beta. *Biol Pharm Bull* **33**:1105–1111.
- Kaneko T, Kanno C, Ichikawa-Tomikawa N, Kashiwagi K, Yaginuma N, Ohkoshi C, Tanaka M, Sugino T, Imura T, Hasegawa H, et al. (2015) Liver X receptor reduces proliferation of human oral cancer cells by promoting cholesterol efflux via up-regulation of ABCA1 expression. *Oncotarget* **6**:33345–33357.
- Kang MH, Zhang LH, Wijesekara N, de Haan W, Butland S, Bhattacharjee A, and Hayden MR (2013) Regulation of ABCA1 protein expression and function in hepatic and pancreatic islet cells by miR-145. *Arterioscler Thromb Vasc Biol* **33**:2724–2732.
- Kannisto K, Gäfvels M, Jiang Z-Y, Slätis K, Hu X, Jorns C, Steffensen KR, and Eggertsen G (2014) LXR driven induction of HDL-cholesterol is independent of intestinal cholesterol absorption and ABCA1 protein expression. *Lipids* **49**:71–83.
- Karalis DG (2009) Intensive lowering of low-density lipoprotein cholesterol levels for primary prevention of coronary artery disease. *Mayo Clin Proc* **84**:345–352.
- Karalis I and Jukema JW (2018) HDL mimetics infusion and regression of atherosclerosis: is it still considered a valid therapeutic option? *Curr Cardiol Rep* **20**:66.
- Kasada R, Tsuchida Y, Yang HC, Yancey PG, Zhong J, Tao H, Bian A, Fogo AB, Linton MRF, Fazio S, et al. (2018) Chronic kidney disease alters lipid trafficking and inflammatory responses in macrophages: effects of liver X receptor agonism. *BMC Nephrol* **19**:17.
- Katz A, Umeta C, Ott E, Hickey L, Burczynski ME, Burghart P, Vesterqvist O, and Meng X (2009) Safety, pharmacokinetics, and pharmacodynamics of single doses of LXR-623, a novel liver X-receptor agonist, in healthy participants. *J Clin Pharmacol* **49**:643–649.
- Keech A, Simes RJ, Barter P, Best J, Scott R, Taskiran MR, Forder P, Pillai A, Davis T, Glasziou P, et al.; FIELD Study Investigators (2005) Effects of long-term fenofibrate therapy on cardiovascular events in 9795 people with type 2 diabetes mellitus (the FIELD study): randomised controlled trial [published correction appears in *Lancet* (2006) 368:1420]. *Lancet* **366**:1849–1861.
- Kellner-Weibel G, Luke SJ, and Rothblat GH (2003) Cytotoxic cellular cholesterol is selectively removed by apoA-I via ABCA1. *Atherosclerosis* **171**:235–243.
- Kemmerer M, Wittig I, Richter F, Brüne B, and Namgaladze D (2016) AMPK activates LXRα and ABCA1 expression in human macrophages. *Int J Biochem Cell Biol* **78**:1–9.
- Kennedy MA, Barrera GC, Nakamura K, Baldán A, Tarr P, Fishbein MC, Frank J, Francone OL, and Edwards PA (2005) ABCG1 has a critical role in mediating cholesterol efflux to HDL and preventing cellular lipid accumulation. *Cell Metab* **1**:121–131.

- Kennedy MA, Venkateswaran A, Tarr PT, Xenarios I, Kudoh J, Shimizu N, and Edwards PA (2001) Characterization of the human ABCG1 gene: liver X receptor activates an internal promoter that produces a novel transcript encoding an alternative form of the protein. *J Biol Chem* **276**:39438–39447.
- Kerr ID, Haider AJ, and Gelissen IC (2011) The ABCG family of membrane-associated transporters: you don't have to be big to be mighty. *Br J Pharmacol* **164**: 1767–1779.
- Khera AV, Cuchel M, de la Llera-Moya M, Rodrigues A, Burke MF, Jafri K, French BC, Phillips JA, Mucksavage ML, Wilensky RL, et al. (2011) Cholesterol efflux capacity, high-density lipoprotein function, and atherosclerosis. *N Engl J Med* **364**: 127–135.
- Kick E, Martin R, Xie Y, Flatt B, Schweiger E, Wang TL, Busch B, Nyman M, Gu XH, Yan G, et al. (2015) Liver X receptor (LXR) partial agonists: biaryl pyrazoles and imidazoles displaying a preference for LXR β . *Bioorg Med Chem Lett* **25**:372–377.
- Kido K, Evans RA, Gopinath A, and Flynn JD (2018) Severe hypertriglyceridemia induced by sirolimus treated with medical management without plasmapheresis: a case report. *J Pharm Pract* **31**:104–106.
- Kim J, Yoon H, Ramirez CM, Lee SM, Hoe HS, Fernández-Hernando C, and Kim J (2012) MiR-106b impairs cholesterol efflux and increases A β levels by repressing ABCA1 expression. *Exp Neurol* **235**:476–483.
- Kim MS, Sweeney TR, Shigenaga JK, Chui LG, Moser A, Grunfeld C, and Feingold KR (2007) Tumor necrosis factor and interleukin 1 decrease RXR α , PPAR α , PPAR γ , LXR α , and the coactivators SRC-1, PGC-1 α , and PGC-1 β in liver cells. *Metabolism* **56**:267–279.
- Kingwell BA, Chapman MJ, Kontush A, and Miller NE (2014) HDL-targeted therapies: progress, failures and future. *Nat Rev Drug Discov* **13**:445–464.
- Kirchgeessner TG, Martin R, Sleph P, Grimm D, Liu X, Lupisella J, Smalley J, Narayanan R, Xie Y, Ostrowski J, et al. (2015) Pharmacological characterization of a novel liver X receptor agonist with partial LXR α activity and a favorable window in nonhuman primates. *J Pharmacol Exp Ther* **352**:305–314.
- Kobayashi A, Takanezawa Y, Hirata T, Shimizu Y, Misasa K, Kioka N, Arai H, Ueda K, and Matsuo M (2006) Efflux of sphingomyelin, cholesterol, and phosphatidylcholine by ABCG1. *J Lipid Res* **47**:1791–1802.
- Kobayashi M, Gouda K, Chisaki I, Ochiai M, Itagaki S, and Iseki K (2011) Regulation mechanism of ABCA1 expression by statins in hepatocytes. *Eur J Pharmacol* **662**:9–14.
- Koldamova RP, Lefterov IM, Ikonovic MD, Skoko J, Lefterov PI, Isanski BA, DeKosky ST, and Lazo JS (2003) 22R-hydroxycholesterol and 9-cis-retinoic acid induce ATP-binding cassette transporter A1 expression and cholesterol efflux in brain cells and decrease amyloid β secretion. *J Biol Chem* **278**:13244–13256.
- Kontush A, Lindahl M, Lhomme M, Calabresi L, Chapman MJ, and Davidson WS (2015) Structure of HDL: particle subclasses and molecular components. *Handb Exp Pharmacol* **224**:3–51.
- Kooistra T, Verschuren L, de Vries-van der Weij J, Koenig W, Toet K, Princen HM, and Kleemann R (2006) Fenofibrate reduces atherogenesis in ApoE $^{-3}$ Leiden mice: evidence for multiple antiatherogenic effects besides lowering plasma cholesterol. *Arterioscler Thromb Vasc Biol* **26**:2322–2330.
- Kratzer A, Buchebner M, Pfeiffer T, Becker TM, Uray G, Miyazaki M, Miyazaki-Anzai S, Ebner B, Chandak PG, Kadam RS, et al. (2009) Synthetic LXR agonist attenuates plaque formation in apoE $^{-3}$ mice without inducing liver steatosis and hypertriglyceridemia. *J Lipid Res* **50**:312–326.
- Kühnast S, Fiocco M, van der Hoorn JW, Princen HM, and Jukema JW (2015) Innovative pharmaceutical interventions in cardiovascular disease: focusing on the contribution of non-HDL-C/LDL-C-lowering versus HDL-C-raising: a systematic review and meta-analysis of relevant preclinical studies and clinical trials. *Eur J Pharmacol* **763**:48–63.
- Kuntz M, Candela P, Saint-Pol J, Lamartinière Y, Boucau MC, Sevin E, Fenart L, and Gosselet F (2015) Bexarotene promotes cholesterol efflux and restricts apical-to-basolateral transport of amyloid- β peptides in an in vitro model of the human blood-brain barrier. *J Alzheimers Dis* **48**:849–862.
- Kwiterovich PO Jr (2000) The metabolic pathways of high-density lipoprotein, low-density lipoprotein, and triglycerides: a current review. *Am J Cardiol* **86**: 51L–10L.
- LaClair KD, Manaye KF, Lee DL, Allard JS, Savonenko AV, Troncoso JC, and Wong PC (2013) Treatment with bexarotene, a compound that increases apolipoprotein-E, provides no cognitive benefit in mutant APP/PS1 mice. *Mol Neurodegener* **8**:18.
- Lagrost L, Gamber P, Dangremont V, Athias A, and Lallemand C (1990) Role of cholesteryl ester transfer protein (CETP) in the HDL conversion process as evidenced by using anti-CETP monoclonal antibodies. *J Lipid Res* **31**:1569–1575.
- Langmann T, Klucken J, Reil M, Liebisch G, Luciani MF, Chimini G, Kaminski WE, and Schmitz G (1999) Molecular cloning of the human ATP-binding cassette transporter 1 (hABCT1): evidence for sterol-dependent regulation in macrophages. *Biochem Biophys Res Commun* **257**:29–33.
- Larrede S, Quinn CM, Jessup W, Frisdal E, Olivier M, Hsieh V, Kim M-J, Van Eck M, Couvert P, Carrie A, et al. (2009) Stimulation of cholesterol efflux by LXR agonists in cholesterol-loaded human macrophages is ABCA1-dependent but ABCG1-independent. *Arterioscler Thromb Vasc Biol* **29**:1930–1936.
- Lauer ME, Graff-Meyer A, Rufer AC, Maugeais C, von der Mark E, Matile H, D'Arcy B, Magg C, Ringler P, Müller SA, et al. (2016) Cholesteryl ester transfer between lipoproteins does not require a ternary tunnel complex with CETP. *J Struct Biol* **194**:191–198.
- Lee J, Hong EM, Byun HW, Choi MH, Jang HJ, Eun CS, Kae SH, and Choi HS (2008) The effect of PPAR α and PPAR γ ligands on inflammation and ABCA1 expression in cultured gallbladder epithelial cells. *Dig Dis Sci* **53**:1707–1715.
- Lee J, Shirk A, Oram JF, Lee SP, and Kuver R (2002) Polarized cholesterol and phospholipid efflux in cultured gall-bladder epithelial cells: evidence for an ABCA1-mediated pathway. *Biochem J* **364**:475–484.
- Lee JY, Karwatsky J, Ma L, and Zha X (2011) ABCA1 increases extracellular ATP to mediate cholesterol efflux to ApoA-I. *Am J Physiol Cell Physiol* **301**:C886–C894.
- Lee JY, Kinch LN, Borek DM, Wang J, Wang J, Urbatsch IL, Xie XS, Grishin NV, Cohen JC, Otwinowski Z, et al. (2016) Crystal structure of the human sterol transporter ABCG5/ABCG8. *Nature* **533**:561–564.
- Levine GN, Keaney JF Jr, and Vita JA (1995) Cholesterol reduction in cardiovascular disease: clinical benefits and possible mechanisms. *N Engl J Med* **332**:512–521.
- Li AC, Binder CJ, Gutierrez A, Brown KK, Plotkin CR, Pattison JW, Vallerod AF, Davis RA, Willson TM, Witztum JL, et al. (2004) Differential inhibition of macrophage foam-cell formation and atherosclerosis in mice by PPAR α , β / δ , and γ . *J Clin Invest* **114**:1564–1576.
- Li AC and Glass CK (2004) PPAR- and LXR-dependent pathways controlling lipid metabolism and the development of atherosclerosis. *J Lipid Res* **45**:2161–2173.
- Li C, Tu Y, Liu T-R, Guo Z-G, Xie D, Zhong J-K, Fan Y-Z, and Lai W-Y (2015) Rosiglitazone attenuates atherosclerosis and increases high-density lipoprotein function in atherosclerotic rabbits. *Int J Mol Med* **35**:715–723.
- Li D, Wang D, Wang Y, Ling W, Feng X, and Xia M (2010) Adenosine monophosphate-activated protein kinase induces cholesterol efflux from macrophage-derived foam cells and alleviates atherosclerosis in apolipoprotein E-deficient mice. *J Biol Chem* **285**:33499–33509.
- Li N, Wang X, Liu P, Lu D, Jiang W, Xu Y, and Si S (2016) E17110 promotes reverse cholesterol transport with liver X receptor β agonist activity in vitro. *Acta Pharm Sin B* **6**:198–204.
- Li N, Xu Y, Feng T, Liu C, Li Y, Wang X, and Si S (2014) Identification of a selective agonist for liver X receptor α (LXR α) via screening of a synthetic compound library. *J Biomol Screen* **19**:566–574.
- Li Y, Wong K, Walsh K, Gao B, and Zang M (2013) Retinoic acid receptor β stimulates hepatic induction of fibroblast growth factor 21 to promote fatty acid oxidation and control whole-body energy homeostasis in mice. *J Biol Chem* **288**:10490–10504.
- Liang B, Wang X, Song X, Bai R, Yang H, Yang Z, Xiao C, and Bian Y (2017) MicroRNA-20a/b regulates cholesterol efflux through post-transcriptional repression of ATP-binding cassette transporter A1. *Biochim Biophys Acta Mol Cell Biol Lipids* **1862**:929–938.
- Libby P (2005) The forgotten majority: unfinished business in cardiovascular risk reduction. *J Am Coll Cardiol* **46**:1225–1228.
- Libby P, Ridker PM, and Hansson GK (2011) Progress and challenges in translating the biology of atherosclerosis. *Nature* **473**:317–325.
- Liebson PR (2010) The thiazolidinedione controversy in cardiovascular risk. *Prev Cardiol* **13**:209–212.
- Lin G and Bornfeldt KE (2002) Cyclic AMP-specific phosphodiesterase 4 inhibitors promote ABCA1 expression and cholesterol efflux. *Biochem Biophys Res Commun* **290**:663–669.
- Liu J, Guo K, Hu L, Luo T, Ma Y, Zhang Y, Lai W, and Guo Z (2019) ZAP70 deficiency promotes reverse cholesterol transport through MAPK/ERK pathway in Jurkat cell. *Mol Immunol* **107**:21–28.
- Liu J, Zhang Z, Xu Y, Feng T, Jiang W, Li Z, Hong B, Xie Z, and Si S (2012) IMB2026791, a xanthone, stimulates cholesterol efflux by increasing the binding of apolipoprotein A-I to ATP-binding cassette transporter A1. *Molecules* **17**:2833–2854.
- Liu XY, Lu Q, Ouyang XP, Tang SL, Zhao GJ, Lv YC, He PP, Kuang HJ, Tang YY, Fu Y, et al. (2013) Apelin-13 increases expression of ATP-binding cassette transporter A1 via activating protein kinase C α signaling in THP-1 macrophage-derived foam cells. *Atherosclerosis* **226**:398–407.
- Liu Z, Wang J, Huang E, Gao S, Li H, Lu J, Tian K, Little PJ, Shen X, Xu S, et al. (2014) Tanshinone IIA suppresses cholesterol accumulation in human macrophages: role of heme oxygenase-1. *J Lipid Res* **55**:201–213.
- Ilaviers G, Rebollo A, Pou J, Vázquez-Carrera M, Sánchez RM, Laguna JC, and Alegret M (2006) Effects of rosiglitazone and atorvastatin on the expression of genes that control cholesterol homeostasis in differentiating monocytes. *Biochem Pharmacol* **71**:605–614.
- Long-Term Intervention with Pravastatin in Ischaemic Disease (LIPID) Study Group (1998) Prevention of cardiovascular events and death with pravastatin in patients with coronary heart disease and a broad range of initial cholesterol levels. *N Engl J Med* **339**:1349–1357.
- Lorenzi I, von Eckardstein A, Cavelier C, Radosavljevic S, and Rohrer L (2008) Apolipoprotein A-I but not high-density lipoproteins are internalised by RAW macrophages: roles of ATP-binding cassette transporter A1 and scavenger receptor BI. *J Mol Med (Berl)* **86**:171–183.
- Lu Q, Tang SL, Liu XY, Zhao GJ, Ouyang XP, Lv YC, He PP, Yao F, Chen WJ, Tang YY, et al. (2013) Tertiary-butylhydroquinone upregulates expression of ATP-binding cassette transporter A1 via nuclear factor E2-related factor 2/heme oxygenase-1 signaling in THP-1 macrophage-derived foam cells. *Circ J* **77**:2399–2408.
- Lu R, Arakawa R, Ito-Osumi C, Iwamoto N, and Yokoyama S (2008) ApoA-I facilitates ABCA1 recycle/accumulation to cell surface by inhibiting its intracellular degradation and increases HDL generation. *Arterioscler Thromb Vasc Biol* **28**: 1820–1824.
- Lu R, Tsuboi T, Okumura-Noji K, Iwamoto N, and Yokoyama S (2016) Caveolin-1 facilitates internalization and degradation of ABCA1 and prooxidative products interfere with this reaction to increase HDL biogenesis. *Atherosclerosis* **253**:54–60.
- Lv YC, Tang YY, Peng J, Zhao GJ, Yang J, Yao F, Ouyang XP, He PP, Xie W, Tan YL, et al. (2014) MicroRNA-19b promotes macrophage cholesterol accumulation and aortic atherosclerosis by targeting ATP-binding cassette transporter A1. *Atherosclerosis* **236**:215–226.
- Lv YC, Yang J, Yao F, Xie W, Tang YY, Ouyang XP, He PP, Tan YL, Li L, Zhang M, et al. (2015) Diosgenin inhibits atherosclerosis via suppressing the MiR-19b-induced downregulation of ATP-binding cassette transporter A1. *Atherosclerosis* **240**:80–89.
- Ma AZS, Song ZY, and Zhang Q (2014) Cholesterol efflux is LXR α isoform-dependent in human macrophages. *BMC Cardiovasc Disord* **14**:80.
- Ma KL, Ruan XZ, Powis SH, Moorhead JF, and Varghese Z (2007) Anti-atherosclerotic effects of sirolimus on human vascular smooth muscle cells. *Am J Physiol Heart Circ Physiol* **292**:H2721–H2728.
- Ma X, Hu YW, Mo ZC, Li XX, Liu XH, Xiao J, Yin WD, Liao DF, and Tang CK (2009) NO-1886 up-regulates Niemann-Pick C1 protein (NPC1) expression through liver X receptor alpha signaling pathway in THP-1 macrophage-derived foam cells. *Cardiovasc Drugs Ther* **23**:199–206.
- Ma X, Li SF, Qin ZS, Ye J, Zhao ZL, Fang HH, Yao ZW, Gu MN, and Hu YW (2015) Propofol up-regulates expression of ABCA1, ABCG1, and SR-B1 through the

- PPAR γ /LXR α signaling pathway in THP-1 macrophage-derived foam cells. *Cardiovasc Pathol* **24**:230–235.
- Machado JT, Iborra RT, Fusco FB, Castilho G, Pinto RS, Machado-Lima A, Nakanakare ER, Seguro AC, Shimizu MH, Catanozi S, et al. (2014) N-acetylcysteine prevents endoplasmic reticulum stress elicited in macrophages by serum albumin drawn from chronic kidney disease rats and selectively affects lipid transporters, ABCA-1 and ABCG-1. *Atherosclerosis* **237**:343–352.
- Maehjima T, Sugano T, Yamazaki H, Yoshinaka Y, Doi T, Tanabe S, and Nishimaki-Mogami T (2011) Pitavastatin increases ABCA1 expression by dual mechanisms: SREBP2-driven transcriptional activation and PPAR α -dependent protein stabilization without activating LXR in rat hepatoma McA-RH7777 cells. *J Pharmacol Sci* **116**:107–115.
- Maiolino G, Rossitto G, Caielli P, Bisogni V, Rossi GP, and Calò LA (2013) The role of oxidized low-density lipoproteins in atherosclerosis: the myths and the facts. *Mediators Inflamm* **2013**:714653.
- Mandolini C, Santovito D, Marcantonio P, Buttitta F, Bucci M, Uchino S, Mezzetti A, and Cipollone F (2015) Identification of microRNAs 758 and 33b as potential modulators of ABCA1 expression in human atherosclerotic plaques. *Nutr Metab Cardiovasc Dis* **25**:202–209.
- Manna PR, Sennoune SR, Martinez-Zaguilan R, Slominski AT, and Pruitt K (2015) Regulation of retinoid mediated cholesterol efflux involves liver X receptor activation in mouse macrophages. *Biochem Biophys Res Commun* **464**:312–317.
- Mao M, Lei H, Liu Q, Chen Y, Zhao L, Li Q, Luo S, Zuo Z, He Q, Huang W, et al. (2014) Effects of miR-33a-5p on ABCA1/G1-mediated cholesterol efflux under inflammatory stress in THP-1 macrophages. *PLoS One* **9**:e109722.
- Marinozzi M, Castro Navas FF, Maggioni D, Carosati E, Bocci G, Carloncelli M, Giorgi G, Cruciani G, Fontana R, and Russo V (2017) Side-chain modified ergosterol and stigmastanol derivatives as liver X receptor agonists. *J Med Chem* **60**:6548–6562.
- Matsumoto A, Mizukami H, Mizuno S, Umegaki K, Nishikawa J, Shudo K, Kagechika H, and Inoue M (2007) beta-Cryptoxanthin, a novel natural RAR ligand, induces ATP-binding cassette transporters in macrophages. *Biochem Pharmacol* **74**:256–264.
- Meade T, Zuhrie R, Cook C, and Cooper J (2002) Bezafibrate in men with lower extremity arterial disease: randomised controlled trial. *BMJ* **325**:1139.
- Meiler S, Baumer Y, Toulmin E, Seng K, and Boisvert WA (2015) MicroRNA 302a is a novel modulator of cholesterol homeostasis and atherosclerosis. *Arterioscler Thromb Vasc Biol* **35**:323–331.
- Méndez-González J, Sören-Castillo S, Calpe-Berdiel L, Rotllan N, Vázquez-Carrera M, Escalá-Gil JC, and Blanco-Vaca F (2010) Disodium ascorbyl phytostanol phosphate (FM-VP4), a modified phytostanol, is a highly active hypocholesterolaemic agent that affects the enterohepatic circulation of both cholesterol and bile acids in mice. *Br J Nutr* **103**:153–160.
- Miao B, Zondlo S, Gibbs S, Cromley D, Hosagrahara VP, Kirchgessner TG, Billheimer J, and Mukherjee R (2004) Raising HDL cholesterol without inducing hepatic steatosis and hypertriglyceridemia by a selective LXR modulator. *J Lipid Res* **45**:1410–1417.
- Moazzini H, Najafi A, and Khani M (2017) Identification of direct target genes of miR-7, miR-9, miR-96, and miR-182 in the human breast cancer cell lines MCF-7 and MDA-MB-231. *Mol Cell Probes* **34**:45–52.
- Mogilenko DA, Shavva VS, Dizhe EB, Orlov SV, and Perevozchikov AP (2010) PPAR γ activates ABCA1 gene transcription but reduces the level of ABCA1 protein in HepG2 cells. *Biochem Biophys Res Commun* **402**:477–482.
- Mohammadpour AH and Akhlaghi F (2013) Future of cholesterol ester transfer protein (CETP) inhibitors: a pharmacological perspective. *Clin Pharmacokinet* **52**:615–626.
- Monzel JV, Budde T, Meyer Zu Schwabedissen HE, Schwebe M, Bien-Möller S, Lütjohann D, Kroemer HK, Jedlitschky G, and Grube M (2017) Doxorubicin enhances oxysterol levels resulting in a LXR-mediated upregulation of cardiac cholesterol transporters. *Biochem Pharmacol* **144**:108–119.
- Moore KJ, Rayner KJ, Suárez Y, and Fernández-Hernando C (2010) microRNAs and cholesterol metabolism. *Trends Endocrinol Metab* **21**:699–706.
- Moore KJ, Rayner KJ, Suárez Y, and Fernández-Hernando C (2011) The role of microRNAs in cholesterol efflux and hepatic lipid metabolism. *Annu Rev Nutr* **31**:49–63.
- Morgado N, Rigotti A, and Valenzuela A (2005) Comparative effect of fish oil feeding and other dietary fatty acids on plasma lipoproteins, biliary lipids, and hepatic expression of proteins involved in reverse cholesterol transport in the rat. *Ann Nutr Metab* **49**:397–406.
- Morrow MP, Grant A, Mujawar Z, Dubrovsky L, Pushkarsky T, Kiselyeva Y, Jennelle L, Mukhamedova N, Remaley AT, Kashanchi F, et al. (2010) Stimulation of the liver X receptor pathway inhibits HIV-1 replication via induction of ATP-binding cassette transporter A1. *Mol Pharmacol* **78**:215–225.
- Mostafa AM, Hamdy NM, Abdel-Rahman SZ, and El-Mesallamy HO (2016) Effect of lumbalaptin and pravastatin combination on cholesterol efflux in adipocytes. *IUBMB Life* **68**:535–543.
- Mostafa AM, Hamdy NM, El-Mesallamy HO, and Abdel-Rahman SZ (2015) Glucagon-like peptide 1 (GLP-1)-based therapy upregulates LXR-ABCA1/ABCG1 cascade in adipocytes. *Biochem Biophys Res Commun* **468**:900–905.
- Mulay V, Wood P, Manetsch M, Darabi M, Cairns R, Hoque M, Chan KC, Reverter M, Alvarez-Guaita A, Rye KA, et al. (2013) Inhibition of mitogen-activated protein kinase Erk1/2 promotes protein degradation of ATP binding cassette transporters A1 and G1 in CHO and HuH7 cells. *PLoS One* **8**:e62667.
- Mulligan JD, Flowers MT, Tebon A, Bitgood JJ, Wellington C, Hayden MR, and Attie AD (2003) ABCA1 is essential for efficient basolateral cholesterol efflux during the absorption of dietary cholesterol in chickens. *J Biol Chem* **278**:13356–13366.
- Muio DM, MacLean PS, Lang DB, Li S, Houmad JA, Way JM, Winegar DA, Corton JC, Dohm GL, and Kraus WE (2002) Fatty acid homeostasis and induction of lipid regulatory genes in skeletal muscles of peroxisome proliferator-activated receptor (PPAR) alpha knock-out mice: evidence for compensatory regulation by PPAR delta. *J Biol Chem* **277**:26089–26097.
- Murphy AJ, Bijl N, Yvan-Charvet L, Welch CB, Bhagwat N, Reheman A, Wang Y, Shaw JA, Levine RL, Ni H, et al. (2013) Cholesterol efflux in megakaryocyte progenitors suppresses platelet production and thrombocytosis. *Nat Med* **19**:586–594.
- Murthy S, Born E, Mathur SN, and Field FJ (2002) LXR/RXR activation enhances basolateral efflux of cholesterol in CaCo-2 cells. *J Lipid Res* **43**:1054–1064.
- Naik SU, Wang X, Da Silva JS, Jaye M, Macphree CH, Reilly MP, Billheimer JT, Rothblat GH, and Rader DJ (2006) Pharmacological activation of liver X receptors promotes reverse cholesterol transport in vivo. *Circulation* **113**:90–97.
- Nakamura K, Kennedy MA, Baldán A, Bojanic DD, Lyons K, and Edwards PA (2004) Expression and regulation of multiple murine ATP-binding cassette transporter G1 mRNAs/isoforms that stimulate cellular cholesterol uptake to high density lipoprotein. *J Biol Chem* **279**:45980–45989.
- Nakaya K, Ayaori M, Hisada T, Sawada S, Tanaka N, Iwamoto N, Ogura M, Yakushiji E, Kusuhara M, Nakamura H, et al. (2007) Telmisartan enhances cholesterol efflux from THP-1 macrophages by activating PPARgamma. *J Atheroscler Thromb* **14**:133–141.
- Nakaya K, Ayaori M, Uto-Kondo H, Hisada T, Ogura M, Yakushiji E, Takiguchi S, Terao Y, Ozasa H, Sasaki M, et al. (2010) Cilostazol enhances macrophage reverse cholesterol transport in vitro and in vivo. *Atherosclerosis* **213**:135–141.
- Nakaya K, Tohyama J, Naik SU, Tanigawa H, MacPhree C, Billheimer JT, and Rader DJ (2011) Peroxisome proliferator-activated receptor- α activation promotes macrophage reverse cholesterol transport through a liver X receptor-dependent pathway. *Arterioscler Thromb Vasc Biol* **31**:1276–1282.
- Nayyar G, Garber DW, Palgunachari MN, Monroe CE, Keenun TD, Handattu SP, Mishra VK, and Anantharamaiah GM (2012) Apolipoprotein E mimetic is more effective than apolipoprotein A-I mimetic in reducing lesion formation in older female apo E null mice. *Atherosclerosis* **224**:326–331.
- Neufeld EB, O'Brien K, Walts AD, Stonik JA, Demosky SJ, Malide D, Combs CA, and Remaley AT (2014) Cellular localization and trafficking of the human ABCG1 transporter. *Biology (Basel)* **3**:781–800.
- Neufeld EB, Remaley AT, Demosky SJ, Stonik JA, Cooney AM, Comly M, Dwyer NK, Zhang M, Blanchette-Mackie J, Santamarina-Fojo S, et al. (2001) Cellular localization and trafficking of the human ABCA1 transporter. *J Biol Chem* **276**:27584–27590.
- Nicholls SJ, Ray KK, Ballantyne CM, Beacham LA, Miller DL, Ruotolo G, Nissen SE, and Riesmeyer JS; ACCENTUATE Investigators (2017) Comparative effects of cholesterol ester transfer protein inhibition, statin or ezetimibe on lipid factors: the ACCENTUATE trial. *Atherosclerosis* **261**:12–18.
- Nicholls SJ, Ruotolo G, Brewer HB, Kane JP, Wang M-D, Krueger KA, Adelman SJ, Nissen SE, and Rader DJ (2015) Cholesterol efflux capacity and pre-beta-1 HDL concentrations are increased in dyslipidemic patients treated with evacetrapib. *J Am Coll Cardiol* **66**:2201–2210.
- Niesor EJ, Magg C, Ogawa N, Okamoto H, von der Mark E, Matile H, Schmid G, Clerc RG, Chapat E, Blum-Kaelin D, et al. (2010) Modulating cholesterol ester transfer protein activity maintains efficient pre-B-HDL formation and increases reverse cholesterol transport. *J Lipid Res* **51**:3443–3454.
- Nishimaki-Mogami T, Tamehiro N, Sato Y, Okuhira K, Sai K, Kagechika H, Shudo K, Abe-Dohmae S, Yokoyama S, Ohno Y, et al. (2008) The RXR agonists PA024 and HX630 have different abilities to activate LXR/RXR and to induce ABCA1 expression in macrophage cell lines. *Biochem Pharmacol* **76**:1006–1013.
- Ogata M, Tsujita M, Hossain MA, Akita N, Gonzalez FJ, Staels B, Suzuki S, Fukutomi T, Kimura G, and Yokoyama S (2009) On the mechanism for PPAR agonists to enhance ABCA1 gene expression. *Atherosclerosis* **205**:413–419.
- Ohama T, Hirano K, Zhang Z, Aoki R, Tsujii K, Nakagawa-Toyama Y, Tsukamoto K, Ikegami C, Matsuyama A, Ishigami M, et al. (2012) Dominant expression of ATP-binding cassette transporter-1 on basolateral surface of Caco-2 cells stimulated by LXR/RXR ligands. *Biochem Biophys Res Commun* **296**:625–630.
- Oliver WR Jr, Shenk JL, Snaith MR, Russell CS, Plunket KD, Bodkin NL, Lewis MC, Winegar DA, Sznaidman ML, Lambert MH, et al. (2001) A selective peroxisome proliferator-activated receptor delta agonist promotes reverse cholesterol transport. *Proc Natl Acad Sci USA* **98**:5306–5311.
- Ono K (2012) Current concept of reverse cholesterol transport and novel strategy for atheroprotection. *J Cardiol* **60**:339–343.
- Ono K (2016) Functions of microRNA-33a/b and microRNA therapeutics. *J Cardiol* **67**:28–33.
- Oram JF (2000) Tangier disease and ABCA1. *Biochim Biophys Acta* **1529**:321–330.
- Oram JF (2003) HDL apolipoproteins and ABCA1: partners in the removal of excess cellular cholesterol. *Arterioscler Thromb Vasc Biol* **23**:720–727.
- Osei-Hwedie DO, Amar M, Sviridov D, and Remaley AT (2011) Apolipoprotein mimetic peptides: mechanisms of action as anti-atherogenic agents. *Pharmacol Ther* **130**:83–91.
- Ozasa H, Ayaori M, Iizuka M, Terao Y, Uto-Kondo H, Yakushiji E, Takiguchi S, Nakaya K, Hisada T, Uehara Y, et al. (2011) Pioglitazone enhances cholesterol efflux from macrophages by increasing ABCA1/ABCG1 expressions via PPAR γ /LXR α pathway: findings from in vitro and ex vivo studies. *Atherosclerosis* **219**:141–150.
- Panzenboeck U, Kratzer I, Sovic A, Wintersperger A, Bernhart E, Hammer A, Malle E, and Sattler W (2006) Regulatory effects of synthetic liver X receptor- and peroxisome-proliferator activated receptor agonists on sterol transport pathways in polarized cerebrovascular endothelial cells. *Int J Biochem Cell Biol* **38**:1314–1329.
- Patel SB, Graf GA, and Temel RE (2018) ABCG5 and ABCG8: more than a defense against xenosterols. *J Lipid Res* **59**:1103–1113.
- Pedersen TR, Kjekshus J, Berg K, Hagfeldt T, Faergeman O, Faergeman G, Pyörälä K, Miettinen T, Wilhelmsen L, Olsson AG, et al.; Scandinavian Simvastatin Survival Study Group (2004) Randomised trial of cholesterol lowering in 4444 patients with coronary heart disease: the Scandinavian Simvastatin Survival Study (4S). *1994. Atheroscler Suppl* **5**:81–87.

- Phillips MC (2014) Molecular mechanisms of cellular cholesterol efflux. *J Biol Chem* **289**:24020–24029.
- Phillips MC (2018) Is ABCA1 a lipid transfer protein? *J Lipid Res* **59**:749–763.
- Piepoli MF, Hoes AW, Agewall S, Albus C, Brotons C, Catapano AL, Cooney MT, Corrà U, Cosyns B, Deaton C, et al.; Authors/Task Force Members Additional Contributor: Simone Binno (Italy) Document Reviewers (2016) 2016 European Guidelines on cardiovascular disease prevention in clinical practice: the Sixth Joint Task Force of the European Society of Cardiology and Other Societies on Cardiovascular Disease Prevention in Clinical Practice (constituted by representatives of 10 societies and by invited experts): developed with the special contribution of the European Association for Cardiovascular Prevention & Rehabilitation (EACPR). *Eur J Prev Cardiol* **23**:NP1–NP96.
- Pou J, Rebollo A, Roglans N, Sánchez RM, Vázquez-Carrera M, Laguna JC, Pedro-Botet J, and Alegret M (2008) Ritonavir increases CD36, ABCA1 and CYP27 expression in THP-1 macrophages. *Exp Biol Med (Maywood)* **233**: 1572–1582.
- Puigserver P, Vázquez F, Bonet ML, Picó C, and Palou A (1996) In vitro and in vivo induction of brown adipocyte uncoupling protein (thermogenin) by retinoic acid. *Biochem J* **317**:827–833.
- Qian L, Ma L, Wu G, Yu Q, Lin H, Ying Q, Wen D, and Gao C (2017) G004, a synthetic sulfonylurea compound, exerts anti-atherosclerosis effects by targeting SIRT1 in ApoE^{-/-} mice. *Vasc Pharmacol* **89**:49–57.
- Quinet EM, Basso MD, Halpern AR, Yates DW, Steffan RJ, Clerin V, Resmini C, Keith JC, Berrodrin TJ, Feingold I, et al. (2009) LXR ligand lowers LDL cholesterol in primates, is lipid neutral in hamster, and reduces atherosclerosis in mouse. *J Lipid Res* **50**:2358–2370.
- Quinet EM, Savio DA, Halpern AR, Chen L, Miller CP, and Nambi P (2004) Gene-selective modulation by a synthetic oxysterol ligand of the liver X receptor. *J Lipid Res* **45**:1929–1942.
- Quinet EM, Savio DA, Halpern AR, Chen L, Schuster GU, Gustafsson J-A, Basso MD, and Nambi P (2006) Liver X receptor (LXR)-beta regulation in LXRalpha-deficient mice: implications for therapeutic targeting. *Mol Pharmacol* **70**:1340–1349.
- Rakshandehroo M, Knoch B, Müller M, and Kersten S (2010) Peroxisome proliferator-activated receptor alpha target genes. *PPAR Res* **2010**:612089.
- Ramasamy I (2016) Update on the molecular biology of dyslipidemias. *Clin Chim Acta* **454**:143–185.
- Ramírez CM, Dávalos A, Goedeke L, Salerno AG, Warrier N, Cirera-Salinas D, Suárez Y, and Fernández-Hernando C (2011) MicroRNA-758 regulates cholesterol efflux through posttranscriptional repression of ATP-binding cassette transporter A1. *Arterioscler Thromb Vasc Biol* **31**:2707–2714.
- Ramírez CM, Rotllan N, Vlassov AV, Dávalos A, Li M, Goedeke L, Aranda JF, Cirera-Salinas D, Araldi E, Salerno A, et al. (2013) Control of cholesterol metabolism and plasma high-density lipoprotein levels by microRNA-144. *Circ Res* **112**:1592–1601.
- Rayner KJ, Fernandez-Hernando C, and Moore KJ (2012) MicroRNAs regulating lipid metabolism in atherogenesis. *Thromb Haemost* **107**:642–647.
- Rayner KJ and Moore KJ (2014) MicroRNA control of high-density lipoprotein metabolism and function. *Circ Res* **114**:183–192.
- Rayner KJ, Sheedy FJ, Esau CC, Hussain FN, Temel RE, Parathath S, van Gils JM, Rayner AJ, Chang AN, Suarez Y, et al. (2011) Antagonism of miR-33 in mice promotes reverse cholesterol transport and regression of atherosclerosis. *J Clin Invest* **121**:2921–2931.
- Reiss AB, Carsons SE, Anwar K, Rao S, Edelman SD, Zhang H, Fernandez P, Cronstein BN, and Chan ES (2008) Atheroprotective effects of methotrexate on reverse cholesterol transport proteins and foam cell transformation in human THP-1 monocyte/macrophages. *Arthritis Rheum* **58**:3675–3683.
- Reiss AB and Cronstein BN (2012) Regulation of foam cells by adenosine. *Arterioscler Thromb Vasc Biol* **32**:879–886.
- Remaley AT (2007) Old drug, new tricks: the unexpected effect of doxazosin on high-density lipoprotein. *Circ Res* **101**:116–118.
- Repa JJ, Turley SD, Lobaccaro JA, Medina J, Li L, Lustig K, Shan B, Heyman RA, Dietschy JM, and Mangelsdorf DJ (2000) Regulation of absorption and ABC1-mediated efflux of cholesterol by RXR heterodimers. *Science* **289**:1524–1529.
- Ridker PM (2014) LDL cholesterol: controversies and future therapeutic directions. *Lancet* **384**:607–617.
- Rinne P, Rami M, Nuutinen S, Santovito D, van der Vorst EPC, Guilleumat-Prats R, Lyytikäinen LP, Raitoharju E, Oksala N, Ring L, et al. (2017) Melanocortin 1 receptor signaling regulates cholesterol transport in macrophages. *Circulation* **136**: 83–97.
- Roglans N, Vázquez-Carrera M, Alegret M, Novell F, Zambón D, Ros E, Laguna JC, and Sánchez RM (2004) Fibrates modify the expression of key factors involved in bile-acid synthesis and biliary-lipid secretion in gallstone patients. *Eur J Clin Pharmacol* **59**:855–861.
- Rongen GA, van Ingen I, Kok M, Vonkeman H, Janssen M, and Jansen TL (2018) Vasodilator function worsens after cessation of tumour necrosis factor inhibitor therapy in patients with rheumatoid arthritis only if a flare occurs. *Clin Rheumatol* **37**:909–916.
- Rosenson RS, Brewer HB Jr, Davidson WS, Fayad ZA, Fuster V, Goldstein J, Hellerstein M, Jiang XC, Phillips MC, Rader DJ, et al. (2012) Cholesterol efflux and atheroprotection: advancing the concept of reverse cholesterol transport. *Circulation* **125**:1905–1919.
- Rotllan N and Fernández-Hernando C (2012) MicroRNA regulation of cholesterol metabolism. *Cholesterol* **2012**:847849.
- Rotllan N, Llaverias G, Julve J, Jauhainen M, Calpe-Berdiel L, Hernández C, Simó R, Blanco-Vaca F, and Escolà-Gil JC (2011) Differential effects of gemfibrozil and fenofibrate on reverse cholesterol transport from macrophages to feces in vivo. *Biochim Biophys Acta* **1811**:104–110.
- Rotllan N, Price N, Pati P, Goedeke L, and Fernández-Hernando C (2016) microRNAs in lipoprotein metabolism and cardiometabolic disorders. *Atherosclerosis* **246**: 352–360.
- Ruan XZ, Moorhead JF, Fernando R, Wheeler DC, Powis SH, and Varghese Z (2003) PPAR agonists protect mesangial cells from interleukin 1beta-induced intracellular lipid accumulation by activating the ABCA1 cholesterol efflux pathway. *J Am Soc Nephrol* **14**:593–600.
- Rubic T, Trottmann M, and Lorenz RL (2004) Stimulation of CD36 and the key effector of reverse cholesterol transport ATP-binding cassette A1 in monocytoid cells by niacin. *Biochem Pharmacol* **67**:411–419.
- Rubins HB, Robins SJ, Collins D, Fye CL, Anderson JW, Elam MB, Faas FH, Linares E, Schaefer EJ, Schechtman G, et al.; Veterans Affairs High-Density Lipoprotein Cholesterol Intervention Trial Study Group (1999) Gemfibrozil for the secondary prevention of coronary heart disease in men with low levels of high-density lipoprotein cholesterol. *N Engl J Med* **341**:410–418.
- Rust S, Rosier M, Funke H, Real J, Amoura Z, Piette JC, Deleuze JF, Brewer HB, Duverger N, Denèfle P, et al. (1999) Tangier disease is caused by mutations in the gene encoding ATP-binding cassette transporter 1. *Nat Genet* **22**:352–355.
- Rye KA, Clay MA, and Barter PJ (1999) Remodelling of high density lipoproteins by plasma factors. *Atherosclerosis* **145**:227–238.
- Sabatine MS, Giugliano RP, Keech AC, Honarpour N, Wiviott SD, Murphy SA, Kuder JF, Wang H, Liu T, Wasserman SM, et al.; FOURIER Steering Committee and Investigators (2017) Evolocumab and clinical outcomes in patients with cardiovascular disease. *N Engl J Med* **376**:1713–1722.
- Sacks FM, Pfeffer MA, Moye LA, Rouleau JL, Rutherford JD, Cole TG, Brown L, Warnica JW, Arnold JM, Wun CC, et al. (1996) The effect of pravastatin on coronary events after myocardial infarction in patients with average cholesterol levels: Cholesterol and Recurrent Events Trial investigators. *N Engl J Med* **335**: 1001–1009.
- Sakr SW, Williams DL, Stoudt GW, Phillips MC, and Rothblat GH (1999) Induction of cellular cholesterol efflux to lipid-free apolipoprotein A-I by cAMP. *Biochim Biophys Acta* **1438**:85–98.
- Sala F, Aranda JF, Rotllan N, Ramírez CM, Aryal B, Elia L, Condorelli G, Catapano AL, Fernández-Hernando C, and Norata GD (2014) MiR-143/145 deficiency attenuates the progression of atherosclerosis in Ldlr^{-/-} mice. *Thromb Haemost* **112**:796–802.
- Sankaranarayanan S, Oram JF, Asztalos BF, Vaughan AM, Lund-Katz S, Adorni MP, Phillips MC, and Rothblat GH (2009) Effects of acceptor composition and mechanism of ABCG1-mediated cellular free cholesterol efflux. *J Lipid Res* **50**:275–284.
- Santamarina-Fojo S, Peterson K, Knapper C, Qiu Y, Freeman L, Cheng JF, Osorio J, Remaley A, Yang XP, Haudenschild C, et al. (2000) Complete genomic sequence of the human ABCA1 gene: analysis of the human and mouse ATP-binding cassette A promoter. *Proc Natl Acad Sci USA* **97**:7987–7992.
- Sato M, Kawata Y, Erami K, Ikeda I, and Imaizumi K (2008) LXR agonist increases the lymph HDL transport in rats by promoting reciprocally intestinal ABCA1 and apo A-I mRNA levels. *Lipids* **43**:125–131.
- Schirris TJ, Renkema GH, Ritschel T, Voermans NC, Bilos A, van Engelen BG, Brandt U, Koopman WJ, Beyrath JD, Rodenburg RJ, et al. (2015a) Statin-induced myopathy is associated with mitochondrial complex III inhibition. *Cell Metab* **22**: 399–407.
- Schirris TJ, Ritschel T, Bilos A, Smeitink JA, and Russel FG (2015b) Statin lactonization by uridine 5'-diphospho-glucuronosyltransferases (UGTs). *Mol Pharm* **12**: 4048–4055.
- See RH, Caday-Malcolm RA, Singaraja RR, Zhou S, Silverston A, Huber MT, Moran J, James ER, Janoo R, Savill JM, et al. (2002) Protein kinase A site-specific phosphorylation regulates ATP-binding cassette A1 (ABCA1)-mediated phospholipid efflux. *J Biol Chem* **277**:41835–41842.
- Sharifov OF, Nayyar G, Garber DW, Handattu SP, Mishra VK, Goldberg D, Anantharamaiah GM, and Gupta H (2011) Apolipoprotein E mimetics and cholesterol-lowering properties. *Am J Cardiovasc Drugs* **11**:371–381.
- Shen L, Peng H, Peng R, Fan Q, Zhao S, Xu D, Morisseau C, Chiamvimonvat N, and Hammock BD (2015) Inhibition of soluble epoxide hydrolase in mice promotes reverse cholesterol transport and regression of atherosclerosis. *Atherosclerosis* **239**: 557–565.
- Shen L, Peng H, Zhao S, and Xu D (2014) A potent soluble epoxide hydrolase inhibitor, t-AUCB, modulates cholesterol balance and oxidized low density lipoprotein metabolism in adipocytes in vitro. *Biol Chem* **395**:443–451.
- Sherman CB, Peterson SJ, and Frishman WH (2010) Apolipoprotein A-I mimetic peptides: a potential new therapy for the prevention of atherosclerosis. *Cardiol Rev* **18**:141–147.
- Shimizu T, Miura S, Tanigawa H, Kuwano T, Zhang B, Uehara Y, and Saku K (2014) Rosuvastatin activates ATP-binding cassette transporter A1-dependent efflux ex vivo and promotes reverse cholesterol transport in macrophage cells in mice fed a high-fat diet. *Arterioscler Thromb Vasc Biol* **34**:2246–2253.
- Silva JC, de Oliveira EM, Turato WM, Trossini GHG, Maltarollo VG, Pitta MGR, Pitta IR, de Las Heras B, Bosca L, Rudnicki M, et al. (2018) GQ-11: a new PPAR agonist improves obesity-induced metabolic alterations in LDLr^{-/-} mice. *Int J Obes* **42**:1062–1072.
- Singaraja RR, Brunham LR, Visscher H, Kastelein JJ, and Hayden MR (2003) Efflux and atherosclerosis: the clinical and biochemical impact of variations in the ABCA1 gene. *Arterioscler Thromb Vasc Biol* **23**:1322–1332.
- Siripurkpong P and Na-Bangchang K (2009) Effects of niacin and chromium on the expression of ATP-binding cassette transporter A1 and apolipoprotein A-1 genes in HepG2 cells. *J Nutr Biochem* **20**:261–268.
- Small DM (2003) Role of ABC transporters in secretion of cholesterol from liver into bile. *Proc Natl Acad Sci USA* **100**:4–6.
- Solanes G, Pedraza N, Iglesias R, Giral M, and Villarroya F (2000) The human uncoupling protein-3 gene promoter requires MyoD and is induced by retinoic acid in muscle cells. *FASEB J* **14**:2141–2143.
- Song G, Liu J, Zhao Z, Yu Y, Tian H, Yao S, Li G, and Qin S (2011) Simvastatin reduces atherogenesis and promotes the expression of hepatic genes associated with reverse cholesterol transport in apoE-knockout mice fed high-fat diet. *Lipids Health Dis* **10**:8.
- Sparrow CP, Baffic J, Lam MH, Lund EG, Adams AD, Fu X, Hayes N, Jones AB, Macnaul KL, Ondeyka J, et al. (2002) A potent synthetic LXR agonist is more effective than cholesterol loading at inducing ABCA1 mRNA and stimulating cholesterol efflux. *J Biol Chem* **277**:10021–10027.

- Sprecher DL, Massien C, Pearce G, Billin AN, Perlstein I, Willson TM, Hassall DG, Ancellin N, Patterson SD, Lobe DC, et al. (2007) Triglyceride:high-density lipoprotein cholesterol effects in healthy subjects administered a peroxisome proliferator activated receptor delta agonist. *Arterioscler Thromb Vasc Biol* **27**:359–365.
- Steiner G (2005) Fibrates and coronary risk reduction. *Atherosclerosis* **182**:199–207.
- Stoekenbroek RM, Stroes ES, and Hovingh GK (2015) ApoA-I mimetics, in *High Density Lipoproteins: From Biological Understanding to Clinical Exploitation* (von Eckardstein A and Kardassis D eds) pp 631–648, Springer International Publishing, Cham, Switzerland.
- Strobach D and Lorenz RL (2003) The bisphosphonate ibandronate stimulates reverse cholesterol transport out of monocyte-derived cells by enhanced ABCA1 transcription. *Biochem Biophys Res Commun* **307**:23–30.
- Stroes ES, Thompson PD, Corsini A, Vladutiu GD, Raal FJ, Ray KK, Roden M, Stein E, Tokgözoğlu L, Nordestgaard BG, et al.; European Atherosclerosis Society Consensus Panel (2015) Statin-associated muscle symptoms: impact on statin therapy. European Atherosclerosis Society Consensus Panel Statement on Assessment, Aetiology and Management. *Eur Heart J* **36**:1012–1022.
- Sun D, Zhang J, Xie J, Wei W, Chen M, and Zhao X (2012) MiR-26 controls LXR-dependent cholesterol efflux by targeting ABCA1 and ARL7. *FEBS Lett* **586**:1472–1479.
- Sun Y, Fan J, Zhu Z, Guo X, Zhou T, Duan W, and Shen X (2015) Small molecule TBTC as a new selective retinoid X receptor α agonist improves behavioral deficit in Alzheimer's disease model mice. *Eur J Pharmacol* **762**:202–213.
- Suzuki S, Nishimaki-Mogami T, Tamehiro N, Inoue K, Arakawa R, Abe-Dohmae S, Tanaka AR, Ueda K, and Yokoyama S (2004) Verapamil increases the apolipoprotein-mediated release of cellular cholesterol by induction of ABCA1 expression via liver X receptor-independent mechanism. *Arterioscler Thromb Vasc Biol* **24**:519–525.
- Tachibana M, Shinohara M, Yamazaki Y, Liu C-C, Rogers J, Bu G, and Kanekiyo T (2016) Rescuing effects of RXR agonist bexarotene on aging-related synapse loss depend on neuronal LRP1. *Exp Neurol* **277**:1–9.
- Takahashi Y and Smith JD (1999) Cholesterol efflux to apolipoprotein AI involves endocytosis and resecretion in a calcium-dependent pathway. *Proc Natl Acad Sci USA* **96**:11358–11363.
- Talbot CPJ, Plat J, Ritsch A, and Mensink RP (2018) Determinants of cholesterol efflux capacity in humans. *Prog Lipid Res* **69**:21–32.
- Tall AR and Rader DJ (2018) Trials and tribulations of CETP inhibitors. *Circ Res* **122**:106–112.
- Tall AR and Yvan-Charvet L (2015) Cholesterol, inflammation and innate immunity. *Nat Rev Immunol* **15**:104–116.
- Tall AR, Yvan-Charvet L, Terasaka N, Pagler T, and Wang N (2008) HDL, ABC transporters, and cholesterol efflux: implications for the treatment of atherosclerosis. *Cell Metab* **7**:365–375.
- Tamehiro N, Park MH, Hawthurst V, Nagpal K, Adams ME, Zannis VI, Golenbock DT, and Fitzgerald ML (2015) LXR agonism upregulates the macrophage ABCA1/syntaxin protein complex that can bind ApoA-I and stabilized ABCA1 protein, but complex loss does not inhibit lipid efflux. *Biochemistry* **54**:6931–6941.
- Tanabe J, Tamasawa N, Yamashita M, Matsuki K, Murakami H, Matsui J, Sugimoto K, Yasujima M, and Suda T (2008) Effects of combined PPARgamma and PPARalpha agonist therapy on reverse cholesterol transport in the Zucker diabetic fatty rat. *Diabetes Obes Metab* **10**:772–779.
- Tang C and Oram JF (2009) The cell cholesterol exporter ABCA1 as a protector from cardiovascular disease and diabetes. *Biochim Biophys Acta* **1791**:563–572.
- Tang H, Shi W, Fu S, Wang T, Zhai S, Song Y, and Han J (2018) Pioglitazone and bladder cancer risk: a systematic review and meta-analysis. *Coglitazone and Med* **7**:1070–1080.
- Tarling EJ and Edwards PA (2011) ATP binding cassette transporter G1 (ABCG1) is an intracellular sterol transporter. *Proc Natl Acad Sci USA* **108**:19719–19724.
- Tauscher A and Kuver R (2003) ABCG5 and ABCG8 are expressed in gallbladder epithelial cells. *Biochem Biophys Res Commun* **307**:1021–1028.
- Taylor F, Ward K, Moore TH, Burke M, Davey Smith G, Casas JP, and Ebrahim S (2011) Statins for the primary prevention of cardiovascular disease. *Cochrane Database Syst Rev* (1):CD004816.
- Teng JJ, Tsai MC, Shih SF, Tsuei BF, Chang H, Chuang YP, Lin CS, Chern CY, and Chen SJ (2018) Chalcone derivatives enhance ATP-binding cassette transporters A1 in human THP-1 macrophages. *Molecules* **23**.
- Terasaka N, Hiroshima A, Koieyama T, Ubukata N, Morikawa Y, Nakai D, and Inaba T (2003) T-0901317, a synthetic liver X receptor ligand, inhibits development of atherosclerosis in LDL receptor-deficient mice. *FEBS Lett* **536**:6–11.
- Thomas J, Bramlett KS, Montrose C, Foxworthy P, Eacho PI, McCann D, Cao G, Kiefer A, McCowan J, Yu KL, et al. (2003) A chemical switch regulates fibrate specificity for peroxisome proliferator-activated receptor alpha (PPARalpha) versus liver X receptor. *J Biol Chem* **278**:2403–2410.
- Tresguerres M, Levin LR, and Buck J (2011) Intracellular cAMP signaling by soluble adenylyl cyclase. *Kidney Int* **79**:1277–1288.
- Tsai JY, Su KH, Shyue SK, Kou YR, Yu YB, Hsiao SH, Chiang AN, Wu YL, Ching LC, and Lee TS (2010) EGB761 ameliorates the formation of foam cells by regulating the expression of SR-A and ABCA1: role of haem oxygenase-1. *Cardiovasc Res* **88**:415–423.
- Tsui CY, Chen CY, Zhao JF, Su KH, Lee HT, Lin SJ, Shyue SK, Hsiao SH, and Lee TS (2014) Activation of soluble guanylyl cyclase prevents foam cell formation and atherosclerosis. *Acta Physiol (Oxf)* **210**:799–810.
- Tsunemi A, Ueno T, Fukuda N, Watanabe T, Tahira K, Haketa A, Hatanaka Y, Tanaka S, Matsumoto T, Matsumoto Y, et al. (2014) A novel gene regulator, pyrrole-imidazole polyamide targeting ABCA1 gene increases cholesterol efflux from macrophages and plasma HDL concentration. *J Mol Med (Berl)* **92**:509–521.
- Uehara Y and Saku K (2014) High-density lipoprotein and atherosclerosis: roles of lipid transporters. *World J Cardiol* **6**:1049–1059.
- Uehara Y, Saku K, and Zhang B (2011) Tangier disease, in *Advances in the Study of Genetic Disorders* (Ikehara K ed) pp 239–254, Intech Open, London, United Kingdom.
- Valasek MA, Clarke SL, and Repa JJ (2007) Fenofibrate reduces intestinal cholesterol absorption via PPARalpha-dependent modulation of NPC1L1 expression in mouse. *J Lipid Res* **48**:2725–2735.
- van Rooij E and Kauppinen S (2014) Development of microRNA therapeutics is coming of age. *EMBO Mol Med* **6**:851–864.
- Vaughan AM and Oram JF (2003) ABCA1 redistributes membrane cholesterol independent of apolipoprotein interactions. *J Lipid Res* **44**:1373–1380.
- Vaughan AM and Oram JF (2005) ABCG1 redistributes cell cholesterol to domains removable by high density lipoprotein but not by lipid-depleted apolipoproteins. *J Biol Chem* **280**:30150–30157.
- Vedhachalam C, Duong PT, Nickel M, Nguyen D, Dhanasekaran P, Saito H, Rothblat GH, Lund-Katz S, and Phillips MC (2007a) Mechanism of ATP-binding cassette transporter A1-mediated cellular lipid efflux to apolipoprotein A-I and formation of high density lipoprotein particles. *J Biol Chem* **282**:25123–25130.
- Vedhachalam C, Ghering AB, Davidson WS, Lund-Katz S, Rothblat GH, and Phillips MC (2007b) ABCA1-induced cell surface binding sites for ApoA-I. *Arterioscler Thromb Vasc Biol* **27**:1603–1609.
- Verschuren L, de Vries-van der Weij J, Zadelaar S, Kleemann R, and Kooistra T (2009) LXR agonist suppresses atherosclerotic lesion growth and promotes lesion regression in apoE*3Leiden mice: time course and mechanisms. *J Lipid Res* **50**:301–311.
- Viñals M, Bermúdez I, Llaverias G, Alegret M, Sanchez RM, Vázquez-Carrera M, and Laguna JC (2005) Aspirin increases CD36, SR-BI, and ABCA1 expression in human THP-1 macrophages. *Circulation* **112**:141–149.
- Voight BF, Peloso GM, Orho-Melander M, Frikke-Schmidt R, Barbalic M, Jensen MK, Hindy G, Holm H, Ding EL, Johnson T, et al. (2012) Plasma HDL cholesterol and risk of myocardial infarction: a mendelian randomisation study. *Lancet* **380**:572–580.
- Voloshyna I, Carsons S, Littlefield MJ, Rieger JM, Figler R, and Reiss AB (2013) Adenosine A2A receptor activation supports an atheroprotective cholesterol balance in human macrophages and endothelial cells. *Biochim Biophys Acta* **1831**:407–416.
- Voloshyna I, Seshadri S, Anwar K, Littlefield MJ, Belilos E, Carsons SE, and Reiss AB (2014) Infliximab reverses suppression of cholesterol efflux proteins by TNF- α : a possible mechanism for modulation of atherogenesis. *BioMed Res Int* **2014**:312647.
- von Eckardstein A and Rohrer L (2009) Transendothelial lipoprotein transport and regulation of endothelial permeability and integrity by lipoproteins. *Curr Opin Lipidol* **20**:197–205.
- Wagschal A, Najafi-Shoushtari SH, Wang L, Goedeke L, Sinha S, deLemos AS, Black JC, Ramirez CM, Li Y, Tewhey R, et al. (2015) Genome-wide identification of microRNAs regulating cholesterol and triglyceride homeostasis. *Nat Med* **21**:1290–1297.
- Wang D, Xia M, Yan X, Li D, Wang L, Xu Y, Jin T, and Ling W (2012) Gut microbiota metabolism of anthocyanin promotes reverse cholesterol transport in mice via repressing miRNA-10b. *Circ Res* **111**:967–981.
- Wang D, Yan X, Xia M, Yang Y, Li D, Li X, Song F, and Ling W (2014a) Coenzyme Q10 promotes macrophage cholesterol efflux by regulation of the activator protein-1/miR-378/ATP-binding cassette transporter G1-signaling pathway. *Arterioscler Thromb Vasc Biol* **34**:1860–1870.
- Wang J, Mitsche MA, Lütjohann D, Cohen JC, Xie XS, and Hobbs HH (2015a) Relative roles of ABCG5/ABCG8 in liver and intestine. *J Lipid Res* **56**:319–330.
- Wang J-M, Wang D, Tan Y-Y, Zhao G, and Ji ZL (2014b) 22(R)-hydroxycholesterol and pioglitazone synergistically decrease cholesterol ester via the PPAR γ -LXR-ABCA1 pathway in cholesteroles of the gallbladder. *Biochim Biophys Res Commun* **447**:152–157.
- Wang JM, Wang D, Tan YY, Zhao G, and Ji ZL (2015b) Pioglitazone reduces lipid droplets in cholesteroles of the gallbladder by increasing ABCA1 and NCEH1 expression. *Mol Cell Biochem* **399**:7–15.
- Wang L, Palme V, Rotter S, Schilcher N, Cukaj M, Wang D, Ladurner A, Heiss EH, Stangl H, Dirsch VM, et al. (2017) Piperine inhibits ABCA1 degradation and promotes cholesterol efflux from THP-1-derived macrophages. *Mol Nutr Food Res* **61**:1500960.
- Wang N, Lan D, Chen W, Matsuura F, and Tall AR (2004) ATP-binding cassette transporters G1 and G4 mediate cellular cholesterol efflux to high-density lipoproteins. *Proc Natl Acad Sci USA* **101**:9774–9779.
- Wang N, Ranalletta M, Matsuura F, Peng F, and Tall AR (2006) LXR-induced redistribution of ABCG1 to plasma membrane in macrophages enhances cholesterol mass efflux to HDL. *Arterioscler Thromb Vasc Biol* **26**:1310–1316.
- Wang N, Silver DL, Thiele C, and Tall AR (2001) ATP-binding cassette transporter A1 (ABCA1) functions as a cholesterol efflux regulatory protein. *J Biol Chem* **276**:23742–23747.
- Wang N, Yvan-Charvet L, Lütjohann D, Mulder M, Vanmierlo T, Kim TW, and Tall AR (2008) ATP-binding cassette transporters G1 and G4 mediate cholesterol and desmosterol efflux to HDL and regulate sterol accumulation in the brain. *FASEB J* **22**:1073–1082.
- Wang X, Luo J, Li N, Liu L, Han X, Liu C, Zuo X, Jiang X, Li Y, Xu Y, et al. (2018) E3317 promotes cholesterol efflux in macrophage cells via enhancing ABCA1 expression. *Biochem Biophys Res Commun* **504**:68–74.
- Wang Y, Moser AH, Shigenaga JK, Grunfeld C, and Feingold KR (2005) Down-regulation of liver X receptor-alpha in mouse kidney and HK-2 proximal tubular cells by LPS and cytokines. *J Lipid Res* **46**:2377–2387.
- Wang Y-H, Chen Y-F, Chen S-R, Chen X, Chen J-W, Shen X-Y, Mou Y-G, and Liu P-Q (2010) Aspirin increases apolipoprotein A-I-mediated cholesterol efflux via enhancing expression of ATP-binding cassette transporter A1. *Pharmacology* **86**:320–326.
- Westertorp M, Bochem AE, Yvan-Charvet L, Murphy AJ, Wang N, and Tall AR (2014) ATP-binding cassette transporters, atherosclerosis, and inflammation. *Circ Res* **114**:157–170.
- White CR, Garber DW, and Anantharamaiah GM (2014) Anti-inflammatory and cholesterol-reducing properties of apolipoprotein mimetics: a review. *J Lipid Res* **55**:2007–2021.

- WHO (2008) *Cardiovascular Disease*, World Health Organization, Geneva, Switzerland.
- WHO (2014) *Global Status Report on Noncommunicable Diseases 2014*, World Health Organization, Geneva, Switzerland.
- Wielinga PR, van der Heijden I, Reid G, Beijnen JH, Wijnholds J, and Borst P (2003) Characterization of the MRP4- and MRP5-mediated transport of cyclic nucleotides from intact cells. *J Biol Chem* **278**:17664–17671.
- Wilkinson MJ, Laffin LJ, and Davidson MH (2014) Overcoming toxicity and side-effects of lipid-lowering therapies. *Best Pract Res Clin Endocrinol Metab* **28**:439–452.
- Wissel G, Kudryavtsev P, Ghemti L, Tammela P, Wipf P, Yliperttula M, Finel M, Urtti A, Kidron H, and Xhaard H (2015) Exploring the structure-activity relationships of ABCG2 modulators using a screening approach. *Bioorg Med Chem* **23**:3513–3525.
- Wlodarczyk Z, Vitko S, Salmela K, Czajkowski Z, and Margreiter R; TERRA Study Group (2005) Lipid metabolism in renal transplant patients receiving tacrolimus/sirolimus combination therapy. *Transplant Proc* **37**:1871–1873.
- Wu J, Zhang Y, Wang N, Davis L, Yang G, Wang X, Zhu Y, Breyer MD, and Guan Y (2004) Liver X receptor- α mediates cholesterol efflux in glomerular mesangial cells. *Am J Physiol Renal Physiol* **287**:F886–F895.
- Wu Z-H and Zhao S-P (2009) Niacin promotes cholesterol efflux through stimulation of the PPAR γ -LXR α -ABCA1 pathway in 3T3-L1 adipocytes. *Pharmacology* **84**:282–287.
- Xie Q, Zhao SP, and Li F (2010) D-4F, an apolipoprotein A-I mimetic peptide, promotes cholesterol efflux from macrophages via ATP-binding cassette transporter A1. *Tohoku J Exp Med* **220**:223–228.
- Xu Y, Lai F, Xu Y, Wu Y, Liu Q, Li N, Wei Y, Feng T, Zheng Z, Jiang W, et al. (2011) Mycophenolic acid induces ATP-binding cassette transporter A1 (ABCA1) expression through the PPAR γ -LXR α -ABCA1 pathway. *Biochem Biophys Res Commun* **414**:779–782.
- Xu Y, Liu H, Liu M, Li F, Liu L, Du F, Fan D, and Yu H (2016) A human apolipoprotein E mimetic peptide reduces atherosclerosis in aged apolipoprotein E null mice. *Am J Transl Res* **8**:3482–3492.
- Xue XH, Shi FF, Chen T, Wei W, Zhou XM, and Chen LD (2016) Inhibition of ERK1/2 improves lipid balance in rat macrophages via ABCA1/G1 and CD36. *Mol Med Rep* **13**:1533–1540.
- Yakushiji E, Ayaori M, Nishida T, Shiotani K, Takiguchi S, Nakaya K, Uto-Kondo H, Ogura M, Sasaki M, Yogo M, et al. (2016) Probucol-oxidized products, spiroquinone and diphenylquinone, promote reverse cholesterol transport in mice. *Arterioscler Thromb Vasc Biol* **36**:591–597.
- Yan W, Zhang T, Cheng J, Zhou X, Qu X, and Hu H (2010) Liver X receptor agonist methyl- β -hydroxy- α , β -epoxycholesterol attenuates atherosclerosis in apolipoprotein E knockout mice without increasing plasma triglyceride. *Pharmacology* **86**:306–312.
- Yancey PG, Bortnick AE, Kellner-Weibel G, de la Llera-Moya M, Phillips MC, and Rothblat GH (2003) Importance of different pathways of cellular cholesterol efflux. *Arterioscler Thromb Vasc Biol* **23**:712–719.
- Yang Y, Lian YT, Huang SY, Yang Y, Cheng LX, and Liu K (2014) GABA and topiramate inhibit the formation of human macrophage-derived foam cells by modulating cholesterol-metabolism-associated molecules. *Cell Physiol Biochem* **33**:1117–1129.
- Yang Z, Cappello T, and Wang L (2015) Emerging role of microRNAs in lipid metabolism. *Acta Pharm Sin B* **5**:145–150.
- Yin QH, Zhang R, Li L, Wang YT, Liu JP, Zhang J, Bai L, Cheng JQ, Fu P, and Liu F (2016) Exendin-4 ameliorates lipotoxicity-induced glomerular endothelial cell injury by improving ABC transporter A1-mediated cholesterol efflux in diabetic apoE knockout mice. *J Biol Chem* **291**:26487–26501.
- Ying R, Yuan Y, Qin Y-F, Tian D, Feng L, Guo Z-G, Sun Y-X, and Li M-X (2013) The combination of L-4F and simvastatin stimulate cholesterol efflux and related proteins expressions to reduce atherosclerotic lesions in apoE knockout mice. *Lipids Health Dis* **12**:180.
- Yokoyama S (2004) ABCA1-mediated generation of HDL and its non-transcriptional regulation. *Seikagaku* **76**:532–538.
- Yokoyama S, Arakawa R, Wu CA, Iwamoto N, Lu R, Tsujita M, and Abe-Dohmae S (2012) Calpain-mediated ABCA1 degradation: post-translational regulation of ABCA1 for HDL biogenesis. *Biochim Biophys Acta* **1821**:547–551.
- Yu L, Gupta S, Xu F, Liverman AD, Moschetta A, Mangelsdorf DJ, Repa JJ, Hobbs HH, and Cohen JC (2005) Expression of ABCG5 and ABCG8 is required for regulation of biliary cholesterol secretion. *J Biol Chem* **280**:8742–8747.
- Yu L, Hammer RE, Li-Hawkins J, Von Bergmann K, Lutjohann D, Cohen JC, and Hobbs HH (2002) Disruption of Abcg5 and Abcg8 in mice reveals their crucial role in biliary cholesterol secretion. *Proc Natl Acad Sci USA* **99**:16237–16242.
- Yu XH, Qian K, Jiang N, Zheng XL, Cayabyab FS, and Tang CK (2014) ABCG5/ABCG8 in cholesterol excretion and atherosclerosis. *Clin Chim Acta* **428**:82–88.
- Yvan-Charvet L, Kling J, Pagler T, Li H, Hubbard B, Fisher T, Sparrow CP, Taggart AK, and Tall AR (2010a) Cholesterol efflux potential and antiinflammatory properties of high-density lipoprotein after treatment with niacin or anacetrapib. *Arterioscler Thromb Vasc Biol* **30**:1430–1438.
- Yvan-Charvet L, Matsuura F, Wang N, Bamberger MJ, Nguyen T, Rinninger F, Jiang X-C, Shear CL, and Tall AR (2007) Inhibition of cholesteryl ester transfer protein by torcetrapib modestly increases macrophage cholesterol efflux to HDL. *Arterioscler Thromb Vasc Biol* **27**:1132–1138.
- Yvan-Charvet L, Wang N, and Tall AR (2010b) Role of HDL, ABCA1, and ABCG1 transporters in cholesterol efflux and immune responses. *Arterioscler Thromb Vasc Biol* **30**:139–143.
- Zanlungo S, Rigotti A, and Nervi F (2004) Hepatic cholesterol transport from plasma into bile: implications for gallstone disease. *Curr Opin Lipidol* **15**:279–286.
- Zanotti I, Favari E, Sposito AC, Rothblat GH, and Bernini F (2004) Pitavastatin increases ABCA1-mediated lipid efflux from Fu5AH rat hepatoma cells. *Biochem Biophys Res Commun* **321**:670–674.
- Zanotti I, Poti F, Favari E, Steffensen KR, Gustafsson J-A, and Bernini F (2006) Pitavastatin effect on ATP binding cassette A1-mediated lipid efflux from macrophages: evidence for liver X receptor (LXR)-dependent and LXR-independent mechanisms of activation by cAMP. *J Pharmacol Exp Ther* **317**:395–401.
- Zanotti I, Poti F, Pedrelli M, Favari E, Moleri E, Franceschini G, Calabresi L, and Bernini F (2008) The LXR agonist T0901317 promotes the reverse cholesterol transport from macrophages by increasing plasma efflux potential. *J Lipid Res* **49**:954–960.
- Zhang C, Yin W, Liao D, Huang L, Tang C, Tsutsumi K, Wang Z, Liu Y, Li Q, Hou H, et al. (2006) NO-1886 upregulates ATP binding cassette transporter A1 and inhibits diet-induced atherosclerosis in Chinese Bama minipigs. *J Lipid Res* **47**:2055–2063.
- Zhang L, Chen Y, Yang X, Yang J, Cao X, Li X, Li L, Miao QR, Hajjar DP, Duan Y, et al. (2016) MEK1/2 inhibitors activate macrophage ABCG1 expression and reverse cholesterol transport-An anti-atherogenic function of ERK1/2 inhibition. *Biochim Biophys Acta* **1861**:1180–1191.
- Zhang L, Jiang M, Shui Y, Chen Y, Wang Q, Hu W, Ma X, Li X, Liu X, Cao X, et al. (2013a) DNA topoisomerase II inhibitors induce macrophage ABCA1 expression and cholesterol efflux: an LXR-dependent mechanism. *Biochim Biophys Acta* **1831**:1134–1145.
- Zhang L-H, Kamanna VS, Ganji SH, Xiong X-M, and Kashyap ML (2012) Niacin increases HDL biogenesis by enhancing DR4-dependent transcription of ABCA1 and lipidation of apolipoprotein A-I in HepG2 cells. *J Lipid Res* **53**:941–950.
- Zhang M, Wu JF, Chen WJ, Tang SL, Mo ZC, Tang YY, Li Y, Wang JL, Liu XY, Peng J, et al. (2014) MicroRNA-27a/b regulates cellular cholesterol efflux, influx and esterification/hydrolysis in THP-1 macrophages. *Atherosclerosis* **234**:54–64.
- Zhang N, Lei J, Lei H, Ruan X, Liu Q, Chen Y, and Huang W (2015a) MicroRNA-101 overexpression by IL-6 and TNF- α inhibits cholesterol efflux by suppressing ATP-binding cassette transporter A1 expression. *Exp Cell Res* **336**:33–42.
- Zhang Q, Ma AZS, Song ZY, Wang C, and Fu XD (2013b) Nifedipine enhances cholesterol efflux in RAW264.7 macrophages. *Cardiovasc Drugs Ther* **27**:425–431.
- Zhang R, Wang Y, Li R, and Chen G (2015b) Transcriptional factors mediating retinoic acid signals in the control of energy metabolism. *Int J Mol Sci* **16**:14210–14244.
- Zhang Y, Zanotti I, Reilly MP, Glick JM, Rothblat GH, and Rader DJ (2003) Overexpression of apolipoprotein A-I promotes reverse transport of cholesterol from macrophages to feces in vivo. *Circulation* **108**:661–663.
- Zhao C and Dahlman-Wright K (2010) Liver X receptor in cholesterol metabolism. *J Endocrinol* **204**:233–240.
- Zhao R, Feng J, and He G (2014) miR-613 regulates cholesterol efflux by targeting LXR α and ABCA1 in PPAR γ activated THP-1 macrophages. *Biochem Biophys Res Commun* **448**:329–334.
- Zhao W, Haller V, and Ritsch A (2015) The polyphenol PGG enhances expression of SR-BI and ABCA1 in J774 and THP-1 macrophages. *Atherosclerosis* **242**:611–617.
- Zhou W, Lin J, Chen H, Wang J, Liu Y, and Xia M (2015) Retinoic acid induces macrophage cholesterol efflux and inhibits atherosclerotic plaque formation in apoE-deficient mice. *Br J Nutr* **114**:509–518.
- Zhou X, Yin Z, Guo X, Hajjar DP, and Han J (2010) Inhibition of ERK1/2 and activation of liver X receptor synergistically induce macrophage ABCA1 expression and cholesterol efflux. *J Biol Chem* **285**:6316–6326.
- Zhu S and Liu JH (2015) Zerumbone, a natural cyclic sesquiterpene, promotes ABCA1-dependent cholesterol efflux from human THP-1 macrophages. *Pharmacology* **95**:258–263.

SUPPLEMENTARY DATA

Brothers in arms: ABCA1- and ABCG1-mediated cholesterol efflux as promising targets in cardiovascular disease treatment

Sanne J.C.M. Frambach, Ria de Haas, Jan A.M. Smeitink, Gerard.A. Rongen,

Frans G.M. Russel & Tom J.J. Schirris

Supplementary Table 1 | Search strategy EMBASE Overview of search strategy sets and search statements, as performed using the EMBASE database on 12-02-2019

Set	Search statement
1.	lipid metabolism/ or sterol metabolism/
2.	(fat metabolism or lipid metabolism or lipid mobilisation or lipid mobilization or lipid turnover or lipometabolism).ti,ab.
3.	(fat metabolism or lipid metabolism or lipid mobilisation or lipid mobilization or lipid turnover or lipometabolism).kw.
4.	or/1-3
5.	phospholipid/
6.	(phosphatides or phospholipids).ti,ab.
7.	(phosphatides or phospholipids).kw.
8.	or/5-7
9.	triacylglycerol/
10.	(triacylglycerol? or triglycerides).ti,ab.
11.	(triacylglycerol? or triglycerides).kw.
12.	or/9-11
13.	liver cell/
14.	(hepatic cell? or hepatocyte?).ti,ab.
15.	(hepatic cell? or hepatocyte?).kw.
16.	or/13-15
17.	liver/
18.	liver?.ti,ab.
19.	liver?.kw.
20.	or/17-19
21.	macrophages/ or foam cells/
22.	(bone marrow?derived macrophage? or macrophage? or monocyte?derived macrophage? or foam cell?).ti,ab.
23.	(bone marrow?derived macrophage? or macrophage? or monocyte?derived macrophage? or foam cell?).kw.
24.	or/21-23
25.	transcription factor Sp1/
26.	(sp1 transcription factor or specificity protein 1 transcription factor).ti,ab.
27.	(sp1 transcription factor or specificity protein 1 transcription factor).kw.
28.	or/25-27
29.	*cholesterol/ or *cholesterol derivative/ or *cholesterol ester/
30.	(cholesterol or epicholesterol or cholesterol esters or cholesteryl esters).ti.
31.	(cholesterol or epicholesterol or cholesterol esters or cholesteryl esters).kw.
32.	or/29-31
33.	*lipoprotein/ or *apolipoprotein/ or *high density lipoprotein/ or *high density lipoprotein 2/ or *high density lipoprotein 3/ or *Apolipoprotein A1/ or *Apolipoprotein E/ or *pre beta high density lipoprotein/
34.	(hdl lipoproteins or heavy lipoproteins or high?density lipoproteins or lipoproteins, hdl or alpha* lipoprotein? or hdl* cholesterol or high density lipoprotein cholesterol or alpha?lipoprotein cholesterol or hdl lipoprotein, pre?beta or hdl, pre-beta1 or hdl, pre-beta or high density lipoprotein?, pre?beta? or nascent hdl or hdl?2 or high density lipoprotein hdl2 or high density lipoprotein?2 or lipoprotein? hdl2 or hdl?3 or high density lipoprotein?3 or high density lipoprotein hdl3 or lipoprotein?, hdl3 or apolipoprotein? or circulating lipoproteins or lipoprotein? or apo-a or apo a or apolipoproteins a or apo a?i isoproteins or apo a-1 or apo?a?i or apo?a?1 or apolipoprotein a 1 or apolipoprotein a?i isoprotein* or apolipoprotein a i or apolipoprotein a?1 or apolipoprotein a-i or apolipoprotein ai propeptide or apolipoprotein ai or pro?apo a?i or pro apolipoprotein a i or pro?apolipoprotein a?i or apo e isoproteins or apo?e or apolipoprotein e isoproteins or apolipoproteins e or apoprotein ?e? or apoproteins e).ti.
35.	(hdl lipoproteins or heavy lipoproteins or high?density lipoproteins or lipoproteins, hdl or alpha* lipoprotein? or hdl* cholesterol or high density lipoprotein cholesterol or alpha?lipoprotein cholesterol or hdl lipoprotein, pre?beta or hdl, pre-beta1 or hdl, pre-beta or high density lipoprotein?, pre?beta? or nascent hdl or hdl?2 or high density lipoprotein hdl2 or high density lipoprotein?2 or lipoprotein? hdl2 or hdl?3 or high density lipoprotein?3 or high density lipoprotein hdl3 or lipoprotein?, hdl3 or apolipoprotein? or circulating lipoproteins or lipoprotein? or apo-a or apo a or apolipoproteins a or apo a?i isoproteins or apo a-1 or apo?a?i or apo?a?1 or apolipoprotein a 1 or apolipoprotein a?i isoprotein* or apolipoprotein a i or apolipoprotein a?1 or apolipoprotein a-i or apolipoprotein ai propeptide or apolipoprotein ai or pro?apo a?i or pro apolipoprotein a i or pro?apolipoprotein a?i or apo e isoproteins or apo?e or apolipoprotein e isoproteins or apolipoproteins e or apoprotein ?e? or apoproteins e).kw.

36.	or/33-35
37.	4 or 8 or 12 or 16 or 20 or 24 or 28 or 32 or 36
38.	protein transport/
39.	(protein transfer or protein uptake or transport, protein or cellular protein localization process* or protein targeting or protein trafficking? or protein translocation).ti,ab.
40.	(protein transfer or protein uptake or transport, protein or cellular protein localization process* or protein targeting or protein trafficking? or protein translocation).kw.
41.	or/38-40
42.	carrier protein/
43.	(biologic* pump? or (membrane adj3 transport*) or metabolic pump? or permease?).ti,ab.
44.	(biologic* pump? or (membrane adj3 transport*) or metabolic pump? or permease?).kw.
45.	or/42-44
46.	glycoprotein/
47.	(*glycoprotein?/ or (cholester?l ester adj3 protein?)).ti,ab.
48.	(*glycoprotein?/ or (cholester?l ester adj3 protein?)).kw.
49.	or/46-48
50.	scavenger receptor BI/
51.	(scavenger receptor B type 1 or scavenger receptor B1 or scavenger receptor class B type 1 or scavenger receptor class B type I or scavenger receptor class BI or scavenger receptor SR B1 or scavenger receptor SR BI or scavenger receptor type B1 or scavenger receptor type BI or SR B1 protein or protein SR BI).ti,ab.
52.	(scavenger receptor B type 1 or scavenger receptor B1 or scavenger receptor class B type 1 or scavenger receptor class B type I or scavenger receptor class BI or scavenger receptor SR B1 or scavenger receptor SR BI or scavenger receptor type B1 or scavenger receptor type BI or SR B1 protein or protein SR BI).kw.
53.	or/50-52
54.	*abc transporter/ or *abc transporter a1/ or "**carrier proteins and binding proteins"/ or *membrane protein/
55.	(abc* transporter? or Atp* binding cassette* or (abc* adj3 protein) or ABC* or cerp protein or cholesterol* efflux regulatory protein).ti.
56.	(abc* transporter? or Atp* binding cassette* or (abc* adj3 protein) or ABC* or cerp protein or cholesterol* efflux regulatory protein).kw.
57.	or/54-56
58.	transport at the cellular level/ or active transport/
59.	(biologic* transport or (active adj3 transport) or uphill transport or facilitated diffusion).ti,ab.
60.	(biologic* transport or (active adj3 transport) or uphill transport or facilitated diffusion).kw.
61.	cholesterol transport/
62.	(cholesterol uptake or transport,cholesterol).ti,ab.
63.	(cholesterol uptake or transport,cholesterol).kw.
64.	58 or 59 or 60
65.	61 or 62 or 63
66.	41 or 45 or 49 or 53 or 64 or 65
67.	57 and 66
68.	dose response/
69.	(dose* response relationship?,drug or drug augmentation? or drug potentiation? or drug synergism? or receptor up-regulation or up* regulation or upregulation).ti,ab.
70.	(dose* response relationship?,drug or drug augmentation? or drug potentiation? or drug synergism? or receptor up-regulation or up* regulation or upregulation).kw.
71.	or/68-70
72.	cardiovascular agent/
73.	(cardiovascular agent? or cardiovascular drug?).ti,ab.
74.	(cardiovascular agent? or cardiovascular drug?).kw.
75.	or/72-74
76.	drug potentiation/
77.	(drug potentiator or drug synergism or drug synergy or potentiation,drug).ti,ab.
78.	(drug potentiator or drug synergism or drug synergy or potentiation,drug).kw.
79.	or/76-78
80.	drug development/

81.	(development,drug or drug discovery or medication development or medication discovery or medicine development or medicine discovery or therapeutic discovery).ti,ab.
82.	(development,drug or drug discovery or medication development or medication discovery or medicine development or medicine discovery or therapeutic discovery).kw.
83.	or/80-82
84.	agents affecting lipid metabolism/
85.	(lipid regulating agent? or lipid regulating drug? or antihyperlipidemics or antilipemics agents or antilipemic drugs or antilipemics or hypolipidemic agents or hypolipidemic drugs).ti,ab.
86.	(lipid regulating agent? or lipid regulating drug? or antihyperlipidemics or antilipemics agents or antilipemic drugs or antilipemics or hypolipidemic agents or hypolipidemic drugs).kw.
87.	or/84-86
88.	molecularly targeted therapy/
89.	molecular targeted therap*.ti,ab.
90.	molecular targeted therap*.kw.
91.	or/88-90
92.	cholesterol ester transfer protein/ or cholesterol ester transfer protein inhibitor/
93.	(cholesterol ester transfer protein* or cholesteryl ester exchange protein or cholesteryl ester transfer protein or CETP or CETP inhibitor? or cholesteryl ester transfer protein inhibitor?).ti,ab.
94.	(cholesterol ester transfer protein* or cholesteryl ester exchange protein or cholesteryl ester transfer protein or CETP or CETP inhibitor? or cholesteryl ester transfer protein inhibitor?).kw.
95.	or/92-94
96.	peroxisome proliferator activated receptor gamma/ or peroxisome proliferator activated receptor gamma agonist/
97.	(ppar?gamma? or peroxisome proliferator?activated receptor gamma or (thiazolidinedione receptor or mppargamma?) or transcription factor? or peroxisome proliferator activated receptor gamma stimulant or peroxisome proliferator activated receptor gamma stimulating agent or PPAR gamma agonist or peroxisome proliferator activated receptor gamma stimulator or peroxisome proliferator activated receptor agonist or PPAR gamma stimulant or PPAR gamma stimulating agent or PPAR gamma stimulator).ti,ab.
98.	(ppar?gamma? or peroxisome proliferator?activated receptor gamma or (thiazolidinedione receptor or mppargamma?) or transcription factor? or peroxisome proliferator activated receptor gamma stimulant or peroxisome proliferator activated receptor gamma stimulating agent or PPAR gamma agonist or peroxisome proliferator activated receptor gamma stimulator or peroxisome proliferator activated receptor agonist or PPAR gamma stimulant or PPAR gamma stimulating agent or PPAR gamma stimulator).kw.
99.	or/96-98
100.	liver X receptor alpha/
101.	liver X receptor alpha.ti,ab.
102.	liver X receptor alpha.kw.
103.	or/100-102
104.	gene expression regulation/
105.	(gene action regulation or gene expression regulation or regulation of gene expression or receptor up-regulation or up?regulation or upregulation).ti,ab.
106.	(gene action regulation or gene expression regulation or regulation of gene expression or receptor up-regulation or up?regulation or upregulation).kw.
107.	or/104-106
108.	orphan nuclear receptor/
109.	orphan nuclear receptor?.ti,ab.
110.	orphan nuclear receptor?.kw.
111.	or/108-110
112.	retinoic acid receptor/
113.	(proteins, retinoic acid-binding or receptor?, retinoic acid or retinoic acid?binding proteins or retinoic acid receptor? or 9-cis-retinoic acid receptor or rxr protein or receptor?, retinoid x or retinoid x receptor? or Xr78e?f protein).ti,ab.
114.	(proteins, retinoic acid-binding or receptor?, retinoic acid or retinoic acid?binding proteins or retinoic acid receptor? or 9-cis-retinoic acid receptor or rxr protein or receptor?, retinoid x or retinoid x receptor? or Xr78e?f protein).kw.
115.	or/112-114
116.	thyroid hormone receptor/
117.	(c erb A protein or oncogene c erb A protein or receptor,thyroid hormone or dit receptor? or diiodotyrosine receptor? or mit receptors or moniodotyrosine receptors or t3 receptor? or t4 receptor? or thyroid hormone receptor? or thyroxine receptor? or triiodothyronine receptor?).ti,ab.

118.	(c erb A protein or oncogene c erb A protein or receptor,thyroid hormone or dit receptor? or diiodotyrosine receptor? or mit receptors or moniodotyrosine receptors or t3 receptor? or t4 receptor? or thyroid hormone receptor? or thyroxine receptor? or triiodothyronine receptor?).kw.
119.	or/116-118
120.	liver X receptor beta/
121.	liver X receptor beta.ti,ab.
122.	liver X receptor beta.kw.
123.	or/120-122
124.	liver X receptor agonist/
125.	(liver X receptor stimulant or liver X receptor stimulating agent or liver X receptor stimulator).ti,ab.
126.	(liver X receptor stimulant or liver X receptor stimulating agent or liver X receptor stimulator).kw.
127.	or/124-126
128.	hydroxymethylglutaryl coenzyme A reductase inhibitor/
129.	(Hmg?coa reductase inhibitor? or hmg-coa reductase inhibitors or hmg?coa statin? or hydroxymethylglutaryl?coa reductase inhibitor? or hydroxymethylglutaryl-coa inhibitors or hydroxymethylglutaryl-coa reductase inhibitors or hydroxymethylglutaryl-coenzyme a inhibitors).ti,ab.
130.	(Hmg?coa reductase inhibitor? or hmg-coa reductase inhibitors or hmg?coa statin? or hydroxymethylglutaryl?coa reductase inhibitor? or hydroxymethylglutaryl-coa inhibitors or hydroxymethylglutaryl-coa reductase inhibitors or hydroxymethylglutaryl-coenzyme a inhibitors).kw.
131.	or/128-130
132.	fibric acid derivative/
133.	(2 phenoxy?2?methylpropionic acid derivatives or 2?phenoxy isobutyric acids or fibrates or fibric acid derivatives or fibric acids or isobutyric acids, 2-phenoxy or methyl 2?phenoxypropanoic acid derivatives).ti,ab.
134.	(2 phenoxy?2?methylpropionic acid derivatives or 2?phenoxy isobutyric acids or fibrates or fibric acid derivatives or fibric acids or isobutyric acids, 2-phenoxy or methyl 2?phenoxypropanoic acid derivatives).kw.
135.	or/132-134
136.	2,4 thiazolidinedione derivative/
137.	(thiazolidine 2,4 dione derivative or thiazolidinedione or thiazolidinedione derivative).ti,ab.
138.	(thiazolidine 2,4 dione derivative or thiazolidinedione or thiazolidinedione derivative).kw.
139.	or/136-138
140.	terpene/
141.	(monoterpene? or terpene?).ti,ab.
142.	(monoterpene? or terpene?).kw.
143.	or/140-142
144.	propofol/
145.	(2,6?diisopropylphenol or 2,6-bis1-methylethylphenol or anepol or crytol or diisoprofol or abbott brand of propofol or alpha brand of propofol or aquafol or astra brand of propofol or astrazeneca brand of propofol or braun brand of propofol or curamed brand of propofol or diprivan or diprofol or disoprivan or disoprofol or fresofol or fresenius brand of propofol or fresenius kabi brand of propofol or gobbifol or ici 35?868 or pofol or propocam or propofol?ipuro or rapinovel or recofol or safol or ivofol or juste brand of propofol or parnell brand of propofol or pisa brand of propofol or propofol* or recofol or rovi brand of propofol or schering brand of propofol or zeneca brand of propofol).ti,ab.
146.	(2,6?diisopropylphenol or 2,6-bis1-methylethylphenol or anepol or crytol or diisoprofol or abbott brand of propofol or alpha brand of propofol or aquafol or astra brand of propofol or astrazeneca brand of propofol or braun brand of propofol or curamed brand of propofol or diprivan or diprofol or disoprivan or disoprofol or fresofol or fresenius brand of propofol or fresenius kabi brand of propofol or gobbifol or ici 35?868 or pofol or propocam or propofol?ipuro or rapinovel or recofol or safol or ivofol or juste brand of propofol or parnell brand of propofol or pisa brand of propofol or propofol* or recofol or rovi brand of propofol or schering brand of propofol or zeneca brand of propofol).kw.
147.	or/144-146
148.	tannin/
149.	(colloid tannin or gallotannic acid or gallotannin or hydrolyzable tannins or rhulitol or tannate or tannic acid or tannine or pyrogallol tannins).ti,ab.
150.	(colloid tannin or gallotannic acid or gallotannin or hydrolyzable tannins or rhulitol or tannate or tannic acid or tannine or pyrogallol tannins).kw.
151.	or/148-150
152.	benzofuran derivative/ or salvianolic acid b/
153.	(3 arylbenzofuran derivative or benzo?b?furan derivative or benzofuran 2,3 * substituted derivative or benzofuran 3,4,5,6 substituted derivative or benzofuran 7 substituted derivative or benzofurans or coumarones or diphenylbenzofuran).ti,ab.

154.	(3 arylbenzofuran derivative or benzo?b?furan derivative or benzofuran 2,3 * substituted derivative or benzofuran 3,4,5,6 substituted derivative or benzofuran 7 substituted derivative or benzofurans or coumarones or diphenylbenzofuran).kw.
155.	or/152-154
156.	probucol/
157.	(bifenabid or biphenabid or dh?581 or lesterol or lodeco or lopicol or lorelco or lursel* or panesclerina or sinlesta or superlipid).ti,ab.
158.	(bifenabid or biphenabid or dh?581 or lesterol or lodeco or lopicol or lorelco or lursel* or panesclerina or sinlesta or superlipid).kw.
159.	or/156-158
160.	caveolin 1/
161.	(caveolin?1 or vip21 protein or vesicular integral membrane protein 21 kda or alpha?caveolin or beta?caveolin).ti,ab.
162.	(caveolin?1 or vip21 protein or vesicular integral membrane protein 21 kda or alpha?caveolin or beta?caveolin).kw.
163.	or/160-162
164.	ubiquinone/
165.	(coenzyme q or ubiquinone*).ti,ab.
166.	(coenzyme q or ubiquinone*).kw.
167.	or/164-166
168.	caffeic acid derivative/
169.	(3,4 dihydroxycinnamic acid derivative or caffeic acids or caffeine salt).ti,ab.
170.	(3,4 dihydroxycinnamic acid derivative or caffeic acids or caffeine salt).kw.
171.	or/168-170
172.	chlorogenic acid/
173.	(1,3,4,5 tetrahydroxycyclohexanecarboxylic acid 3?3,4 dihydroxycinnamate? or 3?3,4 dihydroxycinnamoyl?quinic acid or 3 caffeoylquinic acid or 3 o caffeoylquinic acid or cis chlorogenic acid or trans chlorogenic acid).ti,ab.
174.	(1,3,4,5 tetrahydroxycyclohexanecarboxylic acid 3?3,4 dihydroxycinnamate? or 3?3,4 dihydroxycinnamoyl?quinic acid or 3 caffeoylquinic acid or 3 o caffeoylquinic acid or cis chlorogenic acid or trans chlorogenic acid).kw.
175.	or/172-174
176.	coumaric acid/
177.	(coumarate or coumaric acid? or cumaric acid or hydroxycinnamic acid).ti,ab.
178.	(coumarate or coumaric acid? or cumaric acid or hydroxycinnamic acid).kw.
179.	or/176-178
180.	gallic acid/
181.	(3,4,5-trihydroxybenzoic acid or gallic acid).ti,ab.
182.	(3,4,5-trihydroxybenzoic acid or gallic acid).kw.
183.	or/180-182
184.	nicotinic acid/
185.	(3?pyridinecarboxylic acid or aluminum salt, niacin or enduracin or hydrochloride, niacin or induracin or lithium nicotinate or niacin* or nico* or nicolar or nicotinate, lithium or nicotinate or nicotinic acid or tartrate, niacin or tosylate, niacin or wampocap).ti,ab.
186.	(3?pyridinecarboxylic acid or aluminum salt, niacin or enduracin or hydrochloride, niacin or induracin or lithium nicotinate or niacin* or nico* or nicolar or nicotinate, lithium or nicotinate or nicotinic acid or tartrate, niacin or tosylate, niacin or wampocap).kw.
187.	or/184-186
188.	calpain/
189.	(ca2+?activated protease or calcium?activated neutral prote*ase or calcium?activated protease or calcium dependent neutral prote*ase or calpain* or desminase or neutral prote*ase, calcium-activated or neutral prote*ase, calcium-dependent).ti,ab.
190.	(ca2+?activated protease or calcium?activated neutral prote*ase or calcium?activated protease or calcium dependent neutral prote*ase or calpain* or desminase or neutral prote*ase, calcium-activated or neutral prote*ase, calcium-dependent).kw.
191.	or/188-190
192.	betulin/
193.	(betuline or betulinol or lup 20?29?ene 3beta,28 diol or lup 20?30?ene 3beta,28 diol or trochol).ti,ab.
194.	(betuline or betulinol or lup 20?29?ene 3beta,28 diol or lup 20?30?ene 3beta,28 diol or trochol).kw.
195.	or/192-194

196.	piperine/
197.	1 piperoylpiperidine.ti,ab.
198.	1 piperoylpiperidine.kw.
199.	or/196-198
200.	71 or 75 or 79 or 83 or 87 or 91 or 95 or 99 or 103 or 107 or 111 or 115 or 119 or 123 or 127 or 131 or 135 or 139 or 143 or 147 or 151 or 155 or 159 or 163 or 167 or 171 or 175 or 179 or 183 or 187 or 191 or 195 or 199
201.	37 and 67 and 200
202.	(abc* transporter? or Atp* binding cassette* or (abc* adj3 protein) or ABC* or cerp protein or cholesterol* efflux regulatory protein).ti,ab.
203.	54 or 56 or 202
204.	203 and 66
205.	37 and 200 and 204

Supplementary Table 2 | Search strategy MEDLINE Overview of search strategy sets and search statements, as performed using the MEDLINE database on 12-02-2019

Set	Search statement
1.	atp-binding cassette transporters/ or atp binding cassette transporter 1/
2.	(abc* transporter? or Atp* binding cassette* or (abc* adj3 protein) or ABC* or cerp protein or cholesterol* efflux regulatory protein).ti,ab.
3.	(abc* transporter? or Atp* binding cassette* or (abc* adj3 protein) or ABC* or cerp protein or cholesterol* efflux regulatory protein).kf.
4.	or/1-3
5.	Multiprotein Complexes/ae, ag, ai, de, me, pk, pd, tu [Adverse Effects, Agonists, Antagonists & Inhibitors, Drug Effects, Metabolism, Pharmacokinetics, Pharmacology, Therapeutic Use]
6.	(macromolecular protein complex* or multiprotein complex*).ti,ab.
7.	(macromolecular protein complex* or multiprotein complex*).kf.
8.	or/5-7
9.	biological transport/ or biological transport, active/ or facilitated diffusion/
10.	(biologic* transport or (active adj3 transport) or uphill transport or facilitated diffusion).ti,ab.
11.	(biologic* transport or (active adj3 transport) or uphill transport or facilitated diffusion).kf.
12.	or/9-11
13.	Protein Transport/de, ph [Drug Effects, Physiology]
14.	(protein transport? or cellular protein localization process* or protein targeting or protein trafficking? or protein translocation).ti,ab.
15.	(protein transport? or cellular protein localization process* or protein targeting or protein trafficking? or protein translocation).kf.
16.	or/13-15
17.	carrier proteins/ or membrane transport proteins/ or atp-binding cassette transporters/ or atp binding cassette transporter 1/
18.	(biologic* pump? or (membrane adj3 transport*) or metabolic pump? or permease?).ti,ab.
19.	(biologic* pump? or (membrane adj3 transport*) or metabolic pump? or permease?).kf.
20.	or/17-19
21.	Scavenger Receptors, Class B/ag, ai, bl, de, me, ph, tu [Agonists, Antagonists & Inhibitors, Blood, Drug Effects, Metabolism, Physiology, Therapeutic Use]
22.	(sr-b protein? or scavenger receptor?, class b).ti,ab.
23.	(sr-b protein? or scavenger receptor?, class b).kf.
24.	or/21-23
25.	cholesterol/ or cholesterol esters/
26.	(cholesterol or epicholesterol or cholesterol esters or cholesteryl esters).ti,ab.
27.	(cholesterol or epicholesterol or cholesterol esters or cholesteryl esters).kf.
28.	or/25-27
29.	exp Lipid Metabolism/de, ph [Drug Effects, Physiology]
30.	lipid metabolism.ti,ab.
31.	lipid metabolism.kf.
32.	or/29-31
33.	Phospholipids/ag, ai, bl, me, pk, pd, ph, tu, th [Agonists, Antagonists & Inhibitors, Blood, Metabolism, Pharmacokinetics, Pharmacology, Physiology, Therapeutic Use, Therapy]
34.	(phosphatides or phospholipids).ti,ab.
35.	(phosphatides or phospholipids).kf.
36.	or/33-35
37.	exp Triglycerides/ag, aa, ai, bl, me, pk, pd, ph, tu [Agonists, Analogs & Derivatives, Antagonists & Inhibitors, Blood, Metabolism, Pharmacokinetics, Pharmacology, Physiology, Therapeutic Use]
38.	(triacylglycerol? or triglycerides).ti,ab.
39.	(triacylglycerol? or triglycerides).kf.
40.	exp Hepatocytes/de, me, ph [Drug Effects, Metabolism, Physiology]
41.	(hepatic cell? or hepatocyte?).ti,ab.
42.	(hepatic cell? or hepatocyte?).kf.

43.	exp Liver/de, dt, me, pd, ph, pp, th [Drug Effects, Drug Therapy, Metabolism, Pharmacology, Physiology, Physiopathology, Therapy]
44.	liver?.ti,ab.
45.	liver?.kf.
46.	or/43-45
47.	exp Homeostasis/de, ph [Drug Effects, Physiology]
48.	(autoregulation or homeostasis).ti,ab.
49.	(autoregulation or homeostasis).kf.
50.	or/47-49
51.	macrophages/ or foam cells/
52.	(one marrow?derived macrophage? or macrophage? or monocyte?derived macrophage? or foam cell?).ti,ab.
53.	(bone marrow?derived macrophage? or macrophage? or monocyte?derived macrophage? or foam cell?).kf.
54.	or/51-53
55.	lipoproteins/ or apolipoproteins/ or apolipoproteins a/ or apolipoprotein a-i/ or apolipoproteins e/ or lipoproteins, hdl/ or cholesterol, hdl/ or high-density lipoproteins, pre-beta/ or lipoproteins, hdl2/ or lipoproteins, hdl3/ or lipoproteins, ldl/
56.	(hdl lipoproteins or heavy lipoproteins or high?density lipoproteins or lipoproteins, hdl or alpha* lipoprotein? or hdl* cholesterol or high density lipoprotein cholesterol or alpha?lipoprotein cholesterol or hdl lipoprotein, pre?beta or hdl, pre-beta1 or hdl, pre-beta or high density lipoprotein?, pre?beta? or nascent hdl or hdl?2 or high density lipoprotein hdl2 or high density lipoprotein?2 or lipoprotein? hdl2 or hdl?3 or high density lipoprotein?3 or high density lipoprotein hdl3 or lipoprotein?, hdl3 or apolipoprotein? or circulating lipoproteins or lipoprotein? or apo-a or apo a or apolipoproteins a or apo a?i isoproteins or apo a-1 or apo?a?i or apo?a?1 or apolipoprotein a 1 or apolipoprotein a?i isoprotein* or apolipoprotein a i or apolipoprotein a?1 or apolipoprotein a-i or apolipoprotein ai propeptide or apolipoprotein ai or pro?apo a?i or pro apolipoprotein a i or pro?apolipoprotein a?i or apo e isoproteins or apo?e or apolipoprotein e isoproteins or apolipoproteins e or apoprotein ?e? or apoproteins e).ti,ab.
57.	(hdl lipoproteins or heavy lipoproteins or high?density lipoproteins or lipoproteins, hdl or alpha* lipoprotein? or hdl* cholesterol or high density lipoprotein cholesterol or alpha?lipoprotein cholesterol or hdl lipoprotein, pre?beta or hdl, pre-beta1 or hdl, pre-beta or high density lipoprotein?, pre?beta? or nascent hdl or hdl?2 or high density lipoprotein hdl2 or high density lipoprotein?2 or lipoprotein? hdl2 or hdl?3 or high density lipoprotein?3 or high density lipoprotein hdl3 or lipoprotein?, hdl3 or apolipoprotein? or circulating lipoproteins or lipoprotein? or apo-a or apo a or apolipoproteins a or apo a?i isoproteins or apo a-1 or apo?a?i or apo?a?1 or apolipoprotein a 1 or apolipoprotein a?i isoprotein* or apolipoprotein a i or apolipoprotein a?1 or apolipoprotein a-i or apolipoprotein ai propeptide or apolipoprotein ai or pro?apo a?i or pro apolipoprotein a i or pro?apolipoprotein a?i or apo e isoproteins or apo?e or apolipoprotein e isoproteins or apolipoproteins e or apoprotein ?e? or apoproteins e).kf.
58.	or/55-57
59.	exp Sp1 Transcription Factor/ag, ai, bl, de, me, pd, ph [Agonists, Antagonists & Inhibitors, Blood, Drug Effects, Metabolism, Pharmacology, Physiology]
60.	(sp1 transcription factor or specificity protein 1 transcription factor).ti,ab.
61.	(sp1 transcription factor or specificity protein 1 transcription factor).kf.
62.	or/59-61
63.	exp Sterol Regulatory Element Binding Proteins/ai, de, me, pd, ph [Antagonists & Inhibitors, Drug Effects, Metabolism, Pharmacology, Physiology]
64.	(srebp proteins or sterol regulatory element binding proteins).ti,ab.
65.	(srebp proteins or sterol regulatory element binding proteins).kf.
66.	or/63-65
67.	4 and 12 and 24 and 28 and 32
68.	4 and 24 and 28 and 58
69.	or/40-42
70.	28 and 46 and 58 and 69
71.	or/37-39
72.	28 and 32 and 50 and 58 and 66 and 71
73.	4 and 12 and 24 and 28 and 46
74.	4 and 28 and 62
75.	28 and 54 and 58
76.	4 and 8 and 20 and 28
77.	28 and 50 and 58
78.	66 and 77
79.	4 or 8 or 12 or 16 or 20 or 24
80.	28 or 32 or 36 or 46 or 50 or 54 or 58 or 62 or 66 or 69 or 71

81.	79 and 80
82.	Multiprotein Complexes/
83.	(macromolecular protein complex* or multiprotein complex*).ti,ab.
84.	(macromolecular protein complex* or multiprotein complex*).kf.
85.	or/82-84
86.	Protein Transport/
87.	(protein transport? or cellular protein localization process* or protein targeting or protein trafficking? or protein translocation).ti,ab.
88.	(protein transport? or cellular protein localization process* or protein targeting or protein trafficking? or protein translocation).kf.
89.	or/86-88
90.	carrier proteins/ or membrane transport proteins/ or atp binding cassette transporter 1/
91.	(biologic* pump? or (membrane adj3 transport*) or metabolic pump? or permease?).ti,ab.
92.	(biologic* pump? or (membrane adj3 transport*) or metabolic pump? or permease?).kf.
93.	or/90-92
94.	glycoproteins/ or cholesterol ester transfer proteins/
95.	(*glycoprotein?/ or (cholester?! ester adj3 protein?)).ti,ab.
96.	(*glycoprotein?/ or (cholester?! ester adj3 protein?)).kf.
97.	or/94-96
98.	Scavenger Receptors, Class B/
99.	(sr-b protein? or scavenger receptor?, class b).ti,ab.
100.	(sr-b protein? or scavenger receptor?, class b).kf.
101.	or/98-100
102.	*atp-binding cassette transporters/ or exp *atp binding cassette transporter 1/
103.	(abc* transporter? or Atp* binding cassette* or (abc* adj3 protein) or ABC* or cerp protein or cholesterol* efflux regulatory protein).ti.
104.	*biological transport/ or exp *biological transport, active/ or exp *facilitated diffusion/
105.	(biologic* transport or (active adj3 transport) or uphill transport or facilitated diffusion).ti.
106.	or/102-103
107.	or/104-105
108.	85 or 89 or 93 or 97 or 101 or 106 or 107
109.	biological transport/ or exp biological transport, active/ or exp facilitated diffusion/
110.	(biologic* transport or (active adj3 transport) or uphill transport or facilitated diffusion).ti,ab.
111.	(biologic* transport or (active adj3 transport) or uphill transport or facilitated diffusion).kf.
112.	109 or 110 or 111
113.	85 or 89 or 93 or 97 or 101 or 112
114.	106 and 113
115.	(abc* transporter? or Atp* binding cassette* or (abc* adj3 protein) or ABC* or cerp protein or cholesterol* efflux regulatory protein).kf.
116.	102 or 103 or 115
117.	113 and 116
118.	Lipid Metabolism/
119.	lipid metabolism.ti,ab.
120.	lipid metabolism.kf.
121.	or/118-120
122.	Phospholipids/
123.	(phosphatides or phospholipids).ti,ab.
124.	(phosphatides or phospholipids).kf.
125.	or/122-124
126.	Triglycerides/
127.	(triacylglycerol? or triglycerides).ti,ab.
128.	triacylglycerol?.ti,ab. or triglycerides.kf.
129.	or/126-128
130.	Hepatocytes/

131.	(hepatic cell? or hepatocyte?).ti,ab.
132.	(hepatic cell? or hepatocyte?).kf.
133.	or/130-132
134.	Liver/
135.	liver?.ti,ab.
136.	liver?.kf.
137.	or/134-136
138.	macrophages/ or foam cells/
139.	(bone marrow?derived macrophage? or macrophage? or monocyte?derived macrophage? or foam cell?).ti,ab.
140.	(bone marrow?derived macrophage? or macrophage? or monocyte?derived macrophage? or foam cell?).kf.
141.	or/138-140
142.	lipoproteins/ or apolipoproteins/ or apolipoproteins a/ or apolipoprotein a-i/ or lipoproteins, hdl/ or cholesterol, hdl/ or high-density lipoproteins, pre-beta/ or lipoproteins, hdl2/ or lipoproteins, hdl3/
143.	(hdl lipoproteins or heavy lipoproteins or high?density lipoproteins or lipoproteins, hdl or alpha* lipoprotein? or hdl* cholesterol or high density lipoprotein cholesterol or alpha?lipoprotein cholesterol or hdl lipoprotein, pre?beta or hdl, pre-beta1 or hdl, pre-beta or high density lipoprotein?, pre?beta? or nascent hdl or hdl?2 or high density lipoprotein hdl2 or high density lipoprotein?2 or lipoprotein? hdl2 or hdl?3 or high density lipoprotein?3 or high density lipoprotein hdl3 or lipoprotein?, hdl3 or apolipoprotein? or circulating lipoproteins or lipoprotein? or apo-a or apoa or apolipoproteins a or apo a?i isoproteins or apo a-1 or apo?a?i or apo?a?1 or apolipoprotein a 1 or apolipoprotein a?i isoprotein* or apolipoprotein a i or apolipoprotein a?1 or apolipoprotein a-i or apolipoprotein ai propeptide or apolipoprotein ai or pro?apo a?i or pro apolipoprotein a i or pro?apolipoprotein a?i or apo e isoproteins or apo?e or apolipoprotein e isoproteins or apolipoproteins e or apoprotein ?e? or apoproteins e).ti,ab.
144.	(hdl lipoproteins or heavy lipoproteins or high?density lipoproteins or lipoproteins, hdl or alpha* lipoprotein? or hdl* cholesterol or high density lipoprotein cholesterol or alpha?lipoprotein cholesterol or hdl lipoprotein, pre?beta or hdl, pre-beta1 or hdl, pre-beta or high density lipoprotein?, pre?beta? or nascent hdl or hdl?2 or high density lipoprotein hdl2 or high density lipoprotein?2 or lipoprotein? hdl2 or hdl?3 or high density lipoprotein?3 or high density lipoprotein hdl3 or lipoprotein?, hdl3 or apolipoprotein? or circulating lipoproteins or lipoprotein? or apo-a or apoa or apolipoproteins a or apo a?i isoproteins or apo a-1 or apo?a?i or apo?a?1 or apolipoprotein a 1 or apolipoprotein a?i isoprotein* or apolipoprotein a i or apolipoprotein a?1 or apolipoprotein a-i or apolipoprotein ai propeptide or apolipoprotein ai or pro?apo a?i or pro apolipoprotein a i or pro?apolipoprotein a?i or apo e isoproteins or apo?e or apolipoprotein e isoproteins or apolipoproteins e or apoprotein ?e? or apoproteins e).kf.
145.	or/142-144
146.	Sp1 Transcription Factor/
147.	(sp1 transcription factor or specificity protein 1 transcription factor).ti,ab.
148.	(sp1 transcription factor or specificity protein 1 transcription factor).kf.
149.	or/146-148
150.	*cholesterol/ or *cholesterol derivative/ or *cholesterol ester/
151.	(cholesterol or epicholesterol or cholesterol esters or cholesteryl esters).ti.
152.	or/150-151
153.	121 or 125 or 129 or 133 or 137 or 141 or 145 or 149 or 152
154.	dose-response relationship, drug/ or drug synergism/ or up-regulation/
155.	(dose* response relationship?,drug or drug augmentation? or drug potentiation? or drug synergism? or receptor up-regulation or up* regulation or upregulation).ti,ab.
156.	(dose* response relationship?,drug or drug augmentation? or drug potentiation? or drug synergism? or receptor up-regulation or up* regulation or upregulation).kf.
157.	or/154-156
158.	Cardiovascular Agents/
159.	(cardiovascular agent? or cardiovascular drug?).ti,ab.
160.	(cardiovascular agent? or cardiovascular drug?).kf.
161.	or/158-160
162.	exp Molecular Targeted Therapy/
163.	molecular targeted therap*.ti,ab.
164.	molecular targeted therap*.kf.
165.	or/162-164
166.	exp Cholesterol Ester Transfer Proteins/
167.	(cholesterol ester transfer protein* or cholesteryl ester exchange protein or cholesteryl ester transfer protein).ti,ab.
168.	(cholesterol ester transfer protein* or cholesteryl ester exchange protein or cholesteryl ester transfer protein).kf.
169.	or/166-168

170.	exp Transcription Factors/ or exp PPAR gamma/
171.	(ppar?gamma? or peroxisome proliferator?activated receptor gamma or thiazolidinedione receptor or mppargamma? or transcription factor?).ti,ab.
172.	(ppar?gamma? or peroxisome proliferator?activated receptor gamma or thiazolidinedione receptor or mppargamma? or transcription factor?).kf.
173.	or/170-172
174.	orphan nuclear receptors/ or receptors, retinoic acid/ or retinoid x receptors/ or receptors, thyroid hormone/
175.	(orphan nuclear receptor? or retinoic acid binding protein? or retinoic acid receptor? or dit receptor? or diiodotyrosine receptor? or mit receptors or monoiodotyrosine receptors or t3 receptor? or t4 receptor? or thyroid hormone receptor? or thyroxine receptor? or triiodothyronine receptor?).ti,ab.
176.	(orphan nuclear receptor? or retinoic acid binding protein? or retinoic acid receptor? or dit receptor? or diiodotyrosine receptor? or mit receptors or monoiodotyrosine receptors or t3 receptor? or t4 receptor? or thyroid hormone receptor? or thyroxine receptor? or triiodothyronine receptor?).kf.
177.	or/174-176
178.	receptors, retinoic acid/ or retinoid x receptors/
179.	(proteins, retinoic acid-binding or receptor?, retinoic acid or retinoic acid?binding proteins or retinoic acid receptor? or 9-cis-retinoic acid receptor or rxr protein or receptor?, retinoid x or retinoid x receptor? or Xr78e?f protein).ti,ab.
180.	(proteins, retinoic acid-binding or receptor?, retinoic acid or retinoic acid?binding proteins or retinoic acid receptor? or 9-cis-retinoic acid receptor or rxr protein or receptor?, retinoid x or retinoid x receptor? or Xr78e?f protein).kf.
181.	or/178-180
182.	gene expression regulation/ or up-regulation/
183.	(gene action regulation or gene expression regulation or regulation of gene expression or receptor up-regulation or up?regulation or upregulation).ti,ab.
184.	(gene action regulation or gene expression regulation or regulation of gene expression or receptor up-regulation or up?regulation or upregulation).kf.
185.	or/182-184
186.	Fibric Acids/
187.	(2 phenoxy?2?methylpropionic acid derivatives or 2?phenoxy isobutyric acids or fibrates or fibric acid derivatives or fibric acids or isobutyric acids, 2-phenoxy or methyl 2?phenoxypropanoic acid derivatives).ti,ab.
188.	(2 phenoxy?2?methylpropionic acid derivatives or 2?phenoxy isobutyric acids or fibrates or fibric acid derivatives or fibric acids or isobutyric acids, 2-phenoxy or methyl 2?phenoxypropanoic acid derivatives).kf.
189.	or/186-188
190.	Thiazolidinediones/
191.	(glitazone or thiazolidinediones).ti,ab.
192.	(glitazone or thiazolidinediones).kf.
193.	or/190-192
194.	Tretinoin/
195.	(acid, vitamin a or acid, all-trans-retinoic or acid, beta-all-trans-retinoic or acid, trans-retinoic or potassium salt, tretinoin or retin?a or retinoic acid or tretinoin* or tretinoin or vesanoid or all?trans retinoic acid or beta?all?trans?retinoic acid or trans?retinoic acid).ti,ab.
196.	(acid, vitamin a or acid, all-trans-retinoic or acid, beta-all-trans-retinoic or acid, trans-retinoic or potassium salt, tretinoin or retin?a or retinoic acid or tretinoin* or tretinoin or vesanoid or all?trans retinoic acid or beta?all?trans?retinoic acid or trans?retinoic acid).kf.
197.	or/194-196
198.	hydrocarbons/ or monoterpenes/
199.	(hydrocarbons or monoterpen*).ti,ab.
200.	(hydrocarbons or monoterpen*).kf.
201.	or/198-200
202.	Propofol/
203.	(2,6?diisopropylphenol or 2,6-bis1-methylethylphenol or abbott brand of propofol or alpha brand of propofol or aquafol or astra brand of propofol or astrazeneca brand of propofol or braun brand of propofol or curamed brand of propofol or diprivan or disoprivan or disoprofol or fresenius brand of propofol or fresenius kabi brand of propofol or fresofol or ici?35,868 or ivofol or juste brand of propofol or parnell brand of propofol or pisa brand of propofol or propofol* or recofol or rovi brand of propofol or schering brand of propofol or zeneca brand of propofol).ti,ab.
204.	(2,6?diisopropylphenol or 2,6-bis1-methylethylphenol or abbott brand of propofol or alpha brand of propofol or aquafol or astra brand of propofol or astrazeneca brand of propofol or braun brand of propofol or curamed brand of propofol or diprivan or disoprivan or disoprofol or fresenius brand of propofol or fresenius kabi brand of propofol or fresofol or ici?35,868 or ivofol or juste brand of propofol or parnell brand of propofol or pisa brand of propofol or propofol* or recofol or rovi brand of propofol or schering brand of propofol or zeneca brand of propofol).kf.

205.	or/202-204
206.	gallic acid/ or hydrolyzable tannins/
207.	(3,4,5-trihydroxybenzoic acid or gallic acid or ellagi?tannins or ellagitannins or gallo?tannins or gallotannins or hydrolyzable tannins or pyrogallol tannins).ti,ab.
208.	(3,4,5-trihydroxybenzoic acid or gallic acid or ellagi?tannins or ellagitannins or gallo?tannins or gallotannins or hydrolyzable tannins or pyrogallol tannins).kf.
209.	or/206-208
210.	Benzofurans/
211.	(benzofurans or coumarones or diphenylbenzofuran).ti,ab.
212.	(benzofurans or coumarones or diphenylbenzofuran).kf.
213.	or/210-212
214.	Probucol/
215.	(almirall brand of probucol or aventis brand of probucol or biphenabid or dh?581 or dh581 or hoechst brand of probucol or lorelco or lurselle or panavir or probucol or superlipid).ti,ab.
216.	(almirall brand of probucol or aventis brand of probucol or biphenabid or dh?581 or dh581 or hoechst brand of probucol or lorelco or lurselle or panavir or probucol or superlipid).kf.
217.	or/214-216
218.	Caveolin 1/
219.	(caveolin?1 or vip21 protein or vesicular integral membrane protein 21 kda or alpha?caveolin or beta?caveolin).ti,ab.
220.	(caveolin?1 or vip21 protein or vesicular integral membrane protein 21 kda or alpha?caveolin or beta?caveolin).kf.
221.	or/218-220
222.	Ubiquinone/
223.	(coenzyme q or ubiquinone).ti,ab.
224.	(coenzyme q or ubiquinone).kf.
225.	or/222-224
226.	caffeic acids/ or chlorogenic acid/ or coumaric acids/
227.	(caffeic acids or 3?caffeoylquinic acid or chlorogenic acid or coumaric acids or hydroxycinnamic acids).ti,ab.
228.	(caffeic acids or 3?caffeoylquinic acid or chlorogenic acid or coumaric acids or hydroxycinnamic acids).kf.
229.	Niacin/
230.	(3?pyridinecarboxylic acid or aluminum salt, niacin or enduracin or hydrochloride, niacin or induracin or lithium nicotinate or niacin* or nico* or nicolar or nicotinate, lithium or nicotinate or nicotinic acid or tartrate, niacin or tosylate, niacin or wampocap).ti,ab.
231.	(3?pyridinecarboxylic acid or aluminum salt, niacin or enduracin or hydrochloride, niacin or induracin or lithium nicotinate or niacin* or nico* or nicolar or nicotinate, lithium or nicotinate or nicotinic acid or tartrate, niacin or tosylate, niacin or wampocap).kf.
232.	or/226-228
233.	or/229-231
234.	Calpain/
235.	(ca2+?activated protease or calcium?activated neutral prote*ase or calcium?activated protease or calcium dependent neutral prote*ase or calpain* or desminase or neutral prote*ase, calcium-activated or neutral prote*ase, calcium-dependent).ti,ab.
236.	(ca2+?activated protease or calcium?activated neutral prote*ase or calcium?activated protease or calcium dependent neutral prote*ase or calpain* or desminase or neutral prote*ase, calcium-activated or neutral prote*ase, calcium-dependent).kf.
237.	or/234-236
238.	157 or 161 or 165 or 169 or 173 or 177 or 181 or 185 or 189 or 193 or 197 or 201 or 205 or 209 or 213 or 217 or 221 or 225 or 232 or 233 or 237
239.	117 and 153 and 238
240.	(abc* transporter? or Atp* binding cassette* or (abc* adj3 protein) or ABC* or cerp protein or cholesterol* efflux regulatory protein).ti,ab.
241.	102 or 115 or 240
242.	113 and 241
243.	153 and 238 and 242