



ASSOCIATE EDITOR: LYNETTE DAWS

Pharmacological Modulation of the Crosstalk between Aberrant Janus Kinase Signaling and Epigenetic Modifiers of the Histone Deacetylase Family to Treat Cancer

 Al-Hassan M. Mustafa and  Oliver H. Krämer

Department of Toxicology, University Medical Center, Mainz, Germany (A.-H.M.M., O.H.K.) and Department of Zoology, Faculty of Science, Aswan University, Aswan, Egypt (A.-H.M.M.)

Abstract	36
Significance Statement	36
I. Introduction	36
II. Relevance of Janus Kinases for Tumorigenesis	38
A. Janus Kinase-1	38
B. Janus Kinase-2	39
C. Janus Kinase-3	41
D. Tyrosine Kinase-2	41
III. Histone Deacetylases Are Valid Pharmacological Targets in Janus Kinase Mutant Tumor Cells	41
A. Janus Kinase–Signal Transducer and Activator of Transcription Pathways Control Cellular Susceptibility to Histone Deacetylase Inhibitors	43
B. Histone Deacetylase Inhibitors Can Correct Aberrant Janus Kinase–Signal Transducer and Activator of Transcription Signaling	45
1. Impact of Histone Deacetylase Inhibitors on Janus Kinase-1	45
2. Impact of Histone Deacetylase Inhibitors on Janus Kinase-2	45
a. Histone deacetylase inhibitors attenuate JAK2 ^{V617F} in blood malignancies	45
b. Histone deacetylase inhibitors modulate aberrant JAK2 signaling in solid tumor cells	47
3. Impact of Histone Deacetylase Inhibitors on Janus Kinase-3	47
4. Impact of Histone Deacetylase Inhibitors on Tyrosine Kinase-2	48
5. Histone Deacetylase Inhibitors Modulate Janus Kinase–Signal Transducer and Activator of Transcription–Dependent Immune Functions	48
6. Advantages of Indirect Pan–Janus Kinase Inhibition by Histone Deacetylase Inhibitors	49
IV. Chemically Coupled Multitargeting Drugs	50
V. Targeting Aberrant Janus Kinase Signaling by Proteolysis Targeting Chimeras and Histone Deacetylase Inhibitors	51
VI. Clinical Trials to Assess the Safety and Efficacy of Combined Pharmacological Targeting of Janus Kinases and Histone Deacetylases	53
VII. Summary	54
References	55

Address correspondence to: Oliver H. Krämer, Department of Toxicology, University Medical Center, Obere Zahlbacher Str. 67, 55131 Mainz, Germany. E-mail: okraemer@uni-mainz.de

We gratefully acknowledge that work done by A.-H.M.M. in the group of O.H.K. is funded by the German Research Foundation/Deutsche Forschungsgemeinschaft (DFG) grant KR2291/12-1, DFG project number 445785155 (to O.H.K.) and was initially made possible by the DAAD. Additional support to O.H.K. is from the DFG project number 393547839 - SFB 1361, subproject 11; KR2291/9-1, DFG project number 427404172; KR2291/14-1, DFG project number 469954457; KR2291/15-1, DFG project number 495271833; KR2291/16-1, DFG project number 496927074; KR2291/17-1, DFG project number 502534123; the Wilhelm Sander-Stiftung (grant 2019.086.1); the Brigitte und Dr. Konstanze Wegener-Stiftung (Projekt 65); and the Walter Schulz Stiftung. We thank all our group members for helpful discussions and input on this work.

No author has an actual or perceived conflict of interest with the contents of this article. O.H.K. declares the patents “Synthesis, pharmacology, and use of new and selective FMS-like tyrosine kinase 3 (FLT3) inhibitors, WO2019/034538”, “Novel HDAC6 inhibitors and their uses, WO2016020369A1”, and “The use of molecular markers for the preclinical and clinical profiling of inhibitors of enzymes having histone deacetylase activity, WO/2004/027418.” These patents cover substance classes that are discussed in this work.

dx.doi.org/10.1124/pharmrev.122.000612.

Abstract—Hyperactivated Janus kinase (JAK) signaling is an appreciated drug target in human cancers. Numerous mutant JAK molecules as well as inherent and acquired drug resistance mechanisms limit the efficacy of JAK inhibitors (JAKi). There is accumulating evidence that epigenetic mechanisms control JAK-dependent signaling cascades. Like JAKs, epigenetic modifiers of the histone deacetylase (HDAC) family regulate the growth and development of cells and are often dysregulated in cancer cells. The notion that inhibitors of histone deacetylases (HDACi) abrogate oncogenic JAK-dependent signaling cascades illustrates an intricate crosstalk between JAKs and HDACs. Here, we summarize how structurally divergent, broad-acting as well as isoenzyme-specific HDACi, hybrid fusion pharmacophores containing JAKi and HDACi, and proteolysis targeting chimeras for JAKs inactivate the four JAK proteins JAK1, JAK2, JAK3, and tyrosine kinase-2. These agents suppress aberrant JAK activity through specific transcription-dependent processes and mechanisms that

alter the phosphorylation and stability of JAKs. Pharmacological inhibition of HDACs abrogates allosteric activation of JAKs, overcomes limitations of ATP-competitive type 1 and type 2 JAKi, and interacts favorably with JAKi. Since such findings were collected in cultured cells, experimental animals, and cancer patients, we condense preclinical and translational relevance. We also discuss how future research on acetylation-dependent mechanisms that regulate JAKs might allow the rational design of improved treatments for cancer patients.

Significance Statement—Reversible lysine- ϵ -N acetylation and deacetylation cycles control phosphorylation-dependent Janus kinase–signal transducer and activator of transcription signaling. The intricate crosstalk between these fundamental molecular mechanisms provides opportunities for pharmacological intervention strategies with modern small molecule inhibitors. This could help patients suffering from cancer.

I. Introduction

The Janus kinases (JAK) family consists of four cytosolic nonreceptor tyrosine kinases (TKs). These are tyrosine kinase-2 (TYK2), JAK1, JAK2, and JAK3 (Babon et al., 2014). These molecules consist of about 1150 amino acids and have a molecular mass of approximately 130 kDa. Genes encoding human JAKs locate in chromosomes 19p13.2 (*TYK2*), 1p31.3 (*JAK1*), 9p24.1 (*JAK2*), and 19p13.11 (*JAK3*) (Leonard and O’Shea, 1998).

All JAKs share a modular structure consisting of seven JAK homology (JH) domains that fall into four regions (Fig. 1). These are the N-terminal four-point-one, ezrin, radixin, moesin domain (FERM, JH5–JH7); the Src homology-2 (SH2)-like domain (JH3 and part of JH4); the inactive pseudokinase (JH2); and the catalytically active kinase (JH1) (Schindler et al., 2007) (Fig. 1). The four-point-one, ezrin, radixin, moesin domains mediate specific interactions of JAKs with plasma membrane-inserted receptors that bind extracellular signaling molecules (Funakoshi-Tago et al., 2006; Babon et al., 2014). The

SH2 domains of JAKs contribute to their receptor binding and their dimerization for activating crossphosphorylation (Gorantla et al., 2010; Glassman et al., 2022). The pseudokinase domains of JAKs function as negative regulators of the tyrosine kinase domains by protein-protein interactions (Ungureanu et al., 2011; Lupardus et al., 2014).

TYK2 received its name from its biochemical activity and from being the second identified tyrosine kinase that can phosphorylate tyrosine residues within proteins. Accordingly, TYK2 is the founding member of the JAK family and has been linked to cytokine and growth factor signaling since 1990 (Firmbach-Kraft et al., 1990; Krolewski et al., 1990; Verma et al., 2003).

JAKs were given the name of the ancient two-headed god *Janus*, who faces the past and the future (Darnell et al., 1994; Ihle et al., 1995; Pellegrini and Dusanter-Fourt, 1997; Leonard and O’Shea, 1998). This two-headed functionality reflects their function as linkers between the receptors for cytokines and growth factors and the signal transducers and activators of transcription (STATs). These inducible transcription factors comprise STAT1, STAT2, STAT3, STAT4,

ABBREVIATIONS: ACK1, activated CDC42-associated kinase-1; AKT, serine-threonine protein kinase B; ALL, acute lymphoblastic leukemia; AML, acute myeloid leukemia; B-ALL, B-cell acute lymphoblastic leukemia; BRD4, bromodomain-containing protein-4; CDK, cyclin-dependent kinase; CMML, chronic myelomonocytic leukemia; CRBN, cereblon; CRLF2, cytokine receptor-like factor-2/thymic stromal lymphopoietin receptor; CTCL, cutaneous T-cell lymphoma; Cys909, cysteine 909; DLBCL, diffuse large B-cell lymphoma; ERK, extracellular signal-regulated kinases; ET, essential thrombocythemia; FDA, Food and Drug Administration; FERM, four-point-one, ezrin, radixin, moesin; FLT3, FMS-like tyrosine kinase-3; GOF, gain of function; GSPT1, G1-to-S-phase-transition-1; HCC, hepatocellular carcinoma; HDAC, histone deacetylase; HDACi, histone deacetylase inhibitor; HSP90, heat shock protein-90; IAP, inhibitor of apoptosis; IFN, interferon; JAK, Janus kinase; JAKi, JAK inhibitor; JH, JAK homology; LIFR, leukemia inhibitory factor receptor; MF, myelofibrosis; MPN, myeloproliferative neoplasm; MYC, myelocytomatosis oncogene; NF- κ B, nuclear factor- κ B; NK, natural killer; NSCLC, non-small-cell lung cancer; OSCC, oral squamous cell carcinoma; PanNET, pancreatic neuroendocrine tumors; PARP1, poly-ADP-ribose polymerase-1; PD-1, programmed death receptor-1; PDAC, pancreatic ductal adenocarcinoma; PD-L1, programmed cell death ligand-1; PG, phenyl glutarimide; PMF, primary myelofibrosis; PROTAC, proteolysis targeting chimera; PV, polycythemia vera; RNAi, RNA interference; SH2, Src homology-2; SIAH2, E3 ubiquitin ligase seven-in-absentia-homologue-2; SIRT, sirtuin; SOCS, suppressor of cytokine signaling; STAT, signal transducer and activator of transcription; T-ALL, T-cell acute lymphoblastic leukemia; TK, tyrosine kinase; TKi, tyrosine kinase inhibitor; T-PLL, T-cell prolymphocytic leukemia; TSA, trichostatin A; TYK2, tyrosine kinase-2; VHL, von Hippel-Lindau.




Fig. 1. Structural representation of the JAK family members JAK1–3 and TYK2. Shown is the organization of JAKs into the domains JH1–7 and frequent mutations of JAKs in tumor cells.

STAT5A, STAT5B, and STAT6. Tyrosine-phosphorylated STATs enter the nucleus, where they bind to cognate STAT consensus motifs in genes and promote a recruitment of transcription machineries (Fig. 2). This consequently modulates gene expression patterns that crucially determine cell growth, proliferation, differentiation, and cell death. These vast biologic implications of JAK-STAT signaling can explain why dysregulation of this pathway is frequently associated with solid and hematologic malignancies, autoimmunity, and immunologic failures (Owen et al., 2019; Bharadwaj et al., 2020; Yang et al., 2020).

Histone deacetylases (HDACs) are epigenetic modifiers that fall into four classes. HDAC1, HDAC2, HDAC3, and HDAC8 form class I; HDAC4, HDAC5, HDAC7, and HDAC9 are class IIa; HDAC6 and

HDAC10 represent class IIb; and HDAC11 is the sole member of class IV. The sirtuins (SIRT)s 1–7 represent class III (Table 1). HDAC classes I, II, and IV use Zn^{2+} to polarize H_2O for a nucleophilic attack on the acetyl moiety in acetylated lysine residues (Krämer et al., 2009). SIRTs use NAD^+ for their deacetylation reaction, which couples their catalytic activities to the energy state of cells (Stein and Imai, 2012; Cantó et al., 2015; Jiang et al., 2017b).

Our review summarizes evidence on how structurally divergent histone deacetylase inhibitors (HDACi), derivatives thereof with tyrosine kinase inhibitor (TKi) activity, and TKi with a functional unit driving proteolysis of JAKs kill cancer cells. Moreover, we discuss how the extraction of JAK mutations and regulatory patterns in tumor cells can propel a rational




Fig. 2. The diagram shows wild-type JAK-STAT (left) and mutant JAK2^{V617F}-STAT (right) signaling. The binding of a cytokine, exemplified for erythropoietin (EPO), promotes the dimerization of its receptor and subsequently JAK2 dimerization and crossphosphorylation. Phosphorylation of the cytokine receptor creates a binding interface for the SH2 domains of preformed STAT dimers. These become phosphorylated by JAKs, form high-avidity dimers via SH2-pY interactions, and enter the nucleus to induce specific gene expression patterns. Cognate STAT consensus motifs in gene promoters are usually the palindromic γ -activated sequences and the interferon-stimulated response elements. Independent of receptor activation and due to structurally facilitated JAK2 dimer formation, JAK2^{V617F} activates STATs. The resulting tonic gene activation promotes tumorigenesis. This article summarizes how HDACi can alter the molecules and mechanisms of these pathways. We exemplarily depict the suppressive effects of HDACi on JAK2^{V617F}. These rely on the acetylation of STATs and their subsequent dephosphorylation and on the induction of SOCS proteins that directly inhibit and promote the degradation of JAKs. HDACi additionally decrease transcription of the *JAK2* gene locus and accelerate the proteasomal degradation of JAK2^{V617F}; see text for details. Created with biorender.com.

development of new pharmacological approaches involving HDACi and TKi. We will first present how the four JAK proteins and the 11 Zn²⁺-dependent HDACs contribute to tumorigenesis.

II. Relevance of Janus Kinases for Tumorigenesis

A. Janus Kinase-1

JAK1 is frequently dysregulated in human hematologic malignancies. For example, 10%–27% of patients with T-cell acute lymphoblastic leukemia (T-ALL) and 1.5% of patients with B-cell acute lymphoblastic leukemia (B-ALL) have tumor cells with activating JAK1 mutations (Flex et al., 2008; Jeong et al., 2008; Mullighan et al., 2009b; Zhang et al., 2012; Li et al., 2017). Up to 2% of acute myeloid leukemia (AML) patients carry cells with JAK1 mutations (Tomasson et al., 2008; Xiang et al., 2008). In 8%–12.5% of T-cell prolymphocytic leukemia (T-PLL) cases, JAK1 mutations occur in the tumor cells (Bellanger et al., 2014; Springuel et al., 2015; Wahnschaffe et al., 2019).

Various residues in JAK1 are mutation hotspots (Fig. 1), and additional mutations are continuously updated in public databases, e.g., Genomic Data Commons of the

National Cancer Institute (Grossman et al., 2016; Jensen et al., 2017), Catalogue of Somatic Mutations in Cancer (Tate et al., 2019), and cBioPortal for Cancer Genomics (Cerami et al., 2012; Gao et al., 2013).

Aberrant JAK1 signaling is linked to dysfunctional lymphopoiesis (Chen et al., 2012). For instance, V658F and S646P mutations occur in the JH2 domain of JAK1 (Fig. 1) and impair its autoinhibitory functions on the kinase domain. This leads to constitutively active STAT3 and MEK–extracellular signal-regulated kinases (ERK) signaling and allows a cytokine-independent proliferation of malignant B-ALL and T-ALL cells in vitro and in mice (Jeong et al., 2008; Hornakova et al., 2009; Mullighan et al., 2009a; Li et al., 2017). The finding that difficult-to-treat adult B-ALL cells are highly vulnerable toward the ATP-competitive JAK1/JAK2 inhibitor ruxolitinib verifies a disease relevance of JAK1^{S646P} (Li et al., 2017).

Certain JAK1 mutations, such as A634D and R724H, are characteristic for T-ALL (Flex et al., 2008), whereas the V623A mutation occurs in AML (Xiang et al., 2008). Tumor cell-specific mutations of JAK1 are not only associated with tumorigenesis but also with therapy outcome. This is exemplified by patients with JAK1 mutations frequently showing resistance to chemotherapy (Li et al., 2017). These findings illustrate

TABLE 1
Human histone deacetylases, their cytogenetic and subcellular localization, and selected targets

Group	Class	HDAC	Cytogenetic Locus ^a	Subcellular Localization ^b	Examples of Targets ^b
Zn ²⁺ -Dependent	I	HDAC1	1p35.2-p35.1	Nucleus	CtIP, DNMT1, E2F1, histones, NF-κB members, p53, PR130, pRB, STAT3
		HDAC2	6q21	Nucleus	GATA2, histones, NF-κB members, p53, PR130, pRB, STAT3
		HDAC3	5q31.3	Nucleus, cytoplasm	DNMT1, GATA1, histones, MYC, NF-κB members, p53, pRB, STAT1, STAT3
	IIa ^c	HDAC8	Xq13.1	Nucleus, cytoplasm	Actin, histones, HSP70, SMC3
		HDAC4	2q37.3	Nucleus, cytoplasm	GATA1, histones, HP1, p53, STAT1
		HDAC5	17q21.31	Nucleus, cytoplasm	ER, histones, HP1, SMAD7
		HDAC7 HDAC9	12q13.11 7p21.1	Nucleus, cytoplasm Nucleus, cytoplasm	HIF1α, histones Histones, STAT5, TRIM29, USF1
	IIb	HDAC6	Xp11.23	Nucleus, cytoplasm	HSP70, HSP90, PRX I and II, α-tubulin
	IV	HDAC10	22q13.33	Nucleus, cytoplasm	MSH2, spermidine
		HDAC11	3p25.1	Nucleus, cytoplasm	BUBR1, CDC25A, E2F1, FOXP3, histones
	NAD ⁺ -Dependent	III	SIRT1	10q21.3	Nucleus, cytoplasm, mitochondria
SIRT2			19q13.2	Nucleus, cytoplasm	FOXO, histones, p300, p53, PGC1α, PPARγ, α-tubulin
SIRT3			11p15.5	Mitochondria	ACS2, FOXO, histones, p53, PGC1α, XRCC6
SIRT4 SIRT5			12q24.23-q24.31 6p23	Mitochondria Nucleus, cytoplasm, mitochondria	Histones, mTORC1 CPS1, cytochrome c, histones
SIRT6			19p13.3	Nucleus, endoplasmic reticulum	HIF1α, histones, TNFα
SIRT7			17q25.3	Nucleus, cytoplasm	Histones, p53, RNA polymerase I

ACS2, acetyl-CoA synthetase-2; BUBR1, BUB1-related protein-1; CDC25A, cell division cycle-25A; CPS1, carbamoyl phosphate synthase-1; CtIP, CtBP-interacting protein; DNMT, DNA methyltransferase; E2F, E2F family of transcription factors; ER, estrogen receptor; FOXO, forkhead box protein O; FOXP3, forkhead box protein P-3; GATA, globin transcription factor; HIF1, hypoxia-inducible factor-1; HP1, heterochromatin protein-1; HSP70, heat shock protein 70 kDa; mTORC1, mechanistic target of rapamycin kinase complex-1; MYC, myelocytomatosis oncogene; PGC1, peroxisome proliferator-activated receptor γ coactivator 1; PPARγ, peroxisome proliferator-activated receptor γ; PR130, protein phosphatase 2A regulatory subunit B' PPP2R3A; PRX, peroxiredoxin; SMAD, suppressor of mothers against decapentaplegic; SMC3, structural maintenance of chromosomes protein-3; TNF, tumor necrosis factor; TRIM29, tripartite motif containing-29; USF1, upstream stimulatory factor-1; XRCC6, X-ray repair cross complementing-6.

^aAccording to Ensembl database (www.ensembl.org) and NCBI (www.ncbi.nlm.nih.gov).

^bAccording to the UniProtKB/Swiss-Prot knowledge database (www.uniprot.org).

^cThere are reports in the literature doubting significant catalytic activities of class IIa HDACs (Lahm et al., 2007). Recent data illustrate that these HDACs have reader function (Zhang et al., 2022).

that JAK1 mutations contribute to both the development and the aggressiveness of blood disorders.

Solid tumors can also carry mutant JAK1 molecules. Fusions, rearrangements, missense mutations, nonsense mutations, silent mutations, frameshift deletions, and insertions in JAK1 occur in endometrial cancers, intestinal cancers, and stomach cancers (Albacker et al., 2017). Moreover, JAK1 is mutated in approximately 1.8% of patients suffering from breast carcinoma, uterine corpus neoplasm, colorectal adenocarcinoma, non-small-cell lung carcinoma, and prostate cancer (AACR Project GENIE Consortium, 2017), as well as in 3.8% of hepatocellular carcinoma cases

and 11.5% of endometrial cancer patients (Cerami et al., 2012; Gao et al., 2013; Kan et al., 2013). Further research is required to comprehensively identify the therapeutic potential of such aberrations in these and other solid tumors (Qureshy et al., 2020).

B. Janus Kinase-2

In 2001 and updated in 2008, the World Health Organization categorized polycythemia vera (PV), essential thrombocythemia (ET), and idiopathic myelofibrosis (MF) together with chronic myeloid leukemia and some seldom leukemia subtypes into myeloproliferative neoplasms (MPNs) (Tefferi and Vardiman, 2008; Tefferi et al., 2009; Vannucchi

et al., 2009; Vardiman et al., 2009; Helbig, 2018; Marcellino et al., 2020). PV, ET, and MF are clonal hematopoietic stem cell malignancies with increased self-renewal capacity and a lack of the leukemia fusion protein BCR-ABL, which defines chronic myeloid leukemia. The somatic point mutation JAK2^{V617F} occurs in over 95% of patients with PV and in 50% of patients with ET or MF. Mutant JAK2 also occurs in 20% of other MPNs, such as refractory anemia with ring sideroblasts and thrombocytosis (Patnaik and Tefferi, 2015; Orazi et al., 2017). PV and ET patients have increased numbers of erythroid or megakaryocyte precursors, respectively. Fibrotic bone marrow, hemorrhage, splenomegaly, and enhanced cytokine-induced JAK1/JAK2 signaling are hallmarks of MF (Thiele et al., 2005; Barosi et al., 2007; Tefferi and Vardiman, 2008; Yönal et al., 2016). JAK2^{V617F} manifests in ~5% of AML and myelodysplastic syndrome cases (Renneville et al., 2006; Steensma et al., 2006; Verstovsek et al., 2006; Levine et al., 2007).

In 2005, JAK2^{V617F} was discovered in tumor cells from patients with PV, ET, and MF (Fig. 1). This aberration stems from a guanine-to-thymidine substitution that replaces valine with phenylalanine at codon 617 within the JH2 domain of JAK2 (Baxter et al., 2005; James et al., 2005; Kralovics et al., 2005; Levine et al., 2005). Without a need for cognate receptor stimulation, dimeric JAK2^{V617F} triggers tyrosine phosphorylation of the prosurvival JAK-STAT, phosphatidylinositol 3-kinase/serine-threonine protein kinase (AKT), and MAPK/ERK proteins as the root cause of aberrant cell growth and survival (James et al., 2005; Kralovics et al., 2005; Levine et al., 2005; Gorantla et al., 2010) (Fig. 2). Accordingly, ruxolitinib produces clinical responses in patients suffering from JAK2^{V617F}-associated neoplasms (Barosi et al., 2014; Bose and Verstovsek, 2017b; Cervantes and Pereira, 2017).

Mutant JAK2 not only alters the initiation of cytoplasmic signaling cascades. JAK2^{V617F}-positive cells show a high frequency of genomic instability, which is associated with an increased risk of transformation into aggressive AML (Theocharides et al., 2007; Beer et al., 2010; Aynardi et al., 2018). This serious adverse outcome is seen in 5% of patients with ET or PV (Finazzi et al., 2005; Wolanskyj et al., 2006) and in 15%–30% of primary MF patients (Dupriez et al., 1996). These incidences markedly increase upon genotoxic treatment (Finazzi et al., 2005; Björkholm et al., 2011). This may be due to the capacity of JAK2^{V617F} to suppress cytotoxic programs that are triggered upon chemotherapy-induced DNA replication stress and DNA damage by an upregulation of DNA repair enzymes (Plo et al., 2008; Kurosu et al., 2013; Ueda et al., 2013; Chen et al., 2015; Karantanos and Moliterno, 2018). Thus, alternative therapies that early and effectively eliminate JAK2^{V617F} could prevent aggressive disease progression.

Mutant JAK2 additionally regulates tumorigenesis via the phosphorylation of the residue tyrosine-41 in histone H3. This displaces heterochromatin protein-1 α and abrogates its tumor-suppressive functions (Shi et al., 2006; Dawson et al., 2009; Griffiths et al., 2011). It needs to be considered, though, that other kinases, such as the cell cycle regulatory kinase WEE1 and activated CDC42-associated kinase-1 (ACK1), also catalyze the phosphorylation of histones and thereby modulate the transcription of multiple genes that are relevant to cellular homeostasis (Mahajan et al., 2012; Mahajan et al., 2017; Kim et al., 2020). For instance, the WEE1-mediated phosphorylation of histone H2B at tyrosine-37 during the S phase of the cell cycle suppresses the transcription of histone genes (Mahajan et al., 2012). ACK1 phosphorylates histone H4 at tyrosine-88 and consequently activates transcription of androgen receptor. This drives the progression of prostate cancer to a castration-resistant state (Mahajan et al., 2017; Sawant et al., 2022).

Tightly controlled JAK2 signaling is also relevant for metabolic integrity. JAK2^{V617F} and the JAK2 exon 12 (N542-E543del) mutation, which drive increased erythropoiesis, create a systemically dysregulated metabolic profile with highly elevated mRNA and protein levels of the key glycolytic enzyme 6-phosphofructo-2-kinase/fructose-2,6-biphosphatase-3. Accordingly, this enzyme is a druggable vulnerability of cells with mutant JAK2 (Rao et al., 2019).

JAK2 mutants are likewise involved in chronic myelomonocytic leukemia (CMML). The JAK2 inhibitor TG101209, which was developed to treat MPN cells with JAK2^{V617F} and MPL^{W515L/K} (Pardanani et al., 2007), halted the spontaneous growth of granulocyte-macrophage colony-forming units of cells from CMML patients (Geissler et al., 2016; Padron et al., 2016).

JAK2 mutations are not limited to myeloid leukemia but likewise occur in 8.5% of high-risk B-ALL and in 18%–28% of B-ALL cases associated with Down syndrome (Gaikwad et al., 2009; Kearney et al., 2009). A majority of JAK2 mutations in the lymphoid lineage affect the R683 residue in JH2 (Fig. 1). Overexpression of cytokine receptor-like factor-2/thymic stromal lymphopoietin receptor (CRLF2) is prominent in JAK2^{I682F} and JAK2^{R683G} cells of patients with clinically unfavorable types of acute lymphoblastic leukemia (ALL). The association between these JAK mutants and CRLF2 propels ligand-independent cell proliferation (Mullighan et al., 2009a; Savino et al., 2017; Chang et al., 2021). Below, we summarize that such therapeutically challenging lesions create a remarkable vulnerability to HDACi.

The disease relevance of JAK2-dependent signaling likewise became evident with the JAK2^{L611S} mutation in the JH2 domain of JAK2. This mutation was

identified in a child with B-ALL (Fig. 1) and not detected in the bone marrow upon remission, suggesting that JAK2^{L611S} drove this cancer (Kratz et al., 2006). Therefore, extended screening for JAK2 mutations in childhood ALL may deliver insights into an actionable drug target.

Mutations in JAK2 are not limited to blood malignancies. JAK2 alterations occur in about 2.65% of all cancers, including non-small-cell lung cancer (NSCLC), breast carcinoma, and colorectal adenocarcinoma (AACR Project GENIE Consortium, 2017). Although JAK inhibitors (JAKi) are effective against cells from various solid tumors in preclinical studies, it is currently not solved if such drugs provide a benefit for patients with solid tumors (Qureshy et al., 2020).

C. Janus Kinase-3

JAK3 mutations occur in 12% of juvenile CMML (Sakaguchi et al., 2013; Springuel et al., 2015) and in 15% of acute megakaryoblastic leukemia cases (Walters et al., 2006; Malinge et al., 2008; Springuel et al., 2015) (Fig. 1). These aberrations lead to constitutively active JAK3 that confers growth factor-independent proliferation, being a hallmark of tumorigenesis (Walters et al., 2006; Malinge et al., 2008; Riera et al., 2011; Vainchenker and Constantinescu, 2013). Genomic profiling of a wide variety of T-cell neoplasms showed that up to one-third of patients had genomic aberrations that gave rise to gain-of-function (GOF) mutations in JAK1 and JAK3 (Springuel et al., 2015; Greenplate et al., 2018). JAK3 was found mutated in 10%–16% of T-ALL patients (Zhang et al., 2012; De Keersmaecker et al., 2013; Vicente et al., 2015; Li et al., 2016; Girardi et al., 2017; Liu et al., 2017; Degryse et al., 2018a; Greenplate et al., 2018) and in 36%–42% of T-PLL cases (Bellanger et al., 2014; Springuel et al., 2015; Wahnschaffe et al., 2019).

Further studies revealed JAK3 mutations in 1.8% of all cancers. NSCLC, colorectal adenocarcinoma, breast carcinoma, melanoma, and uterine corpus neoplasm have the greatest prevalence of genetic alterations in the *JAK3* gene (AACR Project GENIE Consortium, 2017). Further research will define whether they are therapeutic targets.

D. Tyrosine Kinase-2

Similar to the other JAKs, activating point mutations at the *TYK2* locus predominantly accumulate in the JH1 and JH2 domains (Leitner et al., 2017; Hammarén et al., 2019) (Fig. 1). The first reported GOF mutation of *TYK2* was V678F. This aberration is homologous to JAK2^{V617F} but has not yet been reported in patients (Staerk et al., 2005; Gakovic et al., 2008). This finding suggests that members of the JAK family have common as well as very individual functions in tumorigenesis. Such a lack of functional redundancy might explain the highly divergent mutation spectra of JAKs in cancer cells.

Two GOF *TYK2* germline mutations (P760L and G761V) were identified in leukemic pediatric patients (Fig. 1). These mutations are in the JH2 pseudokinase domain of *TYK2* and likely constrict its autoinhibitory function on the JH1 kinase domain (Waanders et al., 2017). Sequence analysis of 17 cell lines and 45 pediatric T-ALL patient samples identified further activating mutations of *TYK2* (G36D, S47N, V731I, E957D, R1027H) (Fig. 1). Such aberrations can promote cell survival through the inducible transcription factors STAT1 and STAT3 and activating effects on the antiapoptotic BCL2 proteins (Sanda et al., 2013; Wingelhofer et al., 2018; Wöss et al., 2019).

Activation of wild-type *TYK2*-STAT1 signaling and ensuing expression of the antiapoptotic BCL2 protein as well as germline *TYK2* mutations (A53T, A81V, R197H, V362F, G363S, I684S, R703W, A928V, A1016S, P1104V) are linked to the development of ALL and AML (Kaminker et al., 2007; Tomasson et al., 2008; Sanda et al., 2013). *TYK2*^{P1104A} was detected in breast, liver, stomach, and poor-prognosis malignant peripheral nerve sheath tumors (Hirbe et al., 2017). Although *TYK2*^{P1104A} might inactivate *TYK2* (Kaminker et al., 2007), increased levels of *TYK2* correlate with cell migration and metastasis of prostate cancer cells (Ide et al., 2008; Santos et al., 2015). These data illustrate context-dependent tumor-promoting activities of wild-type and mutant *TYK2*.

Taken together, the above reports impressively demonstrate that a failure to control JAK-STAT signaling is a hallmark feature of early and late stages of cancer development.

III. Histone Deacetylases Are Valid Pharmacological Targets in Janus Kinase Mutant Tumor Cells

Hematopoietic and solid tumor cells frequently display overactive and overexpressed HDAC levels when compared with normal tissues (Fig. 3A). Consequently, acetylation patterns of histones and nonhistone proteins, chromatin compaction, and gene expression are dysregulated. These processes determine cell proliferation and cell fate at multiple layers, including hallmarks of tumorigenesis (Fig. 3, B and C). For example, aberrant HDAC activity restricts the expression of genes encoding tumor suppressors and cell differentiation proteins (Krämer et al., 2014; Wagner et al., 2014; Ceccacci and Minucci, 2016; Imai et al., 2016).

For decades, these important physiologic functions of HDACs have encouraged the development and testing of small molecules that target HDACs (Figs. 3A and 4A). Initial HDACi were identified by their ability to induce histone acetylation before the discovery of the first HDAC in 1996 (Taunton et al., 1996). These compounds of natural and microbial origins include butyrate,




Fig. 3. HDACs are dysregulated in cancer and control chromatin remodeling with histone acetyl transferases (HATs). (A) Cancer cells display altered HDAC expression and activity. This creates a vulnerability of transformed cells to HDACi. (B) HATs catalyze the acetylation of lysine moieties of histone and nonhistone proteins using acetyl-CoA as a co-substrate. The acetyl groups neutralize the positively charged histone tails and thereby weaken the electrostatic interactions between histones and the negatively charged phosphate backbone of DNA. This causes a relaxed state of chromatin. In a chromatin context-dependent manner, this can render DNA more accessible for transcription (depicted as accelerated speed). Elimination of the acetyl groups by HDAC activity reverses these processes. HDACs catalyze the removal of the acetyl group from the lysine residues, causing the release of the acetate molecule. These alterations in transcription critically affect key cellular processes. (C) Biochemistry of the acetylation/deacetylation reactions. Created with BioRender.com.

trapoxin, and trichostatin A (TSA) (Hagopian et al., 1977; Yoshida et al., 1990; Kijima et al., 1993; Krämer et al., 2001). Crystal structures of HDACs were

determined from 1999 on (Finnin et al., 1999). These allowed the development of an increasingly specific arsenal of HDACi based on structure-activity-relationship.

In the 1990s, butyrate and phenylbutyrate were the first HDACi that were applied in compassionate protocols to patients suffering from leukemia (Krämer et al., 2001). The discovery that the antiepileptic, mood-stabilizing drug valproic acid inhibits class I HDACs in vitro and in experimental animals (Göttlicher et al., 2001) illustrated that an agent that increased protein acetylation could be given safely to patients for decades (Michaelis et al., 2007). The US and China Food and Drug Administration (FDA) agencies have approved HDACi for the treatment of poor-prognosis cutaneous T-cell lymphoma (CTCL) and peripheral T-cell lymphoma. These drugs are vorinostat (SAHA), romidepsin (FK228), belinostat (PXD101), and chidamide/tucidinostat (Piekarz et al., 2009; Food and Drug Administration, 2010; Whittaker et al., 2010; Li et al., 2019; Cappellacci et al., 2020). Clinically validated HDACi target Zn²⁺-dependent HDACs (Nebbio et al., 2017; Beyer et al., 2019; Cappellacci et al., 2020) (www.clinicaltrials.org).

HDACi can be broad and isoenzyme-specific inhibitors (Fig. 4A). For example, the hydroxamic acids vorinostat, belinostat, and panobinostat inhibit HDACs belonging to classes I, IIa/b, and IV (pan-HDACi) at nano- to low micromolar levels. The short-chain fatty acid valproic acid selectively inhibits class I HDACs, whereas butyrate and its derivatives inhibit preferentially classes I and IIa at millimolar levels. The benzamide entinostat (MS-275) blocks the class I HDACs HDAC1, HDAC2, and HDAC3 at low micromolar levels, and the nanomolar HDACi romidepsin additionally targets HDAC8 (Fig. 4A) (Bradner et al., 2010; Lechner et al., 2022). Current approaches often focus on hydroxamic acid-based compounds for an improved design of HDACi. This relies on the ability of hydroxamic acids to block HDACs at low concentrations in vitro and in cells as well as the clinical experience with these HDACi.

The pharmacological warhead of HDACi is a Zn²⁺-binding group (Krämer, 2009), as exemplified for hydroxamic acids in Fig. 4B. On the other end of the drug, a cap group has residues for interactions with the rim of the enzyme. A linker of variable length connects these two moieties (Fig. 4B). This structure makes HDACi amenable for poly-pharmacophore design. The cap group can provide residues to anchor additional pharmacological moieties and exit vectors to attach independent pharmacophores. This can be the attachment of TKi (Fig. 4C). In contrast, the Zn²⁺-binding group must be preserved to maintain inhibitory activity against HDACs.

Individual structures of the HDACs (length and depth of the catalytic cavity, amino acid side chains at the entrance site of the catalytic cavity) have allowed the design of selective and specific HDACi. This has been achieved by varying the cap group, the length of the linker, and the type of warhead (Krämer et al.,

2014). For example, tetrahydro- β -carboline-based hydroxamic acid HDACi with large cap groups specifically block the broad and shallow catalytic groove of the second catalytic domain of HDAC6 (17.5 Å). These HDACi selectively inhibit HDAC6 at nanomolar concentrations (Sellmer et al., 2018) and poorly enter class I HDACs because of their narrower catalytic channel rim (~12.5 Å) (Butler et al., 2010). Researchers highly appreciate these and other HDACi as pharmacological tools to define individual functions of HDACs in vitro and in vivo. In addition, specific HDACi are expected to kill certain tumor cells with less adverse effects on normal cells from the same tissue (Michaelis et al., 2007; Krämer, 2009; Juengel et al., 2011; Falkenberg and Johnstone, 2014; Krämer et al., 2014; Koeneke et al., 2015). It will be interesting to see the full clinical potential of such HDACi.

A. Janus Kinase–Signal Transducer and Activator of Transcription Pathways Control Cellular Susceptibility to Histone Deacetylase Inhibitors

JAK-STAT signaling is not only a paradigm for an intracellular signaling pathway but also a determinant for the cellular susceptibility to changes in protein acetylation patterns. In CTCL and diffuse large B-cell lymphoma (DLBCL) cell lines and ex vivo patient samples, aberrant activation of the JAK-STAT pathway confers resistance to HDACi (Fantin et al., 2008; Smith et al., 2010; McKinney et al., 2014; Cortes et al., 2021). A vorinostat-phase IIb clinical trial revealed by immunohistochemistry of skin biopsies that nonresponding CTCL patients carried higher expression levels and activation of STAT1, STAT3, and STAT5. Consistent herewith, RNA interference (RNAi) against STAT3 or STAT5 synergistically induced apoptosis when combined with vorinostat in cultured lymphoma cells (Fantin et al., 2008). These findings encouraged a synchronized inhibition of JAKs and HDACs. A combination of vorinostat with the broad-range, ATP-competitive JAKi pyridone-6 caused growth arrest and triggered apoptosis in CTCL cells with primary robustness to vorinostat due to activated STATs (Fantin et al., 2008). Similarly, inhibition of JAK1 and JAK2 using ruxolitinib or momelotinib overcame the limited response of a subset of CTCL cells toward romidepsin in vitro and in xenografts (Cortes et al., 2021). These data are in line with findings from DLBCL, where concurrent inhibition of JAK-STAT and HDACs using ruxolitinib and panobinostat showed obvious synergy and could tremendously reduce tumor growth in vitro as well as in a xenograft model (McKinney et al., 2014; Davis et al., 2018).

In JAK2^{V617F}-positive MPN cells, concurrent treatment with ruxolitinib and vorinostat resulted in G1-phase cell cycle arrest, apoptotic cell death, and inhibition of colony-formation capacity. This was due to the attenuated expression of phosphorylated and total JAK2^{V617F} and significantly reduced phosphorylation of STAT3 and AKT





Fig. 4. HDAC classes, HDACi, and mode of action. (A) Schematic representation of different classes of HDACs and their pharmacological inhibitors. (B) Targeting the Zn^{2+} -dependent HDACs by HDACi. The structure of hydroxamic acids is shown as an example of the three typical parts of an HDACi. The cap group binds the HDAC at its rim, and the linker serves to position the Zn^{2+} -binding part toward the coordinated metal in the catalytic cleft of the HDAC. (C) The double-hit approach uses JAK-HDAC dual inhibitors. The cap group provides possibilities to attach a TKi. The TKi must preserve its structures that compete with ATP and the HDACi must retain the Zn^{2+} -binding moiety. Ac-K, acetylated lysine residue in a protein.



(Hao et al., 2020). These effects are linked to the suppressors of cytokine signaling (SOCS) proteins, which are negative regulators of JAK-STAT signaling (Jiang et al., 2017a) (Fig. 2). The inhibition of JAK2^{V617F} by ruxolitinib and vorinostat upregulated *SOCS3*. Furthermore, HDACi suppressed the mRNA levels of the antiapoptotic STAT3 target genes *BCL2*, *BIRC5*, *MYC*, and the cell cycle regulator *CCND1*, which can explain the above-mentioned cell cycle arrest in G1. This data shows that ruxolitinib and vorinostat combinations promise better therapeutic outcomes for patients with MPN (Hao et al., 2020). Below, we discuss clinical trials using HDACi and TKi in combination.

Unlike in the above-named tumor types, combinations of ruxolitinib and the hydroxamic acid-based pan-HDACi givinostat (ITF2357) did not induce more apoptosis than givinostat alone in B-ALL cells with CRLF2 overexpression. This deviation may rely on the

resistance of CRLF2/JAK2-mutated cells to type 1 (e.g., ruxolitinib, baricitinib; bind JAKs in the active conformation) and type 2 (CHZ-868; stabilizes the inactive conformation of JAKs) JAKi. Irrespective thereof, givinostat potently inhibited the growth and survival of such cells (Savino et al., 2017).

As in MPNs, JAKi and HDACi interact favorably against breast cancer cells. Vorinostat increased histone acetylation at the *leukemia inhibitory factor receptor (LIFR)* gene promoter. This subsequently recruited bromodomain-containing protein-4 (BRD4) to induce the expression of *LIFR*. This cytokine activates prosurvival JAK-STAT3 signaling in auto- and paracrine fashions. Inhibition of JAKs or BRD4 sensitized breast cancer to HDACi, indicating that JAKs and BRD4 promote an undesired cytoprotective effect in this setting. Thus, combinations of HDACi with inhibitors of JAKs or BRD4 appear as potential therapies for breast tumors

(Zeng et al., 2016). Further studies will show whether JAK-STAT activation protects additional tumor types from HDACi.

Linear correlations between tonic activation of JAK-dependent STAT3 phosphorylation and HDACi resistance are, though, unlikely. For example, DLBCL cells with constitutively tyrosine-phosphorylated STAT3 were more sensitive to panobinostat than those cells lacking detectable STAT3 phosphorylation. In this system, chronic STAT3 phosphorylation was abrogated by HDACi-induced STAT3 hyperacetylation (Gupta et al., 2012), reminiscent of the inhibitory STAT1 phosphorylation-acetylation switch (Krämer et al., 2009). Furthermore, an unbiased screen with a panel of tumor cells demonstrated that PV cells with hyperactive JAK2^{V617F}-dependent signaling cascades displayed the highest sensitivity to vorinostat (Liang et al., 2022). It is tempting to speculate that context-dependent high tonic levels of JAK-STAT signaling create a vulnerability for pharmacological approaches involving HDACi. Such considerations are relevant because clinically validated markers for the stratification of patients into responders and nonresponders to HDACi are not available.

B. Histone Deacetylase Inhibitors Can Correct Aberrant Janus Kinase–Signal Transducer and Activator of Transcription Signaling

In the following section, we sum up evidence on how HDACi decrease aberrant JAK signaling, the underlying molecular mechanisms for this, and potential therapeutic implications of the requirement of HDAC activity for JAK-dependent signaling.

1. Impact of Histone Deacetylase Inhibitors on Janus Kinase-1. In 2004, it was revealed that pretreatment of colon cancer cells with structurally divergent HDACi suppressed cytokine-induced JAK-STAT signaling. The carboxylic acid butyrate and the hydroxamic acids TSA and vorinostat abrogated the activating phosphorylation of JAK1 on Y1022/Y1023 upon treatment with the type II interferon (IFN)- γ (Klampfer et al., 2004). This impaired the IFN-induced phosphorylation of STAT1 at Y701/S727, subsequently caused a failure of STAT1 to translocate to the nucleus, and blunted IFN γ -dependent gene expression. Both overexpression of HDAC1, HDAC2, and HDAC3, as well as RNAi-mediated depletion of these enzymes, illustrated that they promote JAK-STAT activation and the expression of IFN γ target genes. These effects were not due to a caspase-dependent degradation of STAT1 (Klampfer et al., 2004; Owusu and Klampfer, 2017), which can occur in leukemic cells undergoing apoptosis in response to HDACi (Licht et al., 2014).

In contrast to this antagonistic relationship between HDACi and IFN γ , both drugs synergistically suppressed the survival of colon cancer cells and reduced the antiapoptotic BCL-XL protein (Klampfer

et al., 2004). This finding may point to a therapeutic option based on HDACi and IFNs. Studies illustrating apoptosis induction by HDACi and IFNs in human melanoma cells support this idea (Krämer et al., 2006; Roos et al., 2011). Moreover, butyrate levels of up to ~ 7 mM in the colon can be anti-inflammatory and chemopreventive due to a suppression of STAT1 signaling and prosurvival factors (Klampfer et al., 2004; Stempelj et al., 2007). Unlike normal cells, cancer cells often rely on glycolysis to gain energy, instead of the mitochondrial citrate cycle (Warburg effect). This makes cancer cells unable to fuel butyrate into this metabolic pathway. Consequently, butyrate accumulates and causes protein hyperacetylation and antiproliferative activities (Meyer et al., 2021).

2. Impact of Histone Deacetylase Inhibitors on Janus Kinase-2. The oncogenic role of JAK2^{V617F} spurred the development of small-molecule inhibitors of hyperactive JAK2 (Zhao et al., 2016a; Musumeci et al., 2019). The success and survival benefit, along with its striking efficacy regarding spleen volume and symptom reduction, have made ruxolitinib a meaningful therapy in MPNs. Since 2011, ruxolitinib has become the main drug for the treatment of MPNs refractory to the ribonucleotide reductase inhibitor hydroxyurea (Bose and Verstovsek, 2017a). However, ruxolitinib is not fully curative in most cases. This particularly applies to MF patients, who suffer from short survival, and to patients with JAK2 mutant cells that are insensitive to JAKi (Barosi et al., 2014; Bose and Verstovsek, 2017b; Cervantes and Pereira, 2017; Chang et al., 2021). Accordingly, ongoing research strives to identify improved drugs for the treatment of MF patients.

a. Histone deacetylase inhibitors attenuate JAK2^{V617F} in blood malignancies. An alternative, indirect route to target mutant JAK2 in patients suffering from MPNs relies on HDACi (Bose and Verstovsek, 2017b). Givinostat selectively reduced the protein levels of JAK2^{V617F} but not wild-type JAK2 in cell lines and primary human MPN cells. Congruent herewith, JAK2 wild-type cells became the dominant population in heterogeneous patient samples containing wild-type and mutant JAK2 (Guerini et al., 2008). The HDACi-induced attenuation of JAK2^{V617F} occurred by a yet unclear mechanism and despite unchanged mRNA expression (Guerini et al., 2008). Remarkably, givinostat produced promising results in a clinical trial with 29 patients suffering from JAK2^{V617F}-positive blood disorders. Givinostat decreased the allelic burden of JAK2^{V617F}, proving on-target activity against MPN cells (Rambaldi et al., 2010). The givinostat-induced loss of JAK2^{V617F} tied in with anticancer activity in patients with MPNs that were unresponsive to the maximally tolerated dose of hydroxyurea (Finazzi et al., 2013). It similarly appears promising that in an ongoing 4-year study involving 54

MPN patients (51 PV cases; 42–80 years old; trial #NCT01761968), a maximally tolerated dose of 200 mg orally taken givinostat per day has produced over 80% overall response rates. This treatment has been well tolerated, with at most 10% of grade 3 events that have occurred in patients who have concomitantly received hydroxyurea (Rambaldi et al., 2021). These clinical data support the long-term use of givinostat in patients.

Like givinostat, vorinostat, panobinostat, and a further hydroxamic acid-based pan-HDACi, pracinostat (SB939) (Fig. 4A), produce antimyeloproliferative activities that are molecularly linked to reduced JAK2^{V617F}. Vorinostat significantly hampered the aberrant phosphorylation of JAK2^{V617F} and of its downstream targets STAT5, STAT3, AKT, and ERK1/ERK2 in murine and human PV cells (Akada et al., 2012). This finding agrees with remarkable vorinostat-evoked hematologic responses in a JAK2^{V617F} knock-in mouse PV model and a significantly reduced mutant allele burden. Interestingly, vorinostat was more potent than pyridone-6 against erythroid progenitors in this model (Akada et al., 2012). In a phase II clinical trial involving patients with PV and ET, vorinostat demonstrated high efficacy against JAK2^{V617F}-positive peripheral blood mononuclear cells (Andersen et al., 2013). Likewise, panobinostat was effective against MPNs alone or in combination with JAKi. Panobinostat showed clinical efficacy in patients with wild-type JAK2- or JAK2^{V617F}-positive MF and decreased the JAK2^{V617F} allele burden (DeAngelo et al., 2013a; Mascarenhas et al., 2013; Mascarenhas et al., 2017). Pracinostat decreased JAK2^{V617F}, STAT5, and, more pronouncedly, their phosphorylation in cultured PV and ET cells. This was associated with apoptosis-related processing of poly-ADP-ribose polymerase-1 (PARP1). Both pracinostat and the dual JAK2/FMS-like tyrosine kinase-3 (FLT3) inhibitor pacritinib significantly reduced the growth of such tumor cells in mice. Combined application of pracinostat and pacritinib pronounced these effects against ET xenografts (Novotny-Diermayr et al., 2012).

An observation in clinical trials using HDACi is that lower drug doses over longer time are often well tolerated by patients. Moreover, patients with less advanced disease achieve better therapeutic responses. Another recurrent finding in the clinic is that subgroups of patients are particularly amenable to treatment with HDACi (DeAngelo et al., 2013a, 2013b; Mascarenhas et al., 2013; Mascarenhas et al., 2017). Unraveling the molecular features of their malignancies will show which patients can profit from HDACi.

Noteworthy, wild-type JAK2 status does not preclude leukemic cell killing by JAKi and HDACi. Concurrent treatment with low doses of ruxolitinib and vorinostat acted synergistically against six out of 12 cell lines from hematologic malignancies with no known mutations of JAK2. This drug combination triggered reactive oxygen species-mediated tumor cell death (Civallero et al., 2017).

B-ALL represents about one-third of pediatric leukemia, and among them, 7% have overexpressed CRLF2 due to its translocation to strong promoter elements. As mentioned above, B-ALL cells with increased CRLF2 expression and mutant JAK2 are insensitive to JAK2 inhibitors. Promisingly, givinostat was effective against permanent and fresh human B-ALL cells with mutations in CRLF2 and JAK2 (JAK2^{I682F} and JAK2^{R683G}) (Fig. 1) (Savino et al., 2017). Unlike in PV cells (Guerini et al., 2008), givinostat decreased JAK2^{V617F} mRNA in CRLF2/JAK2 mutant human B-ALL cells. Moreover, givinostat increased the mRNA of PTPN1, which is the founding member of the protein tyrosine phosphatase family and an inhibitor of JAK-STAT signaling (Savino et al., 2017). Although the protein levels of mutant JAK2 and PTPN1 were not determined in this study, it illustrates that the B-ALL-associated cooccurrence of dysregulated CRLF2/JAK2 is a pharmacodynamic marker for the high efficiency of givinostat.

In contrast to the reported specific loss of mutant and not wild-type JAK2 in givinostat-treated leukemic cells (Guerini et al., 2008), the benzamide HDACi chidamide, which selectively inhibits HDAC1, HDAC2, HDAC3, and HDAC10 (Fig. 4A), decreased wild-type JAK2 and JAK2^{V617F} mRNA and protein levels in PV and myelodysplastic syndrome cells. This loss of JAK2^{V617F} was associated with decreased STAT3 phosphorylation, cell cycle arrest, apoptosis, and upregulation of SOCS3 mRNA and protein expression. The authors of this work concluded that SOCS3 accumulation, likely upon hyperacetylation of histones in its gene promoter, attenuated wild-type and mutant JAK2 (Zhao et al., 2016b). The possibility of eliminating JAK2^{V617F} and wild-type JAK2-STAT3-dependent signaling with a class I HDACi might offer a clinically safer treatment due to fewer side effects than with pan-HDACi. It is yet unclear whether pharmacological inactivation of class I HDACs and/or the class IIb enzyme HDAC10 by chidamide decreased JAK2.

Studies with the clinically tested hydroxamic acid citarinstat, which has some preference for HDAC6 (Lechner et al., 2022), did not attenuate total and constitutively phosphorylated JAK2 in a panel of lymphoid tumor cells (Cosenza et al., 2020). This data suggests that JAK2 stability does not depend on the catalytic activity of HDAC6. Curiously, highly synergistic cytotoxic combinations of momelotinib and citarinstat decreased p-JAK2 but spared total JAK2 levels. We speculate that such HDACi/TKi combinations unleash yet undefined molecular mechanisms that selectively target a small pool of p-JAK2. The size of this pool varies between cells, and it is conceivable that only a few of these highly catalytically active kinase molecules can modulate biologic processes. Such a low stoichiometry of activation corresponds to

strong biologic effects that small fractions of activated STAT1 can induce (Göder et al., 2021).

Although these data suggest exploiting HDACi for the treatment of preleukemic conditions and leukemia driven by JAK2^{V617F}, additional research is necessary to fully establish HDACi as a valid pharmacological option to treat MPNs. For example, it is ill-defined through which mechanisms HDACi target JAK2^{V617F}-transformed cells. Is it the preferential degradation of JAK2^{V617F}, or are there further mechanisms involved? If it is the degradation of JAK2^{V617F}, then how do HDACi discriminate between wild-type and mutant JAK2? Which factors regulate such processes? It is likewise unclear if the targeting of a subset of HDACs or even of a single member of the 18 HDAC family members in humans suffices to attack leukemic cells driven by JAK2^{V617F}. Considering that JAK2^{V617F} and HDACs control the replication stress and DNA damage responses of leukemia cells (Plo et al., 2008; Kurosu et al., 2013; Marty et al., 2013; Ueda et al., 2013; Chen et al., 2014; Roos and Krumm, 2016; Nikolova et al., 2017), the impact of HDACi on such mechanisms should additionally be investigated as antitumoral effects.

b. Histone deacetylase inhibitors modulate aberrant JAK2 signaling in solid tumor cells. Although JAK2^{V617F} has never been detected in solid malignancies (Lee et al., 2006; Herreros-Villanueva et al., 2010), aberrant JAK2 signaling and constitutive STAT activation are malignant features of multiple solid tumors. Aberrant growth hormones, epidermal growth factors, and chemokines can trigger a hyperactivation of JAK2/STAT5 in prostate, breast, and colorectal cancers (Harry et al., 2012; Thomas et al., 2015; Buchert et al., 2016; Bharadwaj et al., 2020; Qureshy et al., 2020). In the following section, we summarize evidence that JAK2 is a druggable vulnerability of solid tumors.

In colon cancer cells, TSA induced cell cycle arrest and apoptosis. TSA upregulated SOCS1 and SOCS3 through the hyperacetylation of histones H3 and H4 in their gene promoters. This tied in with the reduction of both phosphorylated and total forms of JAK2 and STAT3 as well as their downstream targets BCL2, survivin, and the cell cycle regulator p16/INK4A (Xiong et al., 2012) (Fig. 2). Since JAK2 activation correlates with poor survival of patients with colorectal cancers or prostate cancers (Li et al., 2004; Mao et al., 2011), HDACi might be an option for particularly aggressive colon cancer types.

Additional cancer cells, such as the highly fatal pancreatic ductal adenocarcinoma (PDAC), are sensitive toward the JAK2/JAK3 inhibitor AZ-960 (Scholz et al., 2003; Corcoran et al., 2011). Since PDAC cells are also vulnerable to HDACi (Schneider et al., 2010; Laschanky et al., 2019), inhibition of JAK2 signaling and HDACs might combine favorably against PDAC cells. This may allow a dose reduction of HDACi and

JAKi and, consequently, reduced toxicity against normal tissues.

The rare and highly deadly pancreatic neuroendocrine tumors (PanNET) display epigenetic alterations, including HDAC overexpression. Clinically achievable nanomolar doses of panobinostat stalled the proliferation and caused apoptosis of primary human PanNET and corresponding rodent tumor cells. Global RNA analyses indicated that panobinostat dysregulated JAK-STAT signaling in PanNET cells. Panobinostat did not alter JAK2 and SOCS3 levels but augmented STAT3 phosphorylation. The authors of this study interpret this as a compensatory mechanism suppressing panobinostat-induced apoptosis (Schmitz et al., 2021). Perhaps such compensatory pathways create a susceptibility for cotreatment with HDACi and drugs blocking STAT3 (Bharadwaj et al., 2020). This hypothesis is supported by the above-described notion that breast cancer cells become vulnerable to HDACi when LIFR-induced JAK-STAT signaling is antagonized by small molecules against JAKs and BRD4 (Zeng et al., 2016).

Epigenetic dysregulations also occur in hepatocellular carcinoma (HCC) and might be a vulnerability of this most common and highly fatal liver cancer (Anestopoulos et al., 2015). Increased activation of JAK2-STAT3 signaling in HCC correlates with reduced SOCS3 expression due to transcriptionally silenced methylated CpG regions in the *SOCS3* promoter. Treatment of HCC cells with the DNA hypomethylating agent 5-aza-2'-deoxycytidine/decitabine successfully restored SOCS3 expression, blocked STAT3 activity, and suppressed tumor cell growth. These findings suggest that epigenetic changes promote tumorigenesis through JAK2-STAT3 signaling (Niwa et al., 2005). Whether HDACi and hypomethylating drugs combine favorably in HCC through JAK-STAT inactivation is yet unresolved. A recent study compared the efficacy of equimolar concentrations of vorinostat and decitabine in HCC cell models. Both compounds induced apoptosis, and vorinostat was more potent than decitabine. Vorinostat decreased mRNAs encoding class I HDACs, JAK2, STAT3, and increased SOCS1 and SOCS3 expression (Sanaei et al., 2021). Additional work can show whether these changes are functionally relevant. Irrespective thereof, evidence collected in different cell systems shows that HDACi augmented SOCS protein levels (Fig. 2).

3. Impact of Histone Deacetylase Inhibitors on Janus Kinase-3. Activating JAK3 mutations in T-ALL and other blood malignancies position dysregulated JAK3 signaling as a potential drug target (Zhang et al., 2012; De Keersmaecker et al., 2013; Vicente et al., 2015; Li et al., 2016; Girardi et al., 2017; Liu et al., 2017; Degryse et al., 2018a, 2018b) (Fig. 1). Unlike other JAKs, JAK3 can be inhibited specifically due to its cysteine 909 residue (Cys909). This gatekeeper position of JAK3 is not present

in JAK1, JAK2, or TYK2. This inspired the development of irreversible, selective JAK3 inhibitors (Goedken et al., 2015; Telliez et al., 2016; Elwood et al., 2017; He et al., 2017; Kempson et al., 2017; He et al., 2018). Docking experiments with the JAK3 kinase domain led to the identification of ATP-competitive irreversible tricyclic covalent compounds that inhibit JAK3 in the low nanomolar range. These drugs contain chloroacetamide or acrylamide-derived Michael acceptors to irreversibly bind Cys909. Such molecules displayed high selectivity against the other JAKs (IC_{50} values above 1 μ M) and potently abrogated the JAK3-dependent activation of STAT5 in lymphocytes (Goedken et al., 2015). The irreversible JAK3 inhibitor PF-06651600 blocked JAK3 activity while sparing JAK1-mediated signaling (Telliez et al., 2016). Another irreversible covalent JAK3 inhibitor potently inhibited JAK3 (0.15 nM) with 4300-fold selectivity toward JAK3 over JAK1 (Elwood et al., 2017). Additional covalent irreversible JAK3 inhibitors preferentially inhibited JAK3 (IC_{50} = 57 nM) over other JAKs (IC_{50} > 10 μ M) (He et al., 2017). Likewise, the compound T29 demonstrated the highest potency and selectivity in picomolar doses for JAK3 (IC_{50} = 0.14 nM) over other JAKs (IC_{50} > 5 μ M) and other kinases with a cysteine residue analogous to Cys909 in JAK3 (He et al., 2018). Further irreversible selective agents impair JAK3 (IC_{50} = 1 nM) with high selectivity over JAK1 (IC_{50} = 1100 nM), JAK2 (IC_{50} = 1300 nM), and TYK2 (IC_{50} > 5000 nM). These molecules contain an acrylamide scaffold as an electrophilic warhead (Kempson et al., 2017). The development and characterization of specific JAK3 inhibitors verified their usefulness in treating autoimmune disorders. The putative activity of such compounds against cancer cells is unexplored.

HDACi are a putative option to kill cancer cells expressing mutant JAK3. In CTCL, JAK3 repressed the tumor-suppressive microRNA *miR-22* via tyrosine phosphorylation of STAT5, leading to its nuclear translocation and binding to the *miR-22* gene promoter. This attenuates the validated *miR-22* target genes *MAX*, *MYCBP*, *HDAC6*, *CDK6*, and *PTEN* in malignant T cells. STAT3 and STAT5 repress such genes through the recruitment of transcriptional corepressors such as EZH2 and HDACs. Vorinostat time- and concentration-dependently upregulated *miR-22* in CTCL cells (Sibbesen et al., 2015). HDACi additionally abrogate the JAK3-STAT5-dependent expression of *miR-21* being associated with CTCL (Lindahl et al., 2016). This impact of HDACi on *miRNAs* may represent an Achilles' heel of CTCL.

Selective JAK3 inhibitors may favorably combine with HDACi against subtypes of lymphoma and potentially other malignancies that particularly rely on mutant JAK3. A key consideration with such combinatorial approaches is at which therapeutic levels the desired

specificity for JAK3 disappears due to the impact of HDACi on JAK1, JAK2, and TYK2. Nonetheless, if given for short periods, JAK3 and HDACi may hit the cancer cells hard and simultaneously.

4. Impact of Histone Deacetylase Inhibitors on Tyrosine Kinase-2. Panobinostat abrogated the tyrosine phosphorylation of JAK2, TYK2, STAT3, and expression of MCL1. This consequently induced apoptosis in cell lines and tumor samples derived from DLBCL patients (Gupta et al., 2010; Gupta et al., 2012). This paved the way for phase I/II clinical trials (#NCT01261247 and #NCT01523834) that assessed the efficacy as well as the safety profile of panobinostat in patients with relapsed/refractory DLBCL. Patients tolerated panobinostat and showed a 21% overall response. Whether patients unresponsive to treatment had persistent JAK signaling is an intriguing but unexplored possibility. Truly, other targets of HDACi might also dictate resistance. For example, a recent study using single-cell sequencing illustrates that c-FOS antagonizes HDACi-induced apoptosis and is upregulated in DLBCL cells surviving HDACi (Krämer and Schneider, 2022; Wang et al., 2022).

Another strategy to combat TYK2 may rely on the HDAC-dependent control of the ubiquitin-proteasome system and the finding that cytokines not only activate TYK2 but can lead to its proteasomal degradation. Upon activation by IFN α , a STAT1-dependent induction of the E2 ubiquitin conjugase UBCH8 accelerates the turnover of TYK2 in complex with the E3 ubiquitin ligase seven-in-absentia-homologue-2 (SIAH2) in NSCLC cells. Accordingly, SIAH2 levels inversely correlate with STAT3 phosphorylation and metastatic gene expression in NSCLC patient samples (Müller et al., 2014). HDACi induce E2 ubiquitin conjugase UBCH8 and its cooperativity with SIAH1/SIAH2 to accelerate the proteasomal degradation of multiple oncogenes (Buchwald et al., 2010; Buchwald et al., 2013; Krämer et al., 2013). Eliminating TYK2 and its overactivity in certain tumors with HDACi might be a promising anticancer strategy.

5. Histone Deacetylase Inhibitors Modulate Janus Kinase-Signal Transducer and Activator of Transcription-Dependent Immune Functions. Recent reports demonstrate that the effects of HDACi on JAK-STAT-dependent immunologic tumor surveillance can stall tumor cell proliferation. The overexpression of B7 homologue-1, which is rather known as programmed cell death ligand-1 (PD-L1), is a key mechanism of immune evasion by tumor cells and hence a modern anticancer target (Deng et al., 2018). Conspicuously, HDACs sustain an IFN γ -mediated expression of PD-L1 in gastric cancer. Pretreatment with butyrate, vorinostat, or TSA, as well as a genetic elimination of HDAC1 and HDAC3 by RNAi, suppressed the IFN γ -induced acetylation of histone H3 at lysine residue 9 at the promoter of the *B7 homologue-1/PD-L1*

gene. This equally occurred upon depletion of JAK2, p-JAK1, p-JAK2, and p-STAT1 and a subsequently impaired nuclear translocation of STAT1. HDACi suppressed IFN γ signaling in gastric cancer cells, which hindered their ability to escape detection and elimination by cytotoxic CD8⁺ T cells (Deng et al., 2018). These findings suggest that the reported negative impact of HDACi on STAT signaling (Krämer et al., 2009; Krämer and Heinzl, 2010; Ginter et al., 2012; Wiczorek et al., 2012; Kaowinn et al., 2017; Kotla et al., 2017; Liu et al., 2020; Yang et al., 2021) and herein summarized negative effects of HDACi on JAKs (Fig. 2) are key immunoregulatory and tumor-suppressive mechanisms of such drugs.

These insights could not only improve the treatment of gastric cancers. In murine models with lung and renal carcinoma, entinostat improved the tumor-suppressive effects of an antibody neutralizing the receptor programmed death receptor-1 (PD-1) (Orillion et al., 2017). The class I/IV HDACi mocetinostat (Fig. 4A) similarly promoted tumor antigen presentation, decreased immune suppressive cell types, and augmented antitumor effects of anti-PD-L1 antibodies in colon carcinoma-bearing mice (Briere et al., 2018).

JAK2^{V617F}, and consequently p-STAT3/p-STAT5-induced transcriptional induction of PD-L1, are higher in primary MPN patient-derived monocytes, megakaryocytes, and platelets than in cells from healthy individuals. Inhibition of JAK2 with ruxolitinib or the JAK2 inhibitor SD-1029 and anti-PD-1 or anti-PD-L1 antibodies prolonged the survival of mice xenografted with human peripheral blood mononuclear cells from MPN patients and of JAK2^{V617F} knock-in mice (Prestipino et al., 2018). Targeting PD-L1 via the depletion of JAK2^{V617F} by HDACi may disable the immune evasion of MPNs.

The above-summarized reports inform that HDACi combine favorably with inhibitors of immune checkpoint molecules. The ability of HDACi to target immune suppressive mechanisms is not restricted to the PD-1/PD-L1 axis. For example, entinostat disrupts immunosuppressive functions of regulatory T cells in the tumor microenvironment through the acetylation of STAT3 and an ensuing downregulation of *Foxp3* expression in renal and prostate murine cancer models (Shen et al., 2012).

6. Advantages of Indirect Pan-Janus Kinase Inhibition by Histone Deacetylase Inhibitors. Based on current evidence, HDACi function as indirect pan-JAKi that cause a slow decline of JAK-STAT signaling through transcriptional effects and an impact on JAK protein stability. Although details on such mechanisms are still a subject of investigation, the ability of HDACs to interfere with multiple members of the JAK family seems to be therapeutically beneficial.

Due to the transcriptional and protein stability-centered mechanisms of HDACi (Fig. 2), they can attenuate JAK mutants irrespective of a particular lesion

(Fig. 1). This should also hold for JAK2 resistance mutations in the ATP-binding site that are acquired during the treatment of B-ALL cells with type 1 and type 2 JAKi (Downes et al., 2021) and in case of a co-operation between concomitant activating mutations in JAK1 and JAK3. This acquired resistance to JAKi favors the selection of TKi-resistant tumor clones and consequently increased risk of relapse in patients suffering from T-ALL, T-PLL, and extranodal natural killer (NK)/T-cell lymphoma (Bains et al., 2012; Koo et al., 2012; Atak et al., 2013; Bellanger et al., 2014; Springuel et al., 2014). In MPN cell models and granulocytes from MPN patients, overexpression of JAK2^{V617F} at the transcriptional and protein levels as well as heterodimer formation between mutant JAK2^{V617F} with JAK1 and TYK2 increases their phosphorylation and TKi resistance (Koppikar et al., 2012). Elimination of multiple JAKs by HDACi could abrogate this adaptation to JAKi. One might envision that HDACi can also be effective against tumor-associated JAK fusion proteins resulting from genomic rearrangements.

In a similar fashion, inhibition of heat shock protein-90 (HSP90) attenuates JAK1 and JAK2^{V617F} in MPN cells (Wang et al., 2009; Fiskus et al., 2011; Koppikar et al., 2012; Weigert et al., 2012). Hyperacetylation of HSP90 in response to HDACi was proposed to promote proteasomal degradation of mutant JAK2 (Wang et al., 2009; Fiskus et al., 2011; DeAngelo et al., 2013a). Recent evidence, though, illustrates that the number of acetylation sites in HSP90 and their stoichiometry and biologic relevance are more complex than anticipated (Krämer et al., 2014; Prus et al., 2019). An advantage of HDACi over HSP90 inhibitors is the FDA approval for HDACi (Food and Drug Administration, 2010; Whittaker et al., 2010; Cappellacci et al., 2020).

Moreover, HDACi-triggered mechanisms that eliminate JAK mutants are not susceptible to an observed persistence of phosphorylated JAK2 in the presence of type 1 TKi (Meyer et al., 2015; Wu et al., 2015). Hyperactive JAK2 even appears to sensitize cells to HDACi (Guerini et al., 2008; Savino et al., 2017). This could be particularly relevant considering the allosteric activation mechanism of JAK2 that shifts the molecule from an autoinhibited toward a molecularly open, uninhibited state. Increased complementarity and hydrophobicity of JAK2^{V617F} and other JAK2 mutants allow their ligand-independent dimerization and activation. HDACi-induced elimination of such JAK2 mutants decreases the likelihood of their dimerization and thus abrogates allostery. This idea is supported by the notion that the specific allosteric TYK2 inhibitors BMS-066/BMS-986165 bind the JH1 pseudokinase domain of TYK2 and can freeze it in its inactive, autoinhibited conformation in normal blood cells (Tokarski et al., 2015; Burke et al., 2019; Glassman et al., 2022).

Another advantage of HDACi and a synergistic killing of MPN cells with JAKi/HDACi combinations may rely on a suppression of the coincidence of B-cell non-

Hodgkin lymphomas during treatment of MPN patients with JAK1/JAK2 inhibitors (Porpaczy et al., 2018). Immunosuppressive effects of such drugs on cytotoxic T lymphocytes and NK cells facilitate the conversion of malignant preexisting B-cell clones into aggressive B-cell lymphoma (Low and Song, 2016; Khandelwal et al., 2017; Kim et al., 2017; Assouan et al., 2018). As discussed above, there are beneficial effects of HDACi on the PD1/PD-L1 system.

In the context of the here-discussed interplay between HDACs and JAK signaling, one should still consider that HDACi can attenuate oncogenic signaling at multiple levels. For example, pan-HDACi and class I HDACi reduce WEE1 and ACK1 expression in cells derived from blood malignancies and solid tumors (Cornago et al., 2014; Zhou et al., 2015; Mahendrarajah et al., 2016; Göder et al., 2018; Hu et al., 2020; Wachholz et al., 2022). In addition, HDACi might abrogate JAK-dependent crossdrug resistance mechanisms. For instance, JAK1^{V658F} (Fig. 1) confers resistance of the FLT3-ITD-positive AML cells to the TKi midostaurin and sorafenib (Rummelt et al., 2021). Indirect inhibition of JAK1 signaling by HDACi (Klampfer et al., 2004; Owusu and Klampfer, 2017) may disable such resistance mechanisms. Furthermore, concomitant degradation of FLT3-ITD by HDACi (Buchwald et al., 2010; Pietschmann et al., 2012; Wachholz et al., 2022) could be a double-hit approach against such aggressive cancer cells.

IV. Chemically Coupled Multitargeting Drugs

Despite their wide use and clinical relevance, combination therapies face challenges and limitations. These include, but are not limited to, undefined drug-drug interactions, the need to establish effective dose regimens for two agents, higher costs of development, extra efforts for clinical trials, and complications of the pharmaceutical formulations (Peters, 2013). To prevail over these drawbacks, multipharmacophore therapeutics combining two or more different pharmacophores are developed as an alternative strategy (Zimmermann et al., 2007; Anighoro et al., 2014; Nepali et al., 2014; Fu et al., 2017). Such a chemical coupling allows hitting two cancer-relevant targets with one molecule at the same time. This might overcome drug-drug interactions, might exert a more potent killing effect, can prevent the advent of drug-resistant cells due to mutations in a single drug target, and offers more predictable pharmacokinetic profiles. In addition, a hybrid molecule can facilitate drug dosing to ensure patient compliance (Morphy et al., 2004; Morphy and Rankovic, 2005). A key condition for the success of such strategies is that both pharmacophores retain specific and highly potent activities against their targets. Hence, critical warheads must be present in the fusion molecules (Fig. 4C).

Kinases were one of the earliest molecules to be co-targeted with HDACs. Examples of such compounds resulted from chemical coupling of vorinostat and the macrocycle templates of zotiraciclib/pacritinib (SB1317/SB1518). The macrocycle templates of these TKi are replaced by the cap group of vorinostat (Ning et al., 2015). This preserves the hydroxamic acid group for complex formation with the catalytically active Zn²⁺ in HDACs (Fig. 4C). These hybrid structures displayed high potency against JAK2, FLT3, and HDACs. The multifunctional hybrid molecule 32 is a nanomolar inhibitor of JAK2, FLT3, and global HDAC activity (IC₅₀ values 686, 87, and 87 nM, respectively), and exhibited remarkable anticancer activities against leukemic cells as well as solid tumors (Ning et al., 2015). An additional JAK-HDAC dual inhibitor termed 20a showed high activity against JAK2 (IC₅₀ = 8 nM) and HDAC6 (IC₅₀ = 46 nM). This agent caused apoptotic cell death of AML cells (Huang et al., 2018a).

Pyrimidine-2-amino-pyrazole/hydroxamate derivatives are dual-acting inhibitors that inactivate JAK2 and HDAC6 at nanomolar doses. A compound named 8m produced higher activities in vitro and in vivo than vorinostat and ruxolitinib against a panel of leukemic cell lines (Liang et al., 2019).

Solid tumor-derived cells are also susceptible to HDACi-TKi fusion molecules. Most recently, a series of pyrrolo-[2,3-d]-pyrimidine-based derivatives emerged as dual-acting JAKi/HDACi. Compounds 15d and 15h block JAK1,2,3 and HDAC1/HDAC6 in nanomolar doses, attenuated tyrosine-phosphorylated STAT3, and suppressed the growth of triple-negative human breast cancer in vitro and in mice (Liang et al., 2022). Despite their ability to induce LIFR expression, these multitargeting drugs combated LIFR-induced STAT3 phosphorylation by tumor-associated fibroblasts. Remarkably, 15d/15h were superior to vorinostat. Despite an equal accumulation of acetylated histones and tubulin, targets of HDAC1 or HDAC6 (Table 1), vorinostat activated STAT3 phosphorylation upon induction of LIFR in breast tumor cells (Liang et al., 2022). These data illustrate that combined inhibition of JAKs and HDACs favorably abrogate cytokine-induced phosphorylation events.

The hybrid molecule 24, a chemical coupling of ruxolitinib and vorinostat, inhibits HDAC1,2,3,6,10, and JAK1/JAK2 with >10-fold selectivity over TYK2, JAK3, and 93 other kinases (Yao et al., 2017). Compound 24 inhibited STAT3 tyrosine phosphorylation, induced an accumulation of acetylated histone H3 at 0.15–7.36 μM, apoptosis-related processing of the DNA repair enzyme PARP1, and growth inhibition in a panel of leukemic cells (ALL, AML, MPN, multiple myeloma) and solid tumor-derived cells (breast, colon, prostate) (Yao et al., 2017). Although compound 24 is a more potent inhibitor of HDAC6 than of HDAC1 (1.4 versus 6.9 nM), experiments with the HDAC6 inhibitor

tubastatin A show that inhibition of HDAC6 cannot explain the processing of PARP1 (Yao et al., 2017). These data correspond to a strong biochemical activity of HDAC6 inhibitors without any antiproliferative or cytotoxic effects in several leukemic and solid tumor cells in vitro and in vivo (Noack et al., 2017; Schäfer et al., 2017; Leonhardt et al., 2018; Sellmer et al., 2018; Bandolik et al., 2019; Beyer et al., 2019; Depetter et al., 2019; Pons et al., 2019; Nawar et al., 2022). Presumably, compounds 15d/15h killed cells through inhibition of HDAC1 (Liang et al., 2022). Work on compound 24 was extended by the same group who developed it. Fusing the pharmacophores of vorinostat and ruxolitinib via a single nitrogen atom yielded subnanomolar inhibitors of JAK2 and HDAC6. These compounds had remarkable selectivity against HDAC1 and improved potency against the individual pathways. Compound 13b presented the highest potency and striking subnanomolar activity against JAK2 with an IC_{50} of 41 pM and an IC_{50} against HDAC6 of 200 pM (Yao et al., 2018). Thus, optimized JAK-HDAC hybrid inhibitors could improve cancer therapy.

Similarly, merging the pharmacophores of ruxolitinib and vorinostat created molecules with high potency against solid and hematologic malignancies. These agents attenuated *JAK2* mRNA expression and increased proteasomal degradation of JAK2. This associated with reactive oxygen species accumulation and tumor growth inhibition by apoptosis (Civallero et al., 2017). These data hold for other JAKi-HDACi fusions. Agents resulting from a chemical coupling of the JAK2-selective inhibitors XL019 or pacritinib with vorinostat produced antiproliferative activity and triggered apoptosis in a panel of cancer cells from ALL, MPN, multiple myeloma, breast, colon, and prostate (Yang et al., 2016; Chu-Farseeva et al., 2018). These findings shed light on new leads for bifunctional molecules that block JAKs and HDACs. Furthermore, these results suggest that inhibiting JAK2 phosphorylation with TKi does not exclude its proteasomal degradation upon HDAC inhibition. From these insights, one can deduce that the preferential loss of mutant JAK2 over wild-type JAK2 is unlikely related to JAK2 phosphorylation (Guerini et al., 2008). It will be interesting to see if such fusion molecules overcome the limited clinical potential of XL019 in MPN (Verstovsek et al., 2014) and whether they are effective against other neoplastic diseases.

An additional dual-acting molecule, Roxyl-zhc-84, is a fusion of vorinostat and the CDK4/CDK6 inhibitor abemaciclib. This dual pharmacophore resolved the problem of limited responses of cells from aggressive breast cancer, high-grade serous ovarian cancer, and NSCLC toward traditional HDACi. The mechanism underlying this benefit is an overcoming of JAK1-STAT3-BCL2-mediated drug resistance. Roxyl-zhc-84 was superior to the combinatorial treatment with HDACi and JAKi. Moreover, Roxyl-zhc-84 had favorable pharmacokinetic profiles

in a rat model and produced low toxicity in a mouse model (Huang et al., 2018b). The same group showed that conjugates of HDACi and CDK inhibitors overcame limited responses of oral squamous cell carcinoma (OSCC) toward HDACi monotherapies (Zhao et al., 2019). Roxyl-ZR, a highly Roxyl-zhc-84-related conjugate of vorinostat and abemaciclib, significantly suppressed the survival of OSCC cells through inhibition of the JAK1-STAT3-BCL2 axis and ensuing apoptosis. In mice, Roxyl-ZR mitigated OSCC growth at low toxicity (Zhao et al., 2019).

These results imply the therapeutic potential of hybrid molecules and provide valuable lead compounds for further assessment and structural drug optimization.

V. Targeting Aberrant Janus Kinase Signaling by Proteolysis Targeting Chimeras and Histone Deacetylase Inhibitors

Proteolysis targeting chimeras (PROTACs) are heterobifunctional chimeras that contain ligands for target proteins, ligands that dock to E3 ubiquitin ligases, and linkers to connect these ligands (Fig. 5A). The formation of a ternary complex leads to K48-linked poly-ubiquitination of the target proteins, which are then recognized and rapidly proteolyzed by proteasomes (Pettersson and Crews, 2019; Vogelmann et al., 2020).

Despite over 600 E3 ligases in the human genome, until now few candidate E3 ligases, including inhibitors of apoptosis (IAPs), von Hippel-Lindau (VHL), cereblon (CRBN), murine double minute-2 homolog, damage-specific DNA binding protein-1 and Cullin-4-associated factor-15/16, and ring finger protein-114 are employed for targeted protein degradation (Liu et al., 2019). This is due to the scarcity of small-molecule ligands that recruit E3s without impairing their functions. The identification of more compounds that recruit E3s is anticipated to improve and refine the PROTAC technology (Belcher et al., 2021).

A key advantage of PROTACs over low-molecular-mass inhibitors is the rapid and sustained elimination of their targets. This is frequently accompanied by high efficacy and, in some cases, increased selectivity for their targets (Nowak et al., 2018; Sun et al., 2018; Khan et al., 2019; Shah et al., 2020; Zhang et al., 2020; Alabi et al., 2021; Chang et al., 2021). Besides, PROTACs targeting receptor tyrosine kinases disable both enzymatic activities and scaffolding functions of these proteins. This is important because scaffolding kinase functions reactivate oncogenic signaling pathways and propel kinome rewiring supporting escape clones with drug resistance (Shah et al., 2018; Murtuza et al., 2019). Moreover, JAK-PROTACs can disengage the crossreactivation of specifically inhibited JAKs by the other JAK family members. This is a major resistance mechanism against JAKi (Koppikar et al., 2012).




Fig. 5. Exploiting VHL-based PROTACs and HDACi to correct aberrant JAK-STAT signaling. The game-changing PROTAC technology, as well as the impact of HDACi on VHL expression, suggests a new, synergistically active therapeutic approach through proteolytic mechanisms and HDAC inhibition. (A) PROTACs can accelerate the proteasomal degradation of JAKs. Shown is a VHL-based PROTAC. (B) The expression of the *VHL* gene is suppressed by HDACs. HDACi augment the *VHL* mRNA and protein levels, which can increase the proteasomal degradation of JAKs by PROTACs. This principle may also apply to targets of VHL-PROTACs beyond JAKs. Created with BioRender.com.

Most recently, the ubiquitin-proteasome machinery was successfully hijacked with PROTACs that degrade JAKs through IAPs, VHL, and CRBN in a variety of leukemic cells (Shah et al., 2020; Chang et al., 2021; Alcock et al., 2022). Guided by the JAKi chemotypes pyrimidine and quinoxaline, IAP, VHL, and CRBN-recruiting PROTACs were designed and tested for

their impact on wild-type JAK1/JAK2 levels in human AML cells. These PROTACs accelerated the ubiquitination and proteasomal degradation of JAK1 and JAK2 in micromolar doses (0.6–10 μ M). In this setting, IAP-based PROTACs induced the degradation of JAK1, JAK2, and JAK3 more effectively than PROTACs recruiting VHL and CRBN (Shah et al., 2020).

The clinically used type 1 JAK1/JAK2 inhibitors ruxolitinib and baricitinib were used to rationally design a new series of CRBN-directed PROTACs (Chang et al., 2021). This was achieved by analysis of the crystal structures of the human JAK2 JH1 domain complexed with baricitinib or ruxolitinib at 1.7 to 1.8 Å resolution. It turned out that the pyrimidine rings of the drugs offer a C2-carbon that points away from the catalytic cleft of JAK2 toward the solvent. Introducing a 4-amino-benzamide allowed the attachment of a linker and the imides thalidomide/lenalidomide/pomalidomide as recruiter elements for CRBN. Some of the obtained PROTACs abrogated aberrant JAK-STAT signaling and killed ruxolitinib/baricitinib-resistant ALL cells with a combined overexpression of CRLF2 (due to the IGH-CRLF2 translocation) and JAK2^{I682F}/JAK2^{R683G}. PROTACs 5, 6, 7, and 8 effectively inhibited aberrant JAK-STAT signaling (IC₅₀ < 100 nM) in a panel of ALL cell lines and patient-derived primary ALL xenografts in mice. An up to 70,000-fold higher activity of the PROTACs compared with their parent compounds was not due to increased inhibition of JAK2 phosphorylation but related to its degradation. Even though these efforts provide a new window to target JAK-mutated malignancies, specificity and pharmacokinetics of the existing JAK-directed PROTACs require optimization. The JAK-PROTACs lost the specificity of ruxolitinib and baricitinib for JAK1/JAK2. In addition, these PROTACs inhibited MAP3K2, MAP3K3, and YSK4. Irrespective thereof, compound 8, a fusion of baricitinib and thalidomide, eliminated mutant JAK2 specifically and dose-dependently in cultured cells. Cells with a genetic elimination of CRBN by CRISPR-Cas9 retained JAK2 in the presence of compound 8, verifying CRBN-induced proteasomal degradation. Proteasomal degradation of the tumor-promoting CRBN neo-substrate G1-to-S-phase-transition-1 (GSPT1) by CRBN ligands can cause false-positive effects of PROTACs. This can blur target-specific effects of PROTACs. Interestingly, compound 8 did not degrade GSPT1 in cultured B-ALL cells with a translocation of CRLF2 and JAK2^{I682F}. Nonetheless, compound-8 attenuated wild-type JAK2 and GSPT1 in primary B-ALL cells with a translocation of CRLF2 (Chang et al., 2021).

Based on ruxolitinib and baricitinib, the same group expanded their work and successfully developed JAK2/JAK3-selective PROTACs using the phenyl glutarimide (PG) ligand as CRBN-recruiting warhead. PG has a phenyl group instead of the phthalimide present in the immunomodulatory imide-based PROTACs. The PG-based PROTAC 11 significantly reduced off-target degradation of GSPT1 and showed higher selectivity for JAK2/JAK3 over the other JAKs in CRLF2-rearranged ALL cell lines. PROTAC 11 was more effective than the parental JAK2 inhibitors and displayed the most potency in xenografted, patient-derived ALL cells

harboring an ATF7IP-JAK2 fusion (IC₅₀ = 24 nM). Higher PROTAC doses were needed to kill ALL cells carrying both JAK2 fusions and CRLF2 rearrangements (IC₅₀ < 120 nM) (Alcock et al., 2022).

Phenotypically similar to the PROTAC-mediated degradation of mutant JAKs, HDACi suppress JAK signaling by genomic and nongenomic mechanisms (Guerini et al., 2008; Rambaldi et al., 2010; Akada et al., 2012; Andersen et al., 2013; Zhao et al., 2016b; Bose and Verstovsek, 2017b; Savino et al., 2017) (Fig. 2). Transcriptional effects of HDACi may augment the efficacy of VHL-based PROTACs. Romidepsin and panobinostat induce the expression of VHL in endothelial cells and B-cell lymphomas (Kwon et al., 2002; Kalac et al., 2011). This might potentiate the effects of VHL-based PROTACs through an upregulation of VHL (Fig. 5B). PROTACs targeting HDACs might also promote the effectiveness of VHL-based PROTACs by increasing the amount of available VHL. PROTACs that decrease HDAC1/HDAC2 (Smalley et al., 2022), HDAC3 (Cao et al., 2020; Xiao et al., 2020), HDAC6 (Wu et al., 2019), HDAC8 (Chotitumnavee et al., 2022), and SIRT2 (Schiedel et al., 2018) in cultured cells as well as in animal models were identified. This implies the potential of PROTACs as a comprehensive approach for cancer treatment.

Combinations of JAK-directed PROTACs with HDACi might likewise be useful from the viewpoint that both drugs decrease mutant JAK isoforms. Particularly, the increased sensitivity of JAK2^{V617F} compared with wild-type JAK2 toward HDACi (Guerini et al., 2008) could preferentially target leukemic cells over healthy cells, allowing sustained immune functions and proper hematopoiesis. In addition, the choice of attracted E3 ubiquitin ligase can prevent damage to normal tissues. For example, VHL- and CRBN-based PROTACs spare thrombocytes and platelets due to their low expression of VHL and CRBN. This could prevent fatal thrombocytopenia being a common cause of chemotherapy-associated death (Khan et al., 2019; Zhang et al., 2020). On the other hand, patients suffering from diseases such as ET may not profit well from VHL- and CRBN-based PROTACs and should obtain PROTAC-based drugs recruiting other E3 ubiquitin ligases.

VI. Clinical Trials to Assess the Safety and Efficacy of Combined Pharmacological Targeting of Janus Kinases and Histone Deacetylases

The FDA has approved 10 JAKi, including ruxolitinib and the JAK1/JAK3 inhibitor tofacitinib, for the treatment of MPNs and rheumatoid arthritis (Qureshy et al., 2020). As mentioned above, the FDA has also authorized HDACi. Table 2 lists a selection of clinical trials using combinations of JAKi and HDACi.

Panobinostat was clinically active against JAK2^{V617F}-positive MPNs as monotherapy or in combination with

ruxolitinib (DeAngelo et al., 2013a, 2013b; Mascarenhas et al., 2013; Ribrag et al., 2013; Mascarenhas et al., 2020). Preclinical studies with panobinostat plus ruxolitinib revealed that such combinations effectively corrected JAK2^{V617F}-associated symptoms, including splenomegaly, as well as bone marrow and spleen histology in a JAK2^{V617F}-driven MPN mouse model. Concurrent administration of panobinostat and ruxolitinib disrupted the JAK2^{V617F}-induced STAT5 phosphorylation and significantly inhibited the growth of MPN cells (Evrot et al., 2013). Similarly, combinations of panobinostat and the JAK2 inhibitor TG101209 significantly disrupted JAK2^{V617F}-driven signaling and synergistically induced apoptosis of primary CD34⁺ MPN cells. These were more affected by such treatment than normal CD34⁺ hemopoietic progenitor cells. This signifies a specific antileukemic effect (Wang et al., 2009). It is tempting to speculate that particularly the mutant JAK2^{V617F} expression created this vulnerability.

These insights inspired multiple clinical trials using panobinostat plus ruxolitinib. When applied at lower doses over time, such drug regimen significantly corrected splenomegaly and ameliorated bone marrow fibrosis (Mascarenhas et al., 2020). In a phase Ib clinical trial (#NCT01433445), the combinatorial therapy of ruxolitinib plus panobinostat demonstrated an appropriate safety profile, with low dose-limiting toxicities and modest rates of grade 3/4 anemia (34.2%) and thrombocytopenia (21.2%) (Ribrag et al., 2013).

The phase I/II clinical study (#NCT01693601) confirmed the efficacy and safety of ruxolitinib administered twice daily at 10–15 mg and panobinostat at 10–20 mg three times biweekly. Anemia occurred as the most common therapy-associated feature in 47% of patients (Ribrag et al., 2013; Harrison et al., 2015; Mascarenhas et al., 2020). The reduction in spleen size in this trial was similarly seen in patients who received ruxolitinib in the phase III trial Controlled Myelofibrosis Study with Oral JAK Inhibitor Treatment I and Controlled Myelofibrosis Study with Oral JAK Inhibitor Treatment II (#NCT00934544) (Harrison et al., 2016).

In MF patients, the combination of ruxolitinib and panobinostat was well tolerated and reduced splenomegaly in 57% and 39% of the patients at weeks 24 and 48, respectively. Of note, some patients experienced a reduction in JAK2^{V617F} allele burden and obvious correction in bone marrow fibrosis. Adverse events were not higher than with ruxolitinib or panobinostat single-agent treatments (Harrison et al., 2015).

As a single, orally given agent, pracinostat exhibited modest potency in MF patients (Zorzi et al., 2013; Chu et al., 2015). In a study with 25 MPN patients, 20 received ruxolitinib and pracinostat (#NCT02267278) (Table 2). Ruxolitinib was administered for three cycles,

whereas pracinostat was given afterward in the fourth cycle. Eighty percent of patients had objective responses, and 74% experienced spleen responses. The National Cancer Institute Common Terminology Criteria for Adverse Events reported that anemia and thrombocytopenia were the most frequent adverse events (Bose et al., 2019). The small patient number requires additional investigations on the clinical validity of ruxolitinib and pracinostat.

An ongoing phase I/II study #NCT03598959 is devoted to exploring the efficacy and evaluating the safety profile of tofacitinib (Nwaogu et al., 2020), combined with chidamide in patients with relapsed and refractory NK/T-cell lymphoma.

The molecular mechanisms behind the beneficial interaction of JAKi and HDACi are not entirely clear yet. In at least some cell systems, ruxolitinib and other type 1 JAKi stabilize the phosphorylated form of JAK2^{V617F} (Meyer et al., 2015; Wu et al., 2015). This TKi-mediated accumulation of mutant JAK2 can cause rebound effects when the drug concentrations decline, and this can cause adaptive processes termed hormesis. Since JAK2^{V617F} is more prone than the largely unphosphorylated wild-type JAK2 to HDACi-induced degradation (Guerini et al., 2008; Savino et al., 2017), the elimination of the inhibited mutant JAK2 by HDACi can prevent and disengage such undesired processes. Of course, other mechanism can also contribute to the beneficial outcome of JAKi and HDACi combinations. A topical issue is whether clinical-grade JAKi-HDACi fusions and JAK-directed PROTACs, when applied alone or with HDACi, are as potent or even superior to combined application schemes of JAKi and HDACi.

Despite such success, not all patients profit from HDACi and JAKi. It needs consideration that leukemic (stem) cells can reside in both the periphery and bone marrow. Coculture systems revealed that bone marrow stromal cells protected MPN-derived JAK2^{V617F}-positive cell lines as well as primary bone marrow-derived post-PV AML cells from lethal effects of ruxolitinib and vorinostat. The cytoprotective effects of bone marrow stromal cells were mediated by the secretion of soluble factors that maintain cellular homeostasis via activation of JAK-STAT, phosphatidylinositol 3-kinase, c-JUN N-terminal kinase, MEK-ERK, and nuclear factor- κ B (NF- κ B) signaling pathways (Cardoso et al., 2015). Such findings shed light on mechanisms of tumor survival and pave the way to novel, targeted therapeutic approaches.

VII. Summary

Members of the HDAC family are valuable targets for novel pharmacological intervention strategies against tumor cells. Modern, clinically relevant, broad-acting and isoenzyme-specific inhibitors of Zn²⁺-dependent HDACs modulate the activity and expression of the four JAK

TABLE 2
Examples of clinical studies that have tested JAKi and HDACi for cancer therapy

Drugs	Phase	Chemotypes	Condition or Disease	Clinical Trial Identifier	Status
Panobinostat (pan-HDACi) + ruxolitinib (JAK1/JAK2 inhibitor)	Ib	Hydroxamic acid+ pyrazole	(N = 61); PMF, post-PV MF, post-ET MF	#NCT01433445	Completed
Panobinostat (pan-HDACi) + ruxolitinib (JAK1/JAK2 inhibitor)	I/II	Hydroxamic acid+ pyrazole	(N = 20); PMF, post-PV MF, post-ET MF	#NCT01693601	Completed
Panobinostat(pan-HDACi) + ruxolitinib (JAK1/JAK2 inhibitor)	IV	Hydroxamic acid+ pyrazole	(N = 410); PMF, post-PV MF, post-ET MF and CIMF	#NCT02386800	Recruiting
Pracinostat (pan-HDACi) + ruxolitinib (JAK1/JAK2 inhibitor)	I/II	Hydroxamic acid+ pyrazole	(N = 25); PMF, post-PV MF, post-ET MF	#NCT02267278	Completed
Chidamide (inhibits HDAC1,2,3,10) + tofacitinib (JAK3 inhibitor)	I/II	Benzamide+ pyrazole	(N = 20); extranodal NK/T-cell lymphoma	#NCT03598959	Not yet recruiting

CIMF, chronic idiopathic myelofibrosis; PMF, primary myelofibrosis.

proteins and consequently their tumor-relevant downstream signaling. ATP-competitive JAK inhibitors, structurally diverse HDACi, JAKi-HDACi fusion pharmacophores, and JAK-directed PROTACs combine favorably against malignant cells in culture, experimental animals, and cancer patients. Future research will fully define how HDACs and acetylation-dependent molecular mechanisms control oncogenic JAK signaling and how clinicians can use this knowledge.

Authorship Contributions

Wrote or contributed to the writing of the manuscript: Mustafa, Krämer.

References

- AACR Project GENIE Consortium (2017) AACR Project GENIE: Powering Precision Medicine through an International Consortium. *Cancer Discov* **7**:818–831.
- Akada H, Akada S, Gajra A, Bair A, Graziano S, Hutchison RE, and Mohi G (2012) Efficacy of vorinostat in a murine model of polycythemia vera. *Blood* **119**:3779–3789.
- Alabi S, Jaime-Figueroa S, Yao Z, Gao Y, Hines J, Samarasinghe KT, Vogt L, Rosen N, and Crews CM (2021) Mutant-selective degradation by BRAF-targeting PROTACs. *Nat Commun* **12**:920.
- Albacker LA, Wu J, Smith P, Warmuth M, Stephens PJ, Zhu P, Yu L, and Chmielecki J (2017) Loss of function JAK1 mutations occur at high frequency in cancers with microsatellite instability and are suggestive of immune evasion. *PLoS One* **12**:e0176181.
- Alcock LJ, Chang Y, Jarusiewicz JA, Actis M, Nithianantham S, Mayasundari A, Min J, Maxwell D, Hunt J, Smart B, et al. (2022) Development of Potent and Selective Janus Kinase 2/3 Directing PG-PROTACs. *ACS Med Chem Lett* **13**:475–482.
- Andersen CL, McMullin MF, Ejerblad E, Zweegman S, Harrison C, Fernandes S, Bareford D, Knapper S, Samuelsson J, Löfvenberg E, et al. (2013) A phase II study of vorinostat (MK-0683) in patients with polycythemia vera and essential thrombocythemia. *Br J Haematol* **162**:498–508.
- Anestopoulos I, Voulgaridou GP, Georgakilas AG, Franco R, Pappa A, and Panayiotidis MI (2015) Epigenetic therapy as a novel approach in hepatocellular carcinoma. *Pharmacol Ther* **145**:103–119.
- Anighoro A, Bajorath J, and Rastelli G (2014) Polypharmacology: challenges and opportunities in drug discovery. *J Med Chem* **57**:7874–7887.
- Assouan D, Lebon D, Charbonnier A, Royer B, Marolleau JP, and Gruson B (2018) Ruxolitinib as a promising treatment for corticosteroid-refractory graft-versus-host disease. *Br J Haematol* **181**:687–689.
- Atak ZK, Gianfelici V, Hulselmans G, De Keersmaecker K, Devasia AG, Geerdens E, Mentens N, Chiaretti S, Durinck K, Uyttebroeck A, et al. (2013) Comprehensive analysis of transcriptome variation uncovers known and novel driver events in T-cell acute lymphoblastic leukemia. *PLoS Genet* **9**:e1003997.
- Aynardi J, Manur R, Hess PR, Chekol S, Morrissette JJD, Babushok D, Hexner E, Rogers HJ, Hsi ED, Margolske E, et al. (2018) JAK2 V617F-positive acute myeloid leukaemia (AML): a comparison between de novo AML and secondary AML transformed from an underlying myeloproliferative neoplasm. A study from the Bone Marrow Pathology Group. *Br J Haematol* **182**:78–85.
- Babon JJ, Lucet IS, Murphy JM, Nicola NA, and Varghese LN (2014) The molecular regulation of Janus kinase (JAK) activation. *Biochem J* **462**:1–13.

- Bains T, Heinrich MC, Loriaux MM, Beadling C, Nelson D, Warrick A, Neff TL, Tyner JW, Dunlap J, Corless CL, et al. (2012) Newly described activating JAK3 mutations in T-cell acute lymphoblastic leukemia. *Leukemia* **26**:2144–2146.
- Bandolik JJ, Hamacher A, Schrenk C, Weishaupt R, and Kassack MU (2019) Class I-Histone Deacetylase (HDAC) Inhibition is Superior to pan-HDAC Inhibition in Modulating Cisplatin Potency in High Grade Serous Ovarian Cancer Cell Lines. *Int J Mol Sci* **20**:3052.
- Barosi G, Bergamaschi G, Marchetti M, Vannucchi AM, Guglielmelli P, Antonioli E, Massa M, Rosti V, Campanelli R, Villani L, et al.; Gruppo Italiano Malattie Ematologiche Maligne dell'Adulto (GIMEMA) Italian Registry of Myelofibrosis (2007) JAK2 V617F mutational status predicts progression to large splenomegaly and leukemic transformation in primary myelofibrosis. *Blood* **110**:4030–4036.
- Barosi G, Zhang MJ, and Gale RP (2014) Does ruxolitinib improve survival of persons with MPN-associated myelofibrosis? Should it? *Leukemia* **28**:2267–2270.
- Baxter EJ, Scott LM, Campbell PJ, East C, Fourouclas N, Swanton S, Vassiliou GS, Bench AJ, Boyd EM, Curtin N, et al.; Cancer Genome Project (2005) Acquired mutation of the tyrosine kinase JAK2 in human myeloproliferative disorders. *Lancet* **365**:1054–1061.
- Beer PA, Delhommeau F, LeCouédic JP, Dawson MA, Chen E, Bareford D, Kusec R, McMullin MF, Harrison CN, Vannucchi AM, et al. (2010) Two routes to leukemic transformation after a JAK2 mutation-positive myeloproliferative neoplasm. *Blood* **115**:2891–2900.
- Belcher BP, Ward CC, and Nomura DK (2021) Ligandability of E3 Ligases for Targeted Protein Degradation Applications. *Biochemistry* DOI: 10.1021/acs.biochem.1c00464 [published ahead of print].
- Bellanger D, Jacquemin V, Chopin M, Pierron G, Bernard OA, Ghysdael J, and Stern MH (2014) Recurrent JAK1 and JAK3 somatic mutations in T-cell prolymphocytic leukemia. *Leukemia* **28**:417–419.
- Beyer M, Romanski A, Mustafa AM, Pons M, Büchler I, Vogel A, Pautz A, Sellmer A, Schneider G, Bug G, et al. (2019) HDAC3 Activity is Essential for Human Leukemic Cell Growth and the Expression of β -catenin, MYC, and WT1. *Cancers (Basel)* **11**:1436.
- Bharadwaj U, Kasembeli MM, Robinson P, and Tweardy DJ (2020) Targeting Janus kinases and signal transducer and activator of transcription 3 to treat inflammation, fibrosis, and cancer: rationale, progress, and caution. *Pharmacol Rev* **72**:486–526.
- Björkholm M, Derolf AR, Hultcrantz M, Kristinsson SY, Ekstrand C, Goldin LR, Andreasson B, Birgegård G, Linder O, Malm C, et al. (2011) Treatment-related risk factors for transformation to acute myeloid leukemia and myelodysplastic syndromes in myeloproliferative neoplasms. *J Clin Oncol* **29**:2410–2415.
- Bose P, Swaminathan M, Pemmaraju N, Ferrajoli A, Jabbour EJ, Daver NG, DiNardo CD, Alvarado Y, Yilmaz M, Huynh-Lu J, et al. (2019) A phase 2 study of pracinostat combined with ruxolitinib in patients with myelofibrosis. *Leuk Lymphoma* **60**:1767–1774.
- Bose P and Verstovsek S (2017) Developmental Therapeutics in Myeloproliferative Neoplasms. *Clin Lymphoma Myeloma Leuk* **17**:S43–S52.
- Bose P and Verstovsek S (2017b) JAK2 inhibitors for myeloproliferative neoplasms: what is next? *Blood* **130**:115–125.
- Bradner JE, Mak R, Tanguturi SK, Mazitschek R, Haggarty SJ, Ross K, Chang CY, Bosco J, West N, Morse E, et al. (2010) Chemical genetic strategy identifies histone deacetylase 1 (HDAC1) and HDAC2 as therapeutic targets in sickle cell disease. *Proc Natl Acad Sci USA* **107**:12617–12622.
- Briere D, Sudhakar N, Woods DM, Hallin J, Engstrom LD, Aranda R, Chiang H, Sodr  AL, Olson P, Weber JS, et al. (2018) The class I/IV HDAC inhibitor mocetinostat increases tumor antigen presentation, decreases immune suppressive cell types and augments checkpoint inhibitor therapy. *Cancer Immunol Immunother* **67**:381–392.
- Buchert M, Burns CJ, and Ernst M (2016) Targeting JAK kinase in solid tumors: emerging opportunities and challenges. *Oncogene* **35**:939–951.
- Buchwald M, Pietschmann K, Brand P, Günther A, Mahajan NP, Heinzl T, and Krämer OH (2013) SHAH ubiquitin ligases target the nonreceptor tyrosine kinase ACK1 for ubiquitinylation and proteasomal degradation. *Oncogene* **32**:4913–4920.
- Buchwald M, Pietschmann K, Müller JP, Böhmer FD, Heinzl T, and Krämer OH (2010) Ubiquitin conjugase UBCH8 targets active FMS-like tyrosine kinase 3 for proteasomal degradation. *Leukemia* **24**:1412–1421.

- Burke JR, Cheng L, Gillyooly KM, Strnad J, Zupa-Fernandez A, Catlett IM, Zhang Y, Heimrich EM, McIntyre KW, Cunningham MD, et al. (2019) Autoimmune pathways in mice and humans are blocked by pharmacological stabilization of the TYK2 pseudokinase domain. *Sci Transl Med* **11**:eaaw1736.
- Butler KV, Kalin J, Brochier C, Vistoli G, Langley B, and Koziowski AP (2010) Rational design and simple chemistry yield a superior, neuroprotective HDAC6 inhibitor, tubastatin A. *J Am Chem Soc* **132**:10842–10846.
- Cantó C, Menzies KJ, and Auwerx J (2015) NAD(+) Metabolism and the Control of Energy Homeostasis: A Balancing Act between Mitochondria and the Nucleus. *Cell Metab* **22**:31–53.
- Cao F, de Weerd S, Chen D, Zwinderman MRH, van der Wouden PE, and Dekker FJ (2020) Induced protein degradation of histone deacetylases 3 (HDAC3) by proteolysis targeting chimera (PROTAC). *Eur J Med Chem* **208**:112800.
- Cappellacci L, Perinelli DR, Maggi F, Grifantini M, and Petrelli R (2020) Recent Progress in Histone Deacetylase Inhibitors as Anticancer Agents. *Curr Med Chem* **27**:2449–2493.
- Cardoso BA, Belo H, Barata JT, and Almeida AM (2015) The Bone Marrow-Mediated Protection of Myeloproliferative Neoplastic Cells to Vorinostat and Ruxolitinib Relies on the Activation of JNK and PI3K Signaling Pathways. *PLoS One* **10**:e0143897.
- Ceccacci E and Minucci S (2016) Inhibition of histone deacetylases in cancer therapy: lessons from leukaemia. *Br J Cancer* **114**:605–611.
- Cerami E, Gao J, Dogrusoz U, Gross BE, Sumer SO, Aksoy BA, Jacobsen A, Byrne CJ, Heuer ML, Larsson E, et al. (2012) The cBio cancer genomics portal: an open platform for exploring multidimensional cancer genomics data. *Cancer Discov* **2**:401–404.
- Cervantes F and Pereira A (2017) Does ruxolitinib prolong the survival of patients with myelofibrosis? *Blood* **129**:832–837.
- Chang Y, Min J, Jarusiewicz JA, Actis M, Yu S, Mayasundari A, Yang L, Chepyala D, Alcock L, Roberts KG, et al. (2021) Degradation of Janus kinases in CRLF2-rearranged acute lymphoblastic leukemia. *Blood* **138**:2313–2326.
- Chen E, Ahn JS, Massie CE, Clynes D, Godfrey AL, Li J, Park HJ, Nangalia J, Silber Y, Mullally A, et al. (2014) JAK2V617F promotes replication fork stalling with disease-restricted impairment of the intra-S checkpoint response. *Proc Natl Acad Sci USA* **111**:15190–15195.
- Chen E, Ahn JS, Sykes DB, Breyfogle LJ, Godfrey AL, Nangalia J, Ko A, DeAngelo DJ, Green AR, and Mullally A (2015) RECQL5 Suppresses Oncogenic JAK2-Induced Replication Stress and Genomic Instability. *Cell Rep* **13**:2345–2352.
- Chen E, Staudt LM, and Green AR (2012) Janus kinase deregulation in leukemia and lymphoma. *Immunity* **36**:529–541.
- Chotitumnavee J, Yamashita Y, Takahashi Y, Takada Y, Iida T, Oba M, Itoh Y, and Suzuki T (2022) Selective degradation of histone deacetylase 8 mediated by a proteolysis targeting chimera (PROTAC). *Chem Commun (Camb)* **58**:4635–4638.
- Chu QS, Nielsen TO, Alcindor T, Gupta A, Endo M, Goytain A, Xu H, Verma S, Tozer R, Knowling M, et al. (2015) A phase II study of SB939, a novel pan-histone deacetylase inhibitor, in patients with translocation-associated recurrent/metastatic sarcomas-NCIC-CTG IND 200. *Ann Oncol* **26**:973–981.
- Chu-Farseeva YY, Mustafa N, Poulsen A, Tan EC, Yen JY, Chng WJ, and Dymock BW (2018) Design and synthesis of potent dual inhibitors of JAK2 and HDAC based on fusing the pharmacophores of XL019 and vorinostat. *Eur J Med Chem* **158**:593–619.
- Civallero M, Cosenza M, Pozzi S, and Sacchi S (2017) Ruxolitinib combined with vorinostat suppresses tumor growth and alters metabolic phenotype in hematological diseases. *Oncotarget* **8**:103797–103814.
- Corcoran RB, Contino G, Deshpande V, Tzatsos A, Conrad C, Benes CH, Levy DE, Sletteman J, Engelman JA, and Bardeesy N (2011) STAT3 plays a critical role in KRAS-induced pancreatic tumorigenesis. *Cancer Res* **71**:5020–5029.
- Cornago M, Garcia-Alberich C, Blasco-Angulo N, Vall-Llaura N, Nager M, Herreros J, Comella JX, Sanchis D, and Llovera M (2014) Histone deacetylase inhibitors promote glioma cell death by G2 checkpoint abrogation leading to mitotic catastrophe. *Cell Death Dis* **5**:e1435.
- Cortes JR, Patrone CC, Quinn SA, Gu Y, Sanchez-Martin M, Mackey A, Cooke AJ, Shih BB, Laurent AP, Trager MH, et al. (2021) Jak-STAT Inhibition Mediates Romidepsin and Mechlorethamine Synergism in Cutaneous T-Cell Lymphoma. *J Invest Dermatol* **141**:2908–2920.e7.
- Cosenza M, Civallero M, Marcheselli L, Sacchi S, and Pozzi S (2020) Citarinostat and Momelotinib co-target HDAC6 and JAK2/STAT3 in lymphoid malignant cell lines: a potential new therapeutic combination. *Apoptosis* **25**:370–387.
- Darnell Jr JE, Kerr IM, and Stark GR (1994) Jak-STAT pathways and transcriptional activation in response to IFNs and other extracellular signaling proteins. *Science* **264**:1415–1421.
- Davis N, McKinney MS, Reddy A, Love C, Smith E, Happ L, and Dave S (2018) Novel Mechanisms for Resistance to Targeted Therapy Identified through Machine Learning Approaches in 1167 RNA-Seq Drug Exposure Profiles in Lymphoma. *Blood* **132**:1370.
- Dawson MA, Bannister AJ, Göttgens B, Foster SD, Bartke T, Green AR, and Kouzarides T (2009) JAK2 phosphorylates histone H3Y41 and excludes HP1alpha from chromatin. *Nature* **461**:819–822.
- De Keersmaecker K, Atak ZK, Li N, Vicente C, Patchett S, Girardi T, Gianfelici V, Geerdens E, Clappier E, Porcu M, et al. (2013) Exome sequencing identifies mutation in CNOT3 and ribosomal genes RPL5 and RPL10 in T-cell acute lymphoblastic leukemia. *Nat Genet* **45**:186–190.
- DeAngelo DJ, Mesa RA, Fiskus W, Tefferi A, Paley C, Wadleigh M, Ritchie EK, Snyder DS, Begna K, Ganguly S, et al. (2013a) Phase II trial of panobinostat, an oral pan-deacetylase inhibitor in patients with primary myelofibrosis, post-essential thrombocythaemia, and post-polycythaemia vera myelofibrosis. *Br J Haematol* **162**:326–335.
- DeAngelo DJ, Spencer A, Bhalla KN, Prince HM, Fischer T, Kindler T, Giles FJ, Scott JW, Parker K, Liu A, et al. (2013b) Phase Ia/II, two-arm, open-label, dose-escalation study of oral panobinostat administered via two dosing schedules in patients with advanced hematologic malignancies. *Leukemia* **27**:1628–1636.
- Degryse S, Bornschein S, de Bock CE, Leroy E, Vanden Bempt M, Demeyer S, Jacobs K, Geerdens E, Gielen O, Soulier J, et al. (2018a) Mutant JAK3 signaling is increased by loss of wild-type JAK3 or by acquisition of secondary JAK3 mutations in T-ALL. *Blood* **131**:421–425.
- Degryse S, de Bock CE, Demeyer S, Govaerts I, Bornschein S, Verbeke D, Jacobs K, Binos S, Skerrett-Byrne DA, Murray HC, et al. (2018b) Mutant JAK3 phosphoproteomic profiling predicts synergism between JAK3 inhibitors and MEK/BCL2 inhibitors for the treatment of T-cell acute lymphoblastic leukemia. *Leukemia* **32**:788–800.
- Deng R, Zhang P, Liu W, Zeng X, Ma X, Shi L, Wang T, Yin Y, Chang W, Zhang P, et al. (2018) HDAC is indispensable for IFN- γ -induced B7-H1 expression in gastric cancer. *Clin Epigenetics* **10**:153.
- Depetter Y, Geurs S, De Vreese R, Goethals S, Vandoorn E, Laevens A, Steenbrugge J, Meyer E, de Tullio P, Bracke M, et al. (2019) Selective pharmacological inhibitors of HDAC6 reveal biochemical activity but functional tolerance in cancer models. *Int J Cancer* **145**:735–747.
- Downes CEJ, McClure BJ, Bruning JB, Page E, Breen J, Rehn J, Yeung DT, and White DL (2021) Acquired JAK2 mutations confer resistance to JAK inhibitors in cell models of acute lymphoblastic leukemia. *NPJ Precis Oncol* **5**:75.
- Dupriez B, Morel P, Demory J, Lai JL, Simon M, Plantier I, and Bauters F (1996) Prognostic factors in agnogenic myeloid metaplasia: a report on 195 cases with a new scoring system. *Blood* **88**:1013–1018.
- Elwood F, Witter DJ, Piesvaux J, Kraybill B, Bays N, Alpert C, Goldenblatt P, Qu Y, Ivanovska I, Lee HH, et al. (2017) Evaluation of JAK3 Biology in Autoimmune Disease Using a Highly Selective, Irreversible JAK3 Inhibitor. *J Pharmacol Exp Ther* **361**:229–244.
- Evrot E, Ebel N, Romanet V, Roelli C, Andraos R, Qian Z, Dölemeyer A, Damassa E, Sterker D, Cozens R, et al. (2013) JAK1/2 and Pan-deacetylase inhibitor combination therapy yields improved efficacy in preclinical mouse models of JAK2V617F-driven disease. *Clin Cancer Res* **19**:6230–6241.
- Falkenberg KJ and Johnstone RW (2014) Histone deacetylases and their inhibitors in cancer, neurological diseases and immune disorders. *Nat Rev Drug Discov* **13**:673–691.
- Fantin VR, Loboda A, Paweletz CP, Hendrickson RC, Pierce JW, Roth JA, Li L, Gooden F, Korenchuk S, Hou XS, et al. (2008) Constitutive activation of signal transducers and activators of transcription predicts vorinostat resistance in cutaneous T-cell lymphoma. *Cancer Res* **68**:3785–3794.
- Finazzi G, Caruso V, Marchioli R, Capnist G, Chisesi T, Finelli C, Gugliotta L, Landolfi R, Kutti J, Gisslinger H, et al.; ECLAP Investigators (2005) Acute leukemia in polycythemia vera: an analysis of 1638 patients enrolled in a prospective observational study. *Blood* **105**:2664–2670.
- Finazzi G, Vannucchi AM, Martinelli V, Ruggeri M, Nobile F, Specchia G, Pogliani EM, Olimpieri OM, Fioritoni G, Musolino C, et al. (2013) A phase II study of Givinostat in combination with hydroxycarbamide in patients with polycythemia vera unresponsive to hydroxycarbamide monotherapy. *Br J Haematol* **161**:688–694.
- Finnin MS, Doniphan JR, Cohen A, Richon VM, Rifkind RA, Marks PA, Breslow R, and Pavlentic NP (1999) Structures of a histone deacetylase homologue bound to the TSA and SAHA inhibitors. *Nature* **401**:188–193.
- Firmbach-Kraft I, Byers M, Shows T, Dalla-Favera R, and Krolewski JJ (1990) tyk2, prototype of a novel class of non-receptor tyrosine kinase genes. *Oncogene* **5**:1329–1336.
- Fiskus W, Verstovsek S, Manshour T, Rao R, Balusu R, Venkannagari S, Rao NN, Ha K, Smith JE, Hembruff SL, et al. (2011) Heat shock protein 90 inhibitor is synergistic with JAK2 inhibitor and overcomes resistance to JAK2-TKI in human myeloproliferative neoplasm cells. *Clin Cancer Res* **17**:7347–7358.
- Flex E, Petrangeli V, Stella L, Chiaretti S, Hornakova T, Knoops L, Ariola C, Fodale V, Clappier E, Paoloni F, et al. (2008) Somatic acquired JAK1 mutations in adult acute lymphoblastic leukemia. *J Exp Med* **205**:751–758.
- Fu RG, Sun Y, Sheng WB, and Liao DF (2017) Designing multi-targeted agents: An emerging anticancer drug discovery paradigm. *Eur J Med Chem* **136**:195–211.
- Funakoshi-Tago M, Pelletier S, Matsuda T, Parganas E, and Ihle JN (2006) Receptor specific downregulation of cytokine signaling by autophosphorylation in the FERM domain of Jak2. *EMBO J* **25**:4763–4772.
- Gaikwad A, Rye CL, Devidas M, Heerema NA, Carroll AJ, Israeli S, Plon SE, Basso G, Pession A, and Rabin KR (2009) Prevalence and clinical correlates of JAK2 mutations in Down syndrome acute lymphoblastic leukaemia. *Br J Haematol* **144**:930–932.
- Gakovik M, Ragimbeau J, Francois V, Constantinescu SN, and Pellegrini S (2008) The Stat3-activating Tyk2 V678F mutant does not up-regulate signaling through the type I interferon receptor but confers ligand hypersensitivity to a homodimeric receptor. *J Biol Chem* **283**:18522–18529.
- Gao J, Aksoy BA, Dogrusoz U, Dresdner G, Gross B, Sumer SO, Sun Y, Jacobsen A, Sinha R, Larsson E, et al. (2013) Integrative analysis of complex cancer genomics and clinical profiles using the cBioPortal. *Sci Signal* **6**:p1.
- Geissler K, Jäger E, Barna A, Sliwa T, Knöbl P, Schwarzwinger I, Gisslinger H, and Valent P (2016) In vitro and in vivo effects of JAK2 inhibition in chronic myelomonocytic leukemia. *Eur J Haematol* **97**:562–567.
- Ginter T, Bier C, Knauer SK, Sughra K, Hildebrand D, Münz T, Liebe T, Heller R, Henke A, Stauber RH, et al. (2012) Histone deacetylase inhibitors block IFN- γ -induced STAT1 phosphorylation. *Cell Signal* **24**:1453–1460.
- Girardi T, Vicente C, Cools J, and De Keersmaecker K (2017) The genetics and molecular biology of T-ALL. *Blood* **129**:1113–1123.
- Glassman CR, Tsutsumi N, Saxton RA, Lupardus PJ, Jude KM, and Garcia KC (2022) Structure of a Janus kinase cytokine receptor complex reveals the basis for dimeric activation. *Science* **376**:163–169.
- Göder A, Emmerich C, Nikolova T, Kiweler N, Schreiber M, Kühl T, Imhof D, Christmann M, Heinzl T, Schneider G, et al. (2018) HDAC1 and HDAC2 integrate

- checkpoint kinase phosphorylation and cell fate through the phosphatase-2A subunit PR130. *Nat Commun* **9**:764.
- Göder A, Ginter T, Heinzl T, Stroh S, Fahrer J, Henke A, and Krämer OH (2021) STAT1 N-terminal domain discriminatively controls type I and type II IFN signaling. *Cytokine* **144**:155552.
- Goedken ER, Argiriadi MA, Banach DL, Fiamengo BA, Foley SE, Frank KE, George JS, Harris CM, Hobson AD, Ihle DC, et al. (2015) Tricyclic covalent inhibitors selectively target Jak3 through an active site thiol. *J Biol Chem* **290**:4573–4589.
- Gorantla SP, Dechow TN, Grundler R, Illert AL, Zum Büschenfelde CM, Kremer M, Peschel C, and Duyster J (2010) Oncogenic JAK2V617F requires an intact SH2-like domain for constitutive activation and induction of a myeloproliferative disease in mice. *Blood* **116**:4600–4611.
- Göttlicher M, Minucci S, Zhu P, Krämer OH, Schimpf A, Giavara S, Sleeman JP, Lo Coco F, Nervi C, Pelicci PG, et al. (2001) Valproic acid defines a novel class of HDAC inhibitors inducing differentiation of transformed cells. *EMBO J* **20**:6969–6978.
- Greenplate A, Wang K, Tripathi RM, Palma N, Ali SM, Stephens PJ, Miller VA, Shyr Y, Guo Y, Reddy NM, et al. (2018) Genomic profiling of T-cell neoplasms reveals frequent JAK1 and JAK3 mutations with clonal evasion from targeted therapies. *JCO Precis Oncol* **2018**:1–16.
- Griffiths DS, Li J, Dawson MA, Trotter MW, Cheng YH, Smith AM, Mansfield W, Liu P, Kouzarides T, Nichols J, et al. (2011) LIF-independent JAK signalling to chromatin in embryonic stem cells uncovered from an adult stem cell disease. *Nat Cell Biol* **13**:13–21.
- Grossman RL, Heath AP, Ferretti V, Varmus HE, Lowy DR, Kibbe WA, and Staudt LM (2016) Toward a shared vision for cancer genomic data. *N Engl J Med* **375**:1109–1112.
- Guerini V, Barbu V, Spinelli O, Salvi A, Dellacasa C, Carobbio A, Introna M, Barbu T, Golay J, and Rambaldi A (2008) The histone deacetylase inhibitor ITF2357 selectively targets cells bearing mutated JAK2(V617F). *Leukemia* **22**:740–747.
- Gupta M, Han JJ, Stenson M, Wellik L, and Witzig TE (2010) HDAC Class I Inhibition Acetylates a Non-Histone Protein STAT3 by Modulating p300-STAT3-HDAC1 Interaction in Activated B-Cell Like (ABC) Diffuse Large B-Cell Lymphoma. *Blood* **116**:115.
- Gupta M, Han JJ, Stenson M, Wellik L, and Witzig TE (2012) Regulation of STAT3 by histone deacetylase-3 in diffuse large B-cell lymphoma: implications for therapy. *Leukemia* **26**:1356–1364.
- Hagopian HK, Riggs MG, Swartz LA, and Ingram VM (1977) Effect of n-butyrate on DNA synthesis in chick fibroblasts and HeLa cells. *Cell* **12**:855–860.
- Hammarén HM, Virtanen AT, Raivola J, and Silvennoinen O (2019) The regulation of JAKs in cytokine signaling and its breakdown in disease. *Cytokine* **118**:48–63.
- Hao X, Xing W, Yuan J, Wang Y, Bai J, Bai J, and Zhou Y (2020) Cotargeting the JAK/STAT signaling pathway and histone deacetylase by ruxolitinib and vorinostat elicits synergistic effects against myeloproliferative neoplasms. *Invest New Drugs* **38**:610–620.
- Harrison CN, Kiladjan J-J, Heidel FH, Vannucchi AM, Passamonti F, Hayat A, Conneally E, Martino B, Kindler T, Lipka DB, et al. (2015) Efficacy, Safety, and Confirmation of the Recommended Phase 2 Starting Dose of the Combination of Ruxolitinib (RUX) and Panobinostat (PAN) in Patients (Pts) with Myelofibrosis (MF). *Blood* **126**:4060.
- Harrison CN, Vannucchi AM, Kiladjan JJ, Al-Ali HK, Gisslinger H, Knoops L, Cervantes F, Jones MM, Sun K, McQuitty M, et al. (2016) Long-term findings from COMFORT-II, a phase 3 study of ruxolitinib vs best available therapy for myelofibrosis. *Leukemia* **30**:1701–1707.
- Harry BL, Eckhardt SG, and Jimeno A (2012) JAK2 inhibition for the treatment of hematologic and solid malignancies. *Expert Opin Investig Drugs* **21**:637–655.
- He L, Pei H, Lan T, Tang M, Zhang C, and Chen L (2017) Design and Synthesis of a Highly Selective JAK3 Inhibitor for the Treatment of Rheumatoid Arthritis. *Arch Pharm (Weinheim)* **350**.
- He L, Shao M, Wang T, Lan T, Zhang C, and Chen L (2018) Design, synthesis, and SAR study of highly potent, selective, irreversible covalent JAK3 inhibitors. *Mol Divers* **22**:343–358.
- Helbig G (2018) Classical Philadelphia-negative myeloproliferative neoplasms: focus on mutations and JAK2 inhibitors. *Med Oncol* **35**:119.
- Herreros-Villanueva M, Garcia-Girón C, and Er TK (2010) No evidence for JAK2 V617F mutation in colorectal cancer. *Br J Biomed Sci* **67**:220–222.
- Hirbe AC, Kaushal M, Sharma MK, Dahiya S, Pekmezci M, Perry A, and Gutmann DH (2017) Clinical genomic profiling identifies TYK2 mutation and overexpression in patients with neurofibromatosis type 1-associated malignant peripheral nerve sheath tumors. *Cancer* **123**:1194–1201.
- Hornakova T, Staerk J, Royer Y, Flex E, Tartaglia M, Constantinescu SN, Knoops L, and Renaud JC (2009) Acute lymphoblastic leukemia-associated JAK1 mutants activate the Janus kinase/STAT pathway via interleukin-9 receptor alpha homodimers. *J Biol Chem* **284**:6773–6781.
- Hu C, Xia H, Bai S, Zhao J, Edwards H, Li X, Yang Y, Lyu J, Wang G, Zhan Y, et al. (2020) CUDC-907, a novel dual PI3K and HDAC inhibitor, in prostate cancer: Antitumor activity and molecular mechanism of action. *J Cell Mol Med* **24**:7239–7253.
- Huang Y, Dong G, Li H, Liu N, Zhang W, and Sheng C (2018a) Discovery of Janus Kinase 2 (JAK2) and Histone Deacetylase (HDAC) Dual Inhibitors as a Novel Strategy for the Combinational Treatment of Leukemia and Invasive Fungal Infections. *J Med Chem* **61**:6056–6074.
- Huang Z, Zhou W, Li Y, Cao M, Wang T, Ma Y, Guo Q, Wang X, Zhang C, Zhang C, et al. (2018b) Novel hybrid molecule overcomes the limited response of solid tumours to HDAC inhibitors by suppressing JAK1-STAT3-BCL2 signalling. *Theranostics* **8**:4995–5011.
- Ide H, Nakagawa T, Terado Y, Kamiyama Y, Muto S, and Horie S (2008) Tyk2 expression and its signaling enhances the invasiveness of prostate cancer cells. *Biochem Biophys Res Commun* **369**:292–296.
- Ihle JN, Witthuhn BA, Quelle FW, Yamamoto K, and Silvennoinen O (1995) Signaling through the hematopoietic cytokine receptors. *Annu Rev Immunol* **13**:369–398.
- Imai Y, Maru Y, and Tanaka J (2016) Action mechanisms of histone deacetylase inhibitors in the treatment of hematological malignancies. *Cancer Sci* **107**:1543–1549.
- James C, Ugo V, Le Couédic JP, Staerk J, Delhommeau F, Lacout C, Garçon L, Raslova H, Berger R, Bennaceur-Griscelli A, et al. (2005) A unique clonal JAK2 mutation leading to constitutive signalling causes polycythaemia vera. *Nature* **434**:1144–1148.
- Jensen MA, Ferretti V, Grossman RL, and Staudt LM (2017) The NCI Genomic Data Commons as an engine for precision medicine. *Blood* **130**:453–459.
- Jeong EG, Kim MS, Nam HK, Min CK, Lee S, Chung YJ, Yoo NJ, and Lee SH (2008) Somatic mutations of JAK1 and JAK3 in acute leukemias and solid cancers. *Clin Cancer Res* **14**:3716–3721.
- Jiang M, Zhang WW, Liu P, Yu W, Liu T, and Yu J (2017a) Dysregulation of SOCS-Mediated Negative Feedback of Cytokine Signaling in Carcinogenesis and Its Significance in Cancer Treatment. *Front Immunol* **8**:70.
- Jiang Y, Liu J, Chen D, Yan L, and Zheng W (2017b) Sirtuin Inhibition: Strategies, Inhibitors, and Therapeutic Potential. *Trends Pharmacol Sci* **38**:459–472.
- Juengel E, Bhasin M, Libermann T, Barth S, Michaelis M, Cinatl Jr J, Jones J, Hudak L, Jonas D, and Blaheta RA (2011) Alterations of the gene expression profile in renal cell carcinoma after treatment with the histone deacetylase-inhibitor valproic acid and interferon-alpha. *World J Urol* **29**:779–786.
- Kalac M, Scotto L, Marchi E, Amengual J, Seshan VE, Bhagat G, Ulahannan N, Leshchenko VV, Temkin AM, Parekh S, et al. (2011) HDAC inhibitors and decitabine are highly synergistic and associated with unique gene-expression and epigenetic profiles in models of DLBCL. *Blood* **118**:5506–5516.
- Kaminker JS, Zhang Y, Waugh A, Haverty PM, Peters B, Sebanovic D, Stinson J, Forrest WF, Bazan JF, Seshagiri S, et al. (2007) Distinguishing cancer-associated missense mutations from common polymorphisms. *Cancer Res* **67**:465–473.
- Kan Z, Zheng H, Liu X, Li S, Barber TD, Gong Z, Gao H, Hao K, Willard MD, Xu J, et al. (2013) Whole-genome sequencing identifies recurrent mutations in hepatocellular carcinoma. *Genome Res* **23**:1422–1433.
- Kaowinn S, Jun SW, Kim CS, Shin DM, Hwang YH, Kim K, Shin B, Kaewpiboon C, Jeong HH, Koh SS, et al. (2017) Increased EGFR expression induced by a novel oncogene, CUG2, confers resistance to doxorubicin through Stat1-HDAC4 signaling. *Clin Oncol (Dordr)* **40**:549–561.
- Karantanos T and Moliterno AR (2018) The roles of JAK2 in DNA damage and repair in the myeloproliferative neoplasms: Opportunities for targeted therapy. *Blood Rev* **32**:426–432.
- Kearney L, Gonzalez De Castro D, Yeung J, Procter J, Horsley SW, Eguchi-Ishimae M, Bateman CM, Anderson K, Chaplin T, Young BD, et al. (2009) Specific JAK2 mutation (JAK2R683) and multiple gene deletions in Down syndrome acute lymphoblastic leukemia. *Blood* **113**:646–648.
- Kempson J, Ovalle D, Guo J, Wroblewski ST, Lin S, Spergel SH, Duan JJ, Jiang B, Lu Z, Das J, et al. (2017) Discovery of highly potent, selective, covalent inhibitors of JAK3. *Bioorg Med Chem Lett* **27**:4622–4625.
- Khan S, Zhang X, Lv D, Zhang Q, He Y, Zhang P, Liu X, Thummuri D, Yuan Y, Wiegand JS, et al. (2019) A selective BCL-X_L PROTAC degrader achieves safe and potent antitumor activity. *Nat Med* **25**:1938–1947.
- Khandelwal P, Teusink-Cross A, Davies SM, Nelson AS, Dandoy CE, El-Bietar J, Marsh RA, Kumar AR, Grimley MS, Jodele S, et al. (2017) Ruxolitinib as salvage therapy in steroid-refractory acute graft-versus-host disease in pediatric hematopoietic stem cell transplant patients. *Biol Blood Marrow Transplant* **23**:1122–1127.
- Kijima M, Yoshida M, Sugita K, Horinouchi S, and Beppu T (1993) Trapoxin, an antitumor cyclic tetrapeptide, is an irreversible inhibitor of mammalian histone deacetylase. *J Biol Chem* **268**:22429–22435.
- Kim EH, Cao D, Mahajan NP, Andriole GL, and Mahajan K (2020) ACK1-AR and AR-HOXB13 signaling axes: epigenetic regulation of lethal prostate cancers. *NAR Cancer* **2**:zcaa018.
- Kim H-J, Ko YH, Kim JE, Lee S-S, Lee H, Park G, Paik JH, Cha HJ, Choi Y-D, Han JH, et al.; Hematopathology Study Group of the Korean Society of Pathologists (2017) Epstein-Barr virus-associated lymphoproliferative disorders: review and update on 2016 WHO classification. *J Pathol Transl Med* **51**:352–358.
- Klampfer L, Huang J, Swaby LA, and Augenlicht L (2004) Requirement of histone deacetylase activity for signaling by STAT1. *J Biol Chem* **279**:30358–30368.
- Koeneke E, Witt O, and Oehme I (2015) HDAC Family Members Intertwined in the Regulation of Autophagy: A Druggable Vulnerability in Aggressive Tumor Entities. *Cells* **4**:135–168.
- Koo GC, Tan SY, Tang T, Poon SL, Allen GE, Tan L, Chong SC, Ong WS, Tay K, Tao M, et al. (2012) Janus kinase 3-activating mutations identified in natural killer/T-cell lymphoma. *Cancer Discov* **2**:591–597.
- Koppikar P, Bhagwat N, Kilpivaara O, Manshoury T, Adli M, Hricik T, Liu F, Saunders LM, Mullally A, Abdel-Wahab O, et al. (2012) Heterodimeric JAK-STAT activation as a mechanism of persistence to JAK2 inhibitor therapy. *Nature* **489**:155–159.
- Kotla S, Singh NK, and Rao GN (2017) ROS via BTK-p300-STAT1-PPAR γ signaling activation mediates cholesterol crystals-induced CD36 expression and foam cell formation. *Redox Biol* **11**:350–364.
- Kralovics R, Passamonti F, Buser AS, Teo SS, Tiedt R, Passweg JR, Tichelli A, Cazzola M, and Skoda RC (2005) A gain-of-function mutation of JAK2 in myeloproliferative disorders. *N Engl J Med* **352**:1779–1790.
- Krämer OH (2009) HDAC2: a critical factor in health and disease. *Trends Pharmacol Sci* **30**:647–655.
- Krämer OH, Baus D, Knauer SK, Stein S, Jäger E, Stauber RH, Grez M, Pfitzner E, and Heinzl T (2006) Acetylation of Stat1 modulates NF-kappaB activity. *Genes Dev* **20**:473–485.
- Krämer OH, Göttlicher M, and Heinzl T (2001) Histone deacetylase as a therapeutic target. *Trends Endocrinol Metab* **12**:294–300.

- Krämer OH and Heinzel T (2010) Phosphorylation-acetylation switch in the regulation of STAT1 signaling. *Mol Cell Endocrinol* **315**:40–48.
- Krämer OH, Knauer SK, Greiner G, Jandt E, Reichardt S, Gührs KH, Stauber RH, Böhrer FD, and Heinzel T (2009) A phosphorylation-acetylation switch regulates STAT1 signaling. *Genes Dev* **23**:223–235.
- Krämer OH, Mahboobi S, and Sellmer A (2014) Drugging the HDAC6-HSP90 interplay in malignant cells. *Trends Pharmacol Sci* **35**:501–509.
- Krämer OH and Schneider G (2022) Single-cell profiling guided combination therapy of c-Fos and histone deacetylase inhibitors in diffuse large B-cell lymphoma. *Clin Transl Med* **12**:e858.
- Krämer OH, Stauber RH, Bug G, Hartkamp J, and Knauer SK (2013) SIAH proteins: critical roles in leukemogenesis. *Leukemia* **27**:792–802.
- Kratz CP, Böll S, Kontny U, Schrappe M, Niemeyer CM, and Stanulla M (2006) Mutational screen reveals a novel JAK2 mutation, L611S, in a child with acute lymphoblastic leukemia. *Leukemia* **20**:381–383.
- Krolewski JJ, Lee R, Eddy R, Shows TB, and Dalla-Favera R (1990) Identification and chromosomal mapping of new human tyrosine kinase genes. *Oncogene* **5**:277–282.
- Kurosu T, Nagao T, Wu N, Oshikawa G, and Miura O (2013) Inhibition of the PI3K/Akt/GSK3 pathway downstream of BCR/ABL, Jak2-V617F, or FLT3-ITD downregulates DNA damage-induced Chk1 activation as well as G2/M arrest and prominently enhances induction of apoptosis. *PLoS One* **8**:e79478.
- Kwon HJ, Kim MS, Kim MJ, Nakajima H, and Kim KW (2002) Histone deacetylase inhibitor FK228 inhibits tumor angiogenesis. *Int J Cancer* **97**:290–296.
- Lahm A, Paolini C, Pallaoro M, Nardi MC, Jones P, Neddermann P, Sambucini S, Bottomley MJ, Lo Surdo P, Carfi A, et al. (2007) Unraveling the hidden catalytic activity of vertebrate class IIa histone deacetylases. *Proc Natl Acad Sci USA* **104**:17335–17340.
- Laschanky RS, Humphrey LE, Ma J, Smith LM, Enke TJ, Shukla SK, Dasgupta A, Singh PK, Howell GM, Brattain MG, et al. (2019) Selective Inhibition of Histone Deacetylases 1/2/6 in Combination with Gemcitabine: A Promising Combination for Pancreatic Cancer Therapy. *Cancers (Basel)* **11**:1327.
- Lechner S, Malgou MIP, Grätz C, Steimbach RR, Baron A, Rütger P, Nadal S, Stumpf C, Loos C, Ku X, et al. (2022) Target deconvolution of HDAC pharmacopoeia reveals MBLAC2 as common off-target. *Nat Chem Biol* **18**:1–9.
- Lee JW, Soung YH, Kim SY, Nam SW, Park WS, Lee JY, Yoo NJ, and Lee SH (2006) Absence of JAK2 V617F mutation in gastric cancers. *Acta Oncol* **45**:222–223.
- Leitner NR, Witalisz-Siepracka A, Strobl B, and Müller M (2017) Tyrosine kinase 2 - Surveillant of tumours and bona fide oncogene. *Cytokine* **89**:209–218.
- Leonard WJ and O'Shea JJ (1998) Jaks and STATs: biological implications. *Annu Rev Immunol* **16**:293–322.
- Leonhardt M, Sellmer A, Krämer OH, Dove S, Elz S, Kraus B, Beyer M, and Mahboobi S (2018) Design and biological evaluation of tetrahydro- β -carboline derivatives as highly potent histone deacetylase 6 (HDAC6) inhibitors. *Eur J Med Chem* **152**:329–357.
- Levine RL, Pardanani A, Tefferi A, and Gilliland DG (2007) Role of JAK2 in the pathogenesis and therapy of myeloproliferative disorders. *Nat Rev Cancer* **7**:673–683.
- Levine RL, Wadleigh M, Cools J, Ebert BL, Wernig G, Huntly BJ, Boggon TJ, Wlodarska I, Clark JJ, Moore S, et al. (2005) Activating mutation in the tyrosine kinase JAK2 in polycythemia vera, essential thrombocythemia, and myeloid metaplasia with myelofibrosis. *Cancer Cell* **7**:387–397.
- Li H, Ahonen TJ, Alanen K, Xie J, LeBaron MJ, Pretlow TG, Ealley EL, Zhang Y, Nurmi M, Singh B, et al. (2004) Activation of signal transducer and activator of transcription 5 in human prostate cancer is associated with high histological grade. *Cancer Res* **64**:4774–4782.
- Li Q, Li B, Hu L, Ning H, Jiang M, Wang D, Liu T, Zhang B, and Chen H (2017) Identification of a novel functional JAK1 S646P mutation in acute lymphoblastic leukemia. *Oncotarget* **8**:34687–34697.
- Li Y, Buijss-Gladdines JG, Canté-Barrett K, Stubbs AP, Vroegindewij EM, Smits WK, van Marion R, Dinjens WN, Horstmann M, Kuiper RP, et al. (2016) IL-7 Receptor Mutations and Steroid Resistance in Pediatric T cell Acute Lymphoblastic Leukemia: A Genome Sequencing Study. *PLoS Med* **13**:e1002200.
- Li Y, Wang F, Chen X, Wang J, Zhao Y, Li Y, and He B (2019) Zinc-dependent Deacetylase (HDAC) Inhibitors with Different Zinc Binding Groups. *Curr Top Med Chem* **19**:223–241.
- Liang X, Tang S, Liu X, Liu Y, Xu Q, Wang X, Saidahmatov A, Li C, Wang J, Zhou Y, et al. (2022) Discovery of Novel Pyrrolo[2,3-d]pyrimidine-based Derivatives as Potent JAK/HDAC Dual Inhibitors for the Treatment of Refractory Solid Tumors. *J Med Chem* **65**:1243–1264.
- Liang X, Zang J, Li X, Tang S, Huang M, Geng M, Chou CJ, Li C, Cao Y, Xu W, et al. (2019) Discovery of Novel Janus Kinase (JAK) and Histone Deacetylase (HDAC) Dual Inhibitors for the Treatment of Hematological Malignancies. *J Med Chem* **62**:3898–3923.
- Licht V, Noack K, Schlott B, Förster M, Schlenker Y, Licht A, Krämer OH, and Heinzel T (2014) Caspase-3 and caspase-6 cleave STAT1 in leukemic cells. *Oncotarget* **5**:2305–2317.
- Lindahl LM, Fredholm S, Joseph C, Nielsen BS, Jønson L, Willerslev-Olsen A, Ghud M, Blümel E, Petersen DL, Sibbesen N, et al. (2016) STAT5 induces miR-21 expression in cutaneous T cell lymphoma. *Oncotarget* **7**:45730–45744.
- Liu J, Jin L, Chen X, Yuan Y, Zuo Y, Miao Y, Feng Q, Zhang H, Huang F, Guo T, et al. (2020) USP12 translocation maintains interferon antiviral efficacy by inhibiting CBP acetyltransferase activity. *PLoS Pathog* **16**:e1008215.
- Liu L, Damerell DR, Koukoulis L, Tong Y, Marsden BD, and Schapira M (2019) UbiHub: a data hub for the explorers of ubiquitination pathways. *Bioinformatics* **35**:2882–2884.
- Liu Y, Easton J, Shao Y, Maciaszek J, Wang Z, Wilkinson MR, McCastlain K, Edmonson M, Pounds SB, Shi L, et al. (2017) The genomic landscape of pediatric and young adult T-lineage acute lymphoblastic leukemia. *Nat Genet* **49**:1211–1218.
- Low LK and Song JY (2016) B-cell lymphoproliferative disorders associated with primary and acquired immunodeficiency. *Surg Pathol Clin* **9**:55–77.
- Lupardus PJ, Ullsch M, Wallweber H, Bir Köhli P, Johnson AR, and Eigenbrot C (2014) Structure of the pseudokinase-kinase domains from protein kinase TYK2 reveals a mechanism for Janus kinase (JAK) autoinhibition. *Proc Natl Acad Sci USA* **111**:8025–8030.
- Mahajan K, Fang B, Koomen JM, and Mahajan NP (2012) H2B Tyr37 phosphorylation suppresses expression of replication-dependent core histone genes. *Nat Struct Mol Biol* **19**:930–937.
- Mahajan K, Malla P, Lawrence HR, Chen Z, Kumar-Sinha C, Malik R, Shukla S, Kim J, Coppola D, Lawrence NJ, et al. (2017) ACK1/TNK2 Regulates Histone H4 Tyr88-phosphorylation and AR Gene Expression in Castration-Resistant Prostate Cancer. *Cancer Cell* **31**:790–803.e8.
- Mahendrarajah N, Paulus R, and Krämer OH (2016) Histone deacetylase inhibitors induce proteolysis of activated CDC42-associated kinase-1 in leukemic cells. *J Cancer Res Clin Oncol* **142**:2263–2273.
- Malinge S, Ragu C, Della-Valle V, Pisani D, Constantinescu SN, Perez C, Villeval JL, Reinhardt D, Landman-Parker J, Michaux L, et al. (2008) Activating mutations in human acute megakaryoblastic leukemia. *Blood* **112**:4220–4226.
- Mao YL, Li ZW, Lou CJ, Pang D, and Zhang YQ (2011) Phospho-STAT5 expression is associated with poor prognosis of human colonic adenocarcinoma. *Pathol Oncol Res* **17**:333–339.
- Marcellino BK, Verstovsek S, and Mascarenhas J (2020) The Myelodepletive Phenotype in Myelofibrosis: Clinical Relevance and Therapeutic Implication. *Clin Lymphoma Myeloma Leuk* **20**:415–421.
- Marty C, Lacout C, Droin N, Le Couédic JP, Ribrag V, Solary E, Vainchenker W, Villeval JL, and Plo I (2013) A role for reactive oxygen species in JAK2 V617F myeloproliferative neoplasm progression. *Leukemia* **27**:2187–2195.
- Mascarenhas J, Lu M, Li T, Petersen B, Hochman T, Najfeld V, Goldberg JD, and Hoffman R (2013) A phase I study of panobinostat (LBH589) in patients with primary myelofibrosis (PMF) and post-polycythemia vera/essential thrombocythemia myelofibrosis (post-PV/ET MF). *Br J Haematol* **161**:68–75.
- Mascarenhas J, Marcellino BK, Lu M, Kremianskaya M, Fabris F, Sandy L, Mehrotra M, Houldsworth J, Najfeld V, El Jamal S, et al. (2020) A phase I study of panobinostat and ruxolitinib in patients with primary myelofibrosis (PMF) and post-polycythemia vera/essential thrombocythemia myelofibrosis (post-PV/ET MF). *Leuk Res* **88**:106272.
- Mascarenhas J, Sandy L, Lu M, Yoon J, Petersen B, Zhang D, Ye F, Newsom C, Najfeld V, Hochman T, et al. (2017) A phase II study of panobinostat in patients with primary myelofibrosis (PMF) and post-polycythemia vera/essential thrombocythemia myelofibrosis (post-PV/ET MF). *Leuk Res* **53**:13–19.
- McKinney MS, Beaven AW, Moffitt A, Smith JL, Lock E, Jima D, Healy J, Li G, Greenough A, Love CL, et al. (2014) Chemical Genomics Reveals JAK STAT Activation As a Mechanism of Resistance to HDAC Inhibitors in B Cell Lymphomas. *Blood* **124**:271.
- Meyer FB, Marx C, Spangel SB, and Thierbach R (2021) Butyrate and Metformin Affect Energy Metabolism Independently of the Metabolic Phenotype in the Tumor Therapy Model. *Biomolecules* **11**:1831.
- Meyer SC, Keller MD, Chiu S, Koppikar P, Guryanova OA, Rapaport F, Xu K, Manova K, Pankov D, O'Reilly RJ, et al. (2015) CHZ868, a Type II JAK2 Inhibitor, Reverses Type I JAK Inhibitor Persistence and Demonstrates Efficacy in Myeloproliferative Neoplasms. *Cancer Cell* **28**:15–28.
- Michaelis M, Doerr HW, and Cinatl Jr J (2007) Valproic acid as anti-cancer drug. *Curr Pharm Des* **13**:3378–3393.
- Morphy R, Kay C, and Rankovic Z (2004) From magic bullets to designed multiple ligands. *Drug Discov Today* **9**:641–651.
- Morphy R and Rankovic Z (2005) Designed multiple ligands. An emerging drug discovery paradigm. *J Med Chem* **48**:6523–6543.
- Müller S, Chen Y, Ginter T, Schäfer C, Buchwald M, Schmitz LM, Klitzsch J, Schütz A, Haitel A, Schmid K, et al. (2014) SLAH2 antagonizes TYK2-STAT3 signaling in lung carcinoma cells. *Oncotarget* **5**:3184–3196.
- Mullighan CG, Collins-Underwood JR, Phillips LA, Loudin MG, Liu W, Zhang J, Ma J, Coustan-Smith E, Harvey RC, Willman CL, et al. (2009a) Rearrangement of CRLF2 in B-progenitor- and Down syndrome-associated acute lymphoblastic leukemia. *Nat Genet* **41**:1243–1246.
- Mullighan CG, Zhang J, Harvey RC, Collins-Underwood JR, Schulman BA, Phillips LA, Tasian SK, Loh ML, Su X, Liu W, et al. (2009b) JAK mutations in high-risk childhood acute lymphoblastic leukemia. *Proc Natl Acad Sci USA* **106**:9414–9418.
- Murtuza A, Bulbul A, Shen JP, Keshavarzian P, Woodward BD, Lopez-Diaz FJ, Lippman SM, and Husain H (2019) Novel third-generation EGFR tyrosine kinase inhibitors and strategies to overcome therapeutic resistance in lung cancer. *Cancer Res* **79**:689–698.
- Musumeci F, Greco C, Giachello I, Fallacara AL, Ibrahim MM, Grossi G, Brullo C, and Schenone S (2019) An Update on JAK Inhibitors. *Curr Med Chem* **26**:1806–1832.
- Nawar N, Bukhari S, Adile AA, Suk Y, Manaswiyoungkul P, Toutah K, Olaoye OO, Raouf YS, Sedighi A, Garcha HK, et al. (2022) Discovery of HDAC6-Selective Inhibitor NN-390 with *in Vitro* Efficacy in Group 3 Medulloblastoma. *J Med Chem* **65**:3193–3217.
- Nebbio A, Carafa V, Conte M, Tambaro FP, Abbondanza C, Martens J, Nees M, Benedetti R, Pallavicini I, Minucci S, et al. (2017) c-Myc Modulation and Acetylation Is a Key HDAC Inhibitor Target in Cancer. *Clin Cancer Res* **23**:2542–2555.
- Nepali K, Sharma S, Sharma M, Bedi PM, and Dhar KL (2014) Rational approaches, design strategies, structure activity relationship and mechanistic insights for anticancer hybrids. *Eur J Med Chem* **77**:422–487.
- Nikolova T, Kieweler N, and Krämer OH (2017) Interstrand Crosslink Repair as a Target for HDAC Inhibition. *Trends Pharmacol Sci* **38**:822–836.

- Ning C-Q, Lu C, Hu L, Bi Y-J, Yao L, He Y-J, Liu L-F, Liu X-Y, and Yu N-F (2015) Macrocyclic compounds as anti-cancer agents: design and synthesis of multi-acting inhibitors against HDAC, FLT3 and JAK2. *Eur J Med Chem* **95**:104–115.
- Niwa Y, Kanda H, Shikauchi Y, Saiura A, Matsubara K, Kitagawa T, Yamamoto J, Kubo T, and Yoshikawa H (2005) Methylation silencing of SOCS-3 promotes cell growth and migration by enhancing JAK/STAT and FAK signalings in human hepatocellular carcinoma. *Oncogene* **24**:6406–6417.
- Noack K, Mahendrarajah N, Hennig D, Schmidt L, Grebien F, Hildebrand D, Christmann M, Kaina B, Sellmer A, Mahboobi S, et al. (2017) Analysis of the interplay between all-trans retinoic acid and histone deacetylase inhibitors in leukemic cells. *Arch Toxicol* **91**:2191–2208.
- Novotny-Diermayr V, Hart S, Goh KC, Cheong A, Ong LC, Hentze H, Pasha MK, Jayaraman R, Ethirajulu K, and Wood JM (2012) The oral HDAC inhibitor pracinostat (SB939) is efficacious and synergistic with the JAK2 inhibitor pacritinib (SB1518) in preclinical models of AML. *Blood Cancer J* **2**:e69.
- Nowak RP, DeAngelo SL, Buckley D, He Z, Donovan KA, An J, Safaei N, Jedrychowski MP, Ponthier CM, Isohey M, et al. (2018) Plasticity in binding confers selectivity in ligand-induced protein degradation. *Nat Chem Biol* **14**:706–714.
- Nwaogu A, Bond A, and Smith PJ (2020) Guideline review: Tofacitinib for adults with moderately to severely active ulcerative colitis - NICE guidance. *Frontline Gastroenterol* **12**:133–136.
- Orazi A, Hasserjian R, Cazzola M, Thiele M, and Malcovati L (2017) *Myelodysplastic/myeloproliferative neoplasm with ring sideroblasts and thrombocytosis. WHO Classification of Tumours of Haematopoietic and Lymphoid Tissues*, revised 4th ed, pp 93–94, International Agency for Research on Cancer, Lyon.
- Orillion A, Hashimoto A, Damayanti N, Shen L, Adelaiye-Ogala R, Arisa S, Chintala S, Ordentlich P, Kao C, Elzey B, et al. (2017) Entinostat Neutralizes Myeloid-Derived Suppressor Cells and Enhances the Antitumor Effect of PD-1 Inhibition in Murine Models of Lung and Renal Cell Carcinoma. *Clin Cancer Res* **23**:5187–5201.
- Owen KL, Brockwell NK, and Parker BS (2019) JAK-STAT signaling: A double-edged sword of immune regulation and cancer progression. *Cancers (Basel)* **11**:2002.
- Owusu BY and Klampfer L (2017) Modulation of STAT1-Driven Transcriptional Activity by Histone Deacetylases. *Methods Mol Biol* **1510**:277–285.
- Padron E, Dezern A, Andrade-Campos M, Vaddi K, Scherle P, Zhang Q, Ma Y, Balasis ME, Tinsley S, Ramadan H, et al.; Myelodysplastic Syndrome Clinical Research Consortium (2016) A Multi-Institution Phase I Trial of Ruxolitinib in Patients with Chronic Myelomonocytic Leukemia (CMML). *Clin Cancer Res* **22**:3746–3754.
- Pardanani A, Hood J, Lasho T, Levine RL, Martin MB, Noronha G, Finke C, Mak CC, Mesa R, Zhu H, et al. (2007) TG101209, a small molecule JAK2-selective kinase inhibitor potently inhibits myeloproliferative disorder-associated JAK2V617F and MPLW515L/K mutations. *Leukemia* **21**:1658–1668.
- Patnaik MM and Tefferi A (2015) Refractory anemia with ring sideroblasts and RARS with thrombocytosis. *Am J Hematol* **90**:549–559.
- Pellegrini S and Dusanter-Fourt I (1997) The structure, regulation and function of the Janus kinases (JAKs) and the signal transducers and activators of transcription (STATs). *Eur J Biochem* **248**:615–633.
- Peters JU (2013) Polypharmacology - foe or friend? *J Med Chem* **56**:8955–8971.
- Pettersson M and Crews CM (2019) Proteolysis Targeting Chimeras (PROTACs) - Past, present and future. *Drug Discov Today Technol* **31**:15–27.
- Piekarz R, Wright J, Frye R, Allen SL, Joske D, Kirschbaum M, Lewis ID, Prince M, Smith S, Jaffe ES, et al. (2009) Final Results of a Phase 2 NCI Multicenter Study of Romidepsin in Patients with Relapsed Peripheral T-Cell Lymphoma (PTCL). *Blood* **114**:1657.
- Pietschmann K, Bolck HA, Buchwald M, Spielberg S, Polzer H, Spiekermann K, Bug G, Heinzl T, Böhmer PD, and Krämer OH (2012) Breakdown of the FLT3-ITD/STAT5 axis and synergistic apoptosis induction by the histone deacetylase inhibitor panobinostat and FLT3-specific inhibitors. *Mol Cancer Ther* **11**:2373–2383.
- Plo I, Nakatake M, Malivert L, de Villartay J-P, Giraudier S, Villeval J-L, Wiesmuller L, and Vainchenker W (2008) JAK2 stimulates homologous recombination and genetic instability: potential implication in the heterogeneity of myeloproliferative disorders. *Blood* **112**:1402–1412.
- Pons M, Nagel G, Zeyn Y, Beyer M, Laguna T, Brachetti T, Sellmer A, Mahboobi S, Conradi R, Butter F, et al. (2019) Human platelet lysate as validated replacement for animal serum to assess chemosensitivity. *ALTEX* **36**:277–288.
- Porpaczy E, Tripolt S, Hoelbl-Kovacic A, Gisslinger B, Bago-Horvath Z, Casanova-Hevia E, Clappier E, Decker T, Fajmann S, Fux DA, et al. (2018) Aggressive B-cell lymphomas in patients with myelofibrosis receiving JAK1/2 inhibitor therapy. *Blood* **132**:694–706.
- Prestipino A, Emhardt AJ, Aumann K, O'Sullivan D, Gorantla SP, Duquesne S, Melchinger W, Braun L, Vuckovic S, Boerries M, et al. (2018) Oncogenic JAK2^{V617F} causes PD-L1 expression, mediating immune escape in myeloproliferative neoplasms. *Sci Transl Med* **10**:eaam7729.
- Prus G, Hoegl A, Weinert BT, and Choudhary C (2019) Analysis and Interpretation of Protein Post-Translational Modification Site Stoichiometry. *Trends Biochem Sci* **44**:943–960.
- Qureshy Z, Johnson DE, and Grandis JR (2020) Targeting the JAK/STAT pathway in solid tumors. *J Cancer Metastasis Treat* **6**:27.
- Rambaldi A, Dellacasa CM, Finazzi G, Carobbio A, Ferrari ML, Guglielmelli P, Gattoni E, Salmoiraghi S, Finazzi MC, Di Tollo S, et al. (2010) A pilot study of the Histone-Deacetylase inhibitor Givinostat in patients with JAK2V617F positive chronic myeloproliferative neoplasms. *Br J Haematol* **150**:446–455.
- Rambaldi A, Iurlo A, Vannucchi AM, Martino B, Guarini A, Ruggeri M, von Bubnoff N, De Muro M, McMullin MF, Luciani S, et al. (2021) Long-term safety and efficacy of givinostat in polycythemia vera: 4-year mean follow up of three phase 1/2 studies and a compassionate use program. *Blood Cancer J* **11**:53.
- Rao TN, Hansen N, Hilfiker J, Rai S, Majewska JM, Leković D, Gezer D, Andina N, Galli S, Cassel T, et al. (2019) JAK2-mutant hematopoietic cells display metabolic alterations that can be targeted to treat myeloproliferative neoplasms. *Blood* **134**:1832–1846.
- Renneville A, Quesnel B, Charpentier A, Terriou L, Crinquette A, Lai JL, Cossement C, Lionne-Huyghe P, Rose C, Bauters F, et al. (2006) High occurrence of JAK2 V617 mutation in refractory anemia with ringed sideroblasts associated with marked thrombocytosis. *Leukemia* **20**:2067–2070.
- Riera L, Lasorsa E, Bonello L, Sismondi F, Tondat F, Di Bello C, Di Celle PF, Chiarle R, Godio L, Pich A, et al. (2011) Description of a novel Janus kinase 3 P132A mutation in acute megakaryoblastic leukemia and demonstration of previously reported Janus kinase 3 mutations in normal subjects. *Leuk Lymphoma* **52**:1742–1750.
- Roos WP, Jöst E, Belohlavek C, Nagel G, Fritz G, and Kaina B (2011) Intrinsic anticancer drug resistance of malignant melanoma cells is abrogated by IFN- β and valproic acid. *Cancer Res* **71**:4150–4160.
- Roos WP and Krumm A (2016) The multifaceted influence of histone deacetylases on DNA damage signalling and DNA repair. *Nucleic Acids Res* **44**:10017–10030.
- Rummelt C, Gorantla SP, Meggendorfer M, Charlet A, Endres C, Döhner K, Heidel FH, Fischer T, Haferlach T, Duyster J, et al. (2021) Activating JAK-mutations confer resistance to FLT3 kinase inhibitors in FLT3-ITD positive AML in vitro and in vivo. *Leukemia* **35**:2017–2029.
- Sakaguchi H, Okuno Y, Muramatsu H, Yoshida K, Shiraiishi Y, Takahashi M, Kon A, Sanada M, Chiba K, Tanaka H, et al. (2013) Exome sequencing identifies secondary mutations of SETBP1 and JAK3 in juvenile myelomonocytic leukemia. *Nat Genet* **45**:937–941.
- Sanaei M, Kavousi F, and Pourahmadi M (2021) Effect of Decitabine (5-aza-2'-deoxycytidine, 5-aza-CdR) in Comparison with Vorinostat (Suberoylanilide Hydroxamic Acid, SAHA) on DNMT1, DNMT3a and DNMT3b, HDAC 1-3, SOCS 1, SOCS 3, JAK2, and STAT3 Gene Expression in Hepatocellular Carcinoma HLE and LCL-P1 Cell Lines. *Asian Pac J Cancer Prev* **22**:2089–2098.
- Sanda T, Tyner JW, Gutierrez A, Ngo VN, Glover J, Chang BH, Yost A, Ma W, Fleischman AG, Zhou W, et al. (2013) TYK2-STAT1-BCL2 pathway dependence in T-cell acute lymphoblastic leukemia. *Cancer Discov* **3**:564–577.
- Santos J, Mesquita D, Barros-Silva JD, Jerónimo C, Henrique R, Morais A, Paulo P, and Teixeira MR (2015) Uncovering potential downstream targets of oncogenic GRPR overexpression in prostate carcinomas harboring ETS rearrangements. *Oncoscience* **2**:497–507.
- Savino AM, Sarno J, Trentin L, Vieri M, Fazio G, Bardini M, Bugarin C, Fossati G, Davis KL, Gaipa G, et al. (2017) The histone deacetylase inhibitor givinostat (ITF2357) exhibits potent anti-tumor activity against CRLF2-rearranged BCP-ALL. *Leukemia* **31**:2365–2375.
- Sawant M, Mahajan K, Renganathan A, Weimholt C, Luo J, Kukshal V, Jeer JM, Jeon MS, Zhang B, Li T, et al. (2022) Chronologically modified androgen receptor in recurrent castration-resistant prostate cancer and its therapeutic targeting. *Sci Transl Med* **14**:eabg4132.
- Schäfer C, Göder A, Beyer M, Kiweler N, Mahendrarajah N, Rauch A, Nikolova T, Stojanovic N, Wiczorek M, Reich TR, et al. (2017) Class I histone deacetylases regulate p53/NF- κ B crosstalk in cancer cells. *Cell Signal* **29**:218–225.
- Schiedel M, Herp D, Hammelmann S, Swyter S, Lehotzky A, Robaa D, Oláh J, Ovádi J, Sippl W, and Jung M (2018) Chemically Induced Degradation of Sirtuin 2 (Sirt2) by a Proteolysis Targeting Chimera (PROTAC) Based on Sirtuin Rearranging Ligands (SirReals). *J Med Chem* **61**:482–491.
- Schindler C, Levy DE, and Decker T (2007) JAK-STAT signaling: from interferons to cytokines. *J Biol Chem* **282**:20059–20063.
- Schmitz RL, Weissbach J, Kleilein J, Bell J, Hüttelmaier S, Viol F, Clauditz T, Grabowski P, Laumen H, Rosendahl J, et al. (2021) Targeting HDACs in Pancreatic Neuroendocrine Tumor Models. *Cells* **10**:1408.
- Schneider G, Krämer OH, Fritsche P, Schüler S, Schmid RM, and Saur D (2010) Targeting histone deacetylases in pancreatic ductal adenocarcinoma. *J Cell Mol Med* **14**(6A):1255–1263.
- Scholz A, Heinze S, Detjen KM, Peters M, Welzel M, Hauff P, Schirner M, Wiedenmann B, and Rosewicz S (2003) Activated signal transducer and activator of transcription 3 (STAT3) supports the malignant phenotype of human pancreatic cancer. *Gastroenterology* **125**:891–905.
- Sellmer A, Stangl H, Beyer M, Grünstein E, Leonhardt M, Pongratz H, Eichhorn E, Elz S, Striegl B, Jenei-Lanzl Z, et al. (2018) Maribostat-100 Defines a New Class of Potent and Selective Antiinflammatory and Antirheumatic Histone Deacetylase 6 Inhibitors. *J Med Chem* **61**:3454–3477.
- Shah B, Zhao X, Silva AS, Shain KH, and Tao J (2018) Resistance to Ibrutinib in B Cell Malignancies: One Size Does Not Fit All. *Trends Cancer* **4**:197–206.
- Shah RR, Redmond JM, Mihut A, Menon M, Evans JP, Murphy JA, Bartholomew MA, and Coe DM (2020) Hi-JAK-ing the ubiquitin system: The design and physicochemical optimisation of JAK PROTACs. *Bioorg Med Chem* **28**:115326.
- Shen L, Ciesielski M, Ramakrishnan S, Miles KM, Ellis L, Sotomayor P, Shrikant P, Fenstermaker R, and Pili R (2012) Class I histone deacetylase inhibitor entinostat suppresses regulatory T cells and enhances immunotherapies in renal and prostate cancer models. *PLoS One* **7**:e30815.
- Shi S, Calhoun HC, Xia F, Li J, Le L, and Li WX (2006) JAK signaling globally counteracts heterochromatin gene silencing. *Nat Genet* **38**:1071–1076.
- Sibbesen NA, Kopp KL, Litvinov IV, Jønson L, Willerslev-Olsen A, Fredholm S, Petersen DL, Nastasi C, Krejsgaard T, Lindahl LM, et al. (2015) Jak3, STAT3, and STAT5 inhibit expression of miR-22, a novel tumor suppressor microRNA, in cutaneous T-Cell lymphoma. *Oncotarget* **6**:20555–20569.
- Smalley JP, Baker IM, Pytel WA, Lin LY, Bowman KJ, Schwabe JWR, Cowley SM, and Hodgkinson JT (2022) Optimization of Class I Histone Deacetylase PROTACs Reveals that HDAC1/2 Degradation is Critical to Induce Apoptosis and Cell Arrest in Cancer Cells. *J Med Chem* **65**:5642–5659.
- Smith J, Walsh KJ, Jacobs CL, Liu Q, Fan S, Patel A, and Dave S (2010) Upregulated JAK/STAT signaling represents a major mode of resistance to

- HDAC inhibition in lymphoma and provides a rationale for novel combination therapy. *Blood* **116**:434.
- Springuel L, Hornakova T, Losdyck E, Lambert F, Leroy E, Constantinescu SN, Flex E, Tartaglia M, Knoops L, and Renaud JC (2014) Cooperating JAK1 and JAK3 mutants increase resistance to JAK inhibitors. *Blood* **124**:3924–3931.
- Springuel L, Renaud JC, and Knoops L (2015) JAK kinase targeting in hematologic malignancies: a sinuous pathway from identification of genetic alterations towards clinical indications. *Haematologica* **100**:1240–1253.
- Staerk J, Kallin A, Demoulin J-B, Vainchenker W, and Constantinescu SN (2005) JAK1 and Tyk2 activation by the homologous polycythemia vera JAK2 V617F mutation: cross-talk with IGF1 receptor. *J Biol Chem* **280**:41893–41899.
- Food and Drug Administration(2010) StatBite: FDA oncology drug product approvals in 2009. *J Natl Cancer Inst* **102**:219.
- Steenma DP, McClure RF, Karp JE, Tefferi A, Lasho TL, Powell HL, DeWald GW, and Kaufmann SH (2006) JAK2 V617F is a rare finding in de novo acute myeloid leukemia, but STAT3 activation is common and remains unexplained. *Leukemia* **20**:971–978.
- Stein LR and Imai S (2012) The dynamic regulation of NAD metabolism in mitochondria. *Trends Endocrinol Metab* **23**:420–428.
- Stempel M, Kedinger M, Augenlicht L, and Klampfer L (2007) Essential role of the JAK/STAT1 signaling pathway in the expression of inducible nitric-oxide synthase in intestinal epithelial cells and its regulation by butyrate. *J Biol Chem* **282**:9797–9804.
- Sun Y, Zhao X, Ding N, Gao H, Wu Y, Yang Y, Zhao M, Hwang J, Song Y, Liu W, et al. (2018) PROTAC-induced BTK degradation as a novel therapy for mutated BTK C481S produced ibrutinib-resistant B-cell malignancies. *Cell Res* **28**:779–781.
- Ribrag V, Harrison CN, Heidel FH, Kiladjan J-J, Acharyya S, Mu S, Liu T, Williams D, Giles FJ, Conneally E, et al. (2013) A Phase 1b, Dose-Finding Study Of Ruxolitinib Plus Panobinostat In Patients With Primary Myelofibrosis (PMF), Post-Polycythemia Vera MF (PPV-MF), Or Post-Essential Thrombocythemia MF (PET-MF): Identification Of The Recommended Phase 2 Dose. *Blood* **122**:4045.
- Tate JG, Bamford S, Jubb HC, Sondka Z, Beare DM, Bindal N, Boutselakis H, Cole CG, Creatore C, Dawson E, et al. (2019) COSMIC: the catalogue of somatic mutations in cancer. *Nucleic Acids Res* **47** (D1):D941–D947.
- Taunton J, Hassig CA, and Schreiber SL (1996) A mammalian histone deacetylase related to the yeast transcriptional regulator Rpd3p. *Science* **272**:408–411.
- Tefferi A, Thiele J, and Vardiman JW (2009) The 2008 World Health Organization classification system for myeloproliferative neoplasms: order out of chaos. *Cancer* **115**:3842–3847.
- Tefferi A and Vardiman JW (2008) Classification and diagnosis of myeloproliferative neoplasms: the 2008 World Health Organization criteria and point-of-care diagnostic algorithms. *Leukemia* **22**:14–22.
- Telliez JB, Dowty ME, Wang L, Jussif J, Lin T, Li L, Moy E, Balbo P, Li W, Zhao Y, et al. (2016) Discovery of a JAK3-Selective Inhibitor: Functional Differentiation of JAK3-Selective Inhibition over pan-JAK or JAK1-Selective Inhibition. *ACS Chem Biol* **11**:3442–3451.
- Theocharides A, Boissinot M, Girodon F, Garand R, Teo SS, Lippert E, Talmant P, Tichelli A, Hermouet S, and Skoda RC (2007) Leukemic blasts in transformed JAK2-V617F-positive myeloproliferative disorders are frequently negative for the JAK2-V617F mutation. *Blood* **110**:375–379.
- Thiele J, Kvasnicka HM, Facchetti F, Franco V, van der Walt J, and Orazi A (2005) European consensus on grading bone marrow fibrosis and assessment of cellularity. *Haematologica* **90**:1128–1132.
- Thomas SJ, Snowden JA, Zeidler MP, and Danson SJ (2015) The role of JAK/STAT signalling in the pathogenesis, prognosis and treatment of solid tumours. *Br J Cancer* **113**:365–371.
- Tokarski JS, Zupa-Fernandez A, Trepud JA, Pike K, Chang C, Xie D, Cheng L, Pedicord D, Muckelbauer J, Johnson SR, et al. (2015) Tyrosine Kinase 2-mediated Signal Transduction in T Lymphocytes Is Blocked by Pharmacological Stabilization of Its Pseudokinase Domain. *J Biol Chem* **290**:11061–11074.
- Tomasson MH, Xiang Z, Walgren R, Zhao Y, Kasai Y, Miner T, Ries RE, Lubman O, Fremont DH, McLellan MD, et al. (2008) Somatic mutations and germline sequence variants in the expressed tyrosine kinase genes of patients with de novo acute myeloid leukemia. *Blood* **111**:4797–4808.
- Ueda F, Sumi K, Tago K, Kasahara T, and Funakoshi-Tago M (2013) Critical role of FANCC in JAK2 V617F mutant-induced resistance to DNA cross-linking drugs. *Cell Signal* **25**:2115–2124.
- Ungureanu D, Wu J, Pekkala T, Niranjani Y, Young C, Jensen ON, Xu CF, Neubert TA, Skoda RC, Hubbard SR, et al. (2011) The pseudokinase domain of JAK2 is a dual-specificity protein kinase that negatively regulates cytokine signaling. *Nat Struct Mol Biol* **18**:971–976.
- Vainchenker W and Constantinescu SN (2013) JAK/STAT signaling in hematological malignancies. *Oncogene* **32**:2601–2613.
- Vannucchi AM, Guglielmelli P, and Tefferi A (2009) Advances in understanding and management of myeloproliferative neoplasms. *CA Cancer J Clin* **59**:171–191.
- Vardiman JW, Thiele J, Arber DA, Brunning RD, Borowitz MJ, Porwit A, Harris NL, Le Beau MM, Hellström-Lindberg E, Tefferi A, et al. (2009) The 2008 revision of the World Health Organization (WHO) classification of myeloid neoplasms and acute leukemia: rationale and important changes. *Blood* **114**:937–951.
- Verma A, Kambhampati S, Parmar S, and Platanius LC (2003) Jak family of kinases in cancer. *Cancer Metastasis Rev* **22**:423–434.
- Verstovsek S, Silver RT, Cross NC, and Tefferi A (2006) JAK2V617F mutational frequency in polycythemia vera: 100%, >90%, less? *Leukemia* **20**:2067.
- Verstovsek S, Tam CS, Wadleigh M, Sokol L, Smith CC, Bui LA, Song C, Clary DO, Olszynski P, Cortes J, et al. (2014) Phase I evaluation of XL019, an oral, potent, and selective JAK2 inhibitor. *Leuk Res* **38**:316–322.
- Vicente C, Schwab C, Broux M, Geerdens E, Degryse S, Demeyer S, Lahortiga I, Elliott A, Chilton L, La Starza R, et al. (2015) Targeted sequencing identifies associations between IL7R-JAK mutations and epigenetic modulators in T-cell acute lymphoblastic leukemia. *Haematologica* **100**:1301–1310.
- Vogelmann A, Robaa D, Sippl W, and Jung M (2020) Proteolysis targeting chimeras (PROTACs) for epigenetics research. *Curr Opin Chem Biol* **57**:8–16.
- Waanders E, Scheijen B, Jongmans MC, Venselaar H, van Reijmersdal SV, van Dijk AH, Pastorczak A, Weren RD, van der Schoot CE, van de Vorst M, et al. (2017) Germline activating TYK2 mutations in pediatric patients with two primary acute lymphoblastic leukemia occurrences. *Leukemia* **31**:821–828.
- Wachholz V, Mustafa AM, Zeyn Y, Henninger SJ, Beyer M, Dzulko M, Pié-Staffa A, Brchetti C, Haehnel PS, Sellmer A, et al. (2022) Inhibitors of class I HDACs and of FLT3 combine synergistically against leukemia cells with mutant FLT3. *Arch Toxicol* **96**:177–193.
- Wagner T, Brand P, Heinzel T, and Krämer OH (2014) Histone deacetylase 2 controls p53 and is a critical factor in tumorigenesis. *Biochim Biophys Acta* **1846**:524–538.
- Wahnschaffe L, Braun T, Timonen S, Giri AK, Schrader A, Wagle P, Almusa H, Johansson P, Bellanger D, López C, et al. (2019) JAK/STAT-Activating Genomic Alterations Are a Hallmark of T-PLL. *Cancers (Basel)* **11**:1833.
- Walters DK, Mercher T, Gu TL, O'Hare T, Tyner JW, Loriaux M, Goss VL, Lee KA, Eide CA, Wong MJ, et al. (2006) Activating alleles of JAK3 in acute megakaryoblastic leukemia. *Cancer Cell* **10**:65–75.
- Wang L, Wu Z, Xia Y, Lu X, Li J, Fan L, Qiao C, Qiu H, Gu D, Xu W, et al. (2022) Single-cell profiling-guided combination therapy of c-Fos and histone deacetylase inhibitors in diffuse large B-cell lymphoma. *Clin Transl Med* **12**:e798.
- Wang Y, Fiskus W, Chong DG, Buckley KM, Natarajan K, Rao R, Joshi A, Balusu R, Koul S, Chen J, et al. (2009) Cotreatment with panobinostat and JAK2 inhibitor TG101209 attenuates JAK2V617F levels and signaling and exerts synergistic cytotoxic effects against human myeloproliferative neoplastic cells. *Blood* **114**:5024–5033.
- Weigert O, Lane AA, Bird L, Kopp N, Chapuy B, van Bodegom D, Toms AV, Marubayashi S, Christie AL, McKeown M, et al. (2012) Genetic resistance to JAK2 enzymatic inhibitors is overcome by HSP90 inhibition. *J Exp Med* **209**:259–273.
- Whittaker SJ, Demierre M-F, Kim EJ, Rook AH, Lerner A, Duvic M, Scarisbrick J, Reddy S, Robak T, Becker JC, et al. (2010) Final results from a multicenter, international, pivotal study of romidepsin in refractory cutaneous T-cell lymphoma. *J Clin Oncol* **28**:4485–4491.
- Wieczorek M, Ginter T, Brand P, Heinzel T, and Krämer OH (2012) Acetylation modulates the STAT signaling code. *Cytokine Growth Factor Rev* **23**:293–305.
- Wingelhofer B, Neubauer HA, Valent P, Han X, Constantinescu SN, Gunning PT, Müller M, and Moriggl R (2018) Implications of STAT3 and STAT5 signaling on gene regulation and chromatin remodeling in hematopoietic cancer. *Leukemia* **32**:1713–1726.
- Wolanskyj AP, Schwager SM, McClure RF, Larson DR, and Tefferi A (2006) Essential thrombocythemia beyond the first decade: life expectancy, long-term complication rates, and prognostic factors. *Mayo Clin Proc* **81**:159–166, Elsevier.
- Wöss K, Simonović N, Strobl B, Macho-Maschler S, and Müller M (2019) TYK2: An Upstream Kinase of STATs in Cancer. *Cancers (Basel)* **11**:1728.
- Wu H, Yang K, Zhang Z, Leisten ED, Li Z, Xie H, Liu J, Smith KA, Novakova Z, Barinka C, et al. (2019) Development of multifunctional histone deacetylase 6 degraders with potent antimyeloma activity. *J Med Chem* **62**:7042–7057.
- Wu SC, Li LS, Kopp N, Montero J, Chapuy B, Yoda A, Christie AL, Liu H, Christodoulou A, van Bodegom D, et al. (2015) Activity of the Type II JAK2 Inhibitor CHZ868 in B Cell Acute Lymphoblastic Leukemia. *Cancer Cell* **28**:29–41.
- Xiang Z, Zhao Y, Mitaksov V, Fremont DH, Kasai Y, Molitoris A, Ries RE, Miner TL, McLellan MD, DiPersio JF, et al. (2008) Identification of somatic JAK1 mutations in patients with acute myeloid leukemia. *Blood* **111**:4809–4812.
- Xiao Y, Wang J, Zhao LY, Chen X, Zheng G, Zhang X, and Liao D (2020) Discovery of histone deacetylase 3 (HDAC3)-specific PROTACs. *Chem Commun (Camb)* **56**:9866–9869.
- Xiong H, Du W, Zhang YJ, Hong J, Su WY, Tang JT, Wang YC, Lu R, and Fang JY (2012) Trichostatin A, a histone deacetylase inhibitor, suppresses JAK2/STAT3 signaling via inducing the promoter-associated histone acetylation of SOCS1 and SOCS3 in human colorectal cancer cells. *Mol Carcinog* **51**:174–184.
- Yang EG, Mustafa N, Tan EC, Poulsen A, Ramanujulu PM, Chng WJ, Yen JJ, and Dymock BW (2016) Design and synthesis of janus kinase 2 (JAK2) and histone deacetylase (HDAC) bispecific inhibitors based on pacritinib and evidence of dual pathway inhibition in hematological cell lines. *J Med Chem* **59**:8233–8262.
- Yang H, Liu X, Zhu X, Li X, Jiang L, Zhong M, Zhang M, Chen T, Ma M, Liang X, et al. (2021) CPVL promotes glioma progression via STAT1 pathway inhibition through interactions with the BTK/p300 axis. *JCI Insight* **6**:e146362.
- Yang L, Shi P, Zhao G, Xu J, Peng W, Zhang J, Zhang G, Wang X, Dong Z, Chen F, et al. (2020) Targeting cancer stem cell pathways for cancer therapy. *Signal Transduct Target Ther* **5**:8.
- Yao L, Mustafa N, Tan EC, Poulsen A, Singh P, Duong-Thi MD, Lee JXT, Ramanujulu PM, Chng WJ, Yen JJY, et al. (2017) Design and Synthesis of Ligand Efficient Dual Inhibitors of Janus Kinase (JAK) and Histone Deacetylase (HDAC) Based on Ruxolitinib and Vorinostat. *J Med Chem* **60**:8336–8357.
- Yao L, Ramanujulu PM, Poulsen A, Ohlson S, and Dymock BW (2018) Merging of ruxolitinib and vorinostat leads to highly potent inhibitors of JAK2 and histone deacetylase 6 (HDAC6). *Bioorg Med Chem Lett* **28**:2636–2640.
- Yonal I, Dağlar-Aday A, Akadam-Teker B, Yılmaz C, Nalçacı M, Yavuz AS, and Sargin FD (2016) Impact of JAK2V617F Mutational Status on Phenotypic Features in Essential Thrombocythemia and Primary Myelofibrosis. *Turk J Haematol* **33**:94–101.
- Yoshida M, Kijima M, Akita M, and Beppu T (1990) Potent and specific inhibition of mammalian histone deacetylase both in vivo and in vitro by trichostatin A. *J Biol Chem* **265**:17174–17179.
- Zeng H, Qu J, Jin N, Xu J, Lin C, Chen Y, Yang X, He X, Tang S, Lan X, et al. (2016) Feedback Activation of Leukemia Inhibitory Factor Receptor Limits

- Response to Histone Deacetylase Inhibitors in Breast Cancer. *Cancer Cell* **30**:459–473.
- Zhang J, Ding L, Holmfeldt L, Wu G, Heatley SL, Payne-Turner D, Easton J, Chen X, Wang J, Rusch M, et al. (2012) The genetic basis of early T-cell precursor acute lymphoblastic leukaemia. *Nature* **481**:157–163.
- Zhang X, Thummuri D, Liu X, Hu W, Zhang P, Khan S, Yuan Y, Zhou D, and Zheng G (2020) Discovery of PROTAC BCL-X_L degraders as potent anticancer agents with low on-target platelet toxicity. *Eur J Med Chem* **192**:112186.
- Zhang Y, Andrade R, Hanna AA, and Pflum MKH (2022) Evidence that HDAC7 acts as an epigenetic “reader” of AR acetylation through NCoR-HDAC3 dissociation. *Cell Chem Biol* **29**:1162–1173.e5.
- Zhao B, Huang Z, Qin Z, Li Y, Wang T, Wang L, Zhou W, Yu C, Wang X, Yang S, et al. (2019) Enhancement of Histone Deacetylase Inhibitor Sensitivity in Combination with Cyclin-Dependent Kinase Inhibition for the Treatment of Oral Squamous Cell Carcinoma. *Cell Physiol Biochem* **53**:141–156.
- Zhao C, Khadka DB, and Cho WJ (2016a) Insights into the Structural Features Essential for JAK2 Inhibition and Selectivity. *Curr Med Chem* **23**:1331–1355.
- Zhao S, Guo J, Zhao Y, Fei C, Zheng Q, Li X, and Chang C (2016b) Chidamide, a novel histone deacetylase inhibitor, inhibits the viability of MDS and AML cells by suppressing JAK2/STAT3 signaling. *Am J Transl Res* **8**:3169–3178.
- Zhou L, Zhang Y, Chen S, Kmiecik M, Leng Y, Lin H, Rizzo KA, Dumur CI, Ferreira-Gonzalez A, Dai Y, et al. (2015) A regimen combining the Wee1 inhibitor AZD1775 with HDAC inhibitors targets human acute myeloid leukemia cells harboring various genetic mutations. *Leukemia* **29**:807–818.
- Zimmermann GR, Lehár J, and Keith CT (2007) Multi-target therapeutics: when the whole is greater than the sum of the parts. *Drug Discov Today* **12**:34–42.
- Zorzi AP, Bernstein M, Samson Y, Wall DA, Desai S, Nicksy D, Wainman N, Eisenhauer E, and Baruchel S (2013) A phase I study of histone deacetylase inhibitor, pracinostat (SB939), in pediatric patients with refractory solid tumors: IND203 a trial of the NCIC IND program/C17 pediatric phase I consortium. *Pediatr Blood Cancer* **60**:1868–1874.