

# International Union of Pharmacology. XL. Compendium of Voltage-Gated Ion Channels: Calcium Channels

WILLIAM A. CATTERALL, JOERG STRIESSNIG, TERRANCE P. SNUTCH, AND EDWARD PEREZ-REYES

*Department of Pharmacology, University of Washington School of Medicine, Seattle, Washington (W.A.C.); Abteilung Pharmakologie und Toxikologie, Institut für Pharmazie, Universität Innsbruck, Innsbruck, Austria (J.S.); Biotechnology Laboratory, University of British Columbia, Vancouver, British Columbia, Canada (T.P.S.); and Department of Pharmacology, University of Virginia, Charlottesville, Virginia (E.P.-R.)*

**Abstract**—This summary article presents an overview of the molecular relationships among the voltage-gated calcium channels and a standard nomenclature for them, which is derived from the *IUPHAR*

*Compendium of Voltage-Gated Ion Channels*.<sup>1</sup> The complete Compendium, including data tables for each member of the calcium channel family can be found at <<http://www.iuphar-db.org/iuphar-ic/>>.

## Calcium Channel Subunits

Voltage-gated calcium channels mediate calcium influx in response to membrane depolarization and regulate intracellular processes such as contraction, secretion, neurotransmission, and gene expression. Their activity is essential to couple electrical signals in the cell surface to physiological events in cells. They are members of a gene superfamily of transmembrane ion channel proteins that includes voltage-gated potassium and sodium channels (Catterall, 1995).

The calcium channels that have been characterized biochemically are complex proteins composed of four or five distinct subunits, which are encoded by multiple genes (Fig. 1) (Catterall, 2000). The  $\alpha_1$  subunit of 190 to 250 kDa is the largest subunit, and it incorporates the conduction pore, the voltage sensor and gating apparatus, and the known sites of channel regulation by second messengers, drugs, and toxins. Like the  $\alpha$  subunits of sodium channels, the  $\alpha_1$  subunit of voltage-gated calcium channels is organized in four homologous domains (I–IV) with six transmembrane segments (S1–S6) in each. The S4 segment serves as the voltage sensor. The pore loop between transmembrane segments S5 and S6 in each domain determines ion conductance and selectivity, and changes of only three amino acids (aa) in the pore loops in domains I, III,

and IV will convert a sodium channel to calcium selectivity. An intracellular  $\beta$  subunit and a transmembrane, disulfide-linked  $\alpha_2\delta$  subunit complex are components of most types of calcium channels. A  $\gamma$  subunit has also been found in skeletal muscle calcium channels and related subunits are expressed in heart and brain. Although these auxiliary subunits modulate the properties of the channel complex, the pharmacological and electrophysiological diversity of calcium channels arises primarily from the existence of multiple  $\alpha_1$  subunits (Hofmann et al., 1994).

## Calcium Currents

Calcium currents recorded in different cell types have diverse physiological and pharmacological properties, and an alphabetical nomenclature has evolved for the distinct classes of calcium currents (Tsein et al., 1995). L-type calcium currents require a strong depolarization for activation, are long-lasting, and are blocked by the organic L-type calcium channel antagonists, including dihydropyridines, phenylalkylamines, and benzothiazepines. They are the main calcium currents recorded in muscle and endocrine cells, where they initiate contraction and secretion. N-type, P/Q-type, and R-type calcium currents also require strong depolarization for activation. They are relatively unaffected by L-type calcium channel antagonist drugs but are blocked by specific polypeptide toxins from snail and spider venoms. They are expressed primarily in neurons, where they initiate neurotransmission at most fast synapses and also mediate calcium entry into cell bodies and dendrites. T-type calcium currents are activated by weak depolarization and are transient. They are resistant to both organic antagonists and to the snake and spider toxins used to define the N- and P/Q-type calcium currents. They are expressed in a wide variety of cell types,

Address correspondence to: Dr. William A. Catterall, Department of Pharmacology, University of Washington School of Medicine, Box 357280, Seattle, WA 98195-7280. E-mail: [wcatt@u.washington.edu](mailto:wcatt@u.washington.edu)

<sup>1</sup> This work was previously published in Catterall WA, Chandy KG, and Gutman GA, eds. (2002) *The IUPHAR Compendium of Voltage-Gated Ion Channels*, International Union of Pharmacology Media, Leeds, UK.

Article, publication date, and citation information can be found at <http://pharmrev.aspetjournals.org>.

DOI: 10.1124/pr.55.4.8.




FIG. 1. Subunits of  $\text{Ca}_v1$  channels. The subunit composition and structure of calcium channels purified from skeletal muscle are illustrated. The model is updated from the original description of the subunit structure of skeletal muscle calcium channels. This model also fits available biochemical and molecular biological results for other  $\text{Ca}_v1$  channels and for  $\text{Ca}_v2$  channels. P, sites of phosphorylation by cAMP-dependent protein kinase;  $\Psi$ , sites of N-linked glycosylation. Predicted alpha helices are depicted as cylinders. The lengths of lines correspond approximately to the lengths of the polypeptide segments represented.

where they are involved in shaping the action potential and controlling patterns of repetitive firing.

### Calcium Channel Genes

Mammalian  $\alpha_1$  subunits are encoded by at least ten distinct genes. Historically, various names had been given to the corresponding gene products, giving rise to distinct and sometimes confusing nomenclatures. In 1994, a unified but arbitrary nomenclature was adopted in which  $\alpha_1$  subunits were referred to as  $\alpha_{1S}$  for the original skeletal muscle isoform and  $\alpha_{1A}$  through  $\alpha_{1E}$  for those discovered subsequently (Birnbaumer et al., 1994). In 2000, a rational nomenclature was adopted based on the well defined potassium channel nomenclature (Chandy and Gutman, 1993; Ertel et al., 2000). Calcium channels were named using the chemical sym-

bol of the principal permeating ion (Ca) with the principal physiological regulator (voltage) indicated as a subscript ( $\text{Ca}_v$ ). The numerical identifier corresponds to the  $\text{Ca}_v$  channel  $\alpha_1$  subunit gene subfamily (1 to 3 at present) and the order of discovery of the  $\alpha_1$  subunit within that subfamily (1 through m). According to this nomenclature, the  $\text{Ca}_v1$  subfamily ( $\text{Ca}_v1.1$  to  $\text{Ca}_v1.4$ ) includes channels containing  $\alpha_{1S}$ ,  $\alpha_{1C}$ ,  $\alpha_{1D}$ , and  $\alpha_{1F}$ , which mediate L-type  $\text{Ca}^{2+}$  currents (Table 1). The  $\text{Ca}_v2$  subfamily ( $\text{Ca}_v2.1$  to  $\text{Ca}_v2.3$ ) includes channels containing  $\alpha_{1A}$ ,  $\alpha_{1B}$ , and  $\alpha_{1E}$ , which mediate P/Q-, N-, and R-type  $\text{Ca}^{2+}$  currents, respectively (Table 1). The  $\text{Ca}_v3$  subfamily ( $\text{Ca}_v3.1$  to  $\text{Ca}_v3.3$ ) includes channels containing  $\alpha_{1G}$ ,  $\alpha_{1H}$ , and  $\alpha_{1I}$ , which mediate T-type  $\text{Ca}^{2+}$  currents.

The complete amino acid sequences of these  $\alpha_1$  subunits are more than 70% identical within a family but less than

TABLE 1  
*Physiological function and pharmacology of calcium channels*

Channel	Current	Localization	Specific Antagonists	Cellular Functions
$\text{Ca}_v1.1$	L	Skeletal muscle transverse tubules	Dihydropyridines, phenylalkylamines, benzothiazepines	Excitation-contraction coupling
$\text{Ca}_v1.2$	L	Cardiac myocytes, endocrine cells, neuronal cell bodies and proximal dendrites	Dihydropyridines, phenylalkylamines, benzothiazepines	Excitation-contraction coupling, hormone release, regulation of transcription, synaptic integration
$\text{Ca}_v1.3$	L	Endocrine cells, neuronal cell bodies and dendrites	Dihydropyridines, phenylalkylamines, benzothiazepines	Hormone release, regulation of transcription, synaptic integration
$\text{Ca}_v1.4$	L	Retina	Not established	Neurotransmitter release from rods and bipolar cells
$\text{Ca}_v2.1$	P/Q	Nerve terminals and dendrites	$\omega$ -agatoxin IVA	Neurotransmitter release, dendritic $\text{Ca}^{2+}$ transients
$\text{Ca}_v2.2$	N	Nerve terminals and dendrites	$\omega$ -CTx-GVIA	Neurotransmitter release, dendritic $\text{Ca}^{2+}$ transients
$\text{Ca}_v2.3$	R	Neuronal cell bodies and dendrites	SNX-482	Repetitive firing
$\text{Ca}_v3.1$	T	Neuronal cell bodies and dendrites, cardiac myocytes	None	Pacemaking, repetitive firing
$\text{Ca}_v3.2$	T	Neuronal cell bodies and dendrites, cardiac myocytes	None	Pacemaking, repetitive firing
$\text{Ca}_v3.3$	T	Neuronal cell bodies and dendrites	None	Pacemaking, repetitive firing




FIG. 2. Sequence similarity of voltage-gated calcium channel  $\alpha_1$  subunits. Phylogenetic representation of the primary sequences of the calcium channels. Only the membrane-spanning segments and the pore loops (~350 amino acids) are compared. First, all sequence pairs were compared, which clearly defines three families with intra-family sequence identities above 80% ( $\text{Ca}_V1$ ,  $\text{Ca}_V2$ ,  $\text{Ca}_V3$ ). Then, a consensus sequence was defined for each family and these three sequences were compared with one another, with inter-family sequence identities of ~52% ( $\text{Ca}_V1$  versus  $\text{Ca}_V2$ ) and 28% ( $\text{Ca}_V3$  versus  $\text{Ca}_V1$  or  $\text{Ca}_V2$ ).

40% identical among families. These family relationships are illustrated for the more conserved transmembrane and pore domains in Fig. 2. Division of calcium channels into these three families is phylogenetically ancient, as representatives of each are found in the *Caenorhabditis elegans* genome. Consequently, the genes for the different  $\alpha_1$  subunits have become widely dispersed in the genome and even the most closely related members of the family are not clustered on single chromosomes.

### Calcium Channel Molecular Pharmacology

The pharmacology of the three families of calcium channel is quite distinct. The  $\text{Ca}_V1$  channels are the molecular targets of the organic calcium channel blockers used widely in the therapy of cardiovascular diseases. These drugs are thought to act at three separate, but allosterically coupled, receptor sites (Table 1). Phenylalkylamines are intracellular pore blockers, which are thought to enter the pore from the cytoplasmic side of the channel and block it. Their receptor site is formed by amino acid residues in the S6 segments in domains III and IV, in close analogy to the local anesthetic receptor site on sodium channels (Hockerman et al., 1997; Hofmann et al., 1999; Striessnig, 1999). Dihydropyridines can be channel activators or inhibitors and therefore are thought to act allosterically to shift the channel toward the open or closed state, rather than by occluding the pore. Their receptor site includes amino acid residues in the S6 segments of domains III and IV and the S5 segment of domain III. The dihydropyridine receptor site is closely apposed to the phenylalkylamine receptor site and shares some common amino acid residues. Diltiazem and related benzothiazepines are thought to bind to a third receptor site, but the amino acid residues that

are required for their binding overlap extensively with those required for phenylalkylamine binding.

The  $\text{Ca}_V2$  family of calcium channels is relatively insensitive to dihydropyridine calcium channel blockers, but these calcium channels are specifically blocked with high affinity by peptide toxins from spiders and marine snails (Miljanich and Ramachandran, 1995). The  $\text{Ca}_V2.1$  channels are blocked specifically by  $\omega$ -agatoxin IVA from funnel web spider venom. The  $\text{Ca}_V2.2$  channels are blocked specifically by  $\omega$ -conotoxin GVIA and related cone snail toxins. The  $\text{Ca}_V2.3$  channels are blocked specifically by the synthetic peptide toxin SNX-482 derived from tarantula venom. These peptide toxins are potent blockers of synaptic transmission because of their specific effects on the  $\text{Ca}_V2$  family of calcium channels.

The  $\text{Ca}_V3$  family of calcium channels is insensitive to both the dihydropyridines that block  $\text{Ca}_V1$  channels and the spider and cone snail toxins that block the  $\text{Ca}_V2$  channels, and there are no widely useful pharmacological agents that block T-type calcium currents (Heady et al., 2001). The organic calcium channel blockers mibefradil is somewhat specific for T-type versus L-type calcium currents (3- to 5-fold). The peptide kurtoxin inhibits the activation gating of  $\text{Ca}_V3.1$  and  $\text{Ca}_V3.2$  channels. Development of more specific and high affinity blockers of the  $\text{Ca}_V3$  family of calcium channels would be useful for therapy and for more detailed analysis of the physiological roles of these channels.

This section of the compendium summarizes the major molecular, physiological, and pharmacological properties for each of the ten calcium channels that have been functionally expressed. Quantitative data are included for voltage dependence of activation and inactivation, single-channel conductance, and binding of drugs and neurotoxins, focusing on those agents that are widely used and are diagnostic of channel identity and function.

### References

- Birnbaumer L, Campbell KP, Catterall WA, Harpold MM, Hofmann F, Horne WA, Mori Y, Schwartz A, Snutch TP, Tanabe T, et al. (1994) The naming of voltage-gated calcium channels. *Neuron* **13**:505–506.
- Catterall WA (1995) Structure and function of voltage-gated ion channels. *Annu Rev Biochem* **65**:493–531.
- Catterall WA (2000) Structure and regulation of voltage-gated  $\text{Ca}^{2+}$  channels. *Annu Rev Cell Dev Biol* **16**:521–555.
- Chandy KG and Gutman GA (1993) Nomenclature for mammalian potassium channel genes. *Trends Pharmacol Sci* **14**:434.
- Ertel EA, Campbell KP, Harpold MM, Hofmann F, Mori Y, Perez-Reyes E, Schwartz A, Snutch TP, Tanabe T, Birnbaumer L, et al. (2000) Nomenclature of voltage-gated calcium channels. *Neuron* **25**:533–535.
- Glossmann H and Striessnig J (1990) Molecular properties of calcium channels. *Rev Physiol Biochem Pharmacol* **114**:1–105.
- Heady TN, Gomora JC, Macdonald TL, and Perez-Reyes E (2001) Molecular pharmacology of T-type  $\text{Ca}^{2+}$  channels. *Jpn J Pharmacol* **85**:339–350.
- Hockerman GH, Peterson BZ, Johnson BD, and Catterall WA (1997) Molecular determinants of drug binding and action on L-type calcium channels. *Annu Rev Pharmacol Toxicol* **37**:361–396.
- Hofmann F, Biel M, and Flockerzi V (1994) Molecular basis for  $\text{Ca}^{2+}$  channel diversity. *Annu Rev Neurosci* **17**:399–418.
- Hofmann F, Lacinová L, and Klugbauer N (1999) Voltage-dependent calcium channels: from structure to function. *Rev Physiol Biochem Pharmacol* **139**:33–87.
- Miljanich GP and Ramachandran J (1995) Antagonists of neuronal calcium channels: structure, function, and therapeutic implications. *Annu Rev Pharmacol Toxicol* **35**:707–734.
- Striessnig J (1999) Pharmacology, structure and function of cardiac L-type calcium channels. *Cell Physiol Biochem* **9**:242–269.
- Tsien RW, Lipscombe D, Madison D, Bley K, and Fox A (1995) Reflections on calcium channel diversity, 1988–1994. *Trends Neurosci* **18**:52–54.