

# International Union of Pharmacology. XLIII. Compendium of Voltage-Gated Ion Channels: Transient Receptor Potential Channels

DAVID E. CLAPHAM, CRAIG MONTELL, GUENTER SCHULTZ, AND DAVID JULIUS

*Howard Hughes Medical Institute, Children's Hospital, Boston, Massachusetts (D.E.C.); Department of Biological Chemistry, The Johns Hopkins University School of Medicine, Baltimore, Maryland C.M.); Pharmakologisches Institut, Freie Universitaet Berlin, Berlin, Germany (G.S.); and Department of Cellular and Molecular Pharmacology, University of California, San Francisco, California (D.J.)*

**Abstract**—The transient receptor potential (TRP) proteins are six transmembrane-containing subunits that combine to form cation-selective ion channels. TRP channels are present in yeast, *Drosophila*, *Caenorhabditis elegans*, and mammals. They are widely distributed and sense local changes in stimuli ranging from light to temperature and osmolarity. Mammals contain at least 22 distinct genes encoding these ion

channels. This summary article presents an overview of the molecular relationships among the TRP channels and a standard nomenclature for them, which is derived from the *IUPHAR Compendium of Voltage-Gated Ion Channels*.<sup>1</sup> The complete Compendium, including data tables for each member of the TRP channel family, can be found at <<http://www.iuphar-db.org/iuphar-ic/>>.

## Introduction

The mammalian TRP<sup>2</sup> ion channels are encoded by at least 22 channel subunit genes, rising to >30 if polycystic kidney (PKD, TRPP) and mucolipins (TRPML) are included. TRP channel primary structures predict six transmembrane (6TM)-spanning domains with a pore domain between the fifth (S5) and sixth (S6) segments and both C and N termini located intracellularly. This architecture is a common theme for hundreds of ion channels present in life forms ranging from bacteria to mammals. The mammalian TRP channel family is united primarily by structural homology within the transmembrane-spanning domains (Fig. 1), but overall

sequence identities between members can be as low as 20%. Other features include a 25-amino acid motif (TRP domain) containing TRP box (EWKFAR) just C-terminal to the sixth transmembrane segment. The TRP domain and box are present in all TRPC channel genes but not in all TRP channel genes. The N-terminal cytoplasmic domain of TRPC, TRPV, and ANKTM (TRPA) channels contain ankyrin repeats, whereas the TRPC and TRPM contain proline-rich regions in the region just C-terminal to the predicted 6TM segment (Fig. 2). At present, no one feature other than overall 6TM architecture and homology define the TRP family. Thus, we expect that the definition of TRP channels will evolve as functions and structures are clarified.

Address correspondence to: Dr. David E. Clapham, Howard Hughes Medical Institute, Children's Hospital, 1309 Enders Building, 320 Longwood Ave., Boston, MA 02115. E-mail: [clapham@enders.tch.harvard.edu](mailto:clapham@enders.tch.harvard.edu)

<sup>1</sup> This work was previously published in Catterall WA, Chandy KG, and Gutman GA, eds. (2002) *The IUPHAR Compendium of Voltage-Gated Ion Channels*, International Union of Pharmacology Media, Leeds, UK.

<sup>2</sup> Abbreviations: TRP, transient receptor potential; Ca<sup>2+</sup>, calcium; CaT, calcium transporter; ER, endoplasmic reticulum; ECaC, epithelial calcium channel transporter; IP<sub>3</sub>, inositol 1,4,5-trisphosphate; IP<sub>3</sub>R, IP<sub>3</sub> receptor; KcsA, prokaryotic potassium-selective channel; P<sub>Ca</sub>/P<sub>Na</sub>, ratio of calcium permeability to sodium permeability; PKD, polycystic kidney disease; PIP<sub>2</sub>, phosphatidylinositol 4,5-bisphosphate; PLC, phospholipase C; S, segment, TM, transmembrane; TRPA, ankyrin-repeat TRP; TRPC, canonical TRP; TRPM, melastatin TRP; TRPV, vanilloid TRP.

Article, publication date, and citation information can be found at <http://pharmrev.aspetjournals.org>.

DOI: 10.1124/pr.55.4.6.

Transient receptor potential (*trp*) ion channel subunit genes were first defined in the *Drosophila* visual system. In the *trp* mutation, the light response (receptor potential) decays during prolonged exposure to bright light. TRP-deficient flies are blinded by intense light because sustained Ca<sup>2+</sup> entry via TRP ion channels and subsequent Ca<sup>2+</sup>-dependent adaptation is disrupted. Three genes (TRP, TRPL, TRP $\gamma$ ) in *Drosophila* encode TRP channels mediating fly vision and other unknown functions. Genetic approaches in flies have not resolved the mechanism of TRP activation, but they confirm the importance of phospholipase C $\beta$  (PLC $\beta$ ) and other components of the phosphatidylinositol pathway (Harteneck et al., 2000; Clapham et al., 2001; Montell et al., 2002; Hardie, 2003).




FIG. 1. Phylogenetic relationship in the TRP protein family. The evolutionary tree is calculated by the neighbor-joining method (Saitou and Nei, 1987). The TRPP and TRPML subfamilies are not detailed in this review.

### Structural Features

The 2TM structure of a bacterial  $K^+$  channel (KcsA) is analogous to the S5 and S6 domains joined by a short pore  $\alpha$  helix of the 6TM architecture (Doyle et al., 1998). KcsA channel is a tetramer of 2TM-spanning  $\alpha$  helices. The helices corresponding to S5 face the lipid membrane whereas the helices corresponding to S6 line the pore. At both inner and outer membrane faces, layers of aromatic amino acids form a cuff around the pore. In KcsA the selectivity filter is a narrow region near the outer face of the membrane lined by the carbonyl backbone of five conserved amino acids. These amino acids are not present as a group in the largely nonselective TRP channels. In KcsA, rings of carbonyl oxygens act as surrogate waters to coordinate the dehydrated  $K^+$  ions in the channel. The rest of the S5- and S6-spanning regions are likely to be analogous to KcsA in which the narrow channel in the selectivity filter rapidly broadens in hour-glass fashion. The four short pore  $\alpha$  helices focus their helix dipole negative electrostatic fields on the cavity to shield the cation from the hydrophobic lipid environment. The S6 base lines the rest of the channel on its way to the cytoplasm. The S6 segment and the C-termi-

nal amino acids extending into the cytoplasm are where the interesting gating features of TRP channels are likely to emerge. The most conserved regions between the three TRP subfamilies are in the S6 domain.

The detailed structure of the S1–S4 segments of the 6TM channels is not available, but mutagenesis data provide some clues about their functions. The S4 segment in voltage-sensitive channels contains at least four charged arginines or lysines that convey voltage changes across the membrane into movement of the helix, somehow gating the pore by moving this S4 helix. The TRP channels are very weakly voltage-dependent and lack the full complement of charged amino acids in the S4 domain.

### TRP Channel Functional Features

All TRP channels are nonselective with  $P_{Ca}/P_{Na} \leq 10$ , with the exception of the monovalent-selective TRPM4, M5, and the  $Ca^{2+}$ -selective ( $P_{Ca}/P_{Na} > 100$ ) TRPV5, V6.

TRP channels do not have the sharp voltage sensitivity of the 24 membrane-spanning  $Ca_v$  or  $Na_v$  families (Fig. 3). Thus, upon opening, they depolarize cells from their resting membrane potentials (roughly  $-70$  mV in most mammalian cells) to around 0 mV. In short, they depolarize cells and raise intracellular  $Ca^{2+}$  and/or  $Na^+$ .

Two common signal transduction pathways regulating the release of intracellular  $Ca^{2+}$  are the G protein-coupled and the tyrosine kinase activation of PLC. PLC hydrolyzes phosphatidylinositol 4,5-bisphosphate ( $PIP_2$ ) to form inositol 1,4,5-trisphosphate ( $IP_3$ ) that opens the  $IP_3$  receptor ( $IP_3R$ ), and liberates  $Ca^{2+}$  from the endoplasmic reticulum (Clapham, 1995). Accompanying these chains of events, and not necessarily linked to  $Ca^{2+}$  store (ER) depletion, is activation of the TRP channels. The details of these mechanisms are incompletely understood at present. The strongest associations between the phosphatidylinositol pathway and TRPs involve  $PLC\beta$  and  $PIP_2$ . Based on *Drosophila* TRPs, elements of these signal transduction pathways are linked by scaffolding proteins (Montell, 1999).

Putney (1977) proposed that emptied  $Ca^{2+}$  stores (ER) somehow gate  $Ca^{2+}$  entry of external  $Ca^{2+}$  to replenish the deficit. The physiological hallmark of the store-operated  $Ca^{2+}$  entry process is a large receptor-mediated transient  $[Ca^{2+}]_i$  increase followed by a prolonged high  $[Ca^{2+}]_i$  plateau phase, dependent on  $[Ca^{2+}]_o$ . A very specific and highly  $Ca^{2+}$ -selective current ( $Ca^{2+}$  release activated current;  $I_{CRAC}$ ) is activated by a variety of store depletion protocols in whole-cell recordings from single blood cells (Hoth and Penner, 1992; Lewis, 1999), but store-operated entry may not be solely through  $I_{CRAC}$  channels. From the start, TRPs have been the major suspects for the store-operated channel(s), including  $I_{CRAC}$ . At odds with this supposition is the high  $Ca^{2+}$  selectivity of  $I_{CRAC}$  compared with the cationic nonselectivity of most of the TRP family.




FIG. 2. Architecture of TRP channels is that of the broader class of six transmembrane-spanning ion channels. S1–S6 are transmembrane domains. A represents ankyrin repeats.




FIG. 3. Current-voltage relationships of various TRP channels; a, TRPC1 + TRPC5 heteromer; b, TRPM5; c, TRPV5, TRPV6.

Here we focus on the mammalian genes and use a nomenclature adopted by a number of workers in the field (Montell et al., 2002). TRPC<sub>x</sub>, TRPV<sub>x</sub>, and TRPM<sub>x</sub> correspond to short (C, canonical), (osm-9 -like, Vanilloid) and long (Melastatin family), respectively.

### TRPC (Short, Canonical TRPC) Family

The TRPC group can be divided into four subgroups (TRPC1; TRPC4,5; TRPC3,6,7; TRPC2) by sequence homology as well as functional similarities. TRPC1 was the first member of the mammalian TRP family reported to form an ion channel (Zhu et al., 1996; Zitt et al., 1996). Given the widespread expression of TRPC1 and its ability to coassemble with other TRPC subunits (Xu et al., 1997; Lintschinger et al., 2000; Strubing et al., 2001), TRPC1 might be a component of different heteromeric

TRP complexes. Whether TRPC1 can form functional channels in the absence of other TRP subunits is not established.

The second TRPC subfamily most closely related to TRPC1 is comprised of TRPC4 and TRPC5. Murine TRPC4 and TRPC5 can form homomeric cation channels that are activated following stimulation of G<sub>q</sub>-coupled receptors (Okada et al., 1998; Schaefer et al., 2000) as well as receptor tyrosine kinases (Schaefer et al., 2000). Coexpression of TRPC1 and TRPC4 or TRPC5 resulted in a novel nonselective cation channel with a voltage dependence similar to *N*-methyl-D-aspartate receptor channels, but unlike that of any reported TRP channel (Strubing et al., 2001). The details of the activation mechanism remain elusive but the two primary products of PLC enzyme activity, IP<sub>3</sub> and diacylglycerol, did not

activate TRPC4 and TRPC5 (Hofmann et al., 1999; Schaefer et al., 2000). Both TRPC4 and TRPC5 contain a C-terminal PDZ-binding motif (VTTRL) not present in other TRPs. PDZ domain scaffolding proteins such as the Na<sup>+</sup>/H<sup>+</sup> exchanger regulatory factor (NHERF) as well as signaling molecules like PLCβ1, coimmunoprecipitate with TRPC4 and TRPC5 (Tang et al., 2000), indicating that the channels may be part of multimolecular signaling complexes similar to the signalplex or transducisome of *Drosophila* photoreceptors.

TRPC3, TRPC6, and TRPC7 are ~75% identical and when expressed constitute a cation nonselective current that rectifies in both the inward (negative voltages) and outward (positive voltages) directions. Similar to TRPC3, TRPC6 and TRPC7 are inwardly and outwardly rectifying, have relatively low selectivity of Ca<sup>2+</sup> over Na<sup>+</sup>, are sensitive to intracellular Ca<sup>2+</sup>, and are activated by diacylglycerol (Hofmann et al., 1999; Okada et al., 1999). Their relatively high expression levels in smooth and heart muscle cells make them promising candidates for the as yet molecularly undefined nonselective cation channels in these muscle cells. In support of this idea is the finding that TRPC6 is an essential part of the α<sub>1</sub>-adrenoreceptor-activated cation channel in rabbit portal vein myocytes (Inoue et al., 2001).

Less information is available about TRPC2, which shares about 30% sequence identity with the TRPC3, TRPC6, and TRPC7 subfamilies. Full-length TRPC2 mRNA and several N-terminal splice variants have been found in mouse and rat tissue, but TRPC2 is a pseudogene in humans (Wes et al., 1995; Liman et al., 1999; Vannier et al., 1999; Hofmann et al., 2000). TRPC2 protein was localized to neuronal microvilli in rat vomeronasal organ (Liman et al., 1999). TRPC2-deficient mice display abnormal mating behavior, consistent with a role for this channel in pheromone signaling (Stowers et al., 2002).

### TRPV (osm-9-Like or Vanilloid Receptor TRP) Family

Currently the TRPV family has six members grouped into three subfamilies. TRPV1 and TRPV2 are the vanilloid receptors and vanilloid-like receptors, VR-1 and VRL-1, respectively. TRPV4 is the osm-9-like OTRPC4, and TRPV5 and TRPV6 are the Ca<sup>2+</sup>-selective channels, ECaC1/CaT2 (Epithelial Calcium Channel/Calcium Transporter) and ECaC2 (also called CaT1).

The vanilloid receptors are the most well understood ion channels in this class (Caterina and Julius, 2001). VR-1 (TRPV1) is activated by the "hot" pepper-derived vanilloid compound capsaicin (Caterina et al., 1997) but is not activated by store depletion. The expressed capsaicin receptor is a relatively Ca<sup>2+</sup>-selective ion channel with an outwardly rectifying I–V relation and exhibits Ca<sup>2+</sup>-dependent desensitization. Endogenous cannabinoids receptor ligands, such as anandamide, are poten-

tial TRPV1 agonists. The exact mechanism of TRPV1 activation is not completely understood, but it is sensitive to heat (>43°C), but the temperature at which it is activated is modulated by PIP<sub>2</sub> (Prescott and Julius, 2003). The size of the current is increased by acid pH and is modulated by intracellular PIP<sub>2</sub>, which appears to inhibit the channel (Chuang et al., 2001). Experiments with TRPV1<sup>-/-</sup> mice confirm a role for TRPV1 in transducing the nociceptive, inflammatory, and hypothermic effects of vanilloid compounds and that it contributes to acute thermal nociception and hyperalgesia following tissue injury (Caterina et al., 2000). Analysis of TRPV1<sup>-/-</sup> mice indicates that TRPV1 is essential for normal mechanically evoked purinergic signaling by the bladder urothelium (Birder et al., 2002).

The vanilloid receptor-like channel (VRL-1, TRPV2) is 50% identical to TRPV1, but is insensitive to capsaicin (Caterina et al., 1999). Like TRPV1 it is more permeable to Ca<sup>2+</sup> than Na<sup>+</sup> (P<sub>Ca</sub>/P<sub>Na</sub> = 3/1) and is outwardly rectifying. It has been proposed to mediate high threshold (>52°C) noxious heat sensation, perhaps in the lightly myelinated Aδ nociceptors, but its presence in nonsensory tissue suggests other functions as well.

TRPV3 and TRPV4 are moderately Ca<sup>2+</sup>-selective channels. Both are sensitive to warmth in the range >31°C (Peier et al., 2002b; Smith et al., 2002; Xu et al., 2002) and >25°C, respectively. TRPV3 is also highly expressed in keratinocytes (Peier et al., 2002b), hair follicles, and on the surface of the primate tongue (Xu et al., 2002). Temperature increases TRPV4-mediated current approximately in the >25°C range, and this is potentiated by hypotonicity (Guler et al., 2002). TRPV4 current is increased by cell swelling (Liedtke et al., 2000; Strotmann et al., 2000). TRPV4<sup>-/-</sup> mice display an increase in antidiuretic hormone secretion in response to hyperosmolarity (Mizuno et al., 2003). Hypotonicity increases TRPV4 current in primary afferent nociceptive nerve fibers and is enhanced by the hyperalgesic inflammatory mediator, prostaglandin E<sub>2</sub> (Alessandri-Haber et al., 2003).

TRPV5 (ECaC, CaT2) (Hoenderop et al., 1999) is only 30% identical to TRPV1 but is similar to TRPV6 (66% identical) and indeed many of its electrophysiological properties are indistinguishable from it. The expressed channel is strongly inwardly rectified and is relatively highly Ca<sup>2+</sup>-selective (P<sub>Ca</sub>/P<sub>Na</sub> > 100) (Nilius et al., 2000; Vennekens et al., 2000). These properties are consistent with proposed mechanisms for Ca<sup>2+</sup>-selective channels in which negatively charged glutamic or aspartic acid residues provide a binding site for divalents within the pore (Tsien et al., 1987). Store-dependent activation of this channel has not been reported. TRPV6 has a wide tissue distribution. TRPV6 is Ca<sup>2+</sup>-selective (P<sub>Ca</sub>/P<sub>Na</sub> > 100), activated by low levels of intracellular [Ca<sup>2+</sup>], and inactivated by higher [Ca<sup>2+</sup>]<sub>i</sub> (Yue et al., 2001). Like TRPV5, TRPV6 displays a steeply inwardly rectifying I–V relation, passing most of its current at

hyperpolarized potentials. It exhibits less  $\text{Ca}^{2+}$  selectivity and has a larger single channel conductance than  $I_{\text{CRAC}}$  and thus is unlikely to be CRAC. Neither channel appears to be operated by store depletion.

### TRPM (Long TRPC, Melastatin) Family

The TRPM (long TRP, melastatin) family has eight members divided into four groups. TRPM1 (melastatin) was initially identified through a screen of human melanoma-correlated mRNAs (Duncan et al., 1998). Although analysis of TRPM1 mRNA indicates wide expression, TRPM1's electrophysiological properties have not been studied.

TRPM2 (Nagamine et al., 1998) is a  $\text{Ca}^{2+}$ -permeant nonselective channel with a linear I–V relation (Perraud et al., 2001). ADP-ribose and potentially NAD (Perraud et al., 2001; Sano et al., 2001) bind a C-terminal NUDT9 Nudix hydrolase family domain to gate the channel. As for many  $\text{Ca}^{2+}$ -permeant channels, it appears to be inactivated by  $[\text{Ca}^{2+}]_i$ . TRPM3 (Grimm et al., 2003; Lee et al., 2003) is a  $\text{Ca}^{2+}$ -permeant nonselective channel with a linear I–V relation that is constitutively active when heterologously expressed. Its activity is increased by hypotonicity (200 mOsm/l) (Grimm et al., 2003), and its expression pattern indicates that TRPM4 functions in kidney and in the central nervous system.

TRPM4 and TRPM5 are the only monovalent-selective ion channels of the TRP family. Both are  $\text{Ca}^{2+}$ -activated  $\sim 20$  to  $30$  pS nonselective channels (Launay et al., 2002; Hofmann et al., 2003). G protein receptors coupled to PLC-dependent endoplasmic reticular  $\text{Ca}^{2+}$  release activate these channels, perhaps by direct  $\text{Ca}^{2+}$  binding (Launay et al., 2002; Hofmann et al., 2003). Although their instantaneous I–Vs are linear, they exhibit voltage dependence with continued depolarization (Hofmann et al., 2003; Nilius et al., 2003). TRPM5 is activated by sweet, umami, and bitter taste G protein-coupled receptor pathways (Zhang et al., 2003).

TRPM6 and TRPM7 contain a functional kinase domain within their polypeptide chain. TRPM7 was identified in a yeast two-hybrid screen as a protein interacting with  $\text{PLC}\beta_1$ , and was the first member of the TRPM group to be expressed as a functional ion channel (Clapham et al., 2001; Runnels et al., 2001). TRPM7 passes little inward current under physiological conditions, is permeant to both  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$ , and is inhibited by  $\sim 0.6$  mM intracellular-free  $\text{Mg}^{2+}$  (Nadler et al., 2001). TRPM7 current increases slowly under whole-cell recording conditions, and it is inactivated by  $\text{PIP}_2$  hydrolysis by  $\text{PLC}\beta$  or  $\text{PLC}\gamma$  (Runnels et al., 2002). The function of the kinase domain is poorly understood but is not required for channel activation (Runnels et al., 2002; Schmitz et al., 2003). The substrates of the kinase, an atypical serine/threonine kinase, have not been identified (Yamaguchi et al., 2001). Familial hypomagnesemia

with secondary hypocalcemia is caused by a mutation in TRPM6 (Schlingmann et al., 2002; Walder et al., 2002).

TRPM8 is up-regulated in prostate and other cancers (Tsavaler et al., 2001). TRPM8 is a nonselective, outwardly rectifying channel that can be activated by cold ( $8$ – $28^\circ\text{C}$ ) and enhanced by “cooling” compounds such as menthol and icilin (McKemy et al., 2002; Peier et al., 2002a). It is therefore likely to contribute to the sensation of cold temperature.

### TRPA Ankyrin-Repeat TRP Channel Family

ANKTM1 is a  $\text{Ca}^{2+}$ -permeant nonselective channel with  $\sim 14$  ankyrin repeats in its N terminus. It is activated by noxious cold (Story et al., 2003). TRPA1 is found in nociceptive sensory DRG, and its homolog in *Drosophila*, *painless*. Many other TRP channels have not been systematically tested for temperature sensitivity, but such a comparison would clarify the field.

### Conclusion

The TRP channels are a family of six transmembrane-spanning domain proteins expressed in low numbers per cell to yield small net inward currents. At this time, there is no unifying theme in their function or mechanism for activation. The TRPV (osm-9 like/vanilloid) subfamily is the most well characterized of the group and includes ion channels that are certainly involved in neuronal pain pathways, perhaps to sense heat and osmolarity. The TRPM (long TRPC, melastatin) subfamily may well be the most novel, with potential roles in  $\text{Ca}^{2+}$ -dependent signaling, control of cell cycle progression, division or migration, and thermosensation. TRP proteins are common in many cell types, making expression of confirmed monomeric channels difficult. Several TRPs are known to form heteromultimers, and their electrophysiological properties depend on the subunit composition. The multipotent phosphatidylinositol pathway is involved in most TRP regulation, but the details of this regulation are just beginning to be elucidated.

### References

- Alessandri-Haber N, Yeh JJ, Boyd AE, Parada CA, Chen X, Reichling DB, and Levine JD (2003) Hypotonicity induces TRPV4-mediated nociception in rat. *Neuron* **39**:497–511.
- Birder LA, Nakamura Y, Kiss S, Nealen ML, Barrick S, Kanai AJ, Wang E, Ruiz G, De Groat WC, Apodaca G, et al. (2002) Altered urinary bladder function in mice lacking the vanilloid receptor TRPV1. *Nat Neurosci* **5**:856–860.
- Caterina MJ and Julius D (2001) The vanilloid receptor: a molecular gateway to the pain pathway. *Annu Rev Neurosci* **24**:487–517.
- Caterina MJ, Leffler A, Malmberg AB, Martin WJ, Trafton J, Petersen-Zeitl KR, Koltzenburg M, Basbaum AI, and Julius D (2000) Impaired nociception and pain sensation in mice lacking the capsaicin receptor. *Science (Wash DC)* **288**:306–313.
- Caterina MJ, Rosen TA, Tominaga M, Brake AJ, and Julius D (1999) A capsaicin-receptor homologue with a high threshold for noxious heat. *Nature (Lond)* **398**:436–441.
- Caterina MJ, Schumacher MA, Tominaga M, Rosen TA, Levine JD, and Julius D (1997) The capsaicin receptor: a heat-activated ion channel in the pain pathway. *Nature (Lond)* **389**:816–824.
- Chuang HH, Prescott ED, Kong H, Shields S, Jordt SE, Basbaum AI, Chao MV, and Julius D (2001) Bradykinin and nerve growth factor release the capsaicin receptor from  $\text{PtdIns}(4,5)\text{P}_2$ -mediated inhibition. *Nature (Lond)* **411**:957–962.
- Clapham DE (1995) Calcium signaling. *Cell* **80**:259–268.
- Clapham DE, Runnels LW, and Strubing C (2001) The TRP ion channel family. *Nat Rev Neurosci* **2**:387–396.
- Doyle DA, Morais CJ, Pfuetzner RA, Kuo A, Gulbis JM, Cohen SL, Chait BT, and

- MacKinnon R (1998) The structure of the potassium channel: molecular basis of  $K^+$  conduction and selectivity. *Science (Wash DC)* **280**:69–77.
- Duncan LM, Deeds J, Hunter J, Shao J, Holmgren LM, Woolf EA, Tepper RI, and Shyjan AW (1998) Down-regulation of the novel gene melastatin correlates with potential for melanoma metastasis. *Cancer Res* **58**:1515–1520.
- Grimm C, Kraft R, Sauerbruch S, Schultz G, and Harteneck C (2003) Molecular and functional characterization of the melastatin-related cation channel TRPM3. *J Biol Chem* **278**:21493–21501.
- Guler AD, Lee H, Iida T, Shimizu I, Tominaga M, and Caterina M (2002) Heat-evoked activation of the ion channel, TRPV4. *J Neurosci* **22**:6408–6414.
- Hardie RC (2003) Regulation of TRP channels via lipid second messengers. *Annu Rev Physiol* **65**:735–759.
- Harteneck C, Plant TD, and Schultz G (2000) From worm to man: three subfamilies of TRP channels. *Trends Neurosci* **23**:159–166.
- Hoenderop JG, van der Kemp AW, Hartog A, van de Graaf SF, van Os CH, Willems PH, and Bindels RJ (1999) Molecular identification of the apical  $Ca^{2+}$  channel in 1,25-dihydroxyvitamin D<sub>3</sub>-responsive epithelia. *J Biol Chem* **274**:8375–8378.
- Hofmann T, Chubanov V, Gudermann T, and Montell C (2003) TRPM5 is a voltage-modulated and  $Ca^{2+}$ -activated monovalent selective cation channel. *Curr Biol* **13**:1153–1158.
- Hofmann T, Obukhov AG, Schaefer M, Harteneck C, Gudermann T, and Schultz G (1999) Direct activation of human TRPC6 and TRPC3 channels by diacylglycerol. *Nature (Lond)* **397**:259–263.
- Hofmann T, Schaefer M, Schultz G, and Gudermann T (2000) Cloning, expression and subcellular localization of two novel splice variants of mouse transient receptor potential channel 2. *Biochem J* **351**:115–122.
- Hoth M and Penner R (1992) Depletion of intracellular calcium stores activates a calcium current in mast cells. *Nature (Lond)* **355**:353–356.
- Inoue R, Okada T, Onoue H, Hara Y, Shimizu S, Naitoh S, Ito Y, and Mori Y (2001) The transient receptor potential homologue TRP6 is the essential component of vascular  $\alpha(1)$ -adrenoceptor-activated  $Ca^{2+}$ -permeable cation channel. *Circ Res* **88**:325–332.
- Launay P, Fleig A, Perraud AL, Scharenberg AM, Penner R, and Kinet JP (2002) TRPM4 is a  $Ca^{2+}$ -activated nonselective cation channel mediating cell membrane depolarization. *Cell* **109**:397–407.
- Lee N, Chen J, Sun L, Wu S, Gray KR, Rich A, Huang M, Lin JH, Feder JN, Janovitz EB, et al. (2003) Expression and characterization of human transient receptor potential melastatin 3 (hTRPM3). *J Biol Chem* **278**:20890–20897.
- Lewis RS (1999) Store-operated calcium channels. *Adv Second Messenger Phosphoprotein Res* **33**:279–307.
- Liedtke W, Choe Y, Marti-Renom MA, Bell AM, Denis CS, Sali A, Hudspeth AJ, Friedman JM, and Heller S (2000) Vanilloid receptor-related osmotically activated channel (VR-OAC), a candidate vertebrate osmoreceptor. *Cell* **103**:525–535.
- Liman ER, Corey DP, and Dulac C (1999) TRP2: a candidate transduction channel for mammalian pheromone sensory signaling. *Proc Natl Acad Sci USA* **96**:5791–5796.
- Lintschinger B, Balzer-Geldsetzer M, Baskaran T, Graier WF, Romanin C, Zhu MX, and Groschner K (2000) Coassembly of trp1 and trp3 proteins generates diacylglycerol- and  $Ca^{2+}$ -sensitive cation channels. *J Biol Chem* **275**:27799–27805.
- McKemy DD, Neuhauss WM, and Julius D (2002) Identification of a cold receptor reveals a general role for TRP channels in thermosensation. *Nature (Lond)* **416**:52–58.
- Mizuno A, Matsumoto N, Imai M, and Suzuki M (2003) Impaired osmotic sensation in mice lacking TRPV4. *Am J Physiol Cell Physiol* **285**:C96–C101.
- Montell C (1999) Visual transduction in *Drosophila*. *Annu Rev Cell Dev Biol* **15**:231–268.
- Montell C, Birnbaumer L, and Flockerzi V (2002) The TRP channels, a remarkably functional family. *Cell* **108**:595–598.
- Nadler MJ, Hermosura MC, Inabe K, Perraud AL, Zhu Q, Stokes AJ, Kurokaki T, Kinet JP, Penner R, Scharenberg AM, and Fleig A (2001) LTRPC7 is a Mg-ATP-regulated divalent cation channel required for cell viability. *Nature (Lond)* **411**:590–595.
- Nagamine K, Kudoh J, Minoshima S, Kawasaki K, Asakawa S, Ito F, and Shimizu N (1998) Molecular cloning of a novel putative  $Ca^{2+}$  channel protein (TRPC7) highly expressed in brain. *Genomics* **54**:124–131.
- Nilius B, Prenen J, Droogmans G, Voets T, Vennekens R, Freichel M, Wissenbach U, and Flockerzi V (2003) Voltage dependence of the  $Ca^{2+}$  activated cation channel TRPM4. *J Biol Chem* **278**:30813–30820.
- Nilius B, Vennekens R, Prenen J, Hoenderop JG, Bindels RJ, and Droogmans G (2000) Whole-cell and single channel monovalent cation currents through the novel rabbit epithelial  $Ca^{2+}$  channel ECaC. *J Physiol* **527**:239–248.
- Okada T, Inoue R, Yamazaki K, Maeda A, Kurokaki T, Yamakuni T, Tanaka I, Shimizu S, Ikenaka K, Imoto K, and Mori Y (1999) Molecular and functional characterization of a novel mouse transient receptor potential homologue TRP7.  $Ca^{2+}$ -permeable cation channel that is constitutively activated and enhanced by stimulation of G protein-coupled receptor. *J Biol Chem* **274**:27359–27370.
- Okada T, Shimizu S, Wakamori M, Maeda A, Kurokaki T, Takada N, Imoto K, and Mori Y (1998) Molecular cloning and functional characterization of a novel receptor-activated TRP  $Ca^{2+}$  channel from mouse brain. *J Biol Chem* **273**:10279–10287.
- Peier AM, Moqrich A, Hergarden AC, Reeve AJ, Andersson DA, Story GM, Earley TJ, Dragoni I, McIntyre P, Bevan S, and Patapoutian A (2002a) A TRP channel that senses cold stimuli and menthol. *Cell* **108**:705–715.
- Peier AM, Reeve AJ, Andersson DA, Moqrich A, Earley TJ, Hergarden AC, Story GM, Colley S, Hogenesch JB, McIntyre P, et al. (2002b) A heat-sensitive TRP channel expressed in keratinocytes. *Science (Wash DC)* **296**:2046–2049.
- Perraud AL, Fleig A, Dunn CA, Bagley LA, Launay P, Schmitz C, Stokes AJ, Zhu Q, Bessman MJ, Penner R, et al. (2001) ADP-ribose gating of the calcium-permeable LTRPC2 channel revealed by Nudix motif homology. *Nature (Lond)* **411**:595–599.
- Prescott ED and Julius D (2003) A modular PIP2 binding site as a determinant of capsaicin receptor sensitivity. *Science (Wash DC)* **300**:1284–1288.
- Putney JW Jr (1977) Muscarinic, alpha-adrenergic and peptide receptors regulate the same calcium influx sites in the parotid gland. *J Physiol* **268**:139–149.
- Runnels LW, Yue L, and Clapham DE (2001) TRP-PLIK, a bifunctional protein with kinase and ion channel activities. *Science (Wash DC)* **291**:1043–1047.
- Runnels LW, Yue L, and Clapham DE (2002) The TRPM7 channel is inactivated by PIP(2) hydrolysis. *Nat Cell Biol* **4**:329–336.
- Saitou N and Nei M (1987) The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol Biol Evol* **4**:406–425.
- Sano Y, Inamura K, Miyake A, Mochizuki S, Yokoi H, Matsushime H, and Furuichi K (2001) Immuncyte  $Ca^{2+}$  influx system mediated by LTRPC2. *Science (Wash DC)* **293**:1327–1330.
- Schaefer M, Plant TD, Obukhov AG, Hofmann T, Gudermann T, and Schultz G (2000) Receptor-mediated regulation of the nonselective cation channels TRPC4 and TRPC5. *J Biol Chem* **275**:17517–17526.
- Schlingmann KP, Weber S, Peters M, Niemann Nejsum L, Vitzthum H, Klingel K, Kratz M, Haddad E, Ristoff E, Dinour D, et al. (2002) Hypomagnesemia with secondary hypocalcemia is caused by mutations in TRPM6, a new member of the TRPM gene family. *Nat Genet* **31**:166–170.
- Schmitz C, Perraud AL, Johnson CO, Inabe K, Smith MK, Penner R, Kurokaki T, Fleig A, and Scharenberg AM (2003) Regulation of vertebrate cellular  $Mg^{2+}$ -homeostasis by TRPM7. *Cell* **114**:191–200.
- Smith GD, Gunthorpe MJ, Kelsell RE, Hayes PD, Reilly P, Facer P, Wright JE, Jerman JC, Walhin JP, Ooi L, et al. (2002) TRPV3 is a temperature-sensitive vanilloid receptor-like protein. *Nature (Lond)* **418**:186–190.
- Story GM, Peier AM, Reeve AJ, Eid SR, Mosbacher J, Hricik TR, Earley TJ, Hergarden AC, Andersson DA, Hwang SW, et al. (2003) ANKTM1, a TRP-like channel expressed in nociceptive neurons, is activated by cold temperatures. *Cell* **112**:819–829.
- Stowers L, Holy TE, Meister M, Dulac C, and Koentges G (2002) Loss of sex discrimination and male-male aggression in mice deficient for TRP2. *Science (Wash DC)* **195**:1493–1500.
- Strotmann R, Harteneck C, Nunnenmacher K, Schultz G, and Plant TD (2000) OTRPC4, a nonselective cation channel that confers sensitivity to extracellular osmolarity. *Nat Cell Biol* **2**:695–702.
- Strubing C, Krapivinsky G, Krapivinsky L, and Clapham DE (2001) TRPC1 and TRPC5 form a novel cation channel in mammalian brain. *Neuron* **29**:646–655.
- Tang Y, Tang J, Chen Z, Trost C, Flockerzi V, Li M, Ramesh V, and Zhu MX (2000) Association of mammalian trp4 and phospholipase C isozymes with a PDZ domain-containing protein, NHERF. *J Biol Chem* **275**:37559–37564.
- Tsavaler L, Shapero MH, Morkowski S, and Laus R (2001) Trp-p8, a novel prostate-specific gene, is up-regulated in prostate cancer and other malignancies and shares high homology with transient receptor potential calcium channel proteins. *Cancer Res* **61**:3760–3769.
- Tsien RW, Hess P, McCleskey EW, and Rosenberg RL (1987) Calcium channels: mechanisms of selectivity, permeation and block. *Annu Rev Biophys Chem* **16**:265–290.
- Vannier B, Peyton M, Boulay G, Brown D, Qin N, Jiang M, Zhu X, and Birnbaumer L (1999) Mouse trp2, the homologue of the human trpc2 pseudogene, encodes mTrp2, a store depletion-activated capacitative  $Ca^{2+}$  entry channel. *Proc Natl Acad Sci USA* **96**:2060–2064.
- Vennekens R, Hoenderop JG, Prenen J, Stuiver M, Willems PH, Droogmans G, Nilius B, and Bindels RJ (2000) Permeation and gating properties of the novel epithelial  $Ca^{2+}$  channel. *J Biol Chem* **275**:3963–3969.
- Walder RY, Landau D, Meyer P, Shalev H, Tsolia M, Borochowitz Z, Boettger MB, Beck GE, Englehardt RK, Carmi R, and Sheffeld VC (2002) Mutation of TRPM6 causes familial hypomagnesemia with secondary hypocalcemia. *Nat Genet* **31**:171–174.
- Wes PD, Chevesich J, Jeromin A, Rosenberg C, Stetten G, and Montell C (1995) TRPC1, a human homolog of a *Drosophila* store-operated channel. *Proc Natl Acad Sci USA* **92**:9652–9656.
- Xu H, Ramsey IS, Kotecha SA, Moran MM, Chong JA, Lawson D, Ge P, Lilly J, Silos-Santiago I, Xie Y, et al. (2002) TRPV3 is a calcium-permeable temperature-sensitive cation channel. *Nature (Lond)* **418**:181–186.
- Xu XZ, Li HS, Guggino WB, and Montell C (1997) Coassembly of TRP and TRPL produces a distinct store-operated conductance. *Cell* **89**:1155–1164.
- Yamaguchi H, Matsushita M, Nairn AC, and Kuriyan J (2001) Crystal structure of the atypical protein kinase domain of a TRP channel with phosphotransferase activity. *Mol Cell* **7**:1047–1057.
- Yue L, Peng JB, Hediger MA, and Clapham DE (2001) CaT1 manifests the pore properties of the calcium-release-activated calcium channel. *Nature (Lond)* **410**:705–709.
- Zhang Y, Hoon MA, Chandrashekar J, Mueller KL, Cook B, Wu D, Zuker CS, and Ryba NJ (2003) Coding of sweet, bitter and umami tastes: different receptor cells sharing similar signaling pathways. *Cell* **112**:293–301.
- Zhu X, Jiang M, Peyton M, Boulay G, Hurst R, Stefani E, and Birnbaumer L (1996) trp, a novel mammalian gene family essential for agonist-activated capacitative  $Ca^{2+}$  entry. *Cell* **85**:661–671.
- Zitt C, Zobel A, Obukhov AG, Harteneck C, Kalkbrenner F, Luckhoff A, and Schultz G (1996) Cloning and functional expression of a human  $Ca^{2+}$ -permeable cation channel activated by calcium store depletion. *Neuron* **16**:1189–1196.