

International Union of Pharmacology. XLVIII. Nomenclature and Structure-Function Relationships of Voltage-Gated Calcium Channels

WILLIAM A. CATTERALL, EDWARD PEREZ-REYES, TERRANCE P. SNUTCH, AND JOERG STRIESSNIG

Department of Pharmacology, University of Washington, Seattle, Washington (W.A.C.); Department of Pharmacology, University of Virginia, Charlottesville, Virginia (E.P.-R.); Biotechnology Laboratory, University of British Columbia, Vancouver, British Columbia, Canada (T.P.S.); and Abteilung Pharmakologie und Toxikologie, Institut für Pharmazie, Universität Innsbruck, Innsbruck, Austria (J.S.)

Abstract—The family of voltage-gated calcium channels serves as the key transducers of cell surface membrane potential changes into local intracellular calcium transients that initiate many different physiological events. There are 10 members of the voltage-gated calcium channel family that have been characterized in

mammals, and they serve distinct roles in cellular signal transduction. This article presents the molecular relationships and physiological functions of these calcium channel proteins and provides comprehensive information on their molecular, genetic, physiological, and pharmacological properties.

Introduction

Voltage-gated calcium channels mediate calcium influx in response to membrane depolarization and regulate intracellular processes such as contraction, secretion, neurotransmission, and gene expression in many different cell types. Their activity is essential to couple electrical signals in the cell surface to physiological events in cells. They are members of a gene superfamily of transmembrane ion channel proteins that includes voltage-gated potassium and sodium channels (Yu and Catterall, 2004). This compendium presents an introduction to their biochemical, molecular, and genetic properties, their physiological roles, and their pharmacological significance. Table 1 and the summary tables that follow the text of this article give comprehensive information on each member of the calcium channel family.

Calcium Channel Subunits

The calcium channels that have been characterized biochemically are complex proteins composed of four or five distinct subunits that are encoded by multiple genes (Fig. 1; Catterall, 2000). The α_1 subunit of 190 to 250 kDa is the largest subunit, and it incorporates the conduction pore, the voltage sensor and gating apparatus, and most of the known sites of channel regulation by second messengers, drugs, and toxins. Like the α sub-

units of sodium channels, the α_1 subunit of voltage-gated calcium channels is organized in four homologous domains (I–IV), with six transmembrane segments (S1–S6) in each. The S4 segment serves as the voltage sensor. The pore loop between transmembrane segments S5 and S6 in each domain determines ion conductance and selectivity, and changes of only three amino acids in the pore loops in domains I, III, and IV will convert a sodium channel to calcium selectivity. An intracellular β subunit and a transmembrane, disulfide-linked $\alpha_2\delta$ subunit complex are components of most types of calcium channels. A γ subunit has also been found in skeletal muscle calcium channels, and related subunits are expressed in heart and brain. Although these auxiliary subunits modulate the properties of the channel complex, the pharmacological and electrophysiological diversity of calcium channels arises primarily from the existence of multiple α_1 subunits (Hofmann et al., 1994).

Calcium Currents

Calcium currents recorded in different cell types have diverse physiological and pharmacological properties, and an alphabetical nomenclature has evolved for the distinct classes of calcium currents (Tsien et al., 1995). L-type calcium currents typically require a strong depolarization for activation, are long-lasting, and are blocked by the organic L-type calcium channel antagonists, including dihydropyridines, phenylalkylamines, and benzothiazepines. They are the main calcium currents recorded in muscle and endocrine cells, where they initiate contraction and secretion. L-type currents activating at lower voltages also exist predominantly in neurons and cardiac pacemaker cells. N-type, P/Q-type, and R-type calcium currents

Address correspondence to: Dr. William A. Catterall, Department of Pharmacology, University of Washington, Mailstop 357280, Seattle, WA 98195-7280. E-mail: wcatt@u.washington.edu

The authors serve as the Subcommittee on Calcium Channels of the Nomenclature Committee of the International Union of Pharmacology.

Article, publication date, and citation information can be found at <http://pharmrev.aspetjournals.org>.

doi:10.1124/pr.57.4.5.

TABLE 1
Physiological function and pharmacology of calcium channels

Channel	Current	Localization	Specific Antagonists	Cellular Functions
Ca _v 1.1	L	Skeletal muscle; transverse tubules	Dihydropyridines; phenylalkylamines; benzothiazepines	Excitation-contraction coupling
Ca _v 1.2	L	Cardiac myocytes; smooth muscle myocytes; endocrine cells; neuronal cell bodies; proximal dendrites	Dihydropyridines; phenylalkylamines; benzothiazepines	Excitation-contraction coupling; hormone release; regulation of transcription; synaptic integration
Ca _v 1.3	L	Endocrine cells; neuronal cell bodies and dendrites; cardiac atrial myocytes and pacemaker cells; cochlear hair cells	Dihydropyridines; phenylalkylamines; benzothiazepines	Hormone release; regulation of transcription; synaptic regulation; cardiac pacemaking; hearing; neurotransmitter release from sensory cells
Ca _v 1.4	L	Retinal rod and bipolar cells; spinal cord; adrenal gland; mast cells	Dihydropyridines; phenylalkylamines; benzothiazepines	Neurotransmitter release from photoreceptors
Ca _v 2.1	P/Q	Nerve terminals and dendrites; neuroendocrine cells	ω -Agatoxin IVA	Neurotransmitter release; dendritic Ca ²⁺ transients; hormone release
Ca _v 2.2	N	Nerve terminals and dendrites; neuroendocrine cells	ω -Conotoxin-GVIA	Neurotransmitter release; dendritic Ca ²⁺ transients; hormone release
Ca _v 2.3	R	Neuronal cell bodies and dendrites	SNX-482	Repetitive firing; dendritic calcium transients
Ca _v 3.1	T	Neuronal cell bodies and dendrites; cardiac and smooth muscle myocytes	None	Pacemaking; repetitive firing
Ca _v 3.2	T	Neuronal cell bodies and dendrites; cardiac and smooth muscle myocytes	None	Pacemaking; repetitive firing
Ca _v 3.3	T	Neuronal cell bodies and dendrites	None	Pacemaking; repetitive firing




FIG. 1. Subunit structure of Ca_v1 channels. The subunit composition and structure of calcium channels purified from skeletal muscle are illustrated. The model is updated from the original description of the subunit structure of skeletal muscle calcium channels. This model fits available biochemical and molecular biological results for other Ca_v1 channels and for Ca_v2 channels. Predicted α helices are depicted as cylinders. The lengths of lines correspond approximately to the lengths of the polypeptide segments represented.

also require strong depolarization for activation. They are relatively unaffected by L-type calcium channel antagonist drugs but are blocked by specific polypeptide toxins from snail and spider venoms. They are expressed primarily in neurons, where they initiate neurotransmission at most fast synapses and mediate calcium entry into cell bodies and dendrites. T-type calcium currents are activated by weak depolarization and are transient. They are resistant to both organic antagonists and to the snake and spider toxins used to

define the N- and P/Q-type calcium currents. They are expressed in a wide variety of cell types, where they are involved in shaping the action potential and controlling patterns of repetitive firing.

Calcium Channel Genes

Mammalian α_1 subunits are encoded by at least 10 distinct genes. Historically, various names have been given to the corresponding gene products, giving rise to

distinct and sometimes confusing nomenclatures. In 1994, a unified but arbitrary nomenclature was adopted in which α_1 subunits were referred to as α_{1S} for the original skeletal muscle isoform and α_{1A} through α_{1E} for those discovered subsequently (Birnbaumer et al., 1994). In 2000, a rational nomenclature was adopted (Ertel et al., 2000) based on the well defined potassium channel nomenclature (Chandy and Gutman, 1993). Calcium channels were named using the chemical symbol of the principal permeating ion (Ca) with the principal physiological regulator (voltage) indicated as a subscript (Ca_V). The numerical identifier corresponds to the Ca_V channel α_1 subunit gene subfamily (1 to 3 at present) and the order of discovery of the α_1 subunit within that subfamily (1 through n). According to this nomenclature, the Ca_V1 subfamily ($\text{Ca}_V1.1$ – $\text{Ca}_V1.4$) includes channels containing α_{1S} , α_{1C} , α_{1D} , and α_{1F} , which mediate L-type Ca^{2+} currents (Table 1). The Ca_V2 subfamily ($\text{Ca}_V2.1$ – $\text{Ca}_V2.3$) includes channels containing α_{1A} , α_{1B} , and α_{1E} , which mediate P/Q-type, N-type, and R-type Ca^{2+} currents, respectively (Table 1). The Ca_V3 subfamily ($\text{Ca}_V3.1$ – $\text{Ca}_V3.3$) includes channels containing α_{1G} , α_{1H} , and α_{1I} , which mediate T-type Ca^{2+} currents.

The complete amino acid sequences of these α_1 subunits are more than 70% identical within a subfamily but less than 40% identical among the three subfamilies. These family relationships are illustrated for the more conserved transmembrane and pore domains in Fig. 2. The division of calcium channels into these three families is phylogenetically ancient, as representatives of each are found in the *Caenorhabditis elegans* genome. Consequently, the genes for the different α_1 subunits have become widely dispersed in the genome, and even the most closely related members of the family are not clustered on single chromosomes in mammals.

Calcium Channel Molecular Pharmacology

The pharmacology of the three subfamilies of calcium channels is quite distinct. The Ca_V1 channels are




FIG. 2. Sequence similarity of voltage-gated calcium channel α_1 subunits. Phylogenetic representation of the primary sequences of the calcium channels. Only the membrane-spanning segments and pore loops (~350 amino acids) are compared. First, all sequence pairs were compared, which clearly defines three subfamilies with intrafamily sequence identities above 80% (Ca_V1 , Ca_V2 , and Ca_V3). Then a consensus sequence was defined for each subfamily, and these three sequences were compared to one another, with intersubfamily sequence identities of ~52% (Ca_V1 vs. Ca_V2) and 28% (Ca_V3 vs. Ca_V1 or Ca_V2).

the molecular targets of the organic calcium channel blockers used widely in the therapy of cardiovascular diseases. These drugs are thought to act at three separate, but allosterically coupled, receptor sites (Table 1; reviewed in Glossmann and Striessnig, 1990). Phenylalkylamines are intracellular pore blockers, which are thought to enter the pore from the cytoplasmic side of the channel and block it. Their receptor site is formed by amino acid residues in the S6 segments in domains III and IV, in close analogy to the local anesthetic receptor site on sodium channels (Hockerman et al., 1997; Hofmann et al., 1999; Striessnig, 1999). Dihydropyridines can be channel activators or inhibitors and therefore are thought to act allosterically to shift the channel toward the open or closed state rather than by occluding the pore. Their receptor site includes amino acid residues in the S6 segments of domains III and IV and the S5 segment of domain III. The dihydropyridine receptor site is closely apposed to the phenylalkylamine receptor site and shares some common amino acid residues. Diltiazem and related benzothiazepines are thought to bind to a third receptor site, but the amino acid residues that are required for their binding overlap extensively with those required for phenylalkylamine binding.

The Ca_V2 subfamily of calcium channels is relatively insensitive to dihydropyridine calcium channel blockers, but these calcium channels are specifically blocked with high affinity by peptide toxins from spiders and marine snails (Miljanich and Ramachandran, 1995). The $\text{Ca}_V2.1$ channels are blocked specifically by ω -agatoxin IVA from funnel web spider venom. The $\text{Ca}_V2.2$ channels are blocked specifically by ω -conotoxin GVIA and related cone snail toxins. The $\text{Ca}_V2.3$ channels are blocked specifically by the synthetic peptide toxin SNX-482 derived from tarantula venom. These peptide toxins are potent blockers of synaptic transmission because of their specific effects on the Ca_V2 family of calcium channels.

The Ca_V3 subfamily of calcium channels are insensitive to both the dihydropyridines that block Ca_V1 channels and the spider and cone snail toxins that block the Ca_V2 channels, and there are no widely useful pharmacological agents that block T-type calcium currents (Perez-Reyes, 2003). The organic calcium channel blocker mibefradil is somewhat selective for T-type versus L-type calcium currents (3- to 5-fold). The peptide kurtotoxin inhibits the activation gating of $\text{Ca}_V3.1$ and $\text{Ca}_V3.2$ channels. Development of more specific and high-affinity blockers of the Ca_V3 family of calcium channels would be useful for therapy and a more detailed analysis of the physiological roles of these channels.

Tables 2 through 11 summarize the major molecular, physiological, and pharmacological properties for each of the 10 calcium channels that have been functionally expressed. Quantitative data are included for

voltage dependence of activation and inactivation, single-channel conductance, and binding of drugs and neurotoxins, focusing on those agents that are widely used and diagnostic of channel identity and function.

REFERENCES

Birnbaumer L, Campbell KP, Catterall WA, Harpold MM, Hofmann F, Horne WA, Mori Y, Schwartz A, Snutch TP, Tanabe T, et al. (1994) The naming of voltage-gated calcium channels. *Neuron* **13**:505–506.
 Catterall WA (2000) Structure and regulation of voltage-gated Ca²⁺ channels. *Annu Rev Cell Dev Biol* **16**:521–555.
 Chandy KG and Gutman GA (1993) Nomenclature for mammalian potassium channel genes. *Trends Pharmacol Sci* **14**:434.
 Ertel EA, Campbell KP, Harpold MM, Hofmann F, Mori Y, Perez-Reyes E, Schwartz A, Snutch TP, Tanabe T, Birnbaumer L, et al. (2000) Nomenclature of voltage-gated calcium channels *Neuron* **25**:533–535.

Glossmann H and Striessnig J (1990) Molecular properties of calcium channels *Rev Physiol Biochem Pharmacol* **114**:1–105.
 Hockerman GH, Peterson BZ, Johnson BD, and Catterall WA (1997) Molecular determinants of drug binding and action on L-type calcium channels. *Annu Rev Pharmacol Toxicol* **37**:361–396.
 Hofmann F, Biel M, and Flockerzi V (1994) Molecular basis for Ca²⁺ channel diversity. *Annu Rev Neurosci* **17**:399–418.
 Hofmann F, Lacinová L, and Klugbauer N (1999) Voltage-dependent calcium channels: from structure to function. *Rev Physiol Biochem Pharmacol* **139**:33–87.
 Miljanich GP and Ramachandran J (1995) Antagonists of neuronal calcium channels: structure, function and therapeutic implications. *Annu Rev Pharmacol Toxicol* **35**:707–734.
 Perez-Reyes E (2003) Molecular physiology of low-voltage-activated t-type calcium channels. *Physiol Rev* **83**:117–161.
 Striessnig J (1999) Pharmacology, structure and function of cardiac L-type calcium channels. *Cell Physiol Biochem* **9**:42–269.
 Tsien RW, Lipscombe D, Madison D, Bley K, and Fox A (1995) Reflections on calcium channel diversity, 1988–1994. *Trends Neurosci* **18**:52–54.
 Yu FH and Catterall WA (2004) The VGL-chanome: a protein superfamily specialized for electrical signaling and ionic homeostasis. *Sci STKE*. **253**:re15.

TABLE 2
Ca_v1.1 channels

Channel name	Ca _v 1.1
Description	Voltage-gated calcium channel α ₁ -subunit
Other names	α _{1s} , skeletal muscle L-type Ca ²⁺ channel, skeletal muscle dihydropyridine receptor
Molecular information	Human: 1873aa, L33798 (PMID: 7713519), chr0.1q32, <i>CACNA1S</i> , LocusID: 779 Rat: 1146aa (partial sequence), L04684 (PMID: 1335956), chr. 13, <i>Cacna1s</i> , LocusID: 116652 Mouse: 1861aa, L06234 (PMID: 1281468), chr. 1, <i>Cacna1s</i> , LocusID: 12292 (see 'Comments')
Associated subunits	α _{2δ} , β, γ ^{1,2}
Functional assays	Patch-clamp (whole-cell, single-channel), calcium imaging, gating charge movement, skeletal muscle contraction
Current	I _{Ca,L}
Conductance	13–17pS (in 90–110 mM Ba ²⁺) ^{3,4}
Ion selectivity	Ca ²⁺ > Sr ²⁺ > Mg ²⁺ > Ba ²⁺ ⁵
Activation	V _a = 8–14 mV, τ _a > 50 ms at +10 mV (10 mM Ca ²⁺) ^{4,6}
Inactivation	V _h = –8 mV, 40% current inactivation after 5 s (–5 mV) ⁴
Activators	BayK8644, dihydropyridine agonists, FPL64176 ^{2,8,9}
Gating modifiers	Dihydropyridine antagonists (e.g., (+)-isradipine; IC ₅₀ = 13 nM at –90 mV and 0.15 nM at –65 mV) ⁹
Blockers	Nonselective: cadmium (IC ₅₀ < 0.5 mM) ⁹ ; selective for Ca _v 1.x: verapamil, devapamil (IC ₅₀ < 1 μM) and other phenylalkylamines, (+)- <i>cis</i> -diltiazem (IC ₅₀ < 80 μM) ⁹
Radioligands	(+)-[³ H]isradipine (K _d = 0.2–0.7 nM) and other dihydropyridines; (–)-[³ H]devapamil (K _d = 2.5 nM), (+)- <i>cis</i> -[³ H]diltiazem (K _d = 50 nM) ²
Channel distribution	Skeletal muscle transverse tubules (tetramers) ¹⁰
Physiological functions	Excitation-contraction coupling and Ca ²⁺ homeostasis in skeletal muscle ¹¹
Mutations and pathophysiology	Point mutations cause hypokalemic periodic paralysis and malignant hyperthermia susceptibility in humans and muscular dysgenesis in mice (<i>mdg/mdg</i>) ^{12,13}
Pharmacological significance	Not established
Comments	The gene for Ca _v 1.1 was first isolated and characterized in rabbit (1873aa, M23919, X05921); several groups reported three-dimensional structures of the purified skeletal muscle calcium channel complex determined using electron cryomicroscopy and single-particle averaging ¹⁴

aa, amino acids; chr., chromosome; Bay K8644, methyl 1,4-dihydro-2,6-dimethyl-3-nitro-4-(2-trifluoromethylphenyl)-pyridine-5-carboxylate; FPL64176, methyl 2,5-dimethyl-4-[2-(phenylmethyl)benzoyl]-1H-pyrrole-3-carboxylate.

1. Takahashi M, Seagar MJ, Jones JF, Reber BF, and Catterall WA (1987) Subunit structure of dihydropyridine-sensitive calcium channels from skeletal muscle. *Proc Natl Acad Sci USA* **84**:5478–5482.
2. Glossmann H and Striessnig J (1990) Molecular properties of calcium channels. *Rev Physiol Biochem Pharmacol* **114**:1–105.
3. Dirksen RT, Nakai J, Gonzales A, Imoto K, and Beam KG (1997) The S5-S6 linker of repeat I is a critical determinant of L-type Ca²⁺ channel conductance. *Biophys J* **73**:1402–1409.
4. Freise D, Held B, Wissenbach U, Pfeifer A, Trost C, Himmerkus N, Schweig U, Freichel M, Biel M, Hoffmann F, et al. (2000) Absence of the γ subunit of the skeletal muscle dihydropyridine receptor increases L-type calcium currents and alters channel inactivation properties. *J Biol Chem* **275**:14476–14481.
5. Pizarro G, Fitts R, Uribe I, and Rios E (1989) The voltage-sensor of excitation-contraction coupling in skeletal muscle: ion dependence and selectivity. *J Gen Physiol* **94**:405–428.
6. Dirksen RT and Beam KG (1995) Single calcium channel behavior in native skeletal muscle. *J Gen Physiol* **105**:227–247.
7. Rovnyak GC, Kimball SD, Beyer B, Cucinotta G, DiMarco JD, Gougoutas J, Hedberg A, Malley M, McCarthy JP, Zhang R, et al. (1995) Calcium entry blockers and activators: conformational and structural determinants of dihydropyridine calcium channel modulators. *J Med Chem* **38**:119–129.
8. Striessnig J (1999) Pharmacology, structure and function of cardiac L-type Ca²⁺ channels. *Cell Physiol Biochem* **9**:242–269.
9. Glossmann H and Striessnig J (1988) Ca²⁺ channels. *Vitam Horm* **44**:155–328.
10. Flucher BE and Franzini-Armstrong C (1996) Formation of junctions involved in excitation-contraction coupling in skeletal and cardiac muscle. *Proc Natl Acad Sci USA* **93**:8101–8106.
11. Rios E, Pizarro G, and Stefani E (1992) Charge movement and the nature of signal transduction in skeletal muscle excitation-contraction coupling. *Annu Rev Physiol* **54**:109–133.
12. *CACNA1S*; Online Mendelian Inheritance in Man (OMIM) no. 114208.
13. Striessnig J, Hoda JC, Koschak A, Zaghetto F, Mullner C, Sinnegger-Brauns MJ, Wild C, Watschinger K, Trockenbacher A, and Pelster G (2004) L-type Ca²⁺ channels in Ca²⁺ channelopathies. *Biochem Biophys Res Commun* **322**:1341–1346.
14. Wang MC, Dolphin A, and Kitmitto A (2004) L-type voltage-gated calcium channels: understanding function through structure. *FEBS Lett* **564**:245–250.

TABLE 3
Ca_v1.2 channels

Channel name	Ca _v 1.2
Description	Voltage-gated calcium channel α_1 subunit
Other names	α_{1C} , cardiac or smooth muscle L-type Ca ²⁺ channel, cardiac or smooth muscle dihydropyridine receptor
Molecular information	Human: 2169aa, L29529 (cardiac; PMID: 8392192), 2138aa, Z34815 (fibroblast; PMID: 1316612); 2138aa, AF465484 (jejunum; PMID: 12176756); chr. 12p13.3, <i>CACNA1C</i> , LocusID: 775 Rat: 2169aa, M59786 (aortic smooth muscle; PMID: 2170396); 2140/2143aa, M67516/M67515 (brain; PMID: 1648941); chr. 4q42, <i>Cacna1c</i> , LocusID: 24239 Mouse: 2139aa, L01776 (brain; PMID: 1385406); chr. 6, <i>Cacna1c</i> , LocusID: 12288 (see 'Comments')
Associated subunits	$\alpha_2\delta$, β , $\gamma^{1,2}$
Functional assays	Patch-clamp (whole-cell, single-channel), calcium imaging, cardiac or smooth muscle contraction hormone secretion
Current	I_{CaL}
Conductance	Ba ²⁺ (25pS) > Sr ²⁺ = Ca ²⁺ (9pS) ³
Ion selectivity	Ca ²⁺ > Sr ²⁺ > Ba ²⁺ >> Mg ²⁺ from permeability ratios
Activation	$V_a = -17$ mV (in 2 mM Ca ²⁺ ; HEK cells) ⁴ ; -4 mV (in 15 mM Ba ²⁺ ; HEK cells) to -18.8 mV (in 5 mM Ba ²⁺ ; HEK cells and <i>Xenopus oocytes</i>) ⁵⁻⁷ ; $\tau_a = 1$ ms at $+10$ mV ⁵
Inactivation	$V_h = -50$ to -60 mV (in 2 mM Ca ²⁺ ; HEK cells), $V_h = -18$ to -42 mV (in 5–15 mM Ba ²⁺ ; HEK cells) ^{5,7,8,9} ; $\tau_{fast} = 150$ ms, $\tau_{slow} = 1100$ ms; 61% inactivated after 250 ms in HEK cells ⁸ (at V_{max} in 15 mM Ba ²⁺) ⁴ ; $\sim 70\%$ inactivation after 1 s (at V_{max} in 2 mM Ca ²⁺) ⁴ ; inactivation is accelerated with Ca ²⁺ as charge carrier (calcium-dependent inactivation: 86% inactivated after 250 ms ^{8,10})
Activators	BayK8644, dihydropyridine agonists, FPL64176 ^{10,11}
Gating modifiers	Dihydropyridine antagonists (e.g., isradipine, IC ₅₀ = 7 nM at -60 mV; nimodipine, IC ₅₀ = 139 nM at -80 mV) ^{6,9}
Blockers	Nonselective: Cd ²⁺ ¹² ; selective for Ca _v 1.x: devapamil (IC ₅₀ = 50 nM in 10 mM Ba ²⁺ at -60 mV) and other phenylalkylamines; diltiazem (IC ₅₀ = 33 μ M in 10 mM Ba ²⁺ at -60 mV and 0.05Hz) ¹²
Radioligands	(+)-[³ H]isradipine ($K_d < 0.1$ nM) and other dihydropyridines; (–)-[³ H]devapamil ($K_d = 2.5$ nM), (+)- <i>cis</i> -[³ H]diltiazem ($K_d = 50$ nM) ¹¹
Channel distribution	Cardiac muscle, smooth muscle (including blood vessels, intestine, lung, uterus); endocrine cells (including pancreatic β -cells, pituitary); neurones ¹³ ; subcellular localization: concentrated on granule-containing side of pancreatic β -cells ¹⁴ ; neurons (preferentially somatodendritic) ¹⁵
Physiological functions	Excitation-contraction coupling in cardiac or smooth muscle, action potential propagation in sinoatrial and atrioventricular node, synaptic plasticity, hormone (e.g., insulin) secretion ^{10,13,16,17}
Mutations and pathophysiology	Required for normal embryonic development (mouse, zebrafish) ^{18,19} ; de novo G406R mutation in alternative exon 8A in 1 allele causes Timothy syndrome ²⁰
Pharmacological significance	Mediates cardiovascular effects of clinically used Ca ²⁺ antagonists ¹⁷ ; high concentrations of dihydropyridines exert antidepressant effects through Ca _v 1.2 inhibition ¹⁷
Comments	Tissue-specific splice variants exist—in addition to cardiac channels, smooth muscle and brain channels have been cloned ^{7,21,22} ; the gene for Ca _v 1.2 was first isolated and characterized in rabbit heart (2171aa, P15381, X15539)

aa, amino acids; chr., chromosome; HEK, human embryonic kidney.

- Cooper CL, Vandaele S, Barhanin J, Fosset M, Lazdunski M, and Hosey MM (1987) Purification and characterization of the dihydropyridine-sensitive voltage-dependent calcium channel from cardiac tissue. *J Biol Chem* **262**:509–512.
- Reimer D, Huber IG, Garcia ML, Haase H, and Striessnig J (2000) Beta subunit heterogeneity of L-type calcium channels in smooth muscle tissues. *FEBS Lett* **467**:65–69.
- Tsien RW, Hess P, McCleskey EW, and Rosenberg RL (1987) Calcium channels: mechanisms of selectivity, permeation, and block. *Annu Rev Biophys Chem* **16**:265–290.
- Hu H and Marban E (1998) Isoform-specific inhibition of L-type calcium channels by dihydropyridines is independent of isoform-specific gating properties. *Mol Pharmacol* **53**:902–907.
- Koschak A, Reimer D, Huber I, Grabner M, Glossmann H, Engel J, and Striessnig J (2001) α_1D (Cav1.3) subunits can form L-type calcium channels activating at negative voltages. *J Biol Chem* **276**:22100–22106
- Xu W and Lipscombe D (2001) Neuronal Cav1.3 α_1 L-type channels activate at relatively hyperpolarized membrane potentials and are incompletely inhibited by dihydropyridines. *J Neurosci* **21**:5944–5951.
- Tang ZZ, Liang MC, Lu S, Yu D, Yu CY, Yue DT, and Soong TW (2004) Transcript scanning reveals novel and extensive splice variations in human l-type voltage-gated calcium channel, Ca_v1.2 α_1 subunit. *J Biol Chem* **279**:44335–44343.
- Koschak A, Reimer D, Walther D, Hoda JC, Heinzle T, Grabner M, and Striessnig J (2003) Ca_v1.4 α_1 subunits can form slowly inactivating dihydropyridine-sensitive L-type Ca²⁺ channels lacking Ca²⁺-dependent inactivation. *J Neurosci* **23**:6041–6049.
- Peterson BZ, Johnson BD, Hockerman GH, Acheson M, Scheuer T, and Catterall WA (1997) Analysis of the dihydropyridine receptor site of L-type calcium channels by alanine-scanning mutagenesis. *J Biol Chem* **272**:18752–18758.
- Striessnig J (1999) Pharmacology, structure and function of cardiac L-type calcium channels. *Cell Physiol Biochem* **9**:242–269.
- Glossmann H and Striessnig J (1990) Molecular properties of calcium channels. *Rev Physiol Biochem Pharmacol* **114**:1–105.
- Hockerman GH, Johnson BD, Abbott MR, Scheuer T, and Catterall WA (1997) Molecular determinants of high affinity phenylalkylamine block of L-type calcium channels in transmembrane segment IIIS6 and the pore region of the α_1 subunit. *J Biol Chem* **272**:18759–18765.
- Catterall WA (2001) Structure and regulation of voltage-gated calcium channels. *Annu Rev Cell Dev Biol* **16**:521–555.
- Bokvist K, Eliasson L, Åmmälä C, Renstrom E, and Rorsman P (1995) Co-localization of L-type calcium channels and insulin-containing secretory granules and its significance for the initiation of exocytosis in mouse pancreatic B-cells. *EMBO J* **14**:50–57.
- Hell JW, Westenbroek RE, Warner C, Ahljanian MK, Prystay W, Gilbert MM, Snutch TP, and Catterall WA (1993) Identification and differential subcellular localization of the neuronal class C and class D L-type calcium channel α_1 subunits. *J Cell Biol* **123**:949–962.
- Sinnegger-Brauns MJ, Hetzenauer A, Huber IG, Renstrom E, Wietzorrek G, Berjukov S, Cavalli M, Walter D, Koschak A, Waldschutz R, et al. (2004) Isoform-specific regulation of mood behavior and pancreatic β -cell and cardiovascular function by L-type Ca²⁺ channels. *J Clin Invest* **113**:1430–1439.
- Schulla V, Renstrom E, Feil R, Feil S, Franklin I, Gjinovci A, Jing XJ, Laux D, Lundquist I, Magnuson MA, et al. (2003) Impaired insulin secretion and glucose tolerance in β -cell-selective Ca_v1.2 Ca²⁺ channel null mice. *EMBO J* **22**:3844–3854.
- Seisenberger C, Specht V, Welling A, Platzer J, Pfeifer A, Kubbandner S, Striessnig J, Klugbauer N, Feil R, and Hofmann F (2000) Functional embryonic cardio-myocytes after disruption of the L-type αC (Cav1.2) calcium channel gene in the mouse. *J Biol Chem* **275**:39193–39199.
- Rottbauer W, Baker K, Wo ZG, Mohideen MA, Cantiello HF, and Fishman MC (2001) Growth and function of the embryonic heart depend upon the cardiac-specific L-type calcium channel α_1 subunit. *Dev Cell* **1**:265–275.
- CACNA1C; OMIM no. 114205.
- Snutch TP, Tomlinson WJ, Leonard JP, and Gilbert MM (1991) Distinct calcium channels are generated by alternative splicing and are differentially expressed in the mammalian CNS. *Neuron* **7**:45–57.
- Welling A, Ludwig A, Zimmer S, Klugbauer N, Flockerzi V, Hofmann F (1997) Alternatively spliced IS6 segments of the α_1C gene determine the tissue-specific dihydropyridine sensitivity of cardiac and vascular smooth muscle L-type calcium channels. *Circ Res* **81**:526–532.

TABLE 4
Ca_v1.3 channels

Channel name	Ca _v 1.3
Description	Voltage-gated calcium channel α ₁ subunit
Other names	α _{1D} , “neuroendocrine” L-type Ca ²⁺ channel
Molecular information	Human: 2161aa, M76558 (brain; PMID: 1309651); 2181aa, M83566 (pancreatic β-cells; PMID: 1309948); chr. 3p14.3, <i>CACNA1D</i> , LocusID: 776 Rat: 1646aa, M57682 (brain; PMID: 1648940); 2203aa, D38101 (pancreatic β-cells; PMID: 7760845); chr. 16p16, <i>Cacna1d</i> , LocusID: 29716 Mouse: 2144aa, AJ437291 (embryonic heart; PMID: 12900400); chr. 14, <i>Cacna1d</i> , LocusID: 12289 (see “Comments”)
Associated subunits	Most likely at least α ₂ , β, and δ subunits
Functional assays	Patch-clamp (whole-cell, single-channel), calcium imaging
Current	I _{Ca,L}
Conductance	Not established
Ion selectivity	Not established
Activation	V _a = −15 to −20 mV (mouse cochlear hair cells; 10 mM Ba ²⁺) ^{1,2} ; −18 mV (in 15 mM Ba ²⁺ ; HEK cells) to −37 mV (5 mM Ba ²⁺ ; 2 mM Ca ²⁺ HEK cells or <i>Xenopus</i> oocytes) ^{3,4} ; τ _a < 1 ms at +10 mV ³
Inactivation	V _h = −36 to −43 mV ^{3,5} ; τ _{fast} = 190 ms, τ _{slow} = 1700 ms (at V _{max} in HEK cells) ³ ; calcium-induced inactivation is observed after expression in HEK cells ³ and in cochlear outer hair cells but not in inner hair cells ²
Activators	BayK8644 ^{1–5}
Gating modifiers	Dihydropyridine antagonists (e.g., isradipine, IC ₅₀ = 30 nM at −50 mV and 300 nM at −90 mV; nimodipine, IC ₅₀ = 3 μM at −80 mV) ^{3,4}
Blockers	Nonselective: Cd ²⁺ ⁵
Radioligands	(+)-[³ H]isradipine (K _d < 0.5 nM) ³ ; in radioreceptor assays, HEK cell-expressed Ca _v 1.2 and Ca _v 1.3 channels bind (+)-[³ H]isradipine with indistinguishable K _D ³ ; in functional experiments, however, Ca _v 1.2 channels show higher DHP sensitivity—this discrepancy is explained by the slower inactivation of Ca _v 1.3 decreasing the availability of inactivated channels for state-dependent DHP block
Channel distribution	Sensory cells (photoreceptors, cochlear hair cells ^{1,2}), endocrine cells (including pancreatic β-cells, pituitary, adrenal chromaffin cells, pinealocytes), ^{7–9} low density in heart (atrial muscle, sinoatrial and atrioventricular node) ^{1,7,10} and vascular smooth muscle ⁷ ; neurones ⁶ ; subcellular localization: on neurones preferentially located on proximal dendrites and cell bodies ⁶
Physiological functions	Neurotransmitter release in sensory cells, control of cardiac rhythm and atrioventricular node conductance at rest, ^{1,10,12} mood behavior, ¹² hormone secretion
Mutations and pathophysiology	Deafness, sinoatrial and atrioventricular node dysfunction, ^{1,10,12} no convincing evidence for contribution to pancreatic β-cell L-type currents and insulin secretion in mouse models ^{1,12,13}
Pharmacological significance	Hypothetical drug targets for modulators of heart rate, ¹ antidepressant drugs ¹⁰ and drugs for hearing disorders ¹
Comments	Tissue-specific and developmental (exon 1b) splice variants exist—in addition to brain, pancreatic β-cell and cochlear variants have been cloned; it is likely that Ca _v 1.3 channels form most of the so-called ‘low-voltage-activated’ L-type currents found in the brain and sinoatrial node, although some splice variants of Ca _v 1.2 can also activate at more negative potentials

aa, amino acids; chr., chromosome; HEK, human embryonic kidney; DHP, dihydropyridine.

1. Platzer J, Engel J, Schrott-Fischer A, Stephan K, Bova S, Chen H, Zheng H, and Striessnig J (2000) Congenital deafness and sinoatrial node dysfunction in mice lacking class D L-type calcium channels. *Cell* **102**:89–97.

2. Michna M, Knirsch M, Hoda JC, Muenkner S, Langer P, Platzer J, Striessnig J, and Engel J (2003) Cav1.3 (α_{1D}) Ca²⁺ currents in neonatal outer hair cells of mice. *J Physiol* **553**:747–758.

3. Koschak A, Reimer D, Huber I, Grabner M, Glossmann H, Engel J, and Striessnig J (2001) α_{1D} (Ca_v1.3) subunits can form L-type calcium channels activating at negative voltages. *J Biol Chem* **276**:22100–22106.

4. Xu W and Lipscombe D (2001) Neuronal Ca_v1.3α₁ L-type channels activate at relatively hyperpolarized membrane potentials and are incompletely inhibited by dihydropyridines. *J Neurosci* **21**:5944–5951.

5. Scholze A, Plant TD, Dolphin AC, and Nürnberg B (2001) Functional expression and characterization of a voltage-gated Ca_v1.3 (α_{1D}) calcium channel subunit from an insulin-secreting cell line. *Mol Endocrinol* **15**:1211–1221.

6. Hell JW, Westenbroek RE, Warner C, Ahljianian MK, Prystay W, Gilbert MM, Snutch TP, and Catterall WA (1993) Identification and differential subcellular localization of the neuronal class C and class D L-type calcium channel α₁ subunits. *J Cell Biol* **123**:949–962.

7. Takimoto K, Li D, Nerbonne JM, and Levitan ES (1997) Distribution, splicing and glucocorticoid-induced expression of cardiac α_{1C} and α_{1D} voltage-gated calcium channel mRNAs. *J Mol Cell Cardiol* **29**:3035–3042.

8. Garcia-Palmero E, Renart J, Andres-Mateos E, Solis-Garrido LM, Matute C, Herrero CJ, Garcia AG, and Montiel C (2001) Differential expression of calcium channel subtypes in the bovine adrenal medulla. *Neuroendocrinology* **74**:251–261.

9. Chik CL, Liu QY, Li B, Klein DC, Zylka M, Kim DS, Chin H, Karpinski E, and Ho AK (1997) Alpha 1D L-type Ca²⁺-channel currents: inhibition by a beta-adrenergic agonist and pituitary adenylate cyclase-activating polypeptide (PACAP) in rat pinealocytes. *J Neurochem* **68**:1078–1087.

10. Mangoni ME, Couette B, Bourinet E, Platzer J, Reimer D, Striessnig J, and Nargeot J (2003) Functional role of L-type Cav1.3 Ca²⁺ channels in cardiac pacemaker activity. *Proc Natl Acad Sci USA* **100**:5543–5548.

11. Namkung Y, Skrypnik N, Jeong MJ, Lee T, Lee MS, Kim HL, Chin H, Suh PG, Kim SS, and Shin HS (2001) Requirement for the L-type calcium channel α_{1D} subunit in postnatal pancreatic beta cell generation. *J Clin Invest* **108**:1015–1022.

12. Sinnegger-Brauns MJ, Hetzenauer A, Huber IG, Renstrom E, Wietzorrek G, Berjukov S, Cavalli M, Walter D, Koschak A, Waldschutz R, et al. (2004) Isoform-specific regulation of mood behavior and pancreatic β-cell and cardiovascular function by L-type Ca²⁺ channels. *J Clin Invest* **113**:1430–1439.

13. Schulla V, Renstrom E, Feil R, Feil S, Franklin I, Gjinovci A, Jing XJ, Laux D, Lundquist I, Magnuson MA, et al. (2003) Impaired insulin secretion and glucose tolerance in beta cell-selective Ca_v1.2 Ca²⁺ channel null mice. *EMBO J* **22**:3844–3854.

TABLE 5
Ca_v1.4 channels

Channel name	Ca _v 1.4
Description	Voltage-gated calcium channel α_1 subunit
Other names	α_{1F}
Molecular information	Human: 1966aa, AJ224874 (PMID: 9662399); chr. Xp11.23, <i>CACNA1F</i> , LocusID: 778 Rat: 1981aa, AF365975 (PMID: 11526344); chr. Xq22, <i>Cacna1f</i> , LocusID: 114493 Mouse: 1985aa, AF192497 (PMID: 10873387); chr. X, <i>Cacna1f</i> , LocusID: 54652
Associated subunits	Not established; preliminary functional evidence for β_2 association in retinal neurons ¹
Functional assays	Patch-clamp (whole-cell, single-channel), calcium imaging
Current	$I_{Ca,L}$
Conductance	Preliminary evidence for very small single channel conductance (less than half of Ca _v 1.2); Ba ²⁺ > Ca ²⁺ ^{2,4,6}
Ion selectivity	Not established
Activation	$V_a = -2.5$ to -12 mV (2–20 mM Ca ²⁺ or 15–20 mM Ba ²⁺ ; HEK cells) ^{3–6} ; $\tau_a < 1$ ms at V_{max} (but slower components were also observed) ^{3,6}
Inactivation	$V_h = -9$ to -27 mV (10–20 mM Ba ²⁺ , HEK cells) ^{4,6} ; inactivation kinetics even slower than those of Ca _v 1.3 with incomplete inactivation during 10-s depolarizations to V_{max} ³ ; calcium-induced inactivation is not observed for Ca _v 1.4 channels expressed in HEK cells ^{3,4,6} but after expression in <i>Xenopus oocytes</i> ²
Activators	BayK8644 ^{2–4,6}
Gating modifiers	Dihydropyridine antagonists: nifedipine ($IC_{50} = 944$ nM at -100 mV, ~ 300 nM at -50 mV ⁴ ; isradipine: $\sim 80\%$ inhibition by 100 nM at -50 mV ^{3,6} and 1 μ M at -90 mV ³ ; D- <i>cis</i> -diltiazem ($IC_{50} = 92$ μ M); verapamil: 69% inhibition at 100 μ M (0.2 Hz, holding potential = -80 mV) ⁶
Blockers	Nonselective: Cd ²⁺ ²
Radioligands	Unlike for Ca _v 1.2 and Ca _v 1.3, no high-affinity (+)-[³ H]isradipine binding detectable (HEK cells) (J. Striessnig, unpublished observations)
Channel distribution	Retinal photoreceptors and bipolar cells, spinal cord, lymphoid tissue (plasma and mast cells) ^{1,4,7–10}
Physiological functions	Neurotransmitter release in retinal cells
Mutations and pathophysiology	Mutations cause X-linked congenital stationary night blindness type 2 ^{7,9,11,12}
Pharmacological significance	Not established
Comments	The biophysical properties of heterologously expressed Ca _v 1.4 channels resemble those recorded in retinal neurons, suggesting that this channel type underlies retinal $I_{Ca,L}$ —however, similar to Ca _v 1.4, Ca _v 1.3 channels also inactivate slowly and activate rapidly and may therefore also contribute to retinal $I_{Ca,L}$

aa, amino acids; chr., chromosome; HEK, human embryonic kidney.

1. Ball SL, Powers PA, Shin HS, Morgans CW, Peachey NS, and Gregg RG (2002) Role of the β_2 subunit of voltage-dependent calcium channels in the retinal outer plexiform layer. *Investig Ophthalmol Vis Sci* **43**:1595–1603.
2. Hoda JC, Zaghetto F, Koschak A, and Striessnig J (2005) CSNB2 mutations S229P, G369D, L1068P, and W1440X alter channel gating or functional expression of Ca_v1.4 L-type Ca²⁺ channels. *J Biol Chem* **25**:252–259.
3. Koschak A, Reimer D, Walter D, Hoda JC, Heinzele T, Grabner M, and Striessnig J (2003) Ca_v1.4 α_1 subunits can form slowly inactivating dihydropyridine-sensitive L-type Ca²⁺ channels lacking Ca²⁺-dependent inactivation. *J Biol Chem* **23**:6041–6049.
4. McRory JE, Hamid J, Doering CJ, Garcia E, Parker R, Hamming K, Chen L, Hildebrand M, Beedle AM, Feldcamp L, et al. (2004) The CACNA1F gene encodes an L-type calcium channel with unique biophysical properties and tissue distribution. *J Biol Chem* **24**:1707–1718.
5. Haeseleer F, Imanishi Y, Maeda T, Possin DE, Maeda A, Lee A, Rieke F, and Palczewski K (2004) Essential role of Ca²⁺-binding protein 4, a Ca_v1.4 channel regulator, in photoreceptor synaptic function. *Nat Neurosci* **7**:1079–1087.
6. Baumann L, Gerstner A, Zong X, Biel M, and Wahl-Schott C (2004) Functional characterization of the L-type Ca²⁺ channel Ca_v1.4 α_1 from mouse retina. *Investig Ophthalmol Vis Sci* **45**:708–713.
7. Bech-Hansen NT, Naylor MJ, Maybaum TA, Pearce WG, Koop B, Fishman GA, Mets M, Musarella MA, and Boycott KM (1998) Loss-of-function mutations in a calcium-channel α_1 -subunit gene in Xp11.23 cause incomplete X-linked congenital stationary night blindness. *Nat Genet* **19**:264–267.
8. Naylor MJ, Rancourt DE, and Bech-Hansen NT (2000) Isolation and characterization of a calcium channel gene, *Cacna1f*, the murine orthologue of the gene for incomplete X-linked congenital stationary night blindness. *Genomics* **66**:324–327.
9. Strom TM, Nyakatura G, Apfelstedt-Sylla E, Hellebrand H, Lorenz B, Weber BH, Wutz K, Gutwillinger N, Ruther K, Drescher B, et al. (1998) An L-type calcium-channel gene mutated in incomplete X-linked congenital stationary night blindness. *Nat Genet* **19**:260–263.
10. Firth SI, Morgan IG, Boelen MK, and Morgans CW (2001) Localization of voltage-sensitive L-type calcium channels in the chicken retina. *Clin Experiment Ophthalmol* **29**:183–187.
11. CACNA1F; OMIM no. 300110.
12. Striessnig J, Hoda JC, Koschak A, Zaghetto F, Mullner C, Sinnegger-Brauns MJ, Wild C, Watschinger K, Trockenbacher A, and Pelster G (2004) L-type Ca²⁺ channels in Ca²⁺ channelopathies. *Biochem Biophys Res Commun* **322**:1341–1346.

TABLE 6
Ca_v2.1 channels

Channel name	Ca _v 2.1
Description	Voltage-gated calcium channel α_1 subunit
Other names	α_{1A} , P-type, Q-type, rbA-I (in rat) ¹ ; BI-1, BI-2 (in rabbit) ²
Molecular information	Human: 2510aa, AF004883, 2662aa, AF004884, chr. 19p13, CACNA1A Rat: 2212aa, M64373 Mouse: 2165aa, NM007578, NP031604 Rabbit: 2273aa, X57476 (see "Comments")
Associated subunits	$\alpha_2\delta$, β , possibly γ
Functional assays	Voltage-clamp, patch-clamp, calcium imaging, neurotransmitter release
Current	I _{Ca,P} , I _{Ca,Q}
Conductance	9, 14, 19pS (P-type, cerebellar Purkinje neurones) ⁴ ; 16–17pS (for $\alpha_{1A}/\alpha_2\delta/\beta$ in <i>Xenopus</i> oocytes) ^{2,5,6}
Ion selectivity	Ba ²⁺ > Ca ²⁺
Activation	V _a = -5 mV for native P-type, V _a = -11 mV for native Q-type (with 5 mM Ba ²⁺ charge carrier) ⁷ V _a = -4.1 mV for rat $\alpha_{1A-a}/\alpha_2\delta/\beta_4$ V _a = +2.1 mV for rat $\alpha_{1A-b}/\alpha_2\delta/\beta_4$ (with 5 mM Ba ²⁺ charge carrier) ⁶ V _a = +9.5 mV; τ_a = 2.2 ms at +10 mV for human $\alpha_{1A-1}/\alpha_2\delta/\beta_{1b}$ in HEK293 cells (with 15 mM Ba ²⁺ charge carrier) ³
Inactivation	V _h = -17.2 mV for $\alpha_{1A-a}/\alpha_2\delta/\beta_4$; V _h = -1.6 mV for $\alpha_{1A-b}/\alpha_2\delta/\beta_4$ (with 5 mM Ba ²⁺ charge carrier); V _h = -17 mV, τ_h = 690 ms at +10 mV human $\alpha_{1A-1}/\alpha_2\delta/\beta_{1b}$ in HEK293 cells (with 15 mM Ba ²⁺ charge carrier) ³ ; τ_h > 1 s at 0 mV native P-type (with 5 mM Ba ²⁺ charge carrier) ⁷ (see "Comments")
Activators	None
Gating modifiers	ω -agatoxin IVA (P-type K _d = 1–3 nM ⁸ ; Q-type K _d ~ 100–200 nM ^{5,9}), ω -agatoxin IVB ⁶
Blockers	ω -conotoxin MVIIC ⁸ ; other blockers include piperidines, substituted diphenylbutylpiperidines, piperazines, volatile anesthetics, gabapentin, mibefradil, and peptide toxins DW13.3 and ω -conotoxin SVIB ^{21–26} (see "Comments")
Radioligands	[¹²⁵ I] ω -conotoxin MVIIC
Channel distribution	Neurons (presynaptic terminals, dendrites, some cell bodies), heart, pancreas, pituitary
Physiological functions	Neurotransmitter release in central neurons and neuromuscular junction; excitation-secretion coupling in pancreatic β -cells
Mutations and pathophysiology	Missense mutations in IS4-IS5, IIS4-IIS6, IIS4-IIIS6, and IVS4-IVS6 cause FHM ²⁷ ; a common feature among FHM mutations is an apparent gain-of-function phenotype as a result of a shift in V _{50act} to more hyperpolarized potentials (an increased probability of opening at the single channel level) ^{28,29} ; other effects include a decrease in maximal current density at the whole-cell level and alterations of synaptic transmission ^{28–31} ; point mutations in IIS1, IIS6-IIIS2, IIIS5-IIIS6, and IVS1-IVS5 cause episodic ataxia type-2, a polyglutamine expansion in the carboxyl region causes spinocerebellar ataxia type-6, and mutation of IS5-IS6 and IVS6 causes episodic and progressive ataxia ^{10–12,27}
Pharmacological significance	Peptide toxins that selectively inhibit Ca _v 2.1 channel block a significant portion of neurotransmission in the mammalian CNS ¹³ ; block of Ca _v 2.1 channels inhibits the late-phase formalin response and inflammatory pain but has no significant effect on mechanical allodynia or thermal hyperalgesia ^{14–17} ; mice lacking a functional Ca _v 2.1 gene exhibit cerebellar atrophy, severe muscle spasms, and ataxia and usually die by 3 to 4 weeks postnatal ^{18,19}
Comments	Rates of inactivation and V _h are differentially affected by coexpression with β_{1b} , β_{2a} , β_3 , or β_4 subunits, as well as by alternative splicing of the α_{1A} subunit; identified regions of alternative splicing include the domain I-II linker, domain II-III linker, IVS3-IVS4, and the carboxyl terminus ^{1,2,6,32–34} ; whole-cell currents with P-type kinetics seem to be conducted by the α_{1A-b} splice variant coexpressed with any of the β subunits or by the α_{1A-a} splice variant coexpressed with the β_{2a} subunit ^{6,7,20} ; whole-cell currents with Q-type kinetics seem to be encoded by α_{1A-a} coexpressed with any of the β_{1b} , β_3 , or β_4 subunits ^{6,20} ; whole-cell currents with Q-type pharmacology seem to be encoded by α_{1A} splice variants containing Asp Pro residues in the domain IV S3-S4 linker, whereas whole-cell currents with P-type pharmacology seem to be conducted by α_{1A} splice variants missing Asp Pro residues in IV S3-S4 linker ^{3,6} ; alternative splicing also alters current density, current-voltage relations, calcium/calmodulin-dependent facilitation, sensitivity to mibefradil, and binding to intracellular synaptic proteins such as Mint1, CASK, syntaxin, and SNAP-25 ^{26,32,36}

aa, amino acids; chr., chromosome; HEK, human embryonic kidney; FHM, familial hemiplegic migraine; CNS, central nervous system.

1. Starr TVB, Prystay W, and Snutch TP (1991) Primary structure of a calcium channel that is highly expressed in the rat cerebellum. *Proc Natl Acad Sci USA* **88**:5621–5625.

2. Mori Y, Friedrich T, Kim MS, Mikami A, Nakai J, Ruth P, Bosse E, Hofmann F, Flockerzi V, Furuichi T, et al. (1991) Primary structure and functional expression from complementary DNA of a brain calcium channel. *Nature (Lond)* **350**:398–402.

3. Hans M, Urrutia A, Deal C, Brust PF, Stauderman K, Ellis SB, Harpold M, Johnson EC, and Williams ME (1999) Structural elements in domain IV that influence biophysical and pharmacological properties of human α_{1A} -containing high-voltage-activated calcium channels. *Biophys J* **76**:1384–1400.

4. Llinas R, Sugimori M, Hillman DE, and Cherksey B (1992) Distribution and functional significance of the P-type voltage-dependent calcium channels in the mammalian central nervous system. *Trends Neurosci* **15**:2995–3012.

5. Sather WA, Tanabe T, Zhang JF, Mori Y, Adams ME, and Tsien RW (1993) Distinctive biophysical and pharmacological properties of class A (BI) calcium channel α_1 subunits. *Neuron* **11**:291–303.

6. Bourinet E, Soong TW, Sutton K, Slaymaker S, Mathews E, Montell A, Zamponi GW, Nargeot J, and Snutch TP (1999) Splicing of α_{1A} subunit gene generates phenotypic variants of P- and Q-type calcium channels. *Nat Neurosci* **2**:407–415.

7. Merlestein PG, Foehring RC, Tkatch T, Song WJ, Baranauskas G, and Surmeier JD (1999) Properties of Q-type calcium channels in neostriatal and cortical neurons are correlated with β subunit expression. *J Biol Chem* **19**:7268–7277.
8. Mintz IM, Venema VJ, Swiderek K, Lee T, Bean BP, and Adams ME (1992) P-type calcium channels blocked by the spider toxin ω -Aga-IVA. *Nature (Lond)* **355**:827–829.
9. Randall A and Tsien RW (1995) Pharmacological dissection of multiple types of calcium channel currents in rat cerebellar granule neurons. *J Biol Chem* **15**:2995–3012.
10. Ducros A, Denier C, Joutel A, Vahedi K, Michel A, Darcel F, Madigand M, Guerouaou D, Tison F, Julien J, et al. (1999) Recurrence of the T666M calcium channel CACNA1A gene mutation in familial hemiplegic migraine with progressive cerebellar ataxia. *Am J Hum Genet* **64**:89–98.
11. Kraus RL, Sinnegger MJ, Koschak A, Glossmann H, Stenirri S, Carrera P, and Striessnig J (2000) Three new familial hemiplegic migraine mutants affect P/Q-type calcium channel kinetics. *J Biol Chem* **275**:9239–9243.
12. Ophoff RA, Terwindt GM, Vergouwe MN, van Eijk R, Oefner PJ, Hoffman SM, Lamerdin JE, Mohrenweiser HW, Bulman DE, Ferrari M, et al. (1996) Familial hemiplegic migraine and episodic ataxia type-2 are caused by mutations in the calcium channel gene CACNL1A4. *Cell* **87**:543–552.
13. Dunlap K, Luebke JI, and Turner TJ (1995) Exocytotic Ca^{2+} channels in mammalian central neurons. *Trends Neurosci* **18**:89–98.
14. Chaplan SR, Pogrel JW, and Yaksh TL (1994) Role of voltage-dependent calcium channel subtypes in experimental tactile allodynia. *J Pharmacol Exp Ther* **269**:1117–1123.
15. Malmberg AB and Yaksh TL (1994) Voltage-sensitive calcium channels in spinal nociceptive processing: blockade of N- and P-type channels inhibits formalin-induced nociception. *J Biol Chem* **14**:4882–4890.
16. Sluka KA (1997) Blockade of calcium channels can prevent the onset of secondary hyperalgesia and allodynia induced by intradermal injection of capsaicin in rats. *Pain* **71**:157–164.
17. Sluka KA (1998) Blockade of N- and P/Q-type calcium channels reduces the secondary heat hyperalgesia induced by acute inflammation. *J Pharmacol Exp Ther* **287**:232–237.
18. Jun K, Piedras-Rentera E, Smith SM, Wheeler DB, Lee SB, Lee TG, Chin H, Adams ME, Scheller RH, Tsien RW, and Shin HS (2000) Ablation of P/Q type calcium channel currents, altered synaptic transmission and progressive ataxia in mice lacking the α_{1A} subunit. *Proc Natl Acad Sci USA* **96**:15245–15250.
19. Fletcher CF, Tottene A, Lennon VA, Wilson SM, Dubel SJ, Paylor R, Hosford DA, Tessarollo L, McEnery MW, Pietrobon D, et al. (2001) Dystonia and cerebellar atrophy in *Caen14* null mice lacking P/Q calcium channel activity. *FASEB J* **15**:1288–1290.
20. Stea A, Tomlinson WJ, Soong TW, Bourinet E, Dubel SJ, Vincent SR, and Snutch TP (1994) Localization and functional properties of a rat brain α_{1A} calcium channel reflect similarities to neuronal Q- and P-type channels. *Proc Natl Acad Sci USA* **91**:10567–10580.
21. Sah DW and Bean BP (1994) Inhibition of P-type and N-type calcium channels by dopamine receptor antagonists. *Mol Pharmacol* **45**:84–92.
22. Sutton K, Siok C, Stea A, Zamponi G, Heck SD, Volkman RA, Ahljiyanian MK, and Snutch TP (1998) Inhibition of neuronal calcium channels by a novel peptide spider toxin, DW13.3. *Mol Pharmacol* **54**:407–418.
23. Oka M, Itoh Y, Wada M, Yamamoto A, and Fujita T (2003) Gabapentin blocks L-type and P/Q-type Ca^{2+} channels involved in depolarization-stimulated nitric oxide synthase activity in primary cultures of neurons from mouse cerebral cortex. *Pharm Res* **20**:897–899.
24. Dooley DJ, Donovan CM, Meder WP, and Whetzel SZ (2002) Preferential action of gabapentin and pregabalin at P/Q-type voltage-sensitive calcium channels: inhibition of K^{+} -evoked [^3H]-norepinephrine release from rat neocortical slices. *Synapse* **45**:171–190.
25. Kamatchi GL, Chan CK, Snutch T, Durieux, and Lynch III C (1999) Volatile anesthetic inhibition of neuronal Ca channel currents expressed in *Xenopus* oocytes. *Brain Res* **831**:85–96.
26. Jimenez C, Bourinet E, Leuranguer V, Richard S, Snutch TP, and Nargeot J (2000) Determinants of voltage-dependent inactivation affect Mibefradil block of calcium channels. *Neuropharmacology* **39**:1–10.
27. Lorenzon NM and Beam K (2005) Calcium channelopathies, in *Voltage-Gated Calcium Channels* (Zamponi G ed) pp 240–261, Kluwer Academic/Plenum Publishers.
28. Tottene A, Fellin T, Pagnutti S, Luvisetto S, Striessnig J, Fletcher C, and Pietrobon D (2002) Familial hemiplegic migraine mutations increase Ca^{2+} influx through single human $\text{Ca}_v2.1$ channels and decrease maximal $\text{Ca}_v2.1$ current density in neurons. *Proc Natl Acad Sci USA* **99**:13284–13289.
29. van den Maagdenberg AM, Pietrobon D, Pizzorusso T, Kaja S, Broos LA, Cesetti T, van de Ven RC, Tottene A, van der Kaa J, Plomp JJ, et al. (2004) A *Caen1a* knock in migraine mouse model with increased susceptibility to cortical spreading depression. *Neuron* **41**:701–710.
30. Cao YQ, Piedras-Renteria ES, Smith GB, Chen G, Harata NC, and Tsien RW (2004) Presynaptic Ca^{2+} channels compete for channel type-preferring slots in altered neurotransmission arising from Ca^{2+} channelopathy. *Neuron* **43**:387–400.
31. Cao Y and Tsien R (2005) Effects of familial hemiplegic migraine type 1 mutations on neuronal P/Q-type Ca^{2+} channel activity and inhibitory synaptic transmission. *Proc Natl Acad Sci USA* **102**:2590–2595.
32. Soong TW, DeMaria CD, Alvania RS, Zweifel LS, Liang MC, Mittman S, Agnew W, and Yue DT (2002) Systematic identification of splice variants in human P/Q-type channel α_{12} subunits: implications for current density and Ca^{2+} -dependent inactivation. *J Neurosci* **22**:10142–10152.
33. Krovetz HS, Helton TD, Crews AL and Horne WA (2000) C-Terminal alternative splicing changes the gating properties of a human spinal cord calcium channel alpha 1A subunit. *J Neurosci* **20**:7564–7570.
34. Chaudhuri D, Chang SY, DeMaria CD, Alvania RS, Soong TW, and Yue DT (2004) Alternative splicing as a molecular switch for Ca^{2+} /calmodulin-dependent facilitation of P/Q-type Ca^{2+} channels. *J Neurosci* **24**:6334–6342.
35. Rettig J, Sheng ZH, Kim DK, Hodson CD, Snutch TP, and Catterall WA (1996) Isoform-specific interaction of the α_{1A} subunits of brain Ca^{2+} channels with the presynaptic proteins syntaxin and SNAP-25. *Proc Natl Acad Sci USA* **93**:7363–7368.
36. Maximov A, Sudhof TC, and Bezprozvany I (1999) Association of neuronal calcium channels with modular adaptor proteins. *J Biol Chem* **274**:24453–24456.

TABLE 7
Ca_v2.2 channels

Channel name	Ca _v 2.2
Description	Voltage-gated calcium channel α_1 subunit
Other names	N-type, α_{1B} ; rbB-I, rbB-II (in rat), ^{1,2} BIII (in rabbit) ³
Molecular information	Human: 2339aa, M94172, 2237aa, M94173, ⁴ chr. 9q34, <i>CACN1B</i> Rat: 2336aa, M92905 ¹ Mouse: 2329aa, NM007579, NP031605
Associated subunits	$\alpha_2\delta/\beta_1$, β_3 , β_4 , ⁵ possibly γ
Functional assays	Voltage-clamp, patch-clamp, calcium imaging, neurotransmitter release, ⁴⁵ Ca uptake into synaptosomes
Current	$I_{Ca,N}$
Conductance	20pS (bullfrog sympathetic neurones) ⁶ ; 14.3pS (rabbit BIII cDNA in skeletal muscle myotubes) ³
Ion selectivity	Ba ²⁺ > Ca ²⁺
Activation	$V_a = +7.8$ mV, $\tau_a = 3$ ms at +10 mV (human $\alpha_{1B}/\alpha_2\delta/\beta_{1-3}$ in HEK293 cells, 15 mM Ba ²⁺ charge carrier) ^{4,7} ; $V_a = +9.7$ mV, $\tau_a = 2.8$ ms at +20 mV (rat α_{1B-I}/β_{1b} , in <i>Xenopus</i> oocytes, 40 mM Ba ²⁺ charge carrier) ²
Inactivation	$V_h = -61$ mV, $\tau_h \sim 200$ ms at +10 mV (human $\alpha_{1B}/\alpha_2\delta/\beta_{1-3}$ in HEK293 cells, 15 mM Ba ²⁺ charge carrier) ^{4,7} ; $V_h = -67.5$ mV; $\tau_h = 112$ ms at +20 mV (rat α_{1B-I}/β_{1b} in <i>Xenopus</i> oocytes, 40 mM Ba ²⁺) ²
Activators	None
Gating modifiers	None
Blockers	ω -conotoxin GVIA (1–2 μ M, irreversible block), ω -conotoxin MVIIA (SNX-111, Ziconotide/Prialt), ω -conotoxin MVIIC ⁸ ; other blockers include piperidines, substituted diphenylbutylpiperidines, long alkyl chain molecules, aliphatic monoamines, tetrandine, gabapentin, peptidylamines, volatile anesthetics, the peptide toxins SNX-325 and DW13.3, as well as the ω -conotoxins SVIA, SVIB, and CVID ^{20–34}
Radioligands	[¹²⁵ I] ω -conotoxin GVIA ($K_d = 55$ pM, human $\alpha_{1B}/\alpha_2\delta/\beta_{1-3}$ in HEK293 cells) ⁴
Channel distribution	Neurons (presynaptic terminals, dendrites, cell bodies) ⁹
Physiological functions	Neurotransmitter release in central and sympathetic neurons ¹⁰ ; sympathetic regulation of the circulatory system ^{11,35} ; activity and vigilance state control ³⁶ ; sensation and transmission of pain (see “Pharmacological significance” and “Comments”)
Mutations and pathophysiology	Differing reports exist: mice lacking a functional Ca _v 2.2 gene exhibit a normal life span and no detectable behavioral modifications compared with wild type but possess an increase in basal mean atrial pressure and other functional alterations to the sympathetic nervous system ¹¹ —however, in a different study, approximately 1/3 of the mice lacking a functional Ca _v 2.2 gene did not survive to weaning, but surviving animals were normal except for a decrease in anxiety-related behavior and a suppression of inflammatory and neuropathic pain responses ¹² ; no point mutations in the native Ca _v 2.2. gene have been reported to date
Pharmacological significance	In rats, intrathecal administration of ω -conotoxin GVIA or ω -conotoxin MVIIA shows strong effects on inflammatory pain, postsurgical pain, thermal hyperalgesia, and mechanical allodynia ^{13–15} ; in humans, intrathecal administration of SNX-111 (Ziconotide/Prialt, synthetic ω -conotoxin MVIIA) to patients unresponsive to intrathecal opiates significantly reduced pain scores and in a number of specific instances resulted in relief after many years of continuous pain ¹⁶
Comments	In case studies, Ziconotide/Prialt has been examined for usefulness in the management of intractable spasticity following spinal cord injury in patients unresponsive to baclofen and morphine ¹⁷ ; side effects of intrathecal administration of Ziconotide/Prialt include nystagmus, sedation, confusion, auditory and visual hallucinations, severe agitation, and unruly behavior ¹⁸ ; intravenous administration of Ziconotide to humans results in significant orthostatic hypotension ¹⁹ ; identified regions of alternative splicing include the domain I-II linker, domain II-III linker, IIIS3-IIIS4, IVS3-IVS4, and the carboxyl terminus ^{1–4,37–39} ; splicing affects a number of channel properties, including current-voltage relations and kinetics, and is associated with cell-specific expression—in particular, expression of the e37a splice isoform in dorsal root ganglia correlates with a subset of nociceptive neurons ^{40–42} ; alternative splicing also alters interactions with intracellular synaptic proteins such as Mint1, CASK, syntaxin, and SNAP-25 ^{43–45}

aa, amino acid; chr., chromosome; HEK, human embryonic kidney.

- Dubel SJ, Starr TV, Hell J, Ahljanian MK, Eneart JJ, Catterall WA, and Snutch TP (1992) Molecular cloning of the α_1 subunit of an ω -conotoxin-sensitive calcium channel. *Proc Natl Acad Sci USA* **89**:5058–5062.
- Stea A, Dubel SJ, and Snutch TP (1999) α_{1B} N-type calcium channel isoforms with distinct biophysical properties. *Ann NY Acad Sci* **868**:118–130.
- Fujita Y, Mynlieff M, Dirksen RT, Kim M, Niidome T, Nakai J, Friedrich T, Iwabe N, Miyata T, Furuichi T, et al. (1993) Primary structure and functional expression of the ω -conotoxin-sensitive N-type calcium channel from rabbit brain. *Neuron* **10**:585–598.
- Williams ME, Brust PF, Feldman DH, Patthi S, Simerson S, Maroufi A, McCue AF, Velicelebi G, Ellis SB, and Harpold M (1992) Structure and functional expression of an ω -conotoxin-sensitive human N-type calcium channel. *Science* **257**:389–395.
- Scott VE, De Waard M, Liu H, Gurnett CA, Venzke DP, Lennon VA, and Campbell KP (1996) β subunit heterogeneity in N-type calcium channels. *J Biol Chem* **271**:3207–3212.
- Elmslie KS (1997) Identification of the single channels that underlie the N-type and L-type calcium currents in bullfrog sympathetic neurons. *J Neurosci* **17**:2658–2668.
- Williams ME, Marubio LM, Deal CR, Hans M, Brust PF, Philipson L, Miller RJ, Johnson EC, Harpold MM, and Ellis SB (1994) Structure and functional characterization of neuronal α_{1B} calcium channel subtypes. *J Biol Chem* **269**:22347–22357.
- Hillyard DR, Monje VD, Mintz IM, Bean BP, Nadasdi L, Ramachandran J, Miljanich G, Azimi-Zoonooz A, McIntosh JM, Cruz LJ, et al. (1992) A new conus peptide ligand for mammalian presynaptic calcium channels. *Neuron* **9**:9–77.

9. Westenbroek RE, Hell JW, Warner C, Dubel SJ, Snutch TP, and Catterall W (1992) Biochemical properties and subcellular distribution of an N-type calcium channel α_1 subunit. *Neuron* **6**:1099–1115.
10. Dunlap K, Luebke JI, and Turner TJ (1995) Exocytotic calcium channels in mammalian central neurons. *Trends Neurosci* **18**:9–18.
11. Ino M, Yoshinaga T, Wakamori M, Miyamoto N, Takahashi E, Sonoda J, Kagaya T, Oki T, Nagasu T, Nishizawa Y, et al. (2001) Functional disorders of the sympathetic nervous system in mice lacking the α_{1B} subunit (Ca_v2.2) of N-type calcium channels. *Proc Natl Acad Sci USA* **98**:323–328.
12. Saegusa H, Kurihara T, Zong S, Kazuno A, Matsuda Y, Nonaka T, Han W, Toriyama H, and Tanabe T (2001) Suppression of inflammatory and neuropathic pain symptoms in mice lacking the N-type calcium channel. *EMBO J* **20**:2349–2356.
13. Malmberg AB and Yaksh TL (1994) Voltage-sensitive calcium channels in spinal nociceptive processing: blockade of N- and P-type channels inhibits formalin-induced nociception. *J Neurosci* **14**:4882–4890.
14. Bowersox SS, Gadbois T, Singh T, Pettus M, Wang Y-X, and Luther RR (1996) Selective N-type neuronal voltage-sensitive calcium channel blocker, SNX-111, produces spinal antinociception in rat models of acute, persistent and neuropathic pain. *J Pharmacol Exp Ther* **279**:1243–1249.
15. Sluka KA (1998) Blockade of N- and P/Q-type calcium channels reduces the secondary heat hyperalgesia induced by acute inflammation. *J Pharmacol Exp Ther* **287**:232–237.
16. Vanegas H and Schaible HG (2000) Effects of antagonists to high-threshold calcium channels upon spinal mechanisms of pain, hyperalgesia and allodynia. *Pain* **85**:9–18.
17. Ridgeway B, Wallace M, and Gerayli A (2000) Ziconotide for the treatment of severe spasticity after spinal cord injury. *Pain* **85**:287–289.
18. Penn RD and Paice JA (2000) Adverse effects associated with the intrathecal administration of Ziconotide. *Pain* **85**:291–296.
19. McGuire D, Bowersox S, Fellmann JD, and Luther RR (1997) Sympatholysis after neuron-specific, N-type, voltage-sensitive calcium channel blockade: first demonstration of N-channel function in humans. *J Cardio Pharm* **30**:400–403.
20. Ramilo CA, Zafaralla GC, Nadasdi L, Hammerland LG, Yoshikami D, Gray WR, Kristipati R, Ramachandran J, Miljanich G, and Olivera BM (1992) Novel α - and ω -conotoxins from *Conus striatus* venom. *Biochemistry* **31**:9919–9926.
21. Grantham CJ, Main MJ, and Cannell MB (1994) Fluspirilene block of N-type calcium current in NGF-differentiated PC12 cells. *Br J Pharmacol* **111**:483–488.
22. Sah DW and Bean BP (1994) Inhibition of P-type and N-type calcium channels by dopamine receptor antagonists. *Mol Pharmacol* **45**:84–92.
23. Weinsberg F, Bickmeyer U, and Wiegand H (1994) Effects of tetrandrine on calcium channel currents of bovine chromaffin cells. *Neuropharmacology* **33**:885–890.
24. Sutton KG, Siok C, Stea A, Zamponi GW, Heck SD, Volkman RA, Ahljianian MK, and Snutch TP (1998) Inhibition of neuronal calcium channels by a novel peptide spider toxin, DW13.3. *Mol Pharmacol* **54**:407–418.
25. Roulet JB, Spaetgens RL, Burlingame T, Feng ZP, and Zamponi GW (1999) Modulation of neuronal voltage-gated calcium channels by farnesol. *J Biol Chem* **274**:25439–25446.
26. Hu LY, Ryder TR, Rafferty MF, Siebers KM, Malone T, Chatterjee A, Feng MR, Lotarski SM, Rock DM, Stoehr SJ, et al. (2000) Neuronal N-type calcium channel blockers: a series of 4-piperidinylaniline analogs with analgesic activity. *Drug Des Discov* **17**:85–93.
27. Hu LY, Ryder TR, Rafferty MF, Taylor CP, Feng MR, Kuo BS, Lotarski SM, Miljanich GP, Millerman E, Siebers KM, et al. (2000) The discovery of [1-(4-dimethylamino-benzyl)-piperidin-4-yl]-[4-(3,3-dimethylbutyl)-phenyl]-[3-methyl-but-2-enyl]-amine, an N-type Ca²⁺ channel blocker with oral activity for analgesia. *Bioorg Med Chem* **8**:1203–1212.
28. Ryder TR, Hu LY, Rafferty MF, Millerman E, Szoke BG, and Tarczy-Hornoch K (1999) Multiple parallel synthesis of N, N-dialkyldepeptidylamines as N-type calcium channel blockers. *Bioorg Med Chem Lett* **9**:1813–1818.
29. Kamachi GL, Chan CK, Snutch T, Durieux ME, and Lynch III C (1999) Volatile anesthetic inhibition of neuronal Ca channel currents expressed in *Xenopus* oocytes. *Brain Res* **831**:85–96.
- 30a. Snutch TP and Zamponi GW (2000) inventors, NeuroMed Technologies, Inc., assignee. Calcium channel blockers. U.S. patent 6,011,035. 2000 Jan 4.
- 30b. Snutch TP and Zamponi GW (2001) inventors, NeuroMed Technologies, Inc., assignee. Calcium channel blockers. U.S. patent 6,294,533. 2001 Sep 25.
- 30c. Snutch TP (2001) inventor, NeuroMed Technologies, Inc., assignee. Fused ring calcium channel blockers. U.S. patent 6,310,059. 2001 Oct 30.
- 30d. Snutch TP (2002) inventor, NeuroMed Technologies, Inc., assignee. Preferentially substituted calcium channel blockers. U.S. patent 6,387,897. 2002 May 14.
- 30e. Snutch TP (2002) inventor, NeuroMed Technologies, Inc., assignee. Partially saturated calcium channel blockers. U.S. patent 6,492,375. 2002 Dec 10.
31. Beedle AM and Zamponi GW (2000) Block of voltage-dependent calcium channels by aliphatic monoamines. *Biophys J* **79**:260–270.
32. Lewis RJ, Nielson KJ, Craik DJ, Loughnan ML, Adams DA, Sharpe IA, Luchian T, Adams DJ, Bond T, Thomas L, et al. (2000) Novel ω -conotoxins from *Conus catus* discriminate among neuronal calcium channel subtypes. *J Biol Chem* **275**:35335–35344.
33. Martin DJ, McClelland D, Herd MB, Sutton KG, Hall MD, Lee K, Pinnock RD, and Scott RH (2002) Gabapentin-mediated inhibition of voltage-activated Ca²⁺ channel currents in cultured sensory neurones is dependent on culture conditions and channel subunit expression. *Neuropharmacology* **42**:353–366.
34. Sutton KG, Martin DJ, Pinnock RD, Lee K, and Scott RH (2002) Gabapentin inhibits high-threshold calcium channel currents in cultured rat dorsal root ganglion neurones. *Br J Pharmacol* **135**:257–265.
35. Mori Y, Nishida M, Shimizu S, Ishii M, Yoshinaga T, Ino M, Sawada K, and Niidome T (2002) Ca²⁺ channel α_{1B} subunit (Ca_v2.2) knockout mouse reveals a predominant role of N-type channels in the sympathetic regulation of the circulatory system. *Trends Cardiovasc Med* **12**:270–275.
36. Beuckmann CT, Sinton CM, Miyamoto N, Ino M, and Yanagisawa M (2003) N-type calcium channel α_{1B} subunit (Ca_v2.2) knock-out mice display hyperactivity and vigilance state differences. *J Neurosci* **23**:6793–6797.
37. Ghasemzadeh MB, Pierce RC, and Kalivas PW (1999) The monoamine neurons of the rat brain preferentially express a splice variant of α_{1B} subunit of the N-type calcium channel. *J Neurochem* **73**:1718–1723.
38. Pan JQ and Lipscombe D (2000) Alternative splicing in the cytoplasmic II-III loop of the N-type Ca channel α_{1B} subunit: functional differences are beta subunit-specific. *J Neurosci* **20**:4769–4775.
39. Lin Y, McDonough SI, and Lipscombe D (2004) Alternative splicing in the voltage-sensitive region of N-type Ca_v2.2 channels modulates channel kinetics. *J Neurophysiol* **92**:2820–2830.
40. Lin Z, Haus S, Edgerton J, and Lipscombe D (1997) Identification of functionally distinct isoforms of the N-type Ca²⁺ channel in rat sympathetic ganglia and brain. *Neuron* **18**:153–166.
41. Lin Z, Lin Y, Schorge S, Pan JQ, Beierlein M, and Lipscombe D (1999) Alternative splicing of a short cassette exon in α_{1B} generates functionally distinct N-type calcium channels in central and peripheral neurons. *J Neurosci* **19**:5322–5331.
42. Bell TJ, Thaler C, Castiglioni AJ, Helton TD, and Lipscombe D (2004) Cell specific alternative splicing increases calcium current density in the pain pathway. *Neuron* **42**:127–138.
43. Rettig J, Sheng ZH, Kim DK, Hodson CD, Snutch TP, and Catterall WA (1996) Isoform-specific interaction of the α_{1A} subunits of brain Ca²⁺ channels with the presynaptic proteins syntaxin and SNAP-25. *Proc Natl Acad Sci USA* **93**:7363–7368.
44. Maximov A, Sudhof TC, and Bezprozvany I (1999) Association of neuronal calcium channels with modular adaptor proteins. *J Biol Chem* **274**:24453–24456.
45. Kaneko S, Cooper CB, Nishioka N, Yamasaki H, Suzuki A, Jarvis SE, Akaike A, Satoh M, and Zamponi GW (2002) Identification and characterization of novel human Ca_v2.2 (alpha 1B) calcium channel variants lacking the synaptic protein interaction site. *J Biol Chem* **277**:82–92.

TABLE 8
Ca_v2.3 channels

Channel name	Ca _v 2.3
Description	Voltage-gated calcium channel α ₁ subunit
Other names	R-type, α _{1E} ; rbE-II (in rat) ¹ ; BII-1, BII-2 (in rabbit) ²
Molecular information	Human: 2251aa, L29384, 2270aa, L29385, ³ chr0.1q25-q31, <i>CACNA1E</i> Rat: 2222aa, ¹ GenBank accession no. L15453 Mouse: 2272aa, Q61290
Associated subunits	α ₂ δ/β, possibly γ
Functional assays	Voltage-clamp, patch-clamp, calcium imaging, neurotransmitter release
Current	I _{Ca,R}
Conductance	Not established
Ion selectivity	Ba ²⁺ ~ Ca ²⁺ (rat) ⁴ ; Ba ²⁺ > Ca ²⁺ (human) ³
Activation	V _a = +3.5 mV, τ _a = 1.3 ms at 0 mV (human α _{1E} /α ₂ δ/β ₁₋₃ , 15 mM Ba ²⁺ charge carrier in HEK293 cells) ³ V _a = -29.1 mV, τ _a = 2.1 ms at -10 mV (rat α _{1E} /α ₂ δ/β _{1b} , 4 mM Ba ²⁺ charge carrier in <i>Xenopus</i> oocytes) ¹
Inactivation	V _h = -71 mV, τ _h = 74 ms at 0 mV (human α _{1E} /α ₂ δ/β ₁₋₃ , 15 mM Ba ²⁺ charge carrier in HEK293 cells) ³ ; V _h = -78.1 mV, τ _h = 100 ms at -10 mV (rat α _{1E} /α ₂ δ/β _{1b} , 4 mM Ba ²⁺ charge carrier in <i>Xenopus</i> oocytes) ¹
Activators	None
Gating modifiers	None
Blockers	SNX-482, Ni ²⁺ (IC ₅₀ = 27 μM), Cd ²⁺ (IC ₅₀ = 0.8 μM), mibefradil (IC ₅₀ = 0.4 μM), ¹⁰ volatile anesthetics ¹¹
Radioligands	None
Channel distribution	Neurons (cell bodies, dendrites, some presynaptic terminals), heart, testes, pituitary
Physiological functions	Neurotransmitter release, repetitive firing, long-term potentiation, post-tetanic potentiation, neurosecretion ¹²⁻¹⁴
Mutations and pathophysiology	No point mutations in the native Ca _v 2.3 gene have been reported; mice deficient for the Ca _v 2.3 gene retain a substantial cerebellar R-type current, ⁵ suggesting that R-type currents actually reflect a heterogeneous mixture of channels; homozygous Ca _v 2.3-null mice survive to adulthood, reproduce, and are apparently behaviorally normal ^{5,6} ; mutant mice exhibit an increased resistance to formalin-induced pain, suggesting an involvement of the Ca _v 2.3 calcium channel in transmitting and/or the development of somatic inflammatory pain ⁶
Pharmacological significance	See "Comments"
Comments	Ca _v 2.3 has been variously reported to encode a novel type of calcium channel with properties shared between both low- and high-threshold calcium channels ^{1,4} or a type of high-threshold channel resistant to DHPs, ω-agatoxin-IVA, and ω-conotoxin-GVIA and called R-type (for "residual") ⁷ The tarantula toxin SNX-482 blocks exogenously expressed Ca _v 2.3 currents ⁸ but is only partially effective on native cerebellar R-type currents, ⁹ suggesting that Ca _v 2.3 does not always conduct a significant portion of the R-type current as originally defined ⁷ ; identified regions of alternative splicing include the domain II-III linker and carboxyl terminus and have been shown to affect channel kinetics and Ca ²⁺ -dependent stimulation ^{1-3,15,16}

aa, amino acids; chr., chromosome; HEK, human embryonic kidney; DHP, dihydropyridine.

1. Soong TW, Stea A, Hodson CD, Dubel SJ, Vincent SR, and Snutch TP (1993) Structure and functional expression of a member of the low voltage-activated calcium channel family. *Science* **260**:1133-1136.
2. Niidome T, Kim MS, Friedrich T, and Mori Y (1992) Molecular cloning and characterization of a novel calcium channel from rabbit brain. *FEBS Lett* **308**:7-13.
3. Williams ME, Marubio LM, Deal CR, Hans M, Brust PF, Philipson LH, Miller RJ, Johnson EC, Harpold MM, and Ellis SB (1994) Structure and functional characterization of neuronal α_{1E} calcium channel subtypes. *J Biol Chem* **269**:22347-22357.
4. Bourinet E, Zamponi GW, Stea A, Soong TW, Lewis BA, Jones LP, Yue DT, and Snutch TP (1996) The α_{1E} calcium channel exhibits permeation properties similar to low-voltage-activated calcium channels. *J Neurosci*, **16**:4983-4993.
5. Wilson SM, Toth PT, Oh SB, Gillard SE, Volsen S, Ren D, Philipson LH, Lee EC, Fletcher CF, Tassarollo L, et al. (2000) The status of voltage dependent calcium channels in α_{1E} knockout mice. *J Neurosci* **20**:8566-8571.
6. Saegusa H, Kurhara T, Zong S, Minowa O, Kazuno A, Han W, Matsuda Y, Yamanaka H, Osanai M, Noda T, et al. (2000) Altered pain responses in mice lacking α_{1E} subunit of the voltage dependent Ca channel. *Proc Natl Acad Sci USA* **97**:6132-6137.
7. Randall A and Tsien RW (1995) Pharmacological dissection of multiple types of calcium channel currents in rat cerebellar granule neurons. *J Neurosci* **15**:2995-3012.
8. Newcombe R, Szoke B, Palma A, Wang G, Chen XH, Hopkins W, Cong R, Miller J, Urge L, Tarczy-Hornoch K, et al. (1998) Selective peptide antagonist of the class E calcium channel from the venom of the tarantula *Hysteroecrates gigas*. *Biochemistry* **37**:15353-15362.
9. Tottene A, Volsen S, and Pietrobon D (2000) α_{1E} subunits form the pore of three cerebellar R-type calcium channels with different pharmacological and permeation properties. *J Neurosci* **20**:171-178.
10. Jimenez C, Bourinet E, Leuranguer V, Richard S, Snutch TP, and Nargeot J (2000) Determinants of voltage-dependent inactivation affect Mibefradil block of calcium channels. *Neuropharmacology* **39**:1-10.
11. Kamachi GL, Chan CK, Snutch T, Durieux ME, and Lynch III C (1999) Volatile anesthetic inhibition of neuronal Ca channel currents expressed in *Xenopus* oocytes. *Brain Res* **831**:85-96.
12. Dietrich D, Kirschstein T, Kukley M, Pereverzev A, von der Brölie C, Schneider T, and Beck H (2003) Functional specialization of presynaptic Ca_v2.3 Ca²⁺ channels. *Neuron* **39**:483-496.
13. Jing X, Li DQ, Olofsson CS, Salehi A, Surve VV, Caballero J, Ivarsson R, Lundquist I, Pereverzev A, Schneider T, et al. (2005) Ca_v2.3 calcium channels control second-phase insulin release. *J Clin Invest* **115**:146-154.
14. Pereverzev A, Salehi A, Mikhna M, Renstrom E, Hescheler J, Weiergraber M, Smyth N, and Schneider T (2005) The ablation of the Ca_v2.3/E-type voltage-gated Ca²⁺ channel causes a mild phenotype despite an altered glucose induced glucagon response in isolated islets of Langerhans. *Eur J Pharmacol* **511**:65-72.
15. Pereverzev A, Leroy J, Krieger A, Malecot CO, Hescheler J, Pfitzer G, Klockner U, and Schneider T (2002) Alternate splicing in the cytosolic II-III loop and the carboxy terminus of human E-type voltage-gated Ca²⁺ channels: electrophysiological characterization of isoforms. *Mol Cell Neurosci* **21**:352-365.
16. Klockner U, Pereverzev A, Leroy J, Krieger A, Vajna R, Pfitzer G, Hescheler J, Malecot CO, and Schneider T (2004) The cytoplasmic loop of Ca_v2.3 provides an essential determinant for the phorbol ester-mediated stimulation of E-type Ca²⁺ channel activity. *Eur J Neurosci* **19**:2659-2668.

TABLE 9
Ca_v3.1 channels

Channel name	Ca _v 3.1
Description	Voltage-gated calcium channel α_1 subunit
Other names	T-type, $\alpha_13.1$, α_{1G}
Molecular information	Human: 2377aa, O43497, NM_018896, chr. 17q22, <i>CACNA1G</i> ¹ Rat: 2254aa, O54898, AF027984 Mouse: 2288aa, CAI25956, NM_009783 (see "Comments")
Associated subunits	No biochemical evidence, small changes induced by $\alpha_2\delta_1$ ² and $\alpha_2\delta_2$ ^{3,4}
Functional assays	Voltage-clamp, calcium imaging
Current	$I_{Ca,T}$
Conductance	7.5pS ¹
Ion selectivity	$Sr^{2+} > Ba^{2+} = Ca^{2+}$
Activation	$V_a = -46$ mV, $\tau_a = 1$ ms at -10 mV ^{5,6}
Inactivation	$V_h = -73$ mV, $\tau_h = 11$ ms at -10 mV ^{5,6}
Activators	Not established
Gating modifiers	Kurtoxin, $IC_{50} = 15$ nM ⁷
Blockers	No subtype-specific blocker ⁸ ; selective for Ca _v 3.x relative to Ca _v 1.x and Ca _v 2.x: mibefradil, ^{9,10} U92032, ¹¹ penfluridol and pimozi ¹² ; nonselective: nickel ($IC_{50} = 250$ μ M), ¹³ amiloride ¹⁴
Radioligands	None
Channel distribution	Brain, especially soma and dendrites of neurons in olfactory bulb, amygdala, cerebral cortex, hippocampus, thalamus, hypothalamus, cerebellum, brain stem (human RNA blots, ^{1,5} rat in situ hybridization ¹⁵ and immunocytochemistry ¹⁶); ovary, placenta, heart (especially sinoatrial node; mouse in situ hybridization ¹⁷)
Physiological functions	Thalamic oscillations ¹⁸
Mutations and pathophysiology	Not established
Pharmacological significance	May mediate effect of absence antiepileptic drugs such as ethosuximide ¹⁹ and other thalamocortical dysrhythmias ²⁰
Comments	Splice variants that differ in their voltage dependence have been cloned ⁵

- aa, amino acids; chr., chromosome; U92032, 7-[[4-bis(fluorophenyl)methyl]-1-piperazinyl]methyl-2-[(2-hydroxyethyl)amino]4-(1-methylethyl)-2,4,6-cycloheptatrien-1-one.
- Perez-Reyes E, Cribbs LL, Daud A, Lacerda AE, Barclay J, Williamson MP, Fox M, Rees M, and Lee J-H (1998) Molecular characterization of a neuronal low voltage-activated T-type calcium channel. *Nature (Lond)* **391**:896–900.
 - Dolphin AC, Wyatt CN, Richards J, Beattie RE, Craig P, Lee J-H, Cribbs LL, Volsen SG, and Perez-Reyes E (1999) The effect of $\alpha_2\delta$ and other accessory subunits on expression and properties of the calcium channel α_{1G} . *J Physiol (Lond)* **519**:1:35–45.
 - Gao B, Sekido Y, Maximov A, Saad M, Forgacs E, Latif F, Lerman M, Lee J-H, Perez-Reyes E, Bezprozvanny I, et al. (2000) Functional properties of a new voltage-dependent calcium channel $\alpha_2\delta$ auxiliary subunit gene (*CACNA2D2*). *J Biol Chem* **275**:12237–12242.
 - Hobom M, Dai S, Marais E, Lacinova L, Hofmann F and Klugbauer N (2000) Neuronal distribution and functional characterization of the calcium channel $\alpha_2\delta_2$ subunit. *Eur J Neurosci* **12**:1217–1226.
 - Monteil A, Chemin J, Bourinet E, Mennessier G, Lory P, and Nargeot J (2000) Molecular and functional properties of the human α_{1G} subunit that forms T-type calcium channels. *J Biol Chem* **275**:6090–6100.
 - Klückner U, Lee JH, Cribbs LL, Daud A, Hescheler J, Pereverzev A, Perez-Reyes E, and Schneider T (1999) Comparison of the Ca²⁺ currents induced by expression of three cloned α_1 subunits, α_{1G} , α_{1H} and α_{1I} , of low-voltage-activated T-type Ca²⁺ channels. *Eur J Neurosci* **11**:4171–4178.
 - Chuang RS, Jaffe H, Cribbs L, Perez-Reyes E, and Swartz KJ (1998) Inhibition of T-type voltage-gated calcium channels by a new scorpion toxin. *Nat Neurosci* **1**:668–674.
 - Heady TN, Gomora JC, Macdonald TL, and Perez-Reyes E (2001) Molecular pharmacology of T-type Ca²⁺ channels. *Jpn J Pharmacol* **85**:339–350.
 - Perchenet L, Bénardeau A, and Ertel EA (2000) Pharmacological properties of Cav3.2, a low voltage-activated Ca²⁺ channel cloned from human heart. *Naunyn Schmiedeberg's Arch Pharmacol* **361**:590–599.
 - Martin RL, Lee JH, Cribbs LL, Perez-Reyes E, and Hanck DA (2000) Mibefradil block of cloned T-type calcium channels. *J Pharmacol Exp Ther* **295**:302–308.
 - Avery RB and Johnston D (1997) Ca²⁺ channel antagonist U-92032 inhibits both T-type Ca²⁺ channels and Na⁺ channels in hippocampal CA1 pyramidal neurons. *J Neurophysiol* **77**:1023–1028.
 - Santi CM, Cayabyab FS, Sutton KG, McRory JE, Mezeyova J, Hamming KS, Parker D, Stea A, and Snutch TP (2002) Differential inhibition of T-type calcium channels by neuroleptics. *J Neurosci* **22**:396–403.
 - Lee JH, Gomora JC, Cribbs LL, and Perez-Reyes E (1999) Nickel block of three cloned T-type calcium channels: low concentrations selectively block α_{1H} . *Biophys J* **77**:3034–3042.
 - Lacinova L, Klugbauer N, and Hofmann F (2000) Regulation of the calcium channel α_{1G} subunit by divalent cations and organic blockers. *Neuropharmacology* **39**:1254–1266.
 - Talley EM, Cribbs LL, Lee JH, Daud A, Perez-Reyes E, and Bayliss DA (1999) Differential distribution of three members of a gene family encoding low voltage-activated (T-type) calcium channels. *J Neurosci* **19**:1895–1911.
 - Craig PJ, Beattie RE, Folly EA, Banerjee MD, Reeves MB, Priestley JV, Carney SL, Sher E, Perez-Reyes E, and Volsen SG (1999) Distribution of the voltage-dependent calcium channel α_{1G} subunit mRNA and protein throughout the mature rat brain. *Eur J Neurosci* **11**:2949–2964.
 - Bohn G, Moosmang S, Conrad H, Ludwig A, Hofmann F, and Klugbauer N (2000) Expression of T- and L-type calcium channel mRNA in murine sinoatrial node. *FEBS Lett* **481**:73–76.
 - Perez-Reyes E (2003) Molecular physiology of low-voltage-activated T-type calcium channels. *Physiol Rev* **83**:117–161.
 - Gomora JC, Daud AN, Weiergräber M, and Perez-Reyes E (2001) Block of cloned human T-type calcium channels by succinimide antiepileptic drugs. *Mol Pharmacol* **60**:1121–1132.
 - Llinás RR, Ribary U, Jeanmonod D, Kronberg E, and Mitra PP (1999) Thalamocortical dysrhythmia: a neurological and neuropsychiatric syndrome characterized by magnetoencephalography. *Proc Natl Acad Sci USA* **96**:15222–15227.

TABLE 10
Ca_v3.2 channels

Channel name	Ca _v 3.2
Description	Voltage-gated calcium channel α_1 subunit
Other names	T-type, $\alpha_13.2$, α_{1H}
Molecular information	Human: 2353aa, O95180, AF051946, chr0.16p13.3, <i>CACNA1H</i> ¹ Rat: 2359aa, AAG35187, AF290213 Mouse: 2365aa, NP_067390, NM_021415
Associated subunits	Not established
Functional assays	Voltage-clamp, calcium imaging
Current	I _{Ca,T}
Conductance	9pS ²
Ion selectivity	Ba ²⁺ = Ca ²⁺
Activation	V _a = -46 mV, τ_a = 2 ms at -10 mV ³
Inactivation	V _h = -72 mV, τ_h = 16 ms at -10 mV ³
Activators	None
Gating modifiers	Kurtoxin ⁴
Blockers	Ca _v 3.2 is more sensitive than Ca _v 3.1 to block by nickel (IC ₅₀ = 12 μ M) ⁵ and possibly phenytoin ⁶ and amiloride ^{2,7} ; selective for Ca _v 3.x relative to Ca _v 1.x and Ca _v 2.x: mibefradil, ^{8,9} U92032, ¹⁰ penfluridol and pimozi ¹¹ and amiloride ¹² ; nonselective: nimodipine, ² anesthetics ⁵
Radioligands	None
Channel distribution	Kidney (human Northern ¹), rat smooth muscle (RT-PCR ¹³), liver (human Northern ¹), adrenal cortex (rat, bovine; in situ hybridization and RT-PCR ¹⁴), brain (especially in olfactory bulb, striatum, cerebral cortex, hippocampus, reticular thalamic nucleus; rat in situ hybridization ¹⁵), and heart (especially sinoatrial node; mouse in situ hybridization ¹⁶)
Physiological functions	Smooth muscle contraction, ¹⁷ smooth muscle proliferation, ¹⁸ aldosterone secretion, ¹⁹ cortisol secretion ²⁰
Mutations and pathophysiology	Single nucleotide polymorphisms associated with childhood absence epilepsy patients in a Chinese population ²¹
Pharmacological significance	May mediate effect of absence antiepileptic drugs such as ethosuximide ²² and other thalamocortical dysrhythmias ²³ ; potential drug target in hypertension and angina pectoris ²⁴
Comments	Splice variation found in the linker connecting repeat 3 and 4 ²⁵

aa, amino acids; chr., chromosome; RT-PCR, reverse-transcriptase-polymerase chain reaction.

- Cribbs LL, Lee JH, Yang J, Satin J, Zhang Y, Daud A, Barclay J, Williamson MP, Fox M, Rees M, et al. (1998) Cloning and characterization of α_{1H} from human heart, a member of the T-type Ca²⁺ channel gene family. *Circ Res* **83**:103–109.
- Williams ME, Washburn MS, Hans M, Urrutia A, Brust PF, Prodanovich P, Harpold MM, and Stauderman KA (1999) Structure and functional characterization of a novel human low-voltage activated calcium channel. *J Neurochem* **72**:791–799.
- Klößner U, Lee JH, Cribbs LL, Daud A, Hescheler J, Pereverzev A, Perez-Reyes E, and Schneider T (1999) Comparison of the Ca²⁺ currents induced by expression of three cloned α_1 subunits, α_{1G} , α_{1H} and α_{1I} , of low-voltage-activated T-type Ca²⁺ channels. *Eur J Neurosci* **11**:4171–4178.
- Chuang RS, Jaffe H, Cribbs L, Perez-Reyes E, and Swartz KJ (1998) Inhibition of T-type voltage-gated calcium channels by a new scorpion toxin. *Nat Neurosci* **1**:668–674.
- Lee JH, Gomora JC, Cribbs LL, and Perez-Reyes E (1999) Nickel block of three cloned T-type calcium channels: low concentrations selectively block α_{1H} . *Biophys J* **77**:3034–3042.
- Todorovic SM, Perez-Reyes E, and Lingle CJ (2000) Anticonvulsants but not general anesthetics have differential blocking effects on different T-type current variants. *Mol Pharmacol* **58**:98–108.
- Lacinova L, Klugbauer N, and Hofmann F (2000) Regulation of the calcium channel α_{1G} subunit by divalent cations and organic blockers. *Neuropharmacology* **39**:1254–1266.
- Perchenet L, Bénardeau A, and Ertel EA (2000) Pharmacological properties of Ca_v3.2, a low voltage-activated Ca²⁺ channel cloned from human heart. *Naunyn-Schmiedeberg's Arch Pharmacol* **361**:590–599.
- Martin RL, Lee JH, Cribbs LL, Perez-Reyes E, and Hanck DA (2000) Mibefradil block of cloned T-type calcium channels. *J Pharmacol Exp Ther* **295**:302–308.
- Avery RB and Johnston D (1997) Ca²⁺ channel antagonist U-92032 inhibits both T-type Ca²⁺ channels and Na⁺ channels in hippocampal CA1 pyramidal neurons. *J Neurophysiol* **77**:1023–1028.
- Santi CM, Cayabyab FS, Sutton KG, McRory JE, Mezeyova J, Hamming KS, Parker D, Stea A, and Snutch TP (2002) Differential inhibition of T-type calcium channels by neuroleptics. *J Neurosci* **22**:396–403.
- Tang C-M, Presser F, and Morad M (1988) Amiloride selectively blocks the low threshold (T) calcium channel. *Science* **240**:213–215.
- Hansen PB, Jensen BL, Andreassen D, and Skøtt O (2001) Differential expression of T- and L-type voltage-dependent calcium channels in renal resistance vessels. *Circ Res* **89**:630–638.
- Schrier AD, Wang H, Talley EM, Perez-Reyes E, and Barrett PQ (2001) α_{1H} T-type Ca²⁺ channel is the predominant subtype expressed in bovine and rat zona glomerulosa. *Am J Physiol Cell Physiol* **280**:C265–C272.
- Talley EM, Cribbs LL, Lee JH, Daud A, Perez-Reyes E, and Bayliss DA (1999) Differential distribution of three members of a gene family encoding low voltage-activated (T-type) calcium channels. *J Neurosci* **19**:1895–1911.
- Bohn G, Moosmang S, Conrad H, Ludwig A, Hofmann F, and Klugbauer N (2000) Expression of T- and L-type calcium channel mRNA in murine sinoatrial node. *FEBS Lett* **481**:73–76.
- Sarsero D, Fujiwara T, Molenaar P, and Angus JA (1998) Human vascular to cardiac tissue selectivity of L- and T-type calcium channel antagonists. *Br J Pharmacol* **125**:109–119.
- Schmitt R, Clozel JP, Iberg N, and Bühler FR (1995) Mibefradil prevents neointima formation after vascular injury in rats. Possible role of the blockade of the T-type voltage-operated calcium channel. *Arterioscler Thromb Vasc Biol* **15**:1161–1165.
- Rossier MF, Ertel EA, Vallotton MB, and Capponi AM (1998) Inhibitory action of mibefradil on calcium signaling and aldosterone synthesis in bovine adrenal glomerulosa cells. *J Pharmacol Exp Ther* **287**:824–831.
- Gomora JC, Xu L, Enyeart JA, and Enyeart JJ (2000) Effect of mibefradil on voltage-dependent gating and kinetics of T-type Ca²⁺ channels in cortisol-secreting cells. *J Pharmacol Exp Ther* **292**:96–103.
- Chen YC, Lu JJ, Pan H, Zhang YH, Wu HS, Xu KM, Liu XY, Jiang YW, Bao XH, Yao ZJ, et al. (2003) Association between genetic variation of CACNA1H and childhood absence epilepsy. *Ann Neurol* **54**:239–243.
- Gomora JC, Daud AN, Weiergräber M, and Perez-Reyes E (2001) Block of cloned human T-type calcium channels by succinimide antiepileptic drugs. *Mol Pharmacol* **60**:1121–1132.
- Linás R, Ribary U, Jeanmonod D, Kronberg E, and Mitra PP (1999) Thalamocortical dysrhythmia: a neurological and neuropsychiatric syndrome characterized by magnetoencephalography. *Proc Natl Acad Sci USA* **96**:15222–15227.
- Ertel SI, Ertel EA, and Clozel JP (1997) T-type Ca²⁺ channels and pharmacological blockade: potential pathophysiological relevance. *Cardiovasc Drugs Ther* **11**:723–739.
- Jagannathan S, Punt EL, Gu Y, Arnoult C, Sakkas D, Barratt CLR, and Publicover SJ (2002) Identification and localization of T-type voltage-operated calcium channel subunits in human male germ cells. Expression of multiple isoforms. *J Biol Chem* **277**:8449–8456.

TABLE 11
Ca_v3.3 channels

Channel name	Ca _v 3.3
Description	Voltage-gated calcium channel α_1 subunit
Other names	T-type, $\alpha_13.3$, α_{11}
Molecular information	Human: 2251aa, AAM67414, AF393329, chr. 22q13.1, <i>CACNA11</i> ¹ Rat: 1835aa, AF086827, AAD17796 Mouse 2753aa: XP_139476, XM_139476
Associated subunits	No biochemical evidence, small changes induced by γ_2 ²
Functional assays	Voltage-clamp, calcium imaging
Current	$I_{Ca,T}$
Conductance	11pS ¹
Ion selectivity	$Ba^{2+} = Ca^{2+}$
Activation	$V_a = -44$ mV, $\tau_a = 7$ ms at -10 mV ⁴
Inactivation	$V_h = -72$ mV, $\tau_h = 69$ ms at -10 mV ⁴
Activators	Not established
Gating modifiers	None
Blockers	No subtype-specific blocker ⁵ ; selective for Ca _v 3.x relative to Ca _v 1.x and Ca _v 2.x: mibefradil, ^{6,7} U92032, ⁸ penfluridol, ⁹ pimoizide ⁹ ; nonselective: nickel ($IC_{50} = 216$ μ M) ¹⁰
Radioligands	None
Channel distribution	Brain, especially olfactory bulb, striatum, cerebral cortex, hippocampus, reticular nucleus, lateral habenula, cerebellum (rat in situ hybridization, ¹¹ human Northern ¹²)
Physiological functions	Thalamic oscillations ¹³
Mutations and pathophysiology	Not established
Pharmacological significance	May mediate effect of absence antiepileptic drugs such as ethosuximide ¹⁴ and other thalamocortical dysrhythmias ¹⁵
Comments	Splice variants have been reported ¹⁶

aa, amino acids; chr., chromosome.

1. Lee JH, Daud AN, Cribbs LL, Lacerda AE, Pereverzev A, Klöckner U, Schneider T, and Perez-Reyes E (1999) Cloning and expression of a novel member of the low voltage-activated T-type calcium channel family. *J Neurosci* **19**:1912–1921.
2. Green PJ, Warre R, Hayes PD, McNaughton NC, Medhurst AD, Pangalos M, Duckworth DM, and Randall AD (2001) Kinetic modification of the α_{11} subunit-mediated T-type Ca^{2+} channel by a human neuronal Ca^{2+} channel γ subunit. *J Physiol (Lond)* **533**:467–478.
3. McRory JE, Santi CM, Hamming KS, Mezeyova J, Sutton KG, Baillie DL, Stea A, and Snutch TP (2001) Molecular and functional characterization of a family of rat brain T-type calcium channels. *J Biol Chem* **276**:3999–4011.
4. Klöckner U, Lee JH, Cribbs LL, Daud A, Hescheler J, Pereverzev A, Perez-Reyes E, and Schneider T (1999) Comparison of the Ca^{2+} currents induced by expression of three cloned α_1 subunits, α_{1G} , α_{1H} and α_{1I} , of low-voltage-activated T-type Ca^{2+} channels. *Eur J Neurosci* **11**:4171–4178.
5. Heady TN, Gomora JC, Macdonald TL, and Perez-Reyes E (2001) Molecular pharmacology of T-type Ca^{2+} channels. *Jpn J Pharmacol* **85**:339–350.
6. Perchenet L, Bénardeau A, and Ertel E. (2000) Pharmacological properties of Ca_v3.2, a low voltage-activated Ca^{2+} channel cloned from human heart. *Naunyn Schmiedeberg's Arch Pharmacol* **361**:590–599.
7. Martin RL, Lee JH, Cribbs LL, Perez-Reyes E, and Hanck DA (2000) Mibefradil block of cloned T-type calcium channels. *J Pharmacol Exp Ther* **295**:302–308.
8. Avery RB and Johnston D (1997) Ca^{2+} channel antagonist U-92032 inhibits both T-type Ca^{2+} channels and Na^+ channels in hippocampal CA1 pyramidal neurons. *J Neurophysiol* **77**:1023–1028.
9. Santi CM, Cayabyab FS, Sutton KG, McRory JE, Mezeyova J, Hamming KS, Parker D, Stea A, and Snutch TP (2002) Differential inhibition of T-type calcium channels by neuroleptics. *J Neurosci* **22**:396–403.
10. Lee JH, Gomora JC, Cribbs LL, and Perez-Reyes E (1999) Nickel block of three cloned T-type calcium channels: low concentrations selectively block α_{1H} . *Biophys J* **77**:3034–3042.
11. Talley EM, Cribbs LL, Lee JH, Daud A, Perez-Reyes E, and Bayliss DA (1999) Differential distribution of three members of a gene family encoding low voltage-activated (T-type) calcium channels. *J Neurosci* **19**:1895–1911.
12. Monteil A, Chemin J, Leuranguer V, Altier C, Mennessier G, Bourinet E, Lory P, and Nargeot J (2000) Specific properties of T-type calcium channels generated by the human α_{11} subunit. *J Biol Chem* **275**:16530–16535.
13. Perez-Reyes E (2003) Molecular physiology of low-voltage-activated T-type calcium channels. *Physiol Rev* **83**:117–161.
14. Gomora JC, Daud AN, Weiergräber M, and Perez-Reyes E (2001) Block of cloned human T-type calcium channels by succinimide antiepileptic drugs. *Mol Pharmacol* **60**:1121–1132.
15. Llinás RR, Ribary U, Jeanmonod D, Kronberg E, and Mitra PP (1999) Thalamocortical dysrhythmia: a neurological and neuropsychiatric syndrome characterized by magnetoencephalography. *Proc Natl Acad Sci USA* **96**:15222–15227.
16. Mittman S, Guo J, Emerick MC, and Agnew WS (1999) Structure and alternative splicing of the gene encoding α_{11} , a human brain T calcium channel α_1 subunit. *Neurosci Lett* **269**:121–124.