

# International Union of Basic and Clinical Pharmacology. LXXIII. Nomenclature for the Formyl Peptide Receptor (FPR) Family

RICHARD D. YE, FRANÇOIS BOULAY, JI MING WANG, CLAES DAHLGREN, CRAIG GERARD, MARC PARMENTIER,  
CHARLES N. SERHAN, AND PHILIP M. MURPHY

*Department of Pharmacology, College of Medicine, University of Illinois, Chicago, Illinois (R.D.Y.); Commissariat à l'Énergie Atomique, Institut de Recherches en Technologies et Sciences pour le Vivant, Laboratoire de Biochimie et de Biophysique des Systèmes Intégrés, Grenoble, France (F.B.); Centre National de la Recherche Scientifique, Unité Mixte de Recherche 5092, Grenoble, France (F.B.); Université Joseph Fourier, Grenoble, France (F.B.); Laboratory of Molecular Immunoregulation, Cancer and Inflammation Program, Center for Cancer Research, National Cancer Institute at Frederick, Frederick, Maryland (J.M.W.); Department of Rheumatology and Inflammation Research, University of Göteborg, Göteborg, Sweden (C.D.); Department of Pediatrics, Harvard Medical School and Children's Hospital, Boston, Massachusetts (C.G.); Institut de Recherche Interdisciplinaire en Biologie Humaine et Moléculaire, Université Libre de Bruxelles, Brussels, Belgium (M.P.); Center for Experimental Therapeutics and Reperfusion Injury, Brigham and Women's Hospital, Harvard Medical School, Boston, Massachusetts (C.N.S.); and Laboratory of Molecular Immunology, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Bethesda, Maryland (P.M.M.)*

Abstract	120
I. Introduction and historical overview	120
A. Discovery of <i>N</i> -formyl peptides as potent chemoattractants for phagocytes	120
B. Biochemical characterization of receptors that bind and internalize formyl peptides	123
II. The expanded family of formyl peptide receptors	125
A. Molecular characterization of FPR1	125
B. Identification of additional receptors of the <i>FPR</i> gene family	126
C. Tissue and cellular distribution of the human <i>FPR</i> gene family members	127
D. Polymorphism of the formyl peptide receptors	128
E. Evolution of the formyl peptide receptor gene family	129
III. Ligands for the formyl peptide receptor family of receptors	132
A. Agonists for the formyl peptide receptor family of receptors	132
1. <i>N</i> -formyl peptides	132
2. Microbe-derived nonformyl peptides	134
3. Host-derived peptides	135
a. Peptides associated with amyloidogenic diseases	135
b. Peptides associated with inflammatory and antibacterial responses	136
c. Annexin A1 and its N-terminal peptides	136
d. F2L, a potent endogenous agonist for FPR3	137
e. Other host-derived peptides	137
4. Host-derived, nonpeptide agonists	137
5. Agonists from peptide library	139
6. Agonists from nonpeptide library	140
B. Antagonists for the formyl peptide receptor family of receptors	140
IV. Structure-function relationship of the formyl peptide receptor family of receptors	142
A. Ligand-binding domains	142
B. G protein-coupling specificities	143
C. G protein-coupling domains	144
V. Regulation of formyl peptide receptors	144
A. Desensitization of FPR1	145
B. Phosphorylation of formyl peptide receptors	145
C. Interaction between formyl peptide receptors and $\beta$ -arrestins	146

Address correspondence to: Richard D. Ye, Department of Pharmacology, University of Illinois College of Medicine, 835 South Wolcott Avenue, M/C 868, Chicago, Illinois 60612. E-mail: yer@uic.edu

This article is available online at <http://pharmrev.aspetjournals.org>.

doi:10.1124/pr.109.001578.

1. Uncoupling of G proteins	146
2. Internalization of formyl peptide receptors	146
VI. Formyl peptide receptor signal transduction and activation of cell functions	147
A. Phagocyte activation for bacterial clearance	147
1. Chemotaxis	147
2. Superoxide generation	148
3. Degranulation	150
B. Other cellular functions	150
1. Transcriptional regulation and anti-inflammatory functions	150
2. Neutrophil apoptosis	151
VII. A new IUPHAR-approved nomenclature for the formyl peptide receptor family	152
A. Nomenclature of FPR1	152
B. Nomenclature of FPR2/ALX	152
C. Nomenclature of FPR3	153
VIII. Unmet challenges and future perspectives	153
A. The functions of the formyl peptide receptor family of receptors in other mammals	153
B. Structural basis for the diversity of the formyl peptide receptor family in ligand recognition	153
C. Mechanisms used by FPR2/ALX to respond to two different classes of ligands for proinflammatory and anti-inflammatory effects	154
Acknowledgments	154
References	155

**Abstract**—Formyl peptide receptors (FPRs) are a small group of seven-transmembrane domain, G protein-coupled receptors that are expressed mainly by mammalian phagocytic leukocytes and are known to be important in host defense and inflammation. The three human FPRs (FPR1, FPR2/ALX, and FPR3) share significant sequence homology and are encoded by clustered genes. Collectively, these receptors bind an extraordinarily numerous and structurally diverse group of agonistic ligands, including *N*-formyl and nonformyl peptides of different composition, that chemoattract and activate phagocytes. *N*-formyl peptides, which are encoded in nature only by bacterial and mitochondrial genes and result from obligatory initiation of bacterial and mitochondrial protein synthesis with *N*-formylmethionine, is the only ligand class common to all three human receptors. Surprisingly, the

endogenous anti-inflammatory peptide annexin 1 and its *N*-terminal fragments also bind human FPR1 and FPR2/ALX, and the anti-inflammatory eicosanoid lipoxin A4 is an agonist at FPR2/ALX. In comparison, fewer agonists have been identified for FPR3, the third member in this receptor family. Structural and functional studies of the FPRs have produced important information for understanding the general pharmacological principles governing all leukocyte chemoattractant receptors. This article aims to provide an overview of the discovery and pharmacological characterization of FPRs, to introduce an International Union of Basic and Clinical Pharmacology (IUPHAR)-recommended nomenclature, and to discuss unmet challenges, including the mechanisms used by these receptors to bind diverse ligands and mediate different biological functions.

## I. Introduction and Historical Overview

### A. Discovery of *N*-Formyl Peptides as Potent Chemoattractants for Phagocytes

The phenomenon of pus accumulation at sites of infection is part of the human condition and was described in the very earliest writings. Experimental evidence suggested that this might be due at least in part to bacterially derived phagocyte chemoattractants, as investigators injected live bacteria into tissues of experimental animals and observed accumulation of neutrophils at the injection site (for review, see Harris, 1954). In subsequent studies, Ward et al. (1968) found that filtrates from both Gram-positive and Gram-negative bacteria contain chemotactic activity for phagocytes in

vitro. Further investigations resulted in the identification of these chemotactic factors as low-molecular-weight peptides with blocked amino termini (Schiffmann et al., 1975b). Because prokaryotes initiate protein synthesis with *N*-formyl methionine, Schiffmann et al. (1975b) chemically synthesized and tested short peptides starting with *N*-formyl methionine (CHO-Met). They found potent chemotactic activities for neutrophils in many of the synthetic *N*-formyl peptides tested, especially peptides containing *N*-formyl-methionyl-leucine and *N*-formyl-methionyl-phenylalanine (Schiffmann et al., 1975a). This first report of *N*-formyl peptides serving as chemoattractants for neutrophils spurred a series of studies by Showell et al. (1976) and Freer et al. (1980, 1982) using rabbit neutrophils on the

TABLE 1

*N*-terminal modifications of selected formyl peptides and effects on agonistic activity

Most of the early studies were conducted using neutrophils from rabbits and humans. Note that the concentration of formyl peptides required for superoxide production and enzyme release is 10- to 50-fold higher than the concentration needed for maximal chemotaxis. Therefore, caution should be taken when comparing potency using different assays. Some variations may also exist between assays conducted in different laboratories.

Ligand	Assay	Effects	Potency	References
<i>N</i> -formyl-Met-Leu-Phe	Chemotaxis	Agonistic	pEC <sub>50</sub> = 10.15	Freer et al. (1980); Showell et al. (1976)
Met-Leu-Phe	Chemotaxis	Agonistic	pEC <sub>50</sub> = 6.17	Showell et al. (1976)
<i>N</i> -acetyl-Met-Leu-Phe	Chemotaxis	Agonistic	pEC <sub>50</sub> = 6.70	Freer et al. (1980)
<i>N</i> - <i>p</i> -tolylurea-Met-Leu-Phe	O <sub>2</sub> production	Agonistic	pEC <sub>50</sub> = 8.70	Higgins et al. (1996)
<i>N</i> - <i>tert</i> -butyloxycarbonyl-Met-Leu-Phe	Enzyme release	Antagonistic	pIC <sub>50</sub> = 6.19	Freer et al. (1980)
<i>N</i> - <i>iso</i> -butyloxycarbonyl-Met-Leu-Phe	O <sub>2</sub> production	Antagonistic	pIC <sub>50</sub> = 6.60	Derian et al. (1996)
<i>N</i> -formyl-Met-Phe-Leu	Chemotaxis	Agonistic	pEC <sub>50</sub> = 7.27	Showell et al. (1976)
Met-Phe-Leu	Chemotaxis	Agonistic	pEC <sub>50</sub> = 3.62	Showell et al. (1976)
<i>N</i> -acetyl-Met-Nle-Leu-Phe-Phe	Ca <sup>2+</sup> flux	Agonistic	pEC <sub>50</sub> = 10.00	Gao et al. (1994)
<i>N</i> -formyl-Met-Nle-Leu-Phe-Phe	Ca <sup>2+</sup> flux	Agonistic	pEC <sub>50</sub> = 10.00	Gao et al. (1994)
Met-Nle-Leu-Phe-Phe	Ca <sup>2+</sup> flux	Agonistic	pEC <sub>50</sub> = 9.00	Gao et al. (1994)

pIC<sub>50</sub>, negative logarithm of the IC<sub>50</sub>; pEC<sub>50</sub>, negative logarithm of the EC<sub>50</sub>; pK<sub>d</sub>, negative logarithm of K<sub>d</sub>.

structure-function relationship of the peptides. These studies resulted in identification of *N*-formyl-methionyl-leucyl-phenylalanine (fMet-Leu-Phe, or fMLF<sup>1</sup>) as the most potent agonist among 24 synthetic peptides tested, with an ED<sub>50</sub> of  $7.0 \pm 1.7 \times 10^{-11}$  M in neutrophil chemotaxis assays (Showell et al., 1976). The high potency and efficacy of fMLF to induce chemotaxis of human blood neutrophils was subsequently confirmed in independent studies by many other laboratories (Wilkinson, 1979). At higher concentrations (ED<sub>50</sub> of  $3.2 \pm 0.8 \times 10^{-10}$  M), the same peptide could stimulate lysosomal enzyme release (Freer et al., 1980). Although the formylated Met at position 1 was optimal for its bioactivity,

<sup>1</sup> Abbreviations: 7TM, 7-transmembrane; A $\beta$ ,  $\beta$  amyloid; A $\beta$ <sub>42</sub>, 42-amino acid form of  $\beta$  amyloid peptide; AG-14, 1,3-benzodioxolane-5-carboxylic acid 4'-benzyloxy-3'-methoxybenzylidene-hydrazide; AhR, aryl hydrocarbon receptor; ALX, receptor for lipoxin A4 and aspirin-triggered lipoxins; ANXA1, annexin A1; ATLS, aspirin-triggered lipoxins; CDCA, chenodeoxycholic acid; CGEN-855A, TIPM-FVPESTSKLQKFTSWFM-amide; CHIPS, chemotaxis inhibitory protein of *S. aureus*; CsA, cyclosporin A; CsH, cyclosporin H; c-suPAR, cleaved soluble uPAR; DC, dendritic cell; DCA, deoxycholic acid; ERK, extracellular signal-regulated kinase; fMLF, *N*-formyl-methionyl-leucyl-phenylalanine; FPR, formyl peptide receptor; FPRH1, formyl peptide receptor-homolog 1; FPRL1, formyl peptide receptor-like 1; G-CSF, granulocyte-colony-stimulating factor; GPCR, G protein-coupled receptor; GRK, G protein-coupled receptor kinase; HEK, human embryonic kidney; HSC, hematopoietic stem cells; IL, interleukin; LTB<sub>4</sub>, leukotriene B<sub>4</sub>; LXA4, lipoxin A4 (5*S*,6*R*,15*S*-trihydroxy-7,9,13-*trans*-11-eicosatetraenoic acid); MAPK, mitogen-activated protein kinase; NF- $\kappa$ B, nuclear factor  $\kappa$ B; PAF, platelet-activating factor; PAMP, pathogen-associated molecular pattern; PI3K, phosphatidylinositol 3-kinase; PI-PLC, phosphatidylinositol-specific phospholipase C; PKC, protein kinase C; PMA, phorbol 12-myristate 13-acetate; PtdIns(3,4,5)P<sub>3</sub>, phosphatidylinositol (3,4,5)-trisphosphate; PTEN, phosphatase and tensin homolog; PTX, pertussis toxin; Quin-C1, 4-butoxy-*N*-[2-(4-methoxy-phenyl)-4-oxo-1,4-dihydro-2*H*-quinazolin-3-yl]-benzamide; RBL, rat basophilic leukemia; SAA, serum amyloid A; SNP, single-nucleotide polymorphism; SOCS, suppressor of cytokine signaling; SR-BI, scavenger receptor class B type I; TA, temporin A; *t*-Boc, *t*-butyloxycarbonyl group; TLR, Toll-like receptor; TM, transmembrane; TNF, tumor necrosis factor; uPA, urokinase-type plasminogen activator; uPAR, urokinase-type plasminogen activator receptor.

replacing the Met with norleucine only slightly impaired the potency of the peptide in enzyme release assays.

The above findings led Freer et al. (1980) to propose that the *N*-formyl group is essential for the bioactivity of these chemotactic peptides. Peptides of the same sequence lacking the *N*-formyl group (e.g., Met-Leu-Phe) display reduced potency by 2 to 5 orders of magnitude, as shown in Table 1. However, replacing the *N*-formyl group with a *t*-butyloxycarbonyl group (*t*-Boc), which provides a urethane linkage instead of a peptide linkage, converted the same peptide to an antagonist (Boc-Met-Leu-Phe, Boc-MLF, or Boc1) (Freer et al., 1980). A number of different sequence combinations were tested in addition to *N*-formylated and nonformylated peptides of the same sequence. Based on these experiments, a model for formyl peptide binding requirements was proposed at the time. In this model, the formyl group is necessary for both ligand binding to a receptor and biological activity, and the Met side chain of position 1 occupies a hydrophobic pocket in the receptor. The Leu side chain may interact with a hydrophobic area of the receptor, whereas the phenylalanine side chain resides in a hydrophilic area. The Phe carbonyl plays a role for the C=O interaction with the receptor (Freer et al., 1982). As discussed below, this working model was later tested on a molecularly defined and cloned formyl peptide receptor, and modifications to the model were made through several studies. For instance, the *N*-formyl group is not absolutely required for potency at the receptor, and an *N*-acetylated peptide, Ac-Met-Nle-Leu-Phe-Phe, was found to be highly potent at the human formyl peptide receptor (FPR1) (Gao et al., 1994). The effects of *N*-terminal modifications on the affinity and potency of these peptides are shown in Table 1.

Although synthetic formyl peptides were used in the initial studies, it was believed that these peptides mimicked the behavior of chemotactic peptides released from bacteria during infection (Bennett et al., 1980). In 1984, Marasco et al. reported the isolation of chemotactic pep-

TABLE 2  
*Binding affinity and potency of bacterial and mitochondrial formyl peptides*

pEC<sub>50</sub> is defined as the negative logarithm of the EC<sub>50</sub>. HL-60 cells transfected to express FPR1 or FPR2/ALX, and Chinese hamster ovary cells transfected to express FPR1 were used in some studies

Ligand	Origin	Assay	Potency	Cells (Receptors)	Reference
<i>N</i> -formyl-Met-Leu-Phe	<i>E. coli</i>	Chemotaxis Lysozyme release O <sub>2</sub> production Binding	pEC <sub>50</sub> = 10.15 pEC <sub>50</sub> = 7.49	Neutrophils Neutrophils	Showell et al. (1976); Freer et al. (1980) Freer et al. (1980)
<i>N</i> -formyl-Met-Ile-Phe-Leu	<i>S. aureus</i>	Chemotaxis Competitive binding O <sub>2</sub> production Ca <sup>2+</sup> flux	pEC <sub>50</sub> = 7.00 pK <sub>d</sub> = 9.28–7.61 pK <sub>d</sub> = 6.37 pEC <sub>50</sub> = 7.51 pIC <sub>50</sub> = 8.01	Neutrophils Neutrophils Transfected cells (FPR2/ALX) Monocytes Monocytes	Boxer et al. (1979) Koo et al. (1982) Quehenberger et al (1993) Rot et al. (1987) Rot et al. (1987)
<i>N</i> -formyl-Met-Ile-Val-Ile-Leu	<i>L. monocytogenes</i>	O <sub>2</sub> production Ca <sup>2+</sup> flux	pEC <sub>50</sub> = 8.00 pEC <sub>50</sub> = 8.66	Neutrophils (mFpr1) HL-60 (FPR1)	Southgate et al. (2008) Rabiet et al. (2005)
<i>N</i> -formyl-Met-Ile-Gly-Trp-Ile	<i>L. monocytogenes</i>	O <sub>2</sub> production Ca <sup>2+</sup> flux	pEC <sub>50</sub> = 6.80 pEC <sub>50</sub> = 7.82 pEC <sub>50</sub> = 7.70	HL-60 (FPR2/ALX) Neutrophils (mFpr1) HL-60 (FPR1)	Rabiet et al. (2005) Southgate et al. (2008) Rabiet et al. (2005)
<i>N</i> -formyl-Met-Ile-Val-Thr-Leu-Phe	<i>L. monocytogenes</i>	Ca <sup>2+</sup> flux	pEC <sub>50</sub> = 6.68 pEC <sub>50</sub> = 6.70	HL-60 (FPR2/ALX) HL-60 (FPR1)	Rabiet et al. (2005) Rabiet et al. (2005)
<i>N</i> -formyl-Met-Ile-Gly-Trp-Ile-Ile	<i>L. monocytogenes</i>	Ca <sup>2+</sup> flux	pEC <sub>50</sub> = 7.40	HL-60 (FPR2/ALX)	Rabiet et al. (2005)
<i>N</i> -formyl-Met-Phe-Glu-Asp-Ala-Val-Ala-Trp-Phy	<i>M. avium</i>	Chemotaxis	pEC <sub>50</sub> = 6.00	CHO (FPR1)	Rabiet et al. (2005)
<i>N</i> -formyl-Met-Met-Tyr-Ala-Leu-Phe	Mitochondria, ND6	Ca <sup>2+</sup> flux	pEC <sub>50</sub> = 7.92	HL-60 (FPR1)	Gripentrog et al. (2008)
<i>N</i> -formyl-Met-Leu-Lys-Leu-Ile-Val	Mitochondria, ND4	Ca <sup>2+</sup> flux	pEC <sub>50</sub> = 7.82 pEC <sub>50</sub> = 7.92	HL-60 (FPR2/ALX) HL-60 (FPR1)	Rabiet et al. (2005) Rabiet et al. (2005)
<i>N</i> -formyl-Met-Tyr-Phe-Ile-Asn-Ile-Leu-Thr-Leu	Mitochondria, ND1	Binding	pEC <sub>50</sub> = 7.26 pK <sub>d</sub> = 9.00	HL-60 (FPR2/ALX) CHO (FPR2/ALX)	Rabiet et al. (2005) Chiang et al. (2000)
<i>N</i> -formyl-Met-Phe-Ala-Asp-Arg-Trp	Cytochrome <i>c</i> oxidase subunit	Ca <sup>2+</sup> flux	pEC <sub>50</sub> = 6.80 pEC <sub>50</sub> = 6.68	HL-60 (FPR1) HL-60 (FPR2/ALX)	Rabiet et al. (2005) Rabiet et al. (2005)

CHO, Chinese hamster ovary; pIC<sub>50</sub>, negative logarithm of the IC<sub>50</sub>; pEC<sub>50</sub>, negative logarithm of the EC<sub>50</sub>.

tides from *Escherichia coli* culture supernatants. Using a lysosomal enzyme release assay and competitive binding with [<sup>3</sup>H]fMLF, these investigators identified several distinct formyl peptides with bioactivity for rabbit neutrophils. Further characterization of these peptides led to the identification of fMLF as the major chemotactic factor in *E. coli* culture supernatants. It should be noted that secretion of chemotactic peptides is not limited to *E. coli* and is probably a general property of bacteria and that other bacterial species have also been experimentally shown to release phagocyte-chemotactic factors. For example, Rot et al. (1987, 1989) showed that culture supernatants of *Staphylococcus aureus* contain several peptides that are chemotactic for human monocytes. Two of these peptides, fMet-Ile-Phe-Leu (fMIFL) and fMet-Leu-Phe-Ile (fMLFI), displayed potent activities in chemotaxis and superoxide generation assays (Rot et al., 1987). More recently, Rabiet et al. (2005) reported that several peptides derived from *Listeria monocytogenes*, known to bind to the nonclassic MHC Class Ib molecule M2-H3 (Pamer et al., 1992), are potent chemoattractants for human leukocytes. These results demonstrate that bacteria release *N*-formyl peptides of various sequence compositions that can be recognized by human and rabbit neutrophils and monocytes, and that which can therefore be collectively considered as a pathogen-associated molecular pattern (PAMP). The PAMP concept has heretofore been mainly associated with Toll-like receptors (Medzhitov and Janeway, 2000). Inhibition of bacterial peptide deformylase (an enzyme that deformylates the methionyl group to generate mature proteins) leads to increased bacterial production of formylated peptides (Fu et al., 2003), suggesting a regulatory mechanism for the generation of formyl peptides. A list of selected formyl peptides derived from bacteria and mitochondria is shown in Table 2.

Unlike prokaryotes that initiate protein synthesis with an *N*-formylmethionine, the synthesis of eukaryotic proteins encoded by nuclear DNA is initiated with a nonformylated methionine. However, mitochondrial protein synthesis is initiated with *N*-formylmethionine. Carp (1982) reported that disrupted mitochondria from several types of cells that were tested could stimulate human neutrophil migration. Chemotaxis was also induced by purified proteins of the mitochondria respiratory chain. These activities were inhibited by *t*-Boc-Phe-D-Leu-Phe-D-Leu-Phe (Boc2), an antagonist of the fMLF receptor, suggesting that these mitochondrial chemotactic factors share the same receptor with fMLF. More recent studies of peptides of mitochondrial origin (Chiang et al., 2000; Rabiet et al., 2005) support the conclusion made by Carp, indicating that mitochondrial proteins bearing *N*-formyl modifications are endogenous ligands for formyl peptide receptors (Table 2). Attraction of phagocytes to sites of inflammation and tissue injury may be facilitated by these endogenous ligands.

### B. Biochemical Characterization of Receptors That Bind and Internalize Formyl Peptides

The findings that *N*-formyl peptides could stimulate neutrophil chemotaxis (Schiffmann et al., 1975a; Zigmond, 1977) and lysosomal enzyme release (Showell et al., 1976) provided strong evidence for the presence of a functional receptor on target cells. Formyl peptide receptors were first defined at the biochemical level using radioisotope- and fluorescence-labeled peptide ligands. In 1977, Aswanikumar et al. demonstrated that rabbit neutrophils express specific and saturable binding sites for [<sup>3</sup>H]fNle-Leu-Leu-Phe, with a  $K_d$  of  $1.5 \times 10^{-9}$  M. A similar binding site for fMLF was reported by Williams et al. (1977). In both cases, the bound radiolabeled peptides could be effectively displaced by unlabeled peptides of the same or similar composition, and the concentrations of the peptides that effectively displaced the radiolabeled ligand matched the concentrations used in functional assays. In a series of binding experiments, Niedel et al. (1979a) studied neutrophil uptake of radiolabeled formyl peptides. They found that iodinated fNle-Leu-Phe-Nle-Tyr-Lys retained full biological activity with an  $EC_{50}$  of 0.4 nM in a human neutrophil chemotaxis assay. Reversibility of ligand binding was lost over time and at the appropriate temperature, suggesting receptor-mediated internalization of the formyl peptides. Because unlabeled formyl peptides of similar compositions could inhibit binding and uptake of the iodinated peptide, it was concluded that these peptides interact with the same receptor on neutrophils (Niedel et al., 1979a). Using tetramethylrhodamine-labeled fNle-Leu-Phe-Nle-Tyr-Lys and time-lapse video microscopy, Niedel et al. (1979b) observed rapid internalization and aggregation of the fluorescent peptide, suggesting receptor binding and receptor-mediated uptake of the labeled ligand. In other studies, the relationship between formyl peptide pre-exposure and cell responsiveness was investigated. Vitkauskas et al. (1980) reported that pretreatment of rabbit neutrophils with formyl peptides reduced the available binding sites for [<sup>3</sup>H]fNle-Leu-Phe as well as cell response to subsequent formyl peptide stimulation, a process that the authors termed *deactivation*. A study conducted by Donabedian and Gallin (1981) identified *preferential* and *nonpreferential deactivation* toward the same and different agonists, respectively. In preferential deactivation, incubation of human neutrophils with fMLF reduced the cell-surface binding sites for the same ligand, resulting in a decrease in chemotaxis toward subsequent fMLF stimulation. In nonpreferential deactivation, treatment of human neutrophils with a high concentration of the activated complement C5 fragment (C5a) caused reduced response of the cells to fMLF stimulation, without reducing (and actually increasing) the cell surface binding sites for fMLF. These published studies were among the earliest reports on G protein-coupled receptor (GPCR)-mediated internalization, al-

though the identity of the formyl peptide receptor at the molecular level was still unknown at the time. In addition, what Donabedian and Gallin called *preferential deactivation* was in fact an early example of heterologous desensitization (Didsbury et al., 1991) and cross-desensitization of chemoattractant GPCRs (Richardson et al., 1995).

The study by Donabedian and Gallin (1981) also showed that agonist-induced decrease in the number of formyl peptide binding sites was transient, and these binding sites could return to the cell surface if the cells were kept at 37°C. The study demonstrated a recycling pool of formyl peptide receptors. When neutrophils were sonicated and fractionated on sucrose density gradients, fMLF binding sites were found in the fractions containing specific granules (Fletcher and Gallin, 1983). Therefore, neutrophils contain an intracellular pool of cryptic formyl peptide receptors that may be mobilized to the cell surface. Using time-resolved flow cytometry, Sklar and colleagues studied the dynamics of formyl peptide ligand interaction with its receptor in neutrophils (Sklar et al., 1981, 1984; Sklar and Finney, 1982; Finney and Sklar, 1983). These studies took advantage of the ability of cytometric and fluorimetric analyses to distinguish between receptor-bound and unbound ligands in real time to determine different states of the receptor. The results not only confirmed internalization of ligand-occupied receptors but also determined key parameters of formyl peptide association and dissociation, demonstrating that the ligand-receptor complex could undergo an alteration in affinity (Sklar et al., 1984). Jesaitis et al. (1984, 1985) initiated studies of formyl peptide receptor interaction with the cytoskeleton and found that a receptor-cytoskeleton complex was formed before receptor internalization and was resistant to Triton X-100. In this ternary complex, the formyl peptide ligand binds to its receptor with high affinity and then slowly dissociates from the receptor (Jesaitis et al., 1984). These studies demonstrate that the formyl peptide receptor interacts with intracellular proteins such as cytoskeleton proteins and this interaction can affect the binding properties of the receptor.

Early studies using radiolabeled fMLF identified one class of binding sites in intact neutrophils. Using membrane binding assays, Koo et al. (1982) reported that human neutrophils contain two classes of formyl peptide binding sites with dissociation constants of 0.53 and 24 nM, respectively. The heterogeneity of receptor binding to fMLF was not due to negative cooperativity, because the rate of dissociation was unaltered with increasing receptor occupancy. This result could be interpreted as evidence for the presence of two distinct, noninterconvertible populations of binding sites for formyl peptides, one responsible for neutrophil chemotaxis, which requires lower concentrations of formyl peptides, and the other mediating additional bactericidal functions such as lysoso-

mal enzyme release and superoxide generation known to require higher agonist concentrations (Lehmeyer et al., 1979; Korchak et al., 1984). Alternatively, the different dissociation constants could indicate the presence of one class of receptors present in two affinity states that are interconvertible. A subsequent study conducted by the same authors found that a nonhydrolyzable derivative of GTP, when added to the membrane preparation in a binding assay, could convert a part of the high-affinity binding site to a low-affinity site without altering the total number of receptors (Koo et al., 1983). This effect was reverted by removal of the GTP analog. Similar guanine nucleotide regulation of receptor affinity was reported in other studies of receptors that couple to G proteins (Lad et al., 1977; De Lean et al., 1980). Thus, these biochemical studies were interpreted to mean that a single class of receptors for fMLF is present in two binding states, and that their affinities for binding formyl peptides are regulated by GTP in membrane binding assays in which the receptors are accessible to the GTP analog but the receptor-cytoskeleton interaction is disrupted. Under these experimental conditions, the G protein-bound receptors exhibit high affinity for fMLF, and the receptors uncoupled from G proteins display low binding affinity. This interpretation was later complicated by discovery at the molecular level of FPR1 and FPR2/ALX (see section II), binding fMLF with high and low affinity, respectively.

The biological activities of formyl peptides were extensively characterized after their initial discoveries. The prototypic formyl peptide, fMLF, possesses full agonistic activity that is similar to that of C5a in stimulating chemotaxis, lysosomal enzyme release, and superoxide generation (Schiffmann et al., 1975a; Showell et al., 1976; Lehmeyer et al., 1979). The ability of an activated receptor to mediate multiple cellular functions is intriguing, and the phenomenon spurred additional efforts to characterize the formyl peptide receptors at the biochemical level. Several radiolabeled formyl peptide derivatives were prepared for photoaffinity cross-linking to cell surface receptors. Using this approach, it was determined that the neutrophil formyl peptide receptor is a  $M_r$  50,000~70,000 glycoprotein (Niedel et al., 1980a; Dolmatch and Niedel, 1983; Allen et al., 1986). Treatment with endoglycosidase F reduces the size of the receptor to  $M_r$  32,000, which retains the ability to bind formyl peptides (Malech et al., 1985). These results indicate that the formyl peptide receptor in neutrophils is an *N*-glycosylated cell surface protein, but ligand binding does not depend on glycosylation of the receptor. Additional efforts in purification of the formyl peptide receptor were met with difficulties. Because of the low abundance of the receptor in neutrophils, only partially purified proteins were obtained.

## II. The Expanded Family of Formyl Peptide Receptors

### A. Molecular Characterization of FPR1

Studies conducted throughout the 1980s led to the identification of the formyl peptide receptor as a GPCR. All major neutrophil functions stimulated by fMLF can be inhibited by treatment of the cells with pertussis toxin (Bokoch and Gilman, 1984; Bokoch et al., 1984; Lad et al., 1985; Sha'afi and Molski, 1988; Gierschik et al., 1989; Snyderman and Uhing, 1992), indicating that the G proteins that couple to the formyl peptide receptors belong to the G<sub>i</sub> family of heterotrimeric G proteins (Simon et al., 1991). At that time, a number of G protein-coupled receptors from other tissues had been pharmacologically characterized, but only a few genes coding for these proteins, including rhodopsin,  $\alpha$ - and  $\beta$ -adrenergic receptors, and two of the muscarinic receptors, had been cloned (Dixon et al., 1986; Kubo et al., 1986; Kobilka et al., 1987). The strategies used in cloning these receptor genes involved mostly purification of the receptor proteins and sequencing of the derived peptides to obtain partial sequence of the GPCR for the design of nucleotide probes, which were used for DNA hybridization with cDNA or gene libraries. In addition to DNA hybridization, an expression cloning approach using oocytes from *Xenopus laevis* was adopted to isolate the gene for a muscarinic receptor. A different approach was taken to clone the human formyl peptide receptor gene. Boulay et al. (1990b), with expertise in photoaffinity labeling techniques and protein chemistry, started a cloning project with the synthesis of a photoaffinity hetero-bifunctional derivative of *N*-formyl-Met-Leu-Phe-Lys (fMLFK), with one moiety that was photoactivatable and another that could bind to streptavidin. The rationale behind the synthesis of this derivative was to photolabel the receptor in dibutyryl cAMP-differentiated HL-60 cells, which express the fMLF receptor based on binding and internalization assays (Niedel et al., 1980b). The photoaffinity-labeled receptor was then purified through the interaction of its biotin moiety with streptavidin-coated beads. However, this initial effort to isolate the receptor gene was unsuccessful.

In addition to screening cDNA or genomic libraries using oligonucleotide probes and *X. laevis* oocyte-based expression cloning, the molecular cloning techniques then available included *E. coli*- or mammalian cell-based expression cloning with antibody detection of the recombinant protein on the cell surface or in cell lysate. The mammalian cell expression cloning approach was successfully used in the identification of genes for several cell surface molecules, including the T cell-specific surface protein (CD28), using antibodies against these proteins (Aruffo and Seed, 1987). Taking this approach, Boulay et al. (1990a,b) screened a cDNA expression library made with mRNA from differentiated HL-60 cells. Because no antibody against human formyl peptide re-

ceptor was available at that time, the approach described by Aruffo and Seed (1987) was not directly applicable. Therefore, the *N*-formyl peptide fMLFK was derivatized with a hydrophilic dodecapeptide (*N*-acetyl-SDQALSFLKDYC) that represents the N-terminal region of the bovine mitochondrial ADP/ATP carrier. The dodecapeptide was coupled via the C-terminal cysteine on the free  $\epsilon$ -amino group of the lysine of fMLFK, yielding a water-soluble heterobifunctional ligand that retained its high affinity for FPR1. Because high-affinity antibodies against this peptide were available, it was anticipated that this ligand could be used as a bridge between the receptor and immunoglobulins coated on Petri dishes. However, this approach was also met with difficulties, prompting a switch to labeling the derivatized peptide with <sup>125</sup>I and using this peptide for quantification of any expressed protein that bound the probe in COS cells transfected with fractions of the cDNA library. To improve the chance of detecting a specific receptor on the cell surface, the primary cDNA library was split into pools with approximately 700 independent bacterial colonies each, an approach used in the cloning of the erythropoietin receptor (D'Andrea et al., 1989). After screening 184 pools of the library, several pools yielded signals 5- to 20-fold higher than that of the background. The pools were further divided, and two cDNA variants were found to produce markedly increased signals, in part due to internalization and intracellular accumulation of the radiolabeled formyl peptide derivative. These two cDNA isolates (FPR-26 and FPR-98) represent the two allelic forms of human FPR1 in HL-60 cells (Boulay et al., 1990a,b), encoding a putative seven-transmembrane domain (7TM) receptor with 350 amino acids (Fig. 1).

Shortly after the completion of the initial FPR1 cloning work, Thomas et al. (1990) reported the isolation of a rabbit cDNA, known as F3R, encoding a protein that was first described as rabbit FPR1. This conclusion was based on the ability of the cDNA to encode a receptor capable of mobilizing calcium when expressed in *X. laevis* oocytes and stimulated with fMLF. Subsequent studies showed that F3R encodes a rabbit receptor for IL-8 (Thomas et al., 1991), and the high-affinity formyl peptide receptor in rabbits was found to be the product of another distantly related gene (Ye et al., 1993).

With an available cDNA for human FPR1, it became possible to conduct pharmacological characterization of the receptor in a defined cellular environment free of similar receptors. Functional expression of FPR1 was accomplished in transfected mammalian cells (Prossnitz et al., 1991) and *X. laevis* oocytes (Murphy et al., 1992). In these studies, the recombinant human FPR1 was shown to mediate fMLF-induced calcium mobilization in a pertussis toxin-sensitive manner. Subsequent studies demonstrated that the recombinant FPR1 is able to mediate fMLF-induced actin polymerization and chemotaxis in transfected HL-60 cells (Prossnitz et al., 1993)




FIG. 1. Predicted transmembrane disposition of the human FPR1. The protein sequence of the FPR-98 isoform (Leu110, Ala346) is shown (Boulay et al., 1990a). The transmembrane domains (TMs) are predicted based on hydrophobicity of the amino acid sequence and on similarities to the rhodopsin structure. The amino acids that form the boundaries of the transmembrane domains are numbered. One-letter amino acid code is used. The square blocks in reverse color represent positions at which amino acid substitutions result from polymorphisms, including amino acids 11 (Ile/Thr), 47 (Val/Ala), 101 (Leu/Val), 190 (Arg/Trp), 192 (Asn/Lys) and 346 (Ala/Glu). The circle blocks in reverse color indicate amino acids with known functions as follows. Arg84, Lys85, and Asp284 are critical for high-affinity binding of fMLF (Mills et al., 1998; Quehenberger et al., 1997). Asp122, Arg123, and Cys124 are the signature sequence for G protein interaction (DRY in many GPCRs). NPMLY in the TM7 are known signature sequence (NPXXY) for receptor internalization (Gripentrog et al., 2000; He et al., 2001). The 11 Ser and Thr residues in the cytoplasmic tail are potential phosphorylation sites for GRK2 and GRK3 (Prossnitz et al., 1995). CHO, carbohydrate, marks the identified and potential (in parenthesis) sites for N-glycosylation. The predicted disulfide bond between Cys98 and Cys176 is marked with double-line (=).

and release of granule contents in transfected rat basophilic leukemia (RBL) cells (Ali et al., 1993). More recently, FPR1-dependent production of superoxide was reconstituted in COS<sup>phox</sup> cells (He et al., 2004), a transgenic COS-7 cell line expressing the four essential phagocyte oxidase (phox) proteins (Price et al., 2002). In mice, loss of function assay has shown that Fpr1 is a major receptor to mediate fMLF-induced NADPH oxidase activation. Another receptor, possibly Fpr2, also contributes to oxidant production when mouse neutrophils are stimulated with fMLF at higher concentrations (e.g., 50  $\mu$ M) (Lavigne et al., 2002). These results demonstrate the ability of FPR1 to activate multiple signaling pathways important for the microbicidal functions of phagocytes.

#### B. Identification of Additional Receptors of the FPR Gene Family

The availability of the FPR1 sequence not only facilitated molecular characterization of the receptor in het-

erologous expression systems but also prompted identification of receptors that share sequence homology with FPR1. Strategies were developed to use low-stringency DNA hybridization (under conditions of reduced temperature and/or increased salt concentration), with the cDNA of human FPR1 as a probe, for isolation of cDNAs or genes of similar sequences. In 1992, several laboratories reported the identification of a cDNA (Murphy et al., 1992; Perez et al., 1992; Ye et al., 1992) and gene (Bao et al., 1992) coding for a putative 7TM receptor that shares significant sequence homology to human FPR1. The gene product was given different names, including FPR2 (formyl peptide receptor 2) for its low-affinity binding of fMLF (Ye et al., 1992), FPRL1 (formyl peptide receptor-like 1) (Murphy et al., 1992), FPRH1 (formyl peptide receptor-homolog 1) (Bao et al., 1992), and "receptor related to formyl peptide receptor" (Perez et al., 1992) based on its sequence homology to the human FPR1. Other names used in the literature include HM63 (No-



mura et al., 1993), and FMLP-related receptor II (GenBank accession no. M76672) (Table 3). Pharmacological characterization has led to the identification of the eicosanoids lipoxin A4 (LXA4) and aspirin-triggered lipoxins (Fiore et al., 1994; Chiang et al., 2006), and numerous peptides (Le et al., 2001b, 2002; Migeotte et al., 2006) as ligands for this receptor. Therefore, in addition to FPRL1, which appears frequently in the literature, the names LXA4R and ALX have been introduced to convey the ability of the receptor to interact with LXA4 and aspirin-triggered lipoxins (Brink et al., 2003). Given that IUPHAR nomenclature of a receptor is based on its agonists and *N*-formyl peptides are the only common ligands for this class of receptors, we will refer to the receptor as FPR2/ALX, a nomenclature now recommended by IUPHAR.

FPR2/ALX is a 7TM receptor with 351 amino acids and shares 69% of its amino acids with human FPR1 (Fig. 2). Despite the relatively high level of sequence homology, FPR2/ALX is a low-affinity receptor for fMLF, with a  $K_d$  of 430 nM (Murphy et al., 1992; Ye et al., 1992; Quehenberger et al., 1993). It has been reported that mitochondria-derived formyl peptides are more potent agonists for FPR2/ALX than fMLF (Rabiet et al., 2005), suggesting that its primary function may be to recognize host-driven mitochondrial peptides or possibly other bacterially derived formyl peptides. As discussed in detail below, in addition to formyl peptides and LXA4, FPR2/ALX is also able to interact with nonformylated peptides.

A second gene with significant sequence homology to human *FPR1* was identified using a similar cloning strategy. This gene, initially named *FPRL2* (formyl peptide receptor-like 2) (Murphy et al., 1992) and *FPRH2* (formyl peptide receptor homolog 2) (Bao et al., 1992) based on sequence homology to FPR1, encodes a putative 7TM receptor of 352 amino acids (Table 3). This receptor was also named *RMLP-related receptor 1* (GenBank accession no. M76673). In this article, we will refer to it as FPR3, an IUPHAR-recommended name based on the ability of the receptor to bind certain *N*-formyl peptides. FPR3 shares 56% sequence identity with human FPR1 at the protein level. Although fMLF is inactive at FPR3 (Migeotte et al., 2005), fMMYALF, a peptide derived from a mitochondrial protein, is an agonist at FPR3 in calcium flux assays, with  $pEC_{50}$  of 6.00 ( $EC_{50}$  of  $10^{-6}$  M;  $pEC_{50}$  is the negative logarithm of  $EC_{50}$ ) (Rabiet et al., 2005). Migeotte et al. (2005) recently reported that F2L, a naturally occurring endogenous acylated

peptide derived from the N-terminal sequence of heme-binding protein, is a potent agonist for FPR3 with a  $pEC_{50}$  of 7.00 in a reporter assay. Additional ligands for FPR3 are discussed in Section III below. A survey of FPR3 expression using monoclonal antibodies found an intracellular pool of the receptor. It is noteworthy that, in one published study, one third of blood donors (8 of 24) had no detectable FPR3 expression on the surface of dendritic cells (Migeotte et al., 2005).

The three members of the human FPR gene family are clustered on chromosome 19q13.3–19q13.4, adjacent to the human C5a receptor gene (Bao et al., 1992; Murphy et al., 1992). This suggests that the genes arose by relatively recent duplication of a common ancestor and that the encoded proteins may have related biological functions.

### C. Tissue and Cellular Distribution of the Human FPR Gene Family Members

FPR1 was initially found in human and rabbit neutrophils through functional characterization (Schiffmann et al., 1975b; Showell et al., 1976; Aswanikumar et al., 1977; Williams et al., 1977; Zigmond, 1977). More extensive investigations were conducted when key reagents such as cDNA probes and antibodies became available. The three receptors of the FPR gene family are primarily found in myeloid cells, but the distribution varies within myeloid cell subsets, as described in the next paragraph. The results of these studies also indicate the presence of formyl peptide receptors in nonmyeloid cells. Using an antibody recognizing the carboxyl terminal 11 amino acids of FPR1, Becker et al. (1998) found immunoreactivity at the immunohistochemical level in multiple organs and tissues that include epithelial cells in organs with secretory functions, endocrine cells (including follicular cells of the thyroid and cortical cells of the adrenal gland), liver hepatocytes and Kupffer cells, smooth muscle cells and endothelial cells, brain, spinal cord, and both motor and sensory neurons. In most cases, this has not yet been verified at the RNA level or by functional studies. Human FPR2/ALX has a tissue distribution pattern similar to that of FPR1. Because of significant sequence homology between FPR1 and FPR2/ALX in their carboxyl termini, it is possible that the rabbit polyclonal antibody used in the immunohistochemical analysis have detected both FPR1 and FPR2/ALX. Another study showed endothelial cell expression of formyl peptide binding sites (Rotrosen et al., 1987). The low affinity of the binding site for fMLF is

TABLE 3  
IUPHAR-recommended nomenclature for human FPRs and previously used names

IUPHAR-Recommended Nomenclature	Other Names Used	Gene Location	GenBank Accession Number
FPR1	FPR, NFPR, FMLP, FMLPR	19q13.41	M60627 (cDNA); NM_002029 (gene)
FPR2/ALX	FPRL1, FPRH1, RFP, LXA4R, ALXR, HM63, FMLPX, FPR2A	19q13.3-q13.4	M88107 (cDNA); NM_001005738 (gene)
FPR3	FPRL2, FPRH2, FMLPY	19q13.3-q13.4	NM_002030 (gene)

FPR1	<b>METNSSLP</b> TNISGGT <b>PAVSAGYLF</b> LDIIIT <b>YLVFAVTFVLGVLGNGLVIWVAGFRM</b> THVTTTISYLN <b>LAVADF</b> CF <b>TSTLPPFFMVRKAMGGH</b> 90
FPR2/ALX	<b>METNFSTPLN</b> EYEEVSY <b>ESAGYTVLRILPLV</b> LG <b>VTFLGVLGNGLVIWVAGFRM</b> TRVTTTIC <b>YLNALALADFS</b> TATL <b>PFLIVSMAMGEK</b> 90
FPR3	<b>METNFSIPLN</b> ETEEV <b>LPEPAGHTVLWIFSL</b> LVH <b>GVTFVFGVLGNGLVIWVAGFRM</b> TRVTTTIC <b>YLNALALADFS</b> SAIL <b>PFPMVSVAMREK</b> 90
FPR1	<b>WPFGWFLCK</b> LF <b>FTIVDINLFGSVFLIALIALD</b> RCV <b>CVLHPVW</b> TQ <b>NHRTVSLAKKVIIGPWV</b> MALL <b>LLTLPVIIRV</b> TTVP <b>PKTGTVACTFNF</b> 180
FPR2/ALX	<b>WPFGWFLCK</b> L <b>LIHVVDINLFGSVFLIGFIALD</b> RCIC <b>CVLHPVWAQ</b> NHRTV <b>SLAMKVIIGPWILALV</b> LTLP <b>VFLFTTVTIP</b> NGDT <b>YCTFNF</b> 180
FPR3	<b>WPFASFLCK</b> L <b>VHVMIDINL</b> FVSV <b>YLITIIALD</b> RCIC <b>CVLHPAWAQ</b> NHRT <b>MSLAKRVMTGLWIF</b> TIV <b>LTLPNFI</b> FWTT <b>ISTTNGD</b> TYC <b>IFNF</b> 180
FPR1	SPW <b>TNDPKERIN</b> VAV <b>AMLTVRGII</b> RF <b>IIIGFSAP</b> MS <b>IVAVSYGLI</b> AT <b>KIHKQGLIK</b> SSR <b>PLRVLSF</b> VAA <b>FFLCW</b> SPY <b>QVVALI</b> AT <b>VRIRE</b> 270
FPR2/ALX	<b>ASWGGTPE</b> ERL <b>KVAITMLTAR</b> GI <b>IRFVIGFSL</b> PMS <b>IVAICYGLI</b> AA <b>KHKGMK</b> SSR <b>PLRVLTAV</b> VAS <b>FFICW</b> FP <b>QLVALL</b> GT <b>VWLKE</b> 270
FPR3	<b>AFWGD</b> TAVER <b>LNVFITMAK</b> V <b>FLILHFI</b> IG <b>FTVPM</b> SI <b>ITVCYGI</b> IA <b>AKIHR</b> NH <b>MKSSR</b> PL <b>RVFAAV</b> VAS <b>FFICW</b> FP <b>YELIG</b> IL <b>LMAYWLKE</b> 270
FPR1	-LL <b>QGM</b> YKE <b>IGIAVD</b> VT <b>SALAFF</b> NS <b>CLNP</b> LV <b>FVMGQD</b> FRER <b>LIALP</b> AS <b>LERALTE</b> -- <b>DSTQ</b> TSD <b>TATNST</b> LP <b>SAEVAL</b> QAK 350
FPR2/ALX	<b>MLFYG</b> KY <b>KIIDIL</b> VN <b>PTSSLA</b> FFNS <b>CLNP</b> LV <b>FVMGQD</b> FRER <b>LIIH</b> SL <b>PTSLER</b> AL <b>SE</b> -- <b>DSAP</b> TND <b>TANCA</b> SP <b>PAETEL</b> QAM 351
FPR3	<b>MLLNG</b> KY <b>KIILV</b> LN <b>PTSSLA</b> FFNS <b>CLNP</b> LV <b>FVMGRN</b> FQER <b>LIR</b> SL <b>PTSLER</b> AL <b>TEVP</b> DSA <b>QTSN</b> THT <b>TASPE</b> ETEL <b>QAM</b> 353

FIG. 2. Alignment of the protein sequences of the human FPRs. The putative transmembrane domains (I to VII) are shaded. Sequence of the FPR-98 isoform is shown. Comparison of the three receptors has identified highly conserved regions, including most of TM-I and TM-II, and the short intracellular loop connecting TM-I and TM-II. The second intracellular loops from these receptors, known for G protein interaction, are nearly identical. TM-VII, including the NPXXY motif and a stretch of ~25 amino acids extending toward the C-terminal tail, are also conserved among these receptors. Major differences are found in the extracellular domains between FPR1 and the other two receptors, especially in the amino termini (~50% different), the second extracellular loops (56% different), and the third extracellular loops (~50% different). The two putative *N*-glycosylation sites in the N-terminal domains are conserved among all three receptors. Most of the serines and threonines in the C-terminal tail, along with charged residues that constitute consensus GRK phosphorylation sites, are also conserved among these receptors.

consistent with the binding property of FPR2/ALX. Functional FPR1 has also been found in hepatocytes, glial cells, astrocytes, and platelets (Lacy et al., 1995; McCoy et al., 1995; Le et al., 2000a; Czapiga et al., 2005), suggesting that formyl peptide receptors may have more complex functions than are presently appreciated.

Unlike FPR1 and FPR2/ALX, FPR3 transcripts are not found in neutrophils (Murphy et al., 1992). Instead, it can be detected together with transcripts for FPR1 and FPR2/ALX in monocytes, although the expression pattern changes with monocyte differentiation. In particular, in the process of monocyte differentiation into immature dendritic cells (DCs), the cellular expression of FPR2/ALX progressively declines (Yang et al., 2001a), whereas FPR2/ALX expression remains unchanged during monocyte differentiation into macrophages. There is a progressive loss of FPR1 during differentiation of immature DC to mature DC, such that FPR3 becomes the predominant human formyl peptide receptor in mature DC (Yang et al., 2002; Migeotte et al., 2005). The biological significance of differential expression of formyl peptide receptors in monocytes, macrophages, and DCs has not yet been clearly delineated.

#### D. Polymorphism of the Formyl Peptide Receptors

The initial cloning study resulted in the identification of two allelic forms of human FPR1 (FPR-26 and FPR-98) (Boulay et al., 1990a). The differences between the two cDNAs are Val101, Glu346 in FPR-26 and Leu101, Ala346 in FPR-98. In a subsequent study, additional single-nucleotide polymorphisms (SNPs) of FPR1 were identified (Sahagun-Ruiz et al., 2001), whereas no polymorphism has been identified for FPR2/ALX. The difference in polymorphism suggests that these two structurally similar receptors have undergone distinct processes

of evolutionary selection. FPR1 SNPs identified to date are either synonymous or nonsynonymous, the latter resulting in amino acid changes at positions 11 (Thr/Ile), 47 (Val/Ala), 97 (Leu/Met), 101 (Leu/Val), 190 (Arg/Trp), 192 (Asn/Lys), and 346 (Ala/Glu) (Fig. 1). At least 30 variations have been identified at the nucleotide level, and these SNPs contribute to the different haplotypes of human FPR1 (Sahagun-Ruiz et al., 2001; Zhang et al., 2003; Gunji et al., 2007; Gripenrot et al., 2008).

There is a reported association between juvenile onset, localized periodontitis, and defects in neutrophil responses to formyl peptides (Clark et al., 1977; Van Dyke et al., 1986; Agarwal et al., 1989; Perez et al., 1991). Consequently, attempts have been made to establish a correlation between FPR1 SNPs and defective neutrophil functions in juvenile-localized periodontitis. Gwinn et al. (1999) used single-strand conformation polymorphism analysis to identify two base substitutions resulting in changes at positions 110 (F110S) and 126 (C126W) in FPR1. When tested as recombinant proteins, these FPR1 variants display defective  $G\alpha_i$  protein coupling in reconstitution assays using Sf9-cell derived mutant receptors (Seifert and Wenzel-Seifert, 2001). The Trp126 form of FPR1 exhibits reduced function in chemotaxis and calcium mobilization assays (Jones et al., 2003; Nanamori et al., 2004b). One study also indicated failure of cell surface expression of the Ser110 variant (Jones et al., 2003), whereas in the other study, the Ser110 variant and Ser110/Trp126 variant of FPR1 were normally expressed on the cell surface, but their ability to take up fMLF was compromised (Nanamori et al., 2004b). Although these studies seem to have identified a structural basis for defective FPR1 in juvenile localized periodontitis, a subsequent and more extensive independent survey failed to identify the same base sub-

stitutions in 111 cases of juvenile localized periodontitis examined (Zhang et al., 2003). Therefore, it is still unclear whether the base substitutions resulting in amino acid changes at amino acids 110 and 126 are responsible for the defective neutrophil response to fMLF observed in localized juvenile periodontitis patients.

Mills (2007) and Gripenrog et al. (2008) found that some of the FPR1 haplotypes are indeed functionally distinct. In extracellular signal-regulated kinase (ERK) phosphorylation, chemotaxis, and receptor down-regulation assays, the three tested FPR1 haplotypes displayed similar responsiveness to fMLF; however, differences were found when these FPR1 haplotypes were activated by formyl peptides from other bacterial strains such as *Mycobacterium avium*. These findings suggest individual differences in FPR1-mediated detection of certain bacterial strains.

As discussed above, no SNP has been identified in the human FPR2/ALX gene. The only variant of FPR3 described to date involves an Asp to His substitution at position 338 (GenBank accession no. AAA58482). An examination of this SNP found no difference in receptor expression or functional response between the FPR3 variants in an F2L-induced cAMP inhibition assay (Migeotte et al., 2005).

### *E. Evolution of the Formyl Peptide Receptor Gene Family*

Since the first reported cDNA cloning of human FPR1 (Boulay et al., 1990a,b), orthologs have been identified at the molecular level in other primates (Alvarez et al., 1996), rabbits (Ye et al., 1993), and mice (Gao and Murphy, 1993). Functional formyl peptide receptors have also been found in rat, guinea pig, and horse, and differences relative to human FPR1 have been noted (Oseas et al., 1980; Snyderman and Pike, 1980; Ward et al., 1984). For instance, horse neutrophils respond to fMLF with granule enzyme release but not chemotaxis (Snyderman and Pike, 1980). These cells produce superoxide only when primed with TNF $\alpha$  or platelet-activating factor (PAF) and then challenged with fMLF (Brazil et al., 1998). Neutrophils from other species such as dog were initially reported not to respond to fMLF (Redl et al., 1983) but were later found to respond to very high concentrations (100  $\mu$ M) of fMLF in a chemotaxis assay (Linnekin et al., 1990). Because most studies have been conducted with fMLF, it will be interesting to determine whether neutrophils from dogs and other mammalian species can respond to the many other naturally occurring bioactive formyl peptides that have been described previously (Freer et al., 1980; Rot et al., 1987; Rabiet et al., 2005; Southgate et al., 2008).

The *FPR* gene family has a complex evolutionary history. The number of genes in the family can vary markedly in different mammalian species, indicating differential gene expansion or extinction and suggesting the

presence of differential selective pressures. In particular, the mouse FPR gene family has at least eight members (Fig. 3) clustered on mouse chromosome 17A3.2, as opposed to only three in human on chromosome 19q13.3 in a region syntenic with the mouse cluster (Gao et al., 1998; for review, see Migeotte et al., 2006). The gene product of *Fpr1* is the mouse ortholog of human FPR1. The gene products of both *Fpr-rs1* and *Fpr-rs2* (Fpr-related sequence 1 and 2) are structurally most similar to human FPR2/ALX (Hartt et al., 1999).

The gene product of mouse *Fpr-rs2*, referred to as mouse Fpr2 (mFpr2), is a low-affinity receptor for fMLF (Hartt et al., 1999). This receptor responds to several peptide agonists that activate human FPR2/ALX, including the amyloidogenic proteins serum amyloid A (Liang et al., 2000) and amyloid  $\beta_{(1-42)}$  (Tiffany et al., 2001). An FPR2/ALX-specific antagonist inhibits activation of mouse neutrophils as well as cells stably expressing this receptor, suggesting that the receptor shares significant structural and pharmacological properties with human FPR2/ALX (Onnheim et al., 2008). A variant of mouse *Fpr-rs1* has been cloned that encodes a protein with a four-residue insertion (ARNV, after Leu141 in the 4th transmembrane helix) relative to the reference sequence (Fig. 3); this variant has been named the mouse Lxa4 receptor (Takano et al., 1997). Another study showed that mouse Fpr2 can mediate phosphoinositide turnover when coupled to  $G\alpha_{16}$  and stimulated with LXA4 (Vaughn et al., 2002). It is noteworthy that mouse Fpr2 has been reported to be also a receptor for F2L (Gao et al., 2007), a potent agonist for human FPR3 (Migeotte et al., 2005). These results indicate that the gene products of *Fpr-rs1* (including its variant, mouse Lxa4r) and mouse Fpr2 share some pharmacological properties with human FPR2/ALX, and mouse Fpr2 also has an overlap function with human FPR3 in the detection of F2L. The biological functions of other mouse *Fpr* gene family members have not been determined (Gao et al., 1998; Wang and Ye, 2002). *Fpr-rs4* encodes a putative 7TM-spanning receptor with 323 amino acids. A stop codon at position 246 in the predicted transmembrane 6 causes an early termination of the *Fpr-rs5* open reading frame, which makes it unlikely to encode a functional receptor, although it does not have the typical features of a pseudogene (Gao et al., 1998). *Fpr-rs6* and *Fpr-rs7* are other orphans in the family also identified by low-stringency DNA hybridization (Wang and Ye, 2002). The complex evolution of the *FPR* gene family is apparent from the high sequence divergence between species orthologs ( $\sim 25$ – $30\%$  between human and mouse) (Table 4), which is very high compared with most proteins but characteristic for immunoregulatory proteins, including other chemoattractant receptors, such as chemokine receptors (Murphy, 1993).

Although mouse Fpr1 shares a sequence identity of 77% with human FPR1 (Table 4), it displays low affinity for fMLF. In functional assays, fMLF concentrations of

FIG. 3. Alignment of the predicted receptor sequence of the mouse *Fpr* genes. The putative transmembrane domains (TM-I–TM-VII) are shaded. Dashes indicate gaps in sequence created for alignment purposes. It is noteworthy that there is an eight-residue insertion in the N-terminal region of Fpr-1, just before TM-I. A three-residue insertion is found in the second extracellular loop in Fpr-1. The positively charged residues Arg84 and Lys85, found in human FPR1 and known for the interaction with fMLF, are missing from Fpr-1 and other mouse Fpr-related sequences. In its place are the noncharged residues Ser92 and Met93. The predicted Fpr-rs5 sequence is truncated at amino acid 246, resulting in a putative protein with only five TMs. Fpr-rs4 encodes a protein of 323 residues with a short C-terminal tail. In Fpr-rs1, there is a four-residue deletion in TM-IV, whereas the cloned mouse LXA4 receptor gene encodes a protein with the sequence of ARNV in its place. Polymorphisms exist in the *Fpr-rs1* gene that result in amino acid substitutions at positions 3 (Thr/Ser), 8 (Pro/His), 13 (Asp/Glu), 16 (Ile/Val), 222 (Thr/Tyr), 236 (Phe/Ser), 296 (Ile/Met), and 318 (Gln/Pro) (Takano et al., 1997; Gao et al., 1998; Wang et al., 2002). The highest sequence identity (81%) is found between Fpr-rs1 and Fpr-rs2, and between Fpr-rs3 and Fpr-rs4.

TABLE 4  
Sequence identity between the human and mouse FPRs

Shown in the table are percent of identical amino acids between the receptors, based on pairwise comparison of the entire sequence of each receptor using the ALIGN program (Scientific & Educational Software, ver. 1.02). The FPR1–98 sequence (Boulay et al. 1990a) is used for the comparison. Note that the gene product of *Fpr-rs5* used in sequence comparison is 246 amino acids. The size of each receptor is indicated in the first column

Size (a.a.)	Name	FPR1 (98)	FPR2 /ALX	FPR3	Fpr1	Fpr-rs1	Fpr2	Fpr-rs3	Fpr-rs4	Fpr-rs5	Fpr-rs6	Fpr-rs7
350	FPR1 (98)	100	68	58	77	59	64	55	51	54	51	51
351	FPR2/ALX		100	72	68	73	76	65	61	64	59	58
353	FPR3			100	56	60	63	54	52	58	52	52
364	Fpr1				100	57	62	53	51	55	51	50
347	Fpr-rs1					100	82	65	61	63	58	59
351	Fpr2						100	66	64	64	59	60
343	Fpr-rs3							100	77	78	74	73
323	Fpr-rs4								100	74	73	73
246	Fpr-rs5									100	64	62
339	Fpr-rs6										100	95
338	Fpr-rs7											100

a.a., amino acids.

100 nM or higher are required for activation of cell functions such as calcium mobilization (Gao and Murphy, 1993). Other neutrophil functions, including degranulation and superoxide production, require even higher fMLF concentrations (5–10 μM) in mice. It is notable that both mouse Fpr1 and human FPR2/ALX

are low-affinity receptors for fMLF. The structural basis for the low-affinity binding is not entirely clear, but a comparison of human FPR1, FPR2/ALX, and mouse Fpr1 has identified major differences in key residues known to define the human FPR1 binding pocket for fMLF (Quehenberger et al., 1997; Mills et al., 1998). A

pair of positively charged residues, Arg84 and Lys85 in human FPR1 (Fig. 1), is replaced with uncharged residues in both FPR2/ALX and mouse Fpr1. Moreover, the negatively charged Asp284 in human FPR1, known to stabilize the receptor structure, is also substituted with an uncharged residue in FPR2/ALX and a positively charged Lys in mouse Fpr1 (Fig. 3). Therefore, mouse Fpr1 shares certain structural features with human FPR2/ALX. Restoration of Arg84 and Lys85 in an FPR1-FPR2/ALX chimeric receptor displaying low affinity for fMLF (Quehenberger et al., 1993) has been shown to significantly improve the binding affinity for the formyl peptide from 105 to 1.6 nM (Quehenberger et al., 1997). In comparison, Lys85 and Asp284 are conserved in rabbit FPR1 that displays high binding affinity for fMLF (Ye et al., 1993). This finding suggests that small changes to key residues in formyl peptide receptors can profoundly affect their binding selectivity and may have interesting implications for the evolution of this receptor family.

The transcripts of *Fpr1* and *Fpr-rs2* are most abundant in mouse neutrophils, suggesting that these two receptors might be primarily responsible for detection of formyl peptides. Mouse Fpr2 is 81% identical in protein sequence to mouse Fpr1 and 58% to human FPR1 (Gao et al., 1998) (Fig. 4). It responds even more weakly to fMLF than mouse Fpr1 and contributes to the second

concentration optimum in mouse neutrophil chemotaxis assays using fMLF at micromolar concentrations (Hartt et al., 1999). As a result, mouse neutrophils exhibit a binding affinity for fMLF that is 100- to 500-fold lower than that of human neutrophils. The absence of high affinity interaction with fMLF in mouse neutrophils raises the question of whether mice can effectively detect bacterially derived formyl peptides and whether FPR1 is important for host defense in mice. To address these questions, Gao et al. (1999) generated *Fpr1* knockout mice and examined the ability of these mice to clear *L. monocytogenes*. They found defective chemotaxis of *Fpr1*(-/-) neutrophils toward fMLF. Moreover, the *Fpr1*(-/-) mice were more susceptible to *L. monocytogenes* infection, showing an increased mortality rate (Gao et al., 1999). These results indicate an important function of mouse Fpr1 in host defense. More recently, Southgate et al. (2008) examined mouse neutrophil responses to formyl peptides derived from *L. monocytogenes* (with the sequence of fMIVIL) and *Staphylococcus aureus* (with the sequence of fMIFL) and found that these peptides were at least 100-fold more potent than fMLF in stimulating mouse neutrophil chemotaxis and superoxide production. Using transfected cell lines that individually express mFpr1 and mFpr2, the authors demonstrated that mFpr1 can respond to low nanomolar concentrations of fMIVIL and fMIFL, whereas mFpr2 requires micromolar concentrations of fMIVIL and fMIFL in calcium mobilization assay. Targeted deletion of *Fpr1* results in compromised neutrophil responses to both of these peptides in assays measuring chemotaxis, degranulation, and superoxide generation (Southgate et al., 2008). Therefore, although mouse neutrophils are inefficient in responding to fMLF, a major chemoattractant from *E. coli* culture (Marasco et al., 1984), these innate immune cells are able to detect low nanomolar concentrations of the two peptides from *L. monocytogenes* and *S. aureus*. It is possible that the increased susceptibility of *Fpr1*(-/-) mice to *L. monocytogenes* infection may result from defective neutrophil detection of the *L. monocytogenes*-derived formyl peptides such as fMIVIL. *Listeria* fMIVIL is also known to bind the mouse Class Ib MHC molecule H2-M3 (Gulden et al., 1996) and to activate human FPR2/ALX (Rabiet et al., 2005). *S. aureus* fMIFL is one of the six oligopeptides purified from an *S. aureus* culture and known to activate human neutrophils and monocytes with high potency (Rot et al., 1987). Differential detection of these peptides by mouse Fpr1 may reflect the environment in which mice live. *E. coli* infection can be lethal in humans but is not known to be a natural mouse pathogen. Thus, it is interesting to speculate that selection pressure has favored the acquisition of additional binding properties for more sensitive detection of the *E. coli*-derived fMLF by human FPR1.




FIG. 4. Sequence homology between the FPR family members and their tissue distribution. The predicted protein sequences of the three human (h) FPR genes, the eight mouse (m) Fpr genes, and the rabbit (r) FPR1 gene were compared (Boulay et al., 1990a,b; Ye et al., 1993; Gao et al., 1998; Wang et al., 2002). Based on sequence homology, the hFPR1, mFpr1, and rFPR1 are in the same cluster. The mFpr-rs1, mFpr-rs2 (also termed mFpr2), and mFpr-rs8 are in another cluster closely related to hFPR2/ALX and hFPR3. The mFpr-rs3, mFpr-rs4, mFpr-rs6, and mFpr-rs7 (and, to a lesser extent, mFpr-rs5) are closely related based on their protein sequences (see Table 4 for sequence identity between the gene products). Note that some of these genes are not expressed in neutrophils and monocytes. The tissue expression profiles for mFpr-rs4, mFpr-rs5, and mFpr-rs8 have not been determined. Mo, monocytes; PMN, polymorphonuclear leukocytes; iDC, immature dendritic cells; astro, astrocytes; T, T lymphocytes.




<p>FPR1-selective agonists</p>	 <p>fMLF (M.W. 437.55)</p> <p>AG-14 (M.W. 404.42)</p>
<p>FPR2/ALX-selective agonists</p>	 <p>Lipoxin A4 (M.W. 352.47)</p> <p>Quin-C1 (M.W. 445.51)</p> <p>Compound 43 (M.W. 384.86)</p>
<p>Antagonists</p>	 <p><i>t</i>-Boc-FLFLF (M.W. 785.51)</p> <p>Cyclosporin H (M.W. 1202.61)</p>

FIG. 5. Chemical structures of selected ligands for the formyl peptide receptors. Despite their abilities to bind to FPR1 and/or FPR2/ALX, these ligands have quite different structures. Of the ligands shown, *t*-Boc-FLFLF and CsH are antagonists and others are agonists. Note that the *N*-formyl group that defines agonistic activities in peptides such as fMLF is replaced with a bulky *t*-butyloxycarbonyl group that defines antagonistic activities in peptides such as *t*-Boc-FLFLF. LXA4, Quin-C1, and compound 43 are highly selective agonists for FPR2/ALX, whereas AG-14 and fMLF are selective for FPR1. *t*-Boc-FLFLF is selective for FPR1 at low concentrations but the selectivity is lost at high micromolar concentrations (e.g., 100  $\mu$ M).

### III. Ligands for the Formyl Peptide Receptor Family of Receptors

Ligand diversity is a prominent and unusual feature of the FPR family of receptors. With the exception of the eicosanoid LXA4, all known FPR family ligands are peptides. More recently, synthetic small-molecular-weight ligands for the formyl peptide receptors have emerged from a number of compound library screens. The structural diversity of these ligands is illustrated with selected agonists and antagonists shown in Fig. 5. The reader is also referred to other recent reviews for a discussion on the agonists and antagonists of FPRs (Le et al., 2001b, 2002; Migeotte et al., 2006). In this article, agonists for the FPRs are listed in separate groups in Table 5, based on their chemical constituents and origins. Whereas many of the agonists for FPRs are identified and purified from living organisms (marked by asterisks in Table 5), a number of peptides are synthesized based on the sequences of known proteins of microbe and host origins. Whether these peptides are present in vivo and have physiological functions has yet to be determined.

#### A. Agonists for the Formyl Peptide Receptor Family of Receptors

1. *N*-Formyl Peptides. The *E. coli*-derived tripeptide fMLF is the most widely used chemotactic peptide for several reasons. It was one of the first characterized synthetic chemotactic peptides and has been extensively studied by the phagocyte research community since its initial discovery in 1975. fMLF is the smallest formyl peptide that displays full agonistic activities. Its potency and efficacy in activating major bactericidal functions of neutrophils equals that of C5a, yet fMLF is readily available from most peptide companies at a fraction of the cost of C5a. fMLF can be easily radiolabeled with tritium, and the radiolabeled fMLF serves as an excellent ligand in direct binding assays.<sup>2</sup>

<sup>2</sup> In early reports and some recent publications, the abbreviation "fMLP" is used for fMet-Leu-Phe. This abbreviation is obsolete and can be confusing, because the letter "P" stands for proline in the now widely used one-letter amino acid code, not phenylalanine, as in fMLF. Therefore, fMLF should be the correct abbreviation for fMet-Leu-Phe.

TABLE 5

## Agonists for the human FPRs

The agonists are listed in the order of their potency within each group. The mitochondrial *N*-formylated peptides, listed in the first group, are also host-derived peptides. A more detailed list of *N*-formyl peptides is given in Table 2. Ligands that have been isolated from living organisms in the forms listed, and those generated by the actions of physiologically relevant enzymes, are indicated with an asterisk (\*).

Ligand	Origin/Description	Potency	Selectivity	Reference
<i>N</i> -formyl peptides fMLF and other bacterial formyl peptides*	Bacteria	(see Table 2)	FPR1 > FPR2/ALX	(see Table 2)
Mitochondrial formyl peptides*	Mitochondria	(see Table 2)	FPR1 ≈ FPR2/ALX	(see Table 2)
<i>N</i> -formyl-Nle-Leu-Phe-Nle-Tyr-Lys	Synthetic	p <i>K</i> <sub>d</sub> = 9.22	FPR1 >> FPR2/ALX	Sklar et al. (1984)
Microbe-derived nonformyl peptides				
T20 (DP178)	HIV-1 gp41 aa. 643–678	pEC <sub>50</sub> = 8.30	FPR1	Su et al. (1999c)
Hp (2–20)	<i>H. pylori</i>	pEC <sub>50</sub> = 6.52	FPR2/ALX >> FPR3	Betten et al. (2001)
T21 (DP107)	HIV-1 gp41 aa. 558–595	pEC <sub>50</sub> = 6.30	FPR2/ALX	Su et al. (1999a)
V3 peptide	HIV-1 gp120, V3 loop	pEC <sub>50</sub> = 5.82	FPR2/ALX	Shen et al. (2000)
N36 peptide	HIV-1 gp41 aa. 546–581	pEC <sub>50</sub> = 5.00	FPR2/ALX	Le et al. (2000b)
F peptide	HIV-1 gp120 aa. 414–434	pEC <sub>50</sub> = 5.00	FPR2/ALX	Deng et al. (1999)
Host-derived peptides				
CK68-1 (human CCL23)*	Chemokine	pEC <sub>50</sub> = 9.00–7.82	FPR2/ALX ≈ CCR1	Elagöz et al. (2004)
SHAAptide*	CCL23 <i>N</i> -terminal 18 aa.	pEC <sub>50</sub> = 7.72	FPR2/ALX > CCR1	Miao et al. (2007)
Humanin*	Neuroprotective peptide	pEC <sub>50</sub> = 8.46	FPR2/ALX	Harada et al. (2004); Ying et al. (2004)
F2L*	Heme binding protein	pEC <sub>50</sub> = 8.00	FPR3 >> FPR2/ALX	Migeotte et al. (2005)
SAA*	Acute-phase protein	pEC <sub>50</sub> = 7.35	FPR2/ALX, others	Su et al. (1999b)
Annexin 1 / lipocortin 1*	Annexin 1	pIC <sub>50</sub> = 6.82	FPR1	Walther et al. (2000)
Ac2–26*	Annexin 1	pEC <sub>50</sub> = 6.05–5.77	FPR1, FPR2/ALX	Perretti et al. (2002); Hayhoe et al. (2006)
Ac9–25	Amyloid precursor	pEC <sub>50</sub> = 4.70	FPR1, FPR2/ALX	Karlsson et al. (2005)
AB (1–42)*	uPAR (88–274)	pEC <sub>50</sub> = 7.00	FPR2/ALX	Le et al. (2001a); Tiffany et al. (2001)
D2D3*	Cathelicidin	pEC <sub>50</sub> = 7.08	FPR2/ALX	Resnati et al. (2002)
LL-37*	Prion protein	pEC <sub>50</sub> = 6.00	FPR2/ALX	Yang et al. (2000)
PrP (106–126)*	Anti-microbial peptide	pEC <sub>50</sub> = 4.60	FPR2/ALX	Le et al. (2001b)
Temporin (from <i>Rana temporaria</i> )*	Pituitary adenylate cyclase activating polypeptide	pEC <sub>50</sub> = 6.60	FPR2/ALX	Chen et al. (2004)
Host-derived nonpeptide agonists				
Lipoxin A4 and aspirin-triggered lipoxins*	Eicosanoids	p <i>K</i> <sub>d</sub> = 8.77	FPR2/ALX, Ahr	Kim et al. (2006)
Agonists from peptide library				
WKYMVm	Peptide library	pEC <sub>50</sub> = 10.13	FPR2/ALX > FPR >> FPR3	Le et al. (1999); Christophe et al. (2001)
WKYMVM	Peptide library	pEC <sub>50</sub> = 8.70	FPR2/ALX >> FPR3	Christophe et al. (2001)
MMK-1	Peptide library	pEC <sub>50</sub> = 8.70	FPR2/ALX	Klein et al. (1998); Hu et al. (2001)
MMWLL, formyl-MMWLL	Peptide library	pEC <sub>50</sub> = 8.96	FPR1	Chen et al. (1995)
Agonists from nonpeptide library				
Quinazolinone derivative (Quin-C1)	Combinatorial library	pEC <sub>50</sub> = 5.72	FPR2/ALX >> FPR1	Nanamori et al. (2004)
Pyrazolone, 4-iodo-substituted, no. 43	Combinatorial library	pIC <sub>50</sub> = 7.36	FPR2/ALX >> FPR1	Büri et al. (2006)
AG-14	Drug-like molecule library	pEC <sub>50</sub> = 7.38	FPR1	Schepletkin et al. (2007)

aa., amino acid; pIC<sub>50</sub>, negative logarithm of the IC<sub>50</sub>; pEC<sub>50</sub>, negative logarithm of the EC<sub>50</sub>; p*K*<sub>d</sub>, negative logarithm of *K*<sub>d</sub>.

Whereas fMLF is by far the most frequently used chemotactic peptide in studies of neutrophil functions, this prototypic formyl peptide should not be taken as the sole standard in judging the presence of functional formyl peptide receptors. Because bacterial protein synthesis starts with an *N*-formyl methionine, formyl peptides released from bacteria can be considered a type of microbe-associated molecular pattern, recognizable by specialized receptors in the innate immune cells of the host, such as the Toll-like receptors (TLRs). These non-rearranging innate receptors have evolved to aid the host in detecting nonself such as bacterial products (Medzhitov and Janeway, 2000). Ample evidence shows that the formyl peptide receptors can detect not only the *E. coli*-derived fMLF but also formyl peptides from other bacteria strains and from mitochondria of the host cells. Table 2 lists selected bacterial and mitochondrial formyl peptides that have been characterized for their bioactivities.

Given the variety of formyl peptides from both bacteria and mitochondria, it is worthwhile to revisit some previous studies conducted with the use of fMLF and determine whether the receptors of interest actually are more selective for formyl peptides of different sequences. For instance, FPR2/ALX was first identified as a low affinity receptor for fMLF ( $K_d = 430$  nM), raising the question of whether its true ligand is a formyl peptide. Rabiet et al. (2005) recently conducted an extensive survey of formyl peptides from various sources, including bacteria and mitochondria of mammalian cells (Rabiet et al., 2005). The results from this study demonstrate the ability of FPR2/ALX to respond to mitochondria-derived formyl peptides (fMYFINILTL, fMLKLIV, and fMMYALF) in calcium mobilization assays, with  $EC_{50}$  values of 10 to 160 nM, showing that these peptides are similarly potent on both FPR1 and FPR2/ALX. Moreover, the study also demonstrates the ability of FPR2/ALX to detect and respond to several *L. monocytogenes*-derived formyl peptides, including fMIVIL (Gulden et al., 1996). These results indicate that FPR2/ALX is able to discriminate between *N*-formyl peptides of different sizes, hydrophobicities, and charges. The biological relevance of this property of the receptor is not yet entirely understood.

Two studies discussed in the previous sections provide additional examples that *N*-formyl peptides other than fMLF are more selective for certain receptors. Mouse Fpr1, for example, prefers fMIVIL and fMIFL, peptides derived from *L. monocytogenes* and *S. aureus*, respectively, over the *E. coli*-derived fMLF (Southgate et al., 2008). One of the FPR1 haplotypes tested in the study by Gripenrog et al. showed reduced ability to respond to a *M. avium*-derived formyl peptide but normal response to fMFADRW, a fragment of cytochrome *c* oxidase subunit I (Gripenrog and Miettinen, 2008). These studies demonstrate that although *N*-formyl peptides are a class of ligands representing a pattern recognized by the FPRs,

as has been the case since the initial discoveries by Schiffmann et al. (1975a), there are important differences in potency and receptor selectivity among the individual peptides.

There are other examples in which addition of an *N*-formyl group increases agonistic activity of the peptides. MMWLL (Met-Met-Trp-Leu-Leu), a synthetic peptide isolated from a library screen, becomes more potent at FPR1 with the addition of an *N*-formyl group (Chen et al., 1995). Humanin is an endogenous peptide (MAPRG-FSCLLLLTSEIDLVPVKRRA) with neuroprotective activity (Hashimoto et al., 2001) that also binds to FPR2/ALX and FPR3 (Ying et al., 2004). If humanin is *N*-formylated, it becomes a more potent agonist than nonformylated humanin for these receptors (Harada et al., 2004). In the latter example, although the primary sequences of FPR2/ALX and FPR3 differ considerably from those of FPR1, especially in the ligand binding domains (Mills et al., 1998), these two receptors seem to have retained the ability to preferentially interact with formylated peptides. Whether these receptors mediate the neuroprotective effects of humanin is not yet known.

It has long been hypothesized that *N*-formyl peptides derived from mitochondrial proteins may attract leukocytes to sites of inflammation and tissue damage. Consistent with this, *N*-formylated hexapeptides corresponding to the N terminus of mitochondrial NADH dehydrogenase subunits 4 (fMLKLIV) and 6 (fMMYALF) and cytochrome *c* oxidase subunit I (fMFADRW) are equally potent at FPR1 and FPR2/ALX (Rabiet et al., 2005). fMMYALF is also a low-affinity agonist at FPR3 (Rabiet et al., 2005). A nonformylated peptide fragment (MYFINILTL) derived from mouse NADH dehydrogenase subunit 1 is also an agonist at FPR2/ALX (Chiang et al., 2000). It is not yet known whether these peptides are produced in vivo and whether they modulate inflammation.

**2. Microbe-Derived Nonformyl Peptides.** A cecropin-like peptide (with the sequence AKKVFKRLEKLF-SKIQNDK) from *Helicobacter pylori*, Hp(2–20), was found to attract monocytes and basophils to the gastric mucosa in response to *H. pylori* infection. Hp(2–20) was identified as an agonist at FPR2/ALX and FPR3 (Betten et al., 2001; de Paulis et al., 2004b). Despite the absence of an *N*-formyl group in this case, Hp(2–20) is a full agonist capable of stimulating superoxide production.

HIV-1 envelope proteins contain peptide sequences capable of interacting with either or both FPR1 and FPR2/ALX, including at least three sequences in gp41 and two in gp120. Although T20/DP178 from gp41 (Ac-YTSLIHSLIEESQNQQEKNEQELLELDKWASLWNWF-NH<sub>2</sub>) specifically activates human FPR1 in vitro (Su et al., 1999c) and mouse Fpr1 in vivo (Hartt et al., 2000), T21/DP107 from gp41 (Ac-NLLRAIEAQQHLLQLTVWGKQLQARILAVEERYLKDQ-NH<sub>2</sub>) uses both FPR1 and FPR2/ALX with higher efficacy at FPR2/ALX (Su et al., 1999a). N36 from gp41 (Ac-SGIVQQQNNLL-



RAIEAQQHLLQLTVWGIKQLQARIL-NH<sub>2</sub>), which partially overlaps with T21/DP107, signals only through FPR2/ALX (Le et al., 2000b). Two peptide domains in HIV-1 gp120 are potent chemoattractants and activators for FPR2/ALX, but not for FPR1, in human phagocytic leukocytes. One of the peptide domains, F peptide, consists of 20 amino acids (Ac-EGSDTITLPCRIKQFIN-MWQE-NH<sub>2</sub>) located in the C4-V4 region of gp120 of the HIV-1 Bru strain (Deng et al., 1999). Another peptide (V3 peptide, Ac-RIHIGPGRAFYTTKN-NH<sub>2</sub>) was derived from a linear sequence of the V3 region of the HIV-1 MN strain (Shen et al., 2000). The existence of these peptides *in vivo* and their biological significance are not known at present. The peptide T20/DP178 is a licensed antiretroviral agent (pentafuside) administered subcutaneously that acts at the level of HIV-target cell fusion.

In addition to HIV-1 proteins, other viral proteins contain sequences that can serve as ligands for formyl peptide receptors when tested in the form of synthetic peptides. In Herpes simplex virus type 2, a gG-2p20 peptide corresponding to amino acids 190 to 205 of the secreted glycoprotein sgG-2 (GLLWVEVGGEGPGPT) activates neutrophils and monocytes via FPR1 (Bellner et al., 2005). The gG-2p20-induced activation of phagocytes releases reactive oxygen species that inhibits NK cell cytotoxicity and accelerates apoptotic cell death. Therefore, gG-2p20 peptide may contribute to the reduced function and viability of NK cells during HSV-2 infection by activating phagocytic cells. In a more recent study, Mills (2006) identified additional peptides from coronavirus 229E and Ebola virus as FPR1 ligands. Most of these peptides are antagonists, including the coronavirus 229E peptide MYVKWPWYVWL. It is noteworthy that an *N*-formylated form of this peptide becomes a potent agonist for FPR1. It is important to determine in future studies 1) whether and how these peptides are generated during viral infection and 2) the functional consequences of phagocyte response to these peptides.

**3. Host-Derived Peptides.** In addition to mitochondrial formyl peptides discussed above, a large number of endogenous peptides of various compositions, often without an *N*-formyl group, have been identified as agonists for the formyl peptide receptors, especially FPR2/ALX (Table 5). Of particular interest are peptides associated with amyloidogenic diseases, peptides associated with inflammatory and antibacterial responses, and a peptide derived from heme-binding protein that serves as a potent endogenous agonist at FPR3.

**a. Peptides Associated with Amyloidogenic Diseases.** At least three amyloidogenic polypeptides, associated with chronic inflammation and amyloidosis, have been identified as agonists for FPR2/ALX. Serum amyloid A (SAA) is an acute-phase protein the serum concentration of which is increased by as much as 1000-fold in response to trauma, acute infection, and other envi-

ronmental stress causing acute-phase responses (Kushner and Rzewnicki, 1999). Studies with recombinant human SAA identified it as the first mammalian host-derived chemotactic peptide ligand for FPR2/ALX (Su et al., 1999b). SAA, acting through FPR2/ALX, is chemotactic for monocytes, neutrophils, mast cells, and T lymphocytes; stimulates production of metalloproteases and cytokines; and increases expression of cytokine receptors. In neutrophils, SAA activates FPR2/ALX and induces CXCL8 (IL-8) secretion (He et al., 2003). In monocytes, SAA shows a peculiar pattern of cytokine induction via FPR2/ALX; i.e., the cells respond to low concentrations of SAA by producing TNF- $\alpha$  while releasing IL-10 in response to high concentrations of SAA (Lee et al., 2006a). The synovial tissues of patients with inflammatory arthritis express high levels of SAA and FPR2/ALX, and SAA induces the expression of matrix metalloproteinase-1 and -3 in fibroblast-like synoviocytes (O'Hara et al., 2004; Sodin-Semrl et al., 2004). Furthermore, SAA promotes synovial hyperplasia and angiogenesis through activation of FPR2/ALX (Lee et al., 2006b). In addition to using the formyl peptide receptors, SAA activates neutrophil NADPH oxidase through a different receptor (Björkman et al., 2008). More recent studies have identified additional receptors for SAA, including the class B type I scavenger receptor (mouse) and its human ortholog CLA-1 (CD36 and LIM-PII Analogous-1) (Baranova et al., 2005; Cai et al., 2005), the receptor for advanced glycation end product (RAGE) (Cai et al., 2007), Toll-like receptor 2 (Cheng et al., 2008), and possibly Toll-like receptor 4 (Sandri et al., 2008). These SAA receptors are involved in SAA regulation of cholesterol metabolism and production of selected cytokines.

Another peptide, the 42-amino acid form of  $\beta$  amyloid peptide (A $\beta$ <sub>42</sub>, DAEFRHDSGYEVHHQKLVFFAEDVGSNKGAIIGLMVGGVVIA), which is a cleavage product of the amyloid precursor protein in the brain and a pathologic protein in Alzheimer's disease, was also found to activate FPR2/ALX (Le et al., 2001a; Tiffany et al., 2001). An additional amyloidogenic disease-associated FPR2/ALX agonist is prion protein fragment PrP(106–126), which is produced in human brains with prion disease (Le et al., 2001c). FPR2/ALX mediates the migration and activation of monocytic phagocytes, including macrophages and brain microglia, induced by  $\beta$  amyloid (A $\beta$ ) (Le et al., 2001a; Chen et al., 2006). Moreover, FPR2/ALX promotes the endocytosis of A $\beta$  by macrophages and microglia *in vitro* in the form of receptor and ligand complexes (Yazawa et al., 2001; Chen et al., 2006). If the exposure of macrophages to A $\beta$ <sub>42</sub> is transient, the internalized A $\beta$ <sub>42</sub> is degraded, and FPR2/ALX is rapidly recycled back to the cell surface. But prolonged exposure results in accumulation of the A $\beta$ <sub>42</sub> and FPR2/ALX complex in macrophages, culminating in progressive fibrillary aggregation of A $\beta$ <sub>42</sub> and macrophage death (Yazawa et al., 2001). Therefore, FPR2/ALX not

only mediates the proinflammatory activity of the peptide agonists associated with amyloidogenic diseases, it may also participate in the regulation of fibrillary peptide formation and deposition, which are pathologic features of the diseases that contribute to tissue and organ destruction (Cui et al., 2002). The *in vivo* significance of this to the pathogenesis of Alzheimer's disease is not yet known.

Humanin is a peptide encoded by a cDNA cloned from a relatively healthy region of an Alzheimer's disease brain (Hashimoto et al., 2001). Both secreted and synthetic humanin peptides protect neuronal cells from damage by  $A\beta_{42}$ . Humanin uses FPR2/ALX, FPR3 and mouse Fpr2 as functional receptors to induce chemotaxis of mononuclear phagocytes (Harada et al., 2004; Ying et al., 2004). In addition, humanin reduces aggregation and fibrillary formation by suppressing the interaction of  $A\beta_{42}$  with mononuclear phagocytes through FPR2/ALX. Human neuroblastoma cell lines express functional FPR2/ALX but not FPR1. In these cells, although humanin and  $A\beta_{42}$  both activate FPR2/ALX, only  $A\beta_{42}$  causes apoptotic death of the cells, a process blocked by humanin. These observations suggest that humanin may exert its neuroprotective effects by competitively inhibiting the access of  $A\beta_{42}$  (Ying et al., 2004).

*b. Peptides Associated with Inflammatory and Antibacterial Responses.* Urokinase-type plasminogen activator (uPA) is a serine protease known for its ability to regulate fibrinolysis. uPA binds to a specific high affinity surface receptor (uPAR). In addition to regulating fibrinolysis, the uPA-uPAR system is crucial for cell adhesion and migration and tissue repair. uPAR contains three domains: D1 (binding uPA), D2, and D3. A D3-bound glycosylphosphatidylinositol anchor links uPAR to the plasma membrane. uPAR can be cleaved by different proteases, including uPA, in the D1-D2 linker region. The cleaved soluble uPAR D2D3<sub>88-274</sub> binds and activates FPR2/ALX in monocytes, inducing cell migration (Resnati et al., 2002). The ability of cleaved soluble uPAR (c-suPAR) to activate other members of the FPR family has been reported. For instance, SRSRY, a peptide corresponding to residues 88 to 92 of uPAR, binds to and activates FPR1 (Gargiulo et al., 2005). uPAR<sub>84-95</sub> induces basophil migration by activating both FPR2/ALX and FPR3 (de Paulis et al., 2004a). Recent studies show that uPAR is involved in granulocyte-colony-stimulating factor (G-CSF)-induced mobilization of CD34<sup>+</sup> hematopoietic stem cells. G-CSF increases the expression of uPAR on circulating CD34<sup>+</sup> and CD14<sup>+</sup> cells. This is associated with increased uPAR shedding, leading to the appearance of serum c-suPAR. c-suPAR mobilizes hematopoietic stem cells by promoting their FPR1-mediated migration and by desensitization of the chemokine receptor CXCR4 (Selleri et al., 2005). Additional studies showed that pretreatment of monocytes with the FPR2/ALX agonist D2D3<sub>88-274</sub> markedly decreases chemokine-induced integrin-dependent rapid

cell adhesion (Furlan et al., 2004), indicating that the FPR family receptors regulate leukocyte chemotaxis at multiple levels, i.e., in addition to being direct mediators of cell migration, they may suppress cell responses to chemokines by desensitizing chemokine receptors.

Formyl peptide receptors interact with several bactericidal peptides contained in human neutrophil granules. LL-37 (LLGDFFRKSKEKIGKEFKRIVQRIKDFLRNLPRTES), an enzymatic cleavage fragment of the neutrophil granule protein cathelicidin (Yang et al., 2000), and its mouse homolog CRAMP (GLLRKGGKIGKGLKIKGNFFQKLVPQPEQ) (Kurosaka et al., 2005) are agonists for FPR2/ALX. LL-37 is expressed by leukocytes and epithelial cells and secreted into wounds and onto the airway surface. In addition to its microbicidal activity, LL-37 induces directional migration of human monocytes, neutrophils, and T lymphocytes, a function mediated by FPR2/ALX (Yang et al., 2000). Recent studies showed that LL-37-induced angiogenesis is mediated by FPR2/ALX in vascular endothelial cells. Decreased vascularization during wound repair observed in mice deficient for CRAMP indicates that cathelicidin-mediated angiogenesis is important for cutaneous wound neovascularization *in vivo* (Koczulla et al., 2003). LL-37 seems to be a multifunctional peptide with a central role in innate immunity against bacterial infection and in the induction of arteriogenesis important for angiogenesis. Another antibacterial granule protein, cathepsin G, which is a serine protease and participates in wound healing, is identified as a specific agonist for FPR1 (Sun et al., 2004).

FPR2/ALX is also known to interact with a chemokine variant that potently activates phagocytic leukocytes. CCL23/MPIF-1 (CK $\beta$ 8-1) uses a typical G protein-coupled receptor CCR1 for its leukocyte chemotactic activity. However, an N-terminally truncated form of the CCL23 splice variant CCL23 $\beta$  (amino acids 22-137) activates myeloid cells and FPR2/ALX transfected cell lines at low nanomolar concentration range, making it one of the most potent FPR2/ALX agonists identified so far (Elagoz et al., 2004). A more recent study identified a mechanism of CCL23 $\beta$  processing that involves sequential cleavage of the chemokine *in vitro* with proinflammatory proteases, generating CCR1-specific chemokine and an 18-amino acid peptide, termed SHAAGtide (MLWRRKIGPQMTLSHAAG), that activates FPR2/ALX at nanomolar concentrations (Miao et al., 2007). This study illustrates a novel mechanism by which protease cleavage of a chemokine produces two peptides acting on two different receptors. It will be interesting to determine whether the cleavage product is generated *in vivo* and how the simultaneous activation of two receptors influences the course of inflammation.

*c. Annexin A1 and Its N-terminal Peptides.* ANXA1 and its N-terminal peptides have interesting properties in activating formyl peptide receptors by playing dual roles in inflammatory host responses. ANXA1 (also

termed lipocortin I) is a glucocorticoid-regulated, phospholipid-binding protein of 37 kDa that possesses both pro- and anti-inflammatory activity, mediated in part by activating FPR1 (Ernst et al., 2004a). Expressed in a variety of cell types, ANXA1 is particularly abundant in neutrophils. The protein is primarily cytosolic, but it may also be secreted through a nonclassic secretory process and found on the outer cell surface, causing leukocyte detachment and thereby inhibiting their transendothelial migration (for review, see Perretti, 2003). At low concentrations, both ANXA1 holoprotein and its N-terminal peptides (Ac2–26, Ac-AMVSEFLKQAWFIENEQEYVQTVK, and Ac9–25) elicit  $\text{Ca}^{2+}$  transients through FPR1 without fully activating the MAPK pathway (Ernst et al., 2004a), causing neutrophil desensitization and inhibition of transendothelial migration induced by other chemoattractants such as the chemokine IL-8 (CXCL8). In contrast, at high concentrations, the ANXA1 peptides fully activate neutrophils *in vitro* and become potent proinflammatory stimulants. The antimigratory activity of exogenous and endogenous ANXA1 has been shown in both acute and chronic models of inflammation (Perretti, 2003). *Fpr1* knockout mice exhibit normal neutrophil accumulation during thioglycolate-elicited peritonitis (Perretti et al., 2001). However, a significant part of the effect in reducing intraperitoneal neutrophil infiltration, observed in ANXA1-treated wild type mice, was abolished in the *Fpr1*( $-/-$ ) mice. Other studies have shown that the ANXA1 N-terminal peptides use FPR2/ALX for its anti-inflammatory actions (Perretti et al., 2002). The peptides are also ligands for FPR3 (Ernst et al., 2004a). The ANXA1 core-derived peptide antiinflammin 2 (amino acids 246–254; HDMNKVDK) activates FPR2/ALX (Kamal et al., 2006). The utilization of the formyl peptide receptors by ANXA1 and its amino terminal peptides for their various functions is a complex issue. One published report demonstrates that Ac9–25 stimulates neutrophil NADPH oxidase activation through FPR1, but its inhibitory effect is mediated through a receptor other than FPR1 or FPR2/ALX (Karlsson et al., 2005), suggesting the presence of additional receptors for ANXA1 and its peptides. ANXA1 has been shown to bind to the  $\alpha 4\beta 1$  integrin on undifferentiated U937 cells that do not express FPR1 or FPR2/ALX and inhibits U937 adhesion to microvascular endothelial cells (Solito et al., 2000).

*d. F2L, a Potent Endogenous Agonist for FPR3.* In 2005, Migeotte et al. (2005) reported the purification of a highly potent and efficacious agonist peptide for human FPR3, termed F2L. F2L is an amino-terminally acetylated peptide (Ac-MLGMIKNSLFFGSVETWPWQVL-NH<sub>2</sub>) resulting from the natural cleavage of human heme-binding protein, an intracellular tetrapyrrole-binding protein. The peptide binds and activates FPR3 in the low nanomolar range, triggering typical G protein-mediated intracellular calcium release, inhibition of cAMP accumulation, and phosphorylation of the ERK 1/2

MAPKs. F2L also chemoattracts and activates monocyte-derived DCs. Thus, F2L seems to be a novel and unique natural chemotactic peptide for FPR3 in DCs and monocytes, in agreement with the selective expression of FPR3 in these cells (Yang et al., 2002). F2L may play a role in linking innate and adaptive immune responses by activating antigen-presenting FPR3<sup>+</sup> DCs, which express little FPR1 and FPR2/ALX. It is interesting to note that in mice, F2L activates the FPR2/ALX homolog mFpr2 (Gao et al., 2007). In bone marrow-derived neutrophils deficient in *Fpr2*, the activity of F2L was totally lost, suggesting that in mice this function of human FPR3 is carried out by *Fpr2*. The overlapping function of mouse *Fpr2* with human FPR3 in binding F2L provides another example for the complexity in studying the FPR family receptors using genetically altered mice.

*e. Other Host-Derived Peptides.* Pituitary adenylate cyclase-activating polypeptide 27 (HSDGIFTDSYSRFRKQMAVKKYLA AVL-NH<sub>2</sub>) is an agonist for FPR2/ALX, stimulating chemotaxis, up-regulation of CD11b, and activation of phagocytes (Kim et al., 2006). Temporin A (TA; FLPLIGRVLSGIL-NH<sub>2</sub>) is a frog-derived antimicrobial peptide found to induce the migration of human monocytes, macrophages, and neutrophils (Chen et al., 2004). Characterization of the signaling properties of TA in monocytes and the use of receptor-transfected human embryonic kidney (HEK) 293 cells revealed that this peptide uses FPR2/ALX as a receptor. TA is also chemotactic *in vivo* because it elicited infiltration of neutrophils and monocytes into the injection site in mice. Another temporin peptide, Ranatuerin-6 (Rana-6; FISAIASMLGKFL) also uses FPR2/ALX for chemotaxis of human phagocytes (Chen et al., 2004). The biological and evolutionary significance of these findings is still unclear because, so far, formyl peptide receptors have not been characterized in frogs, and TA homologs have not been characterized in mammals.

*4. Host-Derived, Nonpeptide Agonists.* LXA4 (5S,6R,15S-trihydroxy-7,9,13-*trans*-11-eicosatetraenoic acid) is a potent mediator biosynthesized from arachidonic acid. It is a small molecule with physical chemical properties that differ from most lipids: it has a unique structure and belongs to a class of conjugated tetraene-containing eicosanoids that display stereoselective and highly potent anti-inflammatory and pro-resolving activity *in vivo* in many mammalian systems (for review, see Serhan, 2005) and articles contained in the special issue). In this regard, LXA4 is unusual in that most other eicosanoids are pro-inflammatory. As an endogenous mediator, LXA4 displays multilevel control of processes relevant in acute inflammation via specific and selective actions on multiple cell types via specific receptors (Serhan, 2007). In particular, LXA4 has been reported to interact directly with both human FPR2/ALX and CysLT<sub>1</sub>. It also induces signals that regulate BLT1, and production of chemokines, cytokines (e.g., TNF), and

growth factor receptors (e.g., vascular endothelial growth factor receptor) in human leukocytes, vascular cell types, and mucosal epithelial cells, each contributing to regulate the resolution of inflammation (Serhan, 2005).

LXA4 is the first identified endogenous ligand for FPR2/ALX. Using freshly isolated radioligand for each experiment, specific LXA4 binding sites were initially characterized with isolated human neutrophils (Fiore et al., 1992) and demonstrated to be responsible for the specific LXA4 functions on neutrophils. LXA4 stimulates rapid (within seconds) phospholipase activation in these cells that directly correlates with the induction time course of specific LXA4 binding (Nigam et al., 1990; Fiore et al., 1993). In parallel experiments, a second approach was undertaken with GPCRs that were known to be induced within the same time frame in the promyelocytic cell line HL-60 when differentiated into neutrophils by treatment with either retinoic acid or dimethylsulfoxide. These cells were screened for [<sup>3</sup>H]LXA4 binding and ligand-dependent increases in GTPase activity. Labeled LXA4 gives high-affinity binding with FPR2/ALX as well as ligand selectivity, compared with other eicosanoids including LXB4, leukotriene B4, leukotriene D4, and PGE2. FPR2/ALX expressed in Chinese hamster ovary cells transfected with the FPR2 cDNA pINF114 gave a  $K_d$  of  $\sim 1.7$  nM for [<sup>3</sup>H]LXA4 determined by Scatchard plot analysis (Fiore et al., 1994). This value for recombinant FPR2/ALX is comparable with those obtained for endogenous LXA4 specific binding sites present on peripheral blood neutrophils (e.g.,  $K_d$  values of 0.7 and 0.8 nM obtained with isolated plasma membrane fractions and granule membrane-enriched fractions, respectively). In Chinese hamster ovary cells expressing the recombinant receptor, LXA4 stimulates both GTPase activity and the release of esterified arachidonic acid, which is inhibited by pertussis toxin (Fiore et al., 1994).

Each of the actions of LXA4 proved to be stereoselective in that changes in potencies are associated with double bond isomerization and alcohol chirality (*R* or *S*) as well as dehydrogenation of alcohols and reduction of double bonds. Elimination of the carbon 15 position alcohol from LXA4, denoted 15-deoxy-LXA4, is essentially inactive in vivo and does not stop either neutrophil transmigration or reduce adhesion (Serhan et al., 1995). In nature, LXA4 is enzymatically inactivated by conversion to 15-oxo-LXA4 and 13,14-dihydro-LXA4 (Serhan et al., 1993). The biologically inactive metabolic products and synthetic compounds tested, including 15-oxo-LXA4, 15-deoxy-LXA4, 11-*trans*-LXA4, and 13,14-dihydro-LXA4, do not effectively bind to FPR2/ALX. This contrasts with the active lipoxin ligands that include native LXA4, which carries a 15*S* configuration alcohol that originates from the lipoxygenase biosynthetic pathway and the aspirin-triggered 15*R*- or 15-*epi*-LXA4 (also denoted ATL or aspirin-triggered LXA4) biosynthesized

in the presence of aspirin, each exhibiting stereospecific receptor binding to both human FPR2/ALX and the murine counterpart encoded by *Fpr-rs1* (Takano et al., 1997).

Evidence supporting the anti-inflammatory and pro-resolving functions of LXA4 comes from a number of in vivo studies, as summarized in several recent reviews (Serhan, 2005; Chiang et al., 2006; O'Meara et al., 2008). These in vivo studies have shown that, in nanogram amounts, LXA4 stops neutrophil infiltration (Takano et al., 1997, 1998; Hachicha et al., 1999) and blocks human neutrophil transmigration across mucosal epithelial cells and vascular endothelial cells (Colgan et al., 1993; Kucharzik et al., 2003). One of the mechanisms by which LXA4 and ATLs inhibit neutrophil infiltration is through induction of NO production, which suppresses leukocyte-endothelial cell interaction (Paul-Clark et al., 2004). LXA4 treatment inhibits proinflammatory cytokine production by synoviocytes (Sodin-Semrl et al., 2000), intestinal epithelial cells (Goh et al., 2001; Kucharzik et al., 2003), and bronchial epithelial cells (Bonnans et al., 2007). Several studies have shown that LXA4 inhibits TNF $\alpha$ -induced production of cytokines such as IL-1 $\beta$  and IL-6 (Wu et al., 2005) and chemokines such as IL-8 (Bonnans et al., 2007), as well as LPS-induced secretion of IL-1 $\beta$ , IL-6, and IL-8 (Wu et al., 2008). Dendritic cell production of the immunomodulatory cytokine IL-12 is also regulated by LXA4, as shown in a model of *Toxoplasma gondii* infection (Aliberti et al., 2002a,b). Part of the anti-inflammatory effect of LXA4 is known to involve the inhibition of transcription factors, including NF- $\kappa$ B and activator protein 1, that are responsible for the expression of many proinflammatory cytokines and chemokines (Gewirtz et al., 2002; József et al., 2002; Sodin-Semrl et al., 2004; Wu et al., 2008). Whereas LXA4 suppresses the expression of these proinflammatory cytokines, it stimulates the expression of the anti-inflammatory molecule IL-10 (Souza et al., 2007) and the antioxidant molecule heme oxygenase-1 (Nascimento-Silva et al., 2005). In synovial fibroblasts, LXA4 inhibits the expression of matrix metalloproteinase-3 but enhances the synthesis of tissue inhibitors of metalloproteinases (Sodin-Semrl et al., 2000). LXA4 has also shown antifibrotic effects as it regulates PDGF-induced genes such as TGF- $\beta$  in renal mesangial cells (Rodgers et al., 2005). In addition to its inhibition of proinflammatory cytokine production, LXA4 plays a role in regulating inflammation-induced pain (Svensson et al., 2007). This function is probably mediated through alteration of spinal nociceptive processing through astrocyte activation.

Besides its anti-inflammatory functions, LXA4 and its stable analog exhibit pro-resolution properties (Serhan, 2007). One of the mechanisms involves overriding the apoptosis-delaying effect of proinflammatory factors such as SAA, thereby promoting neutrophil apoptosis (El Kebir et al., 2007). LXA4 also stimulates nonphlo-

gistic phagocytosis of apoptotic neutrophils, which accelerates the clearance of neutrophils and promotes resolution of inflammation (Godson et al., 2000). These examples show that, in general terms, LXA4 actively inhibits many endogenous processes that can amplify local acute inflammation, leading to potent anti-inflammatory as well as pro-resolving actions in vivo.

The multilevel control by LXA4 on processes relevant to acute inflammation raises the intriguing question of how LXA4 binding to FPR2/ALX might translate into the anti-inflammatory and pro-resolving activities and whether other receptors contribute to these activities. Evidence supporting a role of FPR2/ALX in mediating the anti-inflammatory effects of LXA4 came from studies using Boc2, stereoselective analogs, antibodies, and transgenic approaches. Boc2 is a peptide antagonist for both FPR1 and FPR2/ALX as detailed in the following section. Its effect on blocking FPR2/ALX in inflammation models was first reported by Gavins et al. (2003) and confirmed by others (von der Weid et al., 2004; Nascimento-Silva et al., 2005; Machado et al., 2006; Grumbach et al., 2009). Transgenic overexpression of FPR2/ALX resulted in mice that display reduced neutrophil infiltration when challenged with LTB<sub>4</sub> and PGE<sub>2</sub>. The transgene also enhanced the response to aspirin-triggered lipoxins in a mouse model of zymosan-induced peritonitis (Devchand et al., 2003). Binding of LXA4 to FPR2/ALX is stereo-selective, because other hydroxyl-containing eicosanoids, including LXB<sub>4</sub>, LTB<sub>4</sub>, and LXA4 stereoisomers, did not compete for [<sup>3</sup>H]LXA4 binding specifically (Fiore et al., 1992). However, the finding that LXB<sub>4</sub> also exhibits anti-inflammatory functions (Maddox and Serhan, 1996) suggest the possible presence of another functional receptor that differs from FPR2/ALX for the lipoxins. One observed difference between the two receptors is that LXB<sub>4</sub> does not induce calcium mobilization in the monocytic cell line THP-1, whereas LXA4 does (Romano et al., 1996). It is noteworthy that the aryl hydrocarbon receptor (AhR) has also been identified as a functional receptor for LXA4 (Schalch et al., 1999). AhR is not a GPCR and is located intracellularly, whereas FPR2/ALX is found on the plasma membrane. AhR binding of LXA4 renders the receptor capable of interacting with a dioxin responsive element, which can direct expression of a reporter gene with an EC<sub>50</sub> of 50 nM for LXA4. It was recently shown that LXA4 acts on both FPR2/ALX and AhR to exert its anti-inflammatory effect in dendritic cells through induction of suppressor of cytokine signaling-2 (SOCS-2) (Machado et al., 2006, 2008). Whether AhR is a receptor for LXB<sub>4</sub> and the relative contribution of each of these receptors to the anti-inflammatory activities of the lipoxins has yet to be determined.

**5. Agonists from Peptide Library.** Using combinatorial peptide library screens, a number of peptides have been identified as potent agonists for the formyl peptide receptors. Trp-Lys-Tyr-Met-Val-D-Met-NH<sub>2</sub> (WKYMVm),

a hexapeptide representing a modified sequence isolated from a random peptide library, was found to be a highly efficacious stimulant for human B lymphocytes, monocytic cell lines, and blood neutrophils (Baek et al., 1996). WKYMVm uses FPR1, FPR2/ALX, and FPR3 for activation of human phagocytic cells (Le et al., 1999; Kang et al., 2005). WKYMVm is by far the most potent peptide agonist for FPR2/ALX, with an EC<sub>50</sub> well within the picomolar range in chemotaxis assays. WKYMVm, a derivative of WKYMVm with an L-methionine at the carboxyl terminus, is a highly selective agonist for FPR2/ALX and is also a weaker activator of FPR3 (Christophe et al., 2001). A recent study investigated the relationship between FPR1 and FPR2/ALX, both found in neutrophils, in mediating the WKYMVm-stimulated cellular functions. It was found that WKYMVm activates neutrophils through FPR1 only when the signal through FPR2/ALX is blocked (Karlsson et al., 2006), suggesting its preference for one type of receptor despite the presence of two different types of receptors in neutrophils. WKYMVm also inhibits LPS-induced maturation of human monocyte-derived dendritic cells via FPR1 and FPR3, presumably through interference with signaling pathways activated by Toll-like receptor 4 (Kang et al., 2005).

The peptide MMK-1 (LESIFRSLFRVM) was identified from a library screen in genetically engineered *Saccharomyces cerevisiae* cells designed to couple receptor (FPR2/ALX) activation to histidine prototrophy (Klein et al., 1998). Selection for histidine prototrophs among transformants obtained with a plasmid-based random peptide library identified several agonists, each of which yielded autocrine stimulation of the receptor expressed in yeast. MMK-1 was found to be a highly selective chemotactic agonist for FPR2/ALX (Klein et al., 1998; Hu et al., 2001). In another study, a tethered library was screened based on the activation mechanism of the thrombin receptor PAR-1 (Chen et al., 1995). A peptide with the sequence MMWLL was identified as an FPR1 agonist. This peptide becomes 1000 times more potent when *N*-formylated, consistent with the preferential recognition of *N*-formylmethionine-containing peptides by FPR1.

A recent report shows that a peptide library-derived ligand of 21 amino acids, CGEN-855A (TIPM-FVPESTSKLQKFTSWFM-amide), exhibits anti-inflammatory properties (Hecht et al., 2009). CGEN-855A displaced <sup>125</sup>I-WKYMVm binding to FPR2/ALX with an IC<sub>50</sub> value of 189 nM (pIC<sub>50</sub> = 6.72) and a K<sub>i</sub> value of 54.1 nM (pK<sub>i</sub> = 7.27). CGEN-855A does not affect monocyte secretion of cytokines. It has anti-inflammatory activity in the zymosan-triggered air pouch inflammation model, and displays cardioprotection in a mouse model of ischemia/reperfusion-induced myocardial infarction (Hecht et al., 2009). The study indicates that peptides such as ANXA1 N-terminal fragments and CGEN-855A have functional properties similar to those of LXA4.

6. *Agonists from Nonpeptide Library.* Several laboratories have recently identified ligands for the formyl peptide receptors through screening of combinatorial libraries consisting of synthetic, nonpeptide compounds. These synthetic small molecules are of different structures (see Fig. 5) and are highly selective for either FPR1 or FPR2/ALX, providing useful tools for the characterization of formyl peptide receptors. The pharmacological properties of these compounds also make them potentially useful for therapeutic intervention.

Nanamori et al. (2004a) first reported the identification of a quinazolinone derivative named Quin-C1 as a highly selective agonist for FPR2/ALX. Quin-C1 (4-butoxy-*N*-[2-(4-methoxy-phenyl)-4-oxo-1,4-dihydro-2*H*-quinazolin-3-yl]-benzamide) induces chemotaxis and secretion of  $\beta$ -glucuronidase in peripheral blood neutrophils with a potency of approximately 1/1000 that of the peptide agonist WKYMVm, which is the most potent agonist for FPR2/ALX identified so far. In studies using RBL cells expressing either FPR1 or FPR2/ALX, Quin-C1 induced enzyme release only from cells expressing the latter. Quin-C1 selectively stimulates calcium mobilization in RBL cells expressing recombinant FPR2/ALX, induces phosphorylation of ERK1/2, and promotes internalization of FPR2/ALX fused to an enhanced green fluorescent protein. However, unlike most peptide agonists of FPR2/ALX, Quin-C1 did not induce substantial neutrophil superoxide generation, even at concentrations up to 100  $\mu$ M. The structural basis for this biased agonistic activity is still unknown but may be beneficial for its potential use as a therapeutic agent.

Bürli et al. (2006) characterized and subsequently modified a series of compounds initially identified from a cell-based assay for high-throughput screening. These ligands are also highly selective for FPR2/ALX and exhibit anti-inflammatory properties in a mouse ear swelling assay. Two of the pyrazolones (compounds 24 and 43) tested in their study were able to stimulate calcium flux in transfected cells expressing FPR2/ALX, whereas other structural analogs were ineffective. Compound 43 was formulated for oral administration and was found to significantly reduce inflammation in an ear swelling model in mice.

Using a strategy combining computer model-based virtual screening and high-throughput, no-wash cytometry screening, Edwards et al. (2005) identified 30 lead compounds (from an initial pool of 480,000) that are partial agonists or antagonists for FPR1. The *in silico* virtual screening, based on a bovine rhodopsin crystal structure, effectively boosted the physical screen hit rate by 12-fold, eliminating most of the compounds before physical screen. The pharmacophore model for FPR1 developed in this study may be useful in future identification of agonists and antagonists for this class of receptors.

Schepetkin et al. (2007) used a different approach for compound screening. They started in a neutrophil su-

peroxide production assay, combined with substructure screen, fragment focusing, and structure-activity relationship analyses to identify *t*-butyl benzene and thiophene-2-amide-3-carboxylic ester derivatives as potential agonists for neutrophil chemoattractant receptors. Eleven of the compounds were further analyzed in functional assays and one (AG-14) was found to activate neutrophils at nanomolar concentrations. Based on desensitization and antagonist inhibition data, the investigators concluded that AG-14 is an agonist for FPR1. These investigators also identified arylcarboxylic acid hydrazide derivatives as agonists for FPR2/ALX that induce *de novo* production of TNF $\alpha$  through activation of macrophages (Schepetkin et al., 2008).

### B. Antagonists for the Formyl Peptide Receptor Family of Receptors

Early studies showed that replacing the formyl group of fMLF with tertiary butyloxycarbonyl group (*t*-Boc) renders the peptide antagonistic (Freer et al., 1980). *t*-Boc-Met-Leu-Phe (Boc1) and *t*-Boc-Phe-D-Leu-Phe-D-Leu-Phe (Boc2) are two frequently used antagonists for FPR1, with pIC<sub>50</sub> values of 6.19 and 6.59, respectively (Freer et al., 1980) (Table 6). The *N*-ureido-substituted Phe-D-Leu-Phe-D-Leu-Phe was found to also have antagonistic activity (Higgins et al., 1996). It is noteworthy that whereas *t*-Boc-Phe-D-Leu-Phe-D-Leu-Phe-OMe is a full antagonist, the peptide with the *l*-Leu, *t*-Boc-Phe-Leu-Phe-Leu-Phe-OMe, has agonistic activity (Toniolo et al., 1990). In several recent studies, Boc2 was used at high concentrations (e.g., 100  $\mu$ M) for inhibition of FPR2/ALX (Gavins et al., 2003; Machado et al., 2006). A recent study has shown that, when used at low micromolar concentrations, both Boc1 and Boc2 are selective antagonists for FPR1; at high micromolar concentrations, Boc2 partially inhibits FPR2/ALX in addition to FPR1 (Stenfeldt et al., 2007). Therefore, the antagonistic effect of Boc2 at high concentrations is not specific for FPR2/ALX. In another ligand screen, the peptide WRW-WWW was identified as a highly selective antagonist for FPR2/ALX (Bae et al., 2004), with a pIC<sub>50</sub> of 6.64 in calcium flux assay. As an antagonist, WRWWW is more selective at FPR2/ALX than Boc2 (Stenfeldt et al., 2007).

Cyclosporin H (CsH) is a cyclic undecapeptide produced by fungi. It is an optical isomer of the immunosuppressant CsA and contains at its position 11 the amino acid residue *N*-methyl-D-valine, in place of the *N*-methyl-L-valine as seen in CsA (von Wartburg and Traber, 1986). CsH lacks immunosuppressive activity but displays selective antagonistic activity at human FPR1 (Wenzel-Seifert et al., 1991). Studies have shown that CsH is 14-fold more potent than the tertiary butyloxycarbonyl analogs of formyl peptides such as Boc2 in FPR1 binding assays, and approximately 5-fold more potent than Boc2 in the inhibition of fMLF-induced calcium flux and enzyme release (Wenzel-Seifert and Seif-

TABLE 6  
Antagonists for the human formyl peptide receptors

Antagonists for the FPRs are listed in the order of their approximate potency, except that antagonists of same types are listed together.

Ligand	Assay	Potency	Selectivity	References
Chemotaxis inhibitory protein of <i>S. aureus</i> (CHIPS)	Binding	$pK_d = 7.46$	FPR1	Haas et al. (2004)
FPRL1-inhibitor protein (FLIPr)	Binding, $Ca^{2+}$ flux	N.D.	FPR2/ALX $\gg$ FPR1	Prat et al. (2006)
Trp-Arg-Trp-Trp-Trp-Trp (WRW <sup>4</sup> )	$Ca^{2+}$ flux	$pIC_{50} = 6.64$	FPR2/ALX $\gg$ FPR1 $\approx$ FPR3	Bae et al. (2004)
CsH	Binding	$pK_i = 7.00$	FPR1	Wenzel-Seifert et al. (1991)
CsA	Enzyme release	$pK_i = 6.22$	FPR1	Yan et al. (2006)
<i>i</i> -Boc-Met-Leu-Phe	$O_2^-$ generation	$pIC_{50} = 6.60$	FPR1	Derian et al. (1996)
<i>t</i> -Boc-Met-Leu-Phe	Enzyme release	$pIC_{50} = 6.19$	FPR1	Freer et al. (1980)
<i>t</i> -Boc-Phe-Leu-Phe-Leu-Phe	Enzyme release	$pIC_{50} = 6.59$	FPR1 $\gg$ FPR2/ALX	Freer et al. (1980)
Quin-C7	Binding	$pK_i = 5.19$	FPR2/ALX	Zhou et al. (2007)
CDCA	Binding	$pK_i = 4.76-4.52$	FPR1 > FPR2/ALX	Chen et al. (2000)
DCA	Binding	$pK_i = 4.00$	FPR1	Chen et al. (2002)
Spinorphin	$O_2^-$ generation	$pIC_{50} = 4.30$	FPR1	Yamamoto et al. (1997); Liang et al. (2000)

*i*-Boc, *N*-tert-butoxycarbonyl group; *t*-Boc, -butoxycarbonyl group;  $pIC_{50}$ , negative logarithm of the  $IC_{50}$ ;  $pK_i$ , negative logarithm of  $K_i$ ; N.D., binding affinity or potency was not determined.

ert, 1993). CsH is an inverse agonist (negative antagonist) that suppresses the constitutive activity of FPR1 in a GTPase activity assay (Wenzel-Seifert et al., 1998). The biological significance of constitutive activity for FPR is not established. Both Boc2 and CsH competitively displace FPR1-bound [<sup>3</sup>H]fMLF, indicating that its antagonistic activity is mediated through inhibition of fMLF binding.

A study of CsA on differentiated HL-60 myeloid cells has led to the finding that CsA also inhibits fMLF-induced degranulation, although the inhibitory effect is less potent than CsH (Loor et al., 2002). In a more recent study, Yan et al. (2006) reported that CsA dose-dependently suppresses fMLF- and WKYMVm-induced cell activation and inhibits their binding to FPR1. It has not been determined whether CsA is an inverse agonist. Neither CsA nor CsH displays any detectable inhibitory effect on FPR2/ALX-mediated cellular functions.

Two types of endogenous FPR1 antagonists have been identified. Spinorphin (LVVYPWT), an opioid, is an endogenous peptide antagonist for FPR1 with a  $pIC_{50}$  of 4.30 (Yamamoto et al., 1997; Liang et al., 2001). The bile acids deoxycholic acid (DCA) and chenodeoxycholic acid (CDCA) are two other identified antagonists for FPR1 (Chen et al., 2000, 2002). The  $pIC_{50}$  values for DCA and CDCA are 4 and 3.76, respectively. The physiological functions of DCA- and CDCA-mediated FPR1 antagonism have not been fully understood.

Chemotaxis inhibitory protein of *S. aureus* (CHIPS) is a bacteria-derived protein of 14.1 kDa found in more than half of the clinical strains of *S. aureus*. CHIPS has antagonistic activity for FPR1 and C5aR. An N-terminal peptide (FTFEPFPTNEEIESN) derived from CHIPS is a selective antagonist for FPR1 but not C5aR (Haas et al., 2004). Iodinated CHIPS exhibits a  $pK_d$  of 7.46 at FPR1, which is by far the most potent peptide antagonist for

this receptor. The identification of a bacteria-derived FPR1 antagonist suggests a mechanism used by microorganisms to thwart host defenses.

In a subsequent study, the same group reported the identification from *S. aureus* of a 105-amino acid protein, termed FPRL1 inhibitory protein (FLIPr), that selectively inhibits the binding of and activation by FPR2/ALX agonists including MMK-1, WKYMVM, the prion fragment PrP<sub>106-126</sub>, and amyloid peptide A $\beta$ 1-42 (Prat et al., 2006). At higher concentrations, FLIPr also inhibits fMLF binding to FPR1. FLIPr was found to bind directly to FPR2/ALX and FPR1, but not to FPR3 and C5aR. It does not interfere with LXA4 activity on LTB4. The biological function of this inhibition has not been identified.

Quin-C7 is a synthetic, nonpeptide antagonist of FPR2/ALX, developed through chemical modification of the FPR2/ALX agonist Quin-C1 (Zhou et al., 2007). In binding assays, Quin-C7 inhibited iodinated WKYMVM binding to FPR2/ALX with a  $pK_i$  of 5.18. This antagonist is highly selective for FPR2/ALX, as it does not affect the binding of [<sup>3</sup>H]fMLF to transfected cells expressing FPR1 (Zhou et al., 2007).

In summary, several FPR1 and FPR2/ALX antagonists have been identified and characterized. It is notable that, in most cases, these antagonists differ considerably from the identified FPR1 and FPR2/ALX agonists. The observation that *t*-Boc peptides are antagonistic while *N*-formyl peptides of the same or similar composition are agonistic may be helpful to define the binding pocket of these peptides in FPR1. The synthetic, nonpeptide antagonist, Quin-C7 differs from the agonist Quin-C1 only in the *para* position of the phenyl ring (Zhou et al., 2007). Its antagonistic activity on FPR2/ALX provides a potentially useful tool in study the binding properties of FPR2/ALX.

#### IV. Structure-Function Relationship of the Formyl Peptide Receptor Family of Receptors

##### A. Ligand-Binding Domains

The spectrofluorimetric and flow cytometric techniques developed by Sklar et al. (1981, 1987) and Posner et al. (1994) vastly expanded our knowledge of the dynamics of FPR1 and the properties of its ligand binding pocket. These studies are based on *N*-formylated hexapeptides, modified with fluorescein, and used in conjunction with a high-affinity antibody directed against the fluorescein moiety. The antibody possessed the property to capture free fluoresceinated peptide within a few seconds; the peptide-antibody complex neither stimulates the cells nor inhibits subsequent stimulation by an unlabeled formyl peptide. Using a series of fluoresceinated *N*-formyl peptides with four, five, or six amino acids, Sklar et al. (1990) showed that the chromophore of peptides containing four and five amino acids was not exposed to the extracellular environment, whereas the fluorescein moiety of the hexapeptides was accessible to high concentrations of antibodies. These results indicate that the binding pocket can accommodate no more than five amino acids and, like many GPCRs, that interact with small ligands, the binding pocket for fMLF is believed to involve amino acids inserted into the transmembrane domains.

With the cloning of FPR1 and other chemoattractant receptors, such as human FPR2/ALX, murine and rabbit FPR1, and the C5a receptor C5aR, it became theoretically possible to discriminate residues that are critical for the formation of a high-affinity binding site for these peptides. A number of groups have used chimeric receptor approaches to define the molecular determinants involved in ligand recognition. After the observation that FPR2/ALX binds fMLF with an affinity that is 400-fold lower than that of FPR1 (Ye et al., 1992; Quehenberger et al., 1993), receptor chimeras were constructed by sequential replacements of FPR1 segments with the corresponding regions in FPR2/ALX or vice versa (Gao and Murphy, 1993; Quehenberger et al., 1993). Comparison of the functional differences of the FPR1-FPR2/ALX chimeras has allowed the identification of domains that may be essential for binding of fMLF (Quehenberger et al., 1993) and for functions induced by fMLF (Le et al., 2005). The replacement of the first and third extracellular loops of FPR with those of FPR2/ALX resulted in a dramatic decrease in the affinity of fMLF for the chimeras. Conversely, simultaneous replacement of the corresponding loops in FPR2/ALX by those of FPR1 resulted in a significant increase in ligand-binding affinity with a  $K_d$  that shifted from 400 to 18 nM (Quehenberger et al., 1993). Thus, the first and the third extracellular loops and their adjacent transmembrane domains seem to be essential for high-affinity binding of fMLF. Using a similar approach of domain swapping between FPR1 and C5aR, Perez et al. (1993)

proposed a ligand binding model in which the second, third, and fourth extracellular domains and/or their adjacent transmembrane domains together with the first transmembrane domain form a binding pocket for *N*-formyl peptides. The extracellular N- and C-terminal domains were found to affect ligand binding. These authors proposed that the extracellular N-terminal region provide a "lid" to the pocket. However, the role of the extracellular amino terminus of FPR1 is questionable, because other studies have indicated that the N-terminal region is dispensable for high-affinity binding of fMLF (Malech et al., 1985; Mery and Boulay, 1994).

Based on amino acid differences between the transmembrane domains of FPR1 and FPR2/ALX, Miettinen et al. (1997) prepared FPR1 mutants to identify residues responsible for high-affinity ligand binding. Their study led to the identification of 10 amino acid residues in  $\alpha$  helices (transmembrane domains, TM) II to VII that may participate in fMLF binding. The mutations L78A (TM II), D106N, L109A (TM III), T157A (TM IV), R201A, I204Y, R205A (TM V), W254A, Y257A (TM VI), and F291A (TM VII) resulted in reduced affinities for fMLF ( $K_d = 30$ – $128$  nM). Several of these mutations, (i.e., D106N, R201A, and R205A) were found to affect G protein coupling, suggesting that these residues may also be involved in signal transduction and/or are essential for proper folding of the receptor. Quehenberger et al. (1997) used a "gain-of-function" strategy by selectively replacing the nonconserved region of FPR2/ALX with those of FPR1. This approach has allowed the identification of three clusters of amino acids that are able to restore high-affinity fMLF binding to two distinct chimeras (Quehenberger et al., 1993) that exhibited low-affinity fMLF binding. Introduction of two positively charged amino acids, Arg84 and Lys85, markedly improved binding affinity of one chimeric receptor ( $K_d$  shifted from 105 to 1.6 nM). Likewise, restoration of either Gly89/His90 or Phe102/Thr103 improved the binding affinity of another chimeric receptor from a  $K_d$  of 275 nM to 2.3 or 3.3 nM, respectively. Because a high-affinity binding site for fMLF most likely involves multiple noncontiguous residues that must be positioned by the proper folding of all extracellular and transmembrane domains, it is difficult to locate the ligand binding site precisely or to conclude that these three clusters of amino acids are directly involved in the interaction with fMLF without a crystal structure of the receptor. However, it is worth noting that the identification of amino acids at positions 84 and 85 as key residues for the formation of a high-affinity fMLF binding site is consistent with data derived from an independent photoaffinity labeling approach. Mills et al. (1998) developed an elegant strategy that combined the photolabeling properties of CHO-Met-*p*-benzoyl-L-phenylalanine-Phe-Tyr-*N*<sup>ε</sup>-(fluorescein)-Lys-OH, the ability of an anti-fluorescein antibody to immunoprecipitate the cross-linked fluorescein-labeled peptides, and matrix-assisted laser



desorption ionization mass spectroscopy for characterization of the fMLF binding pocket in FPR1. They identified a major photolabeled cyanogen bromide peptide, Val-Arg-Lys-Ala-Hse, corresponding to residues 83 to 87 of FPR1, as the photolabeled region. In the current three-dimensional model, this peptide lies at the interface between the second transmembrane domain and the first extracellular loop. A model has been proposed in which the NH<sub>2</sub> terminus of an *N*-formyl peptide is hydrogen-bonded to both Asp106 and Arg201, the leucine side chain of fMLF is close to Val-Arg-Lys85, and the COOH-terminal group of fMLF is ion-paired with Arg205 (Mills et al., 2000).

To evaluate the structural contributions of the major domains from the receptor side via interactions with LXA4 or anti-inflammatory and proinflammatory peptide ligands, chimeric receptors were constructed from FPR2/ALX and BLT1 receptors with opposing functions. These chimeras demonstrated that TM7 and adjacent regions of FPR2/ALX are essential for LXA4 recognition, and additional regions, including extracellular loops, are required for high-affinity binding for peptide ligands such as the MMK-1 and MHC peptides. A single GPCR can recognize and function with specific chemotactic peptides as well as nonpeptide ligands such as LXA4. Clearly, however, these ligands act with different affinities and/or at separate interaction sites within the receptor (Chiang et al., 2000). It is noteworthy that conserved *N*-glycosylation sites are present on Asn4 and Asn179 of human FPR2/ALX, and bacterial and viral infection are known to interfere with normal *N*-glycosylation of the host cells (Olofsson et al., 1980; Kim and Cunningham, 1993; Villanueva et al., 1994). In this regard, deglycosylation of FPR2/ALX does not dramatically alter LXA4 recognition but significantly lowers the affinity for peptide ligands (Chiang et al., 2000). Thus, *N*-glycosylation is essential for ligand specificity of this receptor and may play an important role in switching or changing this receptor's functional role at local host defense sites.

### B. G Protein-Coupling Specificities

A series of studies initiated in the mid-1980s revealed that the biochemical and the functional responses of neutrophils to fMLF are largely inhibited by pertussis toxin (PTX), a bacterial toxin that ADP-ribosylates the G<sub>i</sub> class of heterotrimeric G protein  $\alpha$  subunits, thereby preventing their interaction with GPCRs (Bokoch and Gilman, 1984; Bokoch et al., 1984; Lad et al., 1985; Sha'afi and Molski, 1988; Gierschik et al., 1989; Snyderman and Uhing, 1992). With the cloning of FPR1, studies of the structure-function relationships of FPR1 with respect to G protein specificities and G protein coupling became possible. The first such studies involved injection of crude or purified mRNA preparations into *X. laevis* oocytes. This system was tentatively used for the functional cloning of FPR1 (Murphy et al., 1990) before

the expression cloning of FPR1 using COS cells (Boulay et al., 1990a,b). Although *X. laevis* oocytes had been used to demonstrate the functional coupling of a variety of GPCRs, no functional coupling could be detected after the injection of FPR1 mRNA despite appropriate expression of the receptor at the surface of the oocytes (Schultz et al., 1992). This result was unexpected, because all nonchemoattractant GPCRs have been capable of functional coupling in *X. laevis* oocytes. Furthermore, a functional FPR1 could be reconstituted after the injection of crude mRNA preparations from differentiated HL-60 cells but not from undifferentiated cells (Murphy et al., 1990). Surprisingly, *X. laevis* oocytes exhibited a functional response to fMLF when FPR1 mRNA was coinjected with an mRNA fraction of 3 to 3.5 kb from undifferentiated HL-60 cells, which by itself did not confer responsiveness to fMLF (Murphy and McDermott, 1991; Schultz et al., 1992). From these results, it had been concluded that a cofactor was required for FPR1 to function properly in *X. laevis* oocytes. The nature of the cofactor(s) remained unknown until Amatruda et al. (1991) demonstrated that the cofactor might be G $\alpha_{16}$ , a pertussis toxin-insensitive G protein of the G<sub>q</sub> class in which expression is restricted to a subset of hematopoietic cells in the early stage of differentiation; coinjection of FPR1 mRNA with the transcript for G $\alpha_{16}$  was found to be sufficient to enable fMLF-induced ion flux in *X. laevis* oocytes (Burg et al., 1995). The restoration of functional coupling by G $\alpha_{16}$  is most likely due to the fact that G $\alpha_{16}$  is a promiscuous G protein that couples to a wide variety of GPCRs (Offermanns and Simon, 1995; Wu et al., 1995), including FPR1 in transfected mammalian cells (Yang et al., 2001b). However, an additional factor may exist because the size of the mRNA for G $\alpha_{16}$  (2.2 kb) differs from the size of the transcript (3–3.5 kb) for the complementary factor present in the undifferentiated HL-60 cells (Murphy and McDermott, 1991; Schultz et al., 1992).

Another approach used in the study of FPR1 signaling involves the measurement of fMLF-mediated increase in the intracellular concentration of calcium in L-cell mouse fibroblasts (Prossnitz et al., 1991) and human HEK 293 cells (Didsbury et al., 1992) that transiently or stably express FPR1. In contrast to *X. laevis* oocytes, transfected L cells or HEK293 cells responded to fMLF with a rapid and transient calcium flux without any additional cofactor(s) from differentiated HL-60 cells. In contrast to L-cells and HEK293 cells, transfected COS-7 cells were unable to elicit fMLF-stimulated phospholipase C activation unless they were cotransfected with G $\alpha_{16}$  (Amatruda et al., 1991). The deficiency in signaling may be attributable to the fact that COS-7 cells express PI-PLC $_{\beta 1}$  but not the myeloid-specific PI-PLC $_{\beta 2}$  isotype. PI-PLC $_{\beta 1}$  is activated by the  $\alpha$  subunits of PTX-resistant G proteins, such as G $\alpha_{q/11}$  and G $\alpha_{16}$  (Kozasa et al., 1993), whereas PI-PLC $_{\beta 2}$  is activated by the  $\beta\gamma$  subunits of G proteins, such as G $\alpha_{12}$  (Camps et al., 1992a,b). The

$\alpha$  subunits of PTX-sensitive G proteins cannot activate any isoforms of PI-PLC (Hepler et al., 1993), but  $G_{\alpha_q}$  and  $G_{\alpha_{11}}$ , which are resistant to PTX, are able to activate PI-PLC $_{\beta 2}$ .

In leukocytes, the role of  $G_{\alpha_{16}}$  seems to be marginal for several reasons. First, the fMLF-mediated functional responses are largely inhibited by treatment with PTX. Second,  $G_{\alpha_{16}}$  transcript abundance is very low in mature neutrophils, and induction of HL-60 cell differentiation is associated with a progressive decrease in expression of  $G_{\alpha_{16}}$  and an increase in the expression of  $G_{\alpha_2}$  (Amatruda et al., 1991). Third, a large body of evidence indicates that the fMLF-induced responses require physical association of FPR1 with PTX-sensitive  $G_{\alpha_2}$  protein in leukocytes (Bommakanti et al., 1992, 1993, 1994, 1995; Schreiber et al., 1993). However, residual molecules of  $G_{\alpha_{16}}$  may account for a small PTX-resistant activity that has been observed in neutrophils, monocytes, and differentiated HL-60 cells stimulated with fMLF (Verghese et al., 1987). It is noteworthy that mouse Fpr1, along with a number of chemokine receptors known to couple to the  $G_i$  proteins, can also couple to  $G_{\alpha_q}$  (Shi et al., 2007). This property is responsible for an alternative pathway for the activation of chemoattractant and chemokine receptors that leads to chemotaxis.

A quantitative analysis of FPR1 coupling to the three  $G_{\alpha_i}$  isoforms,  $G_{\alpha_{11}}$ ,  $G_{\alpha_{12}}$ , and  $G_{\alpha_{13}}$ , was performed with FPR1 fused to the different  $G_{\alpha_i}$  isoforms to obtain a defined 1:1 stoichiometry of the signaling partners. After expression in Sf9 cells, high-affinity agonist binding, GTP $\gamma$ S binding kinetics, and GTP hydrolysis were analyzed. The results indicate that 1) FPR1 couples to all three  $G_{\alpha_i}$  isoforms with similar efficiency, 2) FPR1 can couple efficiently to these G proteins even in the low-affinity state for agonist binding, and 3) in contrast to what was previously thought, FPR1 activates the  $G_{\alpha_i}$  proteins linearly and not catalytically (Wenzel-Seifert et al., 1999).

### C. G Protein-Coupling Domains

A variety of approaches have been taken to define the contact sites responsible for G protein-FPR1 coupling, including site-directed mutagenesis, construction of chimeric receptors, inhibition of high-affinity G protein-dependent fMLF binding, and peptide "walking" to map the regions of the intracellular loops that are in contact with the G protein (for review, see Savarese and Fraser, 1992). The latter approach is based on the use of synthetic receptor-mimetic peptides to block the receptor-G protein interactions. A competition assay based on the sedimentation of a GTP-sensitive FPR1- $G_{\alpha_{12}}$  complex on sucrose gradients in the presence of octyl glucoside was developed by Bommakanti et al. (1992, 1994) to assess the effect of FPR1-derived peptides on the conversion of FPR1 from a 7S (FPR1- $G_i$  complex) to a 4S species (FPR1 only). Synthetic FPR1-derived peptides were also

used by Schreiber et al. (1994) in a competitive enzyme-linked immunosorbent assay. This assay measures the ability of FPR1-derived peptides to inhibit the binding of an anti- $G_{\alpha_{12}}$  antibody to soluble  $G_{\alpha_{12}}$  (Weingarten et al., 1990).

The results from these assays excluded the third intracellular loop of FPR1 as a significant determinant for G protein coupling. The second intracellular loop (amino acids D<sup>122</sup>RCVCLHPQNHRTVSLAKK<sup>144</sup>) and the entire C-terminal region were identified as the major sites of contact between FPR1 and  $G_{\alpha_i}$  proteins. It is notable that the DRY (Asp-Arg-Tyr) motif conserved in many GPCRs (Rovati et al., 2007) becomes DRC in the FPR family of receptors. Peptides corresponding to different regions of the carboxyl-terminal tail produced variable effects. The sequence extending from Arg322 to Thr336 had the best blocking activity and seemed to be the major structural determinant for the carboxyl-terminal tail to interact with  $G_{\alpha_i}$ . Although several studies on other receptors, namely rhodopsin and the  $\beta$ -adrenergic receptors (König et al., 1989; Münch et al., 1991), have suggested no role for the first intracellular loop in  $G_{\alpha_i}$  interaction, Bommakanti et al. (1995) found that a peptide corresponding to the first intracellular loop of FPR1 was able to disrupt FPR1- $G_{\alpha_i}$  complex formation. A role for the first intracellular loop in G protein coupling was further supported by a study showing that its replacement by the counterpart from C5aR resulted in a constitutively active chimeric receptor (Amatruda et al., 1995). This latter observation may either suggest that the short stretch of amino acids that constitutes the first intracellular loop help to maintain an inactivated state of the receptor or reflect an unexpected conformational change resulting from substitution with the C5aR sequence.

Detailed information on the mechanisms of coupling between FPR1 and G proteins were obtained by site-directed mutagenesis studies that targeted amino acids conserved among the different GPCRs (Prossnitz et al., 1995b; Miettinen et al., 1999). Replacement of a cluster of three amino acids starting at position 309 at the N-terminal part of the cytoplasmic tail (RER to GAG) as well as Ala substitution of either Asp71 in the second transmembrane domain, or Arg123 in the conserved DRC (Asp-Arg-Cys) motif at the N terminus of the second intracellular loop, yielded three interesting mutant receptors with poor coupling to  $G_{\alpha_i}$  proteins and deficient signaling capacity. The Asp71 mutant receptor is not activated or phosphorylated in response to fMLF, whereas the Arg123Gly mutant does not bind G protein but is phosphorylated and able to bind  $\beta$ -arrestins (Gripentrog and Miettinen, 2008).

## V. Regulation of Formyl Peptide Receptors

The activation of FPRs is regulated at the levels of receptor to G protein activation (R $\rightarrow$ G), transduction

and amplification of signals from activated G proteins to effectors (G→E) including kinases and small GTPases, and integration of effector signals leading to phagocyte functions (E→F) such as chemotaxis, degranulation, and superoxide generation. This section is focused on major regulation of FPRs at the receptor level.

#### A. Desensitization of FPR1

After stimulation with fMLF, the cellular responses rapidly decline in intensity and the cells become refractory to subsequent stimulations with the same agonist. This loss of responsiveness, known as *homologous desensitization*, is a feature of G protein-coupled receptors that results from phosphorylation of the agonist-occupied receptor by G protein-coupled receptor kinases (GRKs) (Pitcher et al., 1998). A related phenomenon is that stimulation of one receptor can desensitize the cells to subsequent stimulations by ligands for another receptor. This is referred to as *heterologous desensitization* and is thought to result from the modification and/or inactivation of unliganded receptors after phosphorylation by second messenger-triggered kinases, such as protein kinase A and protein kinase C (PKC) (Dohlman et al., 1991).

Through the expression of different chemoattractant receptors in HEK293 cells and measurement of calcium mobilization induced by different agonists, Didsbury et al. (1991) introduced the concept of *receptor class desensitization*. For instance, C5aR and FPR1 were found to desensitize each other, but they cannot desensitize, or be desensitized by, receptors such as the  $\alpha 1$  adrenergic receptor, which is not closely related to the chemoattractant receptors and has different signaling properties (Didsbury et al., 1991). Furthermore, comparison of the ability of fMLF, C5a, and IL-8 to desensitize one another in calcium mobilization assays has led to the observation of hierarchy in receptor desensitization between these chemoattractant receptors. The strength of peptide chemoattractants to desensitize calcium responses to one another (in the order of fMLF > C5a > IL-8) reflects the ability to generate desensitizing signals for both phosphorylation of an unliganded receptor by PKC and inactivation of a distal component in the calcium mobilization pathway. It is noteworthy that all three peptide chemoattractants can desensitize the responses mediated by lipid mediators such as leukotriene B<sub>4</sub> (LTB<sub>4</sub>) and PAF. However, none of these lipids was able to desensitize the calcium response mediated by fMLF, C5a, or IL-8. Thus, this type of receptor cross-desensitization consists of at least two distinct processes. The first process possibly involves the phosphorylation of the receptor by PKC. FPR1 is resistant to this process because of the lack of PKC phosphorylation sites, yet the fMLF-mediated calcium response can be desensitized. The second process occurs downstream of the receptor-G protein complex and most likely results from the inactivation of a shared effector that is required for the cal-

cium mobilization response mediated by peptide chemoattractants. The hierarchy of desensitization among these agonists may be important for leukocyte signal processing when multiple chemoattractants are present (e.g., in tissues with bacterial infection). Under these conditions, the “end-target chemoattractants,” such as fMLF, that are more capable of activating terminal functions of neutrophils, seem to prevail over “intermediary chemoattractants” such as IL-8 and other chemokines present at the site of inflammation (Heit et al., 2002). It has been shown that the “end-target chemoattractants” use both p38 MAPK and the PI3K pathways to maintain polarity and steer a migrating neutrophil, whereas the “intermediary chemoattractants” only activate the PI3K pathway. Through an as-yet-unknown mechanism, the chemoattractants that activate p38 MAPK can maintain PI3K phosphatase and tensin homolog (PTEN) localization to the rear and sides of the migrating cell, which helps to amplify the intracellular PtdIns(3,4,5)P<sub>3</sub> gradient and resist any PI3K activity initiated by “intermediary chemoattractants” from other directions (Heit et al., 2008).

Studies of FPR1 have revealed the dose-dependent characteristics of receptor desensitization, such that FPR1 exposed to a given concentration of fMLF can no longer respond to fMLF at the same concentration but it can still respond to a higher concentration of fMLF. In a study of Mac-1-dependent neutrophil adhesion, Hughes et al. (1992) demonstrated that sequential, stepwise increases in fMLF concentrations could overcome desensitization and augment Mac-1 mobilization from intracellular stores to the cell's surface as well as Mac-1-mediated cell adhesion. This and other results from calcium mobilization assays indicate that full desensitization of chemoattractant receptors requires sustained exposure to saturating concentrations of agonists.

#### B. Phosphorylation of Formyl Peptide Receptors

It has been shown that FPR1 is rapidly phosphorylated in an agonist concentration- and time-dependent manner (Ali et al., 1993; Tardif et al., 1993). PKC is not involved in the phosphorylation of FPR1 based on several observations. First, the C-terminal region of FPR1 does not contain PKC phosphorylation sites. Second, the agonist-dependent phosphorylation of FPR1 is resistant to the PKC inhibitor staurosporin. Third, phorbol 12-myristate 13-acetate (PMA), an activator of protein kinase C, is unable to induce the phosphorylation of FPR1. The phosphorylation sites are restricted to the carboxyl terminal region, and phosphoamino acid analysis has revealed that only serines and threonines are phosphorylated (Prossnitz et al., 1995a). The carboxyl tail of FPR1 contains a total of 11 serine and threonine residues. Eight of these residues, located between and including Ser328 and Thr339, are found to be critical to FPR1 internalization and desensitization (Maestes et al., 1999). These residues are arranged in two domains,

and the sequences D<sup>328</sup>STQTS<sup>332</sup> and D<sup>333</sup>TATNST<sup>339</sup> have the characteristics of GRK phosphorylation sites (Onorato et al., 1991). GRK2 has been identified as the primary kinase that phosphorylates FPR1, using as substrate a fusion protein consisting of the carboxyl-terminal 47 amino acids of FPR1 and glutathione transferase (Prossnitz et al., 1995a). Other GRKs are either significantly less effective (GRK3) or have no detectable activity (GRK5 and GRK6). Site-specific mutagenesis of the FPR1 carboxyl terminus has shown that acidic residues (Glu326, Asp327, and Asp333) are critical to FPR1 phosphorylation (Prossnitz et al., 1995a). Amino acid substitutions indicate that FPR1 phosphorylation uses a hierarchical mechanism in which the phosphorylation of two clusters of adjacent serine and threonine residues (Ser328/Thr329 or Thr331/Ser332) is required for subsequent phosphorylation of other serine and threonine residues. The phosphorylation of Ser328, Ser332, and Ser338 is critical to FPR1 internalization and desensitization as well as  $\beta$ -arrestin 2 binding (Potter et al., 2006). The phosphorylation status of these two clusters of serines and threonines modulates the affinity of FPR1 for  $\beta$ -arrestins and agonists (Key et al., 2003, 2005). The affinity of FPR1 for  $\beta$ -arrestins is controlled by the phosphorylation of serines and threonines in the sequence D<sup>328</sup>STQTS<sup>332</sup>, whereas the affinity of the  $\beta$ -arrestin-bound receptor for the agonist is regulated by the phosphorylation of the serines and threonines in the sequence D<sup>333</sup>TATNST<sup>339</sup>. Phosphorylation of these residues renders the  $\beta$ -arrestin-bound FPR1 able to form a high-affinity ternary complex with the ligand. Phosphorylation of carboxyl terminal serine and threonine residues may produce a localized concentration of negative charges, which facilitates ionic interactions with the positively charged receptor recognition domain of  $\beta$ -arrestin. This interaction may induce a  $\beta$ -arrestin intramolecular conformational change that then stabilizes a firm interaction between the two proteins and/or exposes a secondary high-affinity binding site (Han et al., 2001; Milano et al., 2002).

Like FPR1, FPR2/ALX is phosphorylated in an agonist-dependent manner, but little is known about the nature of the kinase(s) involved in the process. The determinants responsible for its internalization have not yet been identified. In contrast to FPR1 and FPR2/ALX, FPR3 displays a marked level of phosphorylation in the absence of stimulation (Christophe et al., 2001). Because phosphorylation of these receptors is known to affect their internalization, it will be interesting to know whether constitutive phosphorylation of FPR3 is related to its cell surface expression pattern.

### C. Interaction between Formyl Peptide Receptors and $\beta$ -Arrestins

**1. Uncoupling of G Proteins.** In the current model, liganded GPCRs are phosphorylated and form a high-affinity complex with  $\beta$ -arrestin 1 or  $\beta$ -arrestin 2. The

$\beta$ -arrestins are thought to sterically interfere with G protein coupling and thereby inhibit the signaling capacity of the receptor. However, several studies have suggested that the desensitization of FPR1 may be unique. Using a fluorimetric assay in conjunction with solubilized receptors, Bennett et al. demonstrated that fMLF-induced phosphorylation of FPR1 can block FPR1 interaction with G proteins independently of  $\beta$ -arrestin (Bennett et al., 2001). Based on these findings, it is proposed that phosphorylation alone may be sufficient for uncoupling of the receptors from G proteins in vivo and this may not be the primary function of  $\beta$ -arrestins.

In the case of GPCRs that are coupled to the G<sub>q</sub> or G<sub>s</sub> class of G proteins, such as the angiotensin II type I A receptor and the V2 vasopressin receptor,  $\beta$ -arrestins play the role of adaptors and scaffolding molecules that have recently emerged as signal transducers for the activation of ERK in response to agonist stimulation (Shenoy and Lefkowitz, 2003; Lefkowitz and Whalen, 2004). Two studies have examined whether the binding of  $\beta$ -arrestins to FPR1 and FPR2/ALX, which are coupled to the PTX-sensitive G<sub>i</sub> class of G proteins, plays a role as signal transducer to ERK activation. The first study compared the kinetics of ERK activation in both wild-type and  $\beta$ -arrestin-deficient mouse embryonic fibroblasts that stably expressed FPR2/ALX or in HEK293 cells that coexpressed FPR2/ALX and  $\beta$ -arrestin 1 or 2 (Huet et al., 2007). The second study took advantage of FPR1 mutants, such as the Arg123 mutant with a partial defect in G $\alpha_i$  coupling and the Asn297 mutant with a defect in  $\beta$ -arrestin recruitment (Gripentrog and Miettinen, 2008). Both studies have led to the conclusion that ERK activation occurs through G $\alpha_i$  and is not affected by  $\beta$ -arrestins. The lack of involvement of  $\beta$ -arrestins is thought to be a common characteristic among G $\alpha_i$ -coupled chemoattractant receptors, but additional investigations will be necessary to prove this is the case.

**2. Internalization of Formyl Peptide Receptors.** After prolonged stimulation with fMLF, FPR1 was found to colocalize with  $\beta$ -arrestin 1 and  $\beta$ -arrestin 2 in endocytic vesicles in transfected monocytic U937 cells. It is well accepted that the  $\beta$ -arrestins serve to link phosphorylated receptors to the components of the endocytic machinery, including clathrin and the clathrin adaptor AP2 (Goodman et al., 1996; Laporte et al., 2002). This endocytic pathway seems to be used by the majority of GPCRs (for review, see Marchese et al., 2003; Lefkowitz and Whalen, 2004). However, several studies indicate that the internalization of FPR1 diverges from this common model in certain aspects (Gilbert et al., 2001; Xue et al., 2007). Flow cytometric analyses of HEK293 cells coexpressing FPR1 and dominant-negative constructs of  $\beta$ -arrestin 1, dynamin, and clathrin provided the first evidence that FPR1 is not internalized through clathrin-coated pits (Gilbert et al., 2001). Studies with transfected mouse embryonic fibroblasts (Vines et al., 2003)

that are derived from  $\beta$ -arrestin 1 and  $\beta$ -arrestin 2 knockout mice (Kohout et al., 2001) further established that FPR1 can interact with different components of the endocytic machinery in a  $\beta$ -arrestin-independent manner. However, phosphorylation of liganded FPR1 is a prerequisite for the internalization whether the two  $\beta$ -arrestins are present or not. The ability of FPR1 to be internalized via a  $\beta$ -arrestin-independent pathway in mouse embryonic fibroblasts does not exclude the possibility that the classic arrestin- and clathrin-dependent pathway is involved when arrestins associate with the phosphorylated receptor (Gilbert et al., 2001; Vines et al., 2003).

In contrast to FPR1, several studies indicate that liganded FPR2/ALX undergoes clathrin-mediated and dynamin-dependent endocytosis and that it recycles in a slow pathway involving perinuclear recycling endosomes. First, FPR2/ALX internalization can be inhibited by either siRNA-mediated depletion of cellular clathrin or the expression of a dominant-negative mutant of dynamin in HeLa cells (Ernst et al., 2004b). Second, in HEK293 cells cotransfected with FPR2/ALX and a  $\beta$ -arrestin-enhanced green fluorescent protein, the receptor was rapidly internalized after the addition of the synthetic agonist peptide WKYMVM. It colocalized within a few minutes with  $\beta$ -arrestin-enhanced green fluorescent protein vesicles and gathered in the perinuclear region of the cytoplasm (Huet et al., 2007). Third, the internalization of FPR2/ALX, expressed in mouse embryonic fibroblasts lacking both  $\beta$ -arrestins, was highly compromised (Huet et al., 2007). In comparison, FPR2/ALX was normally internalized when only one of the two  $\beta$ -arrestins was expressed.

## VI. Formyl Peptide Receptor Signal Transduction and Activation of Cell Functions

FPR1, along with the receptor for C5a, were the first identified phagocyte chemoattractant receptors. In comparison with chemokines and other “classic chemoattractants” such as PAF and LTB<sub>4</sub>, fMLF and C5a have several properties that qualify them as “end-target chemoattractants” (Heit et al., 2002). These two chemoattractants are predominant in cross-desensitization of other receptors (Ali et al., 1999), and they are potent agonists for stimulating bactericidal functions such as superoxide generation. This section will focus on major phagocyte activities induced by fMLF and the relevant signaling mechanisms. A more detailed description of signaling through FPR1 and similar chemoattractant receptors is provided in a recent review (Rabiet et al., 2007). It is notable that, although there are many similarities between FPR1 and FPR2/ALX (and possibly also FPR3), major differences exist between these receptors in signaling. These include for example the sensitivity to membrane cholesterol depletion (Tuluc et al., 2003), and to the intracellularly acting peptide (PBP10)

derived from a PIP<sub>2</sub> binding domains of the cytoskeletal protein gelsolin (Fu et al., 2004a). Evidence also suggests that the basic mechanisms by which these receptors trigger a transient increase in intracellular Ca<sup>2+</sup> may be different (Partida-Sánchez et al., 2004).

### A. Phagocyte Activation for Bacterial Clearance

1. *Chemotaxis.* N-Formyl peptides have been extensively studied for their abilities to induce directional migration of neutrophils since their initial discoveries (Schiffmann et al., 1975a; Zigmond, 1977). Human neutrophils are able to detect a chemotactic gradient of subnanomolar concentrations of fMLF (Showell et al., 1976; Freer et al., 1980). In dose-response experiments, fMLF induces a typical bell-shaped curve seen with other leukocyte chemoattractants, indicating that increasing chemoattractant concentrations beyond a certain level actually reduces chemotaxis. At least two mechanisms can contribute to the descending curve past an optimal chemotactic concentration. First, saturation of these receptors with a high concentration of chemoattractant eliminates any meaningful gradient, thereby reducing chemotaxis. Second, chemoattractant receptors are desensitized by exposure to high concentrations of agonists, causing reduced responsiveness to subsequent agonist stimulation. It is unclear what proportion that desensitization contributes to the reduced chemotaxis when cells are exposed to a chemoattractant at high concentrations, but studies have shown that mechanisms other than heterologous desensitization are involved in prioritizing migrating cells exposed to multiple chemoattractants such as fMLF and IL-8 (Heit et al., 2008). Chemotaxis of neutrophils does not require chemoattractant receptor internalization or redistribution to the leading edge (Hsu et al., 1997; Neptune and Bourne, 1997). It requires G $\beta\gamma$  released from the activated G $\alpha_i$ , but not G $\alpha_i$  itself or activation of G $\alpha_q$  or G $\alpha_s$  (Neptune et al., 1999). Signaling downstream of G $\beta\gamma$  is important for chemotaxis because it determines the orientation of the cells. This was shown in elegant studies using video microscopy of fluorescent proteins as sensors for the detection of clustered signaling molecules at the leading edge of moving neutrophils (Servant et al., 1999). The PI3K family of lipid kinases has been found to play important roles in signaling for chemotaxis downstream of chemoattractant receptors (Niggli and Keller, 1997; Hirsch et al., 2000; Li et al., 2000; Sasaki et al., 2000). In neutrophils, the small GTPase Rac2 is also necessary for optimal cell migration (Roberts et al., 1999). Therefore, exposing neutrophils to a chemotactic gradient creates an intracellular gradient of signaling molecules that defines the “frontness” or leading edge as well as the direction of cell migration. The leading edge is characterized with increased accumulation of PtdIns(3,4,5)P<sub>3</sub> and actin polymerization. The “backness” or uropod of a migrating cell is characterized by concentrations of molecules such as PTEN, RhoA, and

the G<sub>12/13</sub> family of G proteins (Funamoto et al., 2002; Xu et al., 2003). Whereas FPR1 is known to activate p38 MAPK, which helps to localize PTEN in the circumference other than the leading edge, exactly how FPR1 couples and activates G $\alpha_{12}$  or G $\alpha_{13}$  remains unclear.

Migrating neutrophils in tissues are exposed to multiple chemoattractants and must sort out different and sometimes conflicting signals. How these chemoattractants influence the migration of neutrophils has been studied in vitro using the “under agarose” chemotaxis assay (Nelson et al., 1975). Campbell et al. (1997) and Foxman et al. (1997, 1999) found that navigation of neutrophils is a multistep process, and a migrating cell not only detects multiple chemoattractants but also integrates these different signals. Therefore, the net result of cell migration is determined by the type of orienting signals, the strength of the signals, and the time sequence in which they appear. Heit et al. (2002) identified a hierarchy in chemotactic signaling that determines the direction of migrating cells in opposing chemotactic gradients. They found that “end-target chemoattractants” such as fMLF and C5a dominate over the “intermediary chemoattractants” such as IL-8 (CXCL8) and LTB4 (Heit et al., 2002). The ability of neutrophils to distinguish between these chemoattractants is crucial for their optimal migration toward bacteria with minimal interference by the “intermediary chemoattractants” that are also present in the inflammatory site. In a recent study, the same group of investigators provided new insights into the molecular mechanisms that allow neutrophils to prioritize chemotactic signals (Heit et al., 2008). As mentioned above, PI3K plays an important role in cell migration through generation of PtdIns(3,4,5)P<sub>3</sub>, which is found in the leading edge of a cell undergoing chemotaxis (Dormann et al., 2002; Wang et al., 2002). PTEN converts PtdIns(3,4,5)P<sub>3</sub> back to phosphatidylinositol (4,5)-bisphosphate, thereby counteracting the effect of PI3K. Contrary to PtdIns(3,4,5)P<sub>3</sub>, PTEN is localized in the rear (uropod) as well as sides of a migrating cell, which helps to amplify the intracellular PtdIns(3,4,5)P<sub>3</sub> gradient (Funamoto et al., 2002). In addition to the PI3K/PTEN pathway, “end-target chemoattractants” such as fMLF and C5a activate p38 MAPK, which is important for neutrophil chemotaxis toward these chemoattractants. Inhibition of p38 MAPK prevents neutrophil migration toward fMLF and C5a but not IL-8 (CXCL8) or LTB4, which do not stimulate p38 MAPK activation in neutrophils (Heit et al., 2002; Khan et al., 2005). *Pten*( $-/-$ ) neutrophils migrate normally to a single gradient of fMLF but randomly to a single gradient of an “intermediary chemoattractant” such as MIP-2 (CXCL2) (Heit et al., 2008). These cells exhibit two or more pseudopods and often lack an identifiable uropod when stimulated with MIP-2. In the presence of opposing chemotactic gradients such as fMLF and MIP-2, the *Pten*( $-/-$ ) neutrophils were unable to prioritize chemoattractants to migrate toward fMLF (Heit et

al., 2008). The results from this study suggest an important function of p38 MAPK in maintaining proper localization of PTEN in cells migrating toward an “end-target chemoattractant,” although how p38 MAPK regulates PTEN localization is still unclear. PTEN that is present in the sides and rear of a moving cell helps the cell to resist any PI3K activity initiated by “intermediary chemoattractants” from other directions. This study raises an intriguing question of how two types of chemoattractant receptors that converge to the same PTX-sensitive G $\alpha_i$  protein can trigger differential activation of downstream pathways such as the p38 MAPK pathway. Further studies will be necessary to determine whether the two types of receptors are localized in different plasma membrane microdomains that may recruit different components of intracellular signaling pathways.

**2. Superoxide Generation.** Stimulation of neutrophils with fMLF at concentrations higher than those required for chemotaxis leads to generation of superoxide (Boxer et al., 1979; Lehmeyer et al., 1979). In most published studies, fMLF concentrations of 50 to 100 nM are required for the induction of superoxide production by human neutrophils in suspension. In studies using mouse neutrophils, an even higher fMLF concentration (5–10  $\mu$ M) is necessary for superoxide production. The response is ameliorated when mouse neutrophils are stimulated with formyl peptides of different sequences such as fMIVIL and fMIFL, which are able to activate this function in mouse neutrophils at concentrations as low as 10 nM (Southgate et al., 2008). The different concentration requirements for chemotaxis and superoxide generation effectively prevent oxidant-mediated tissue injury that may be caused by migrating neutrophils. It is not entirely clear why neutrophils require a 50- to 100-fold higher concentration of fMLF for superoxide production than chemotaxis, but a high signaling strength is probably necessary for simultaneous activation of multiple pathways leading to NADPH oxidase activation in phagocytes.

Superoxide production in neutrophils results from membrane assembly and activation of NADPH oxidase, which is a multicomponent enzyme complex for electron transfer leading to one-electron reduction of molecular oxygen (Babior et al., 2002). In the resting state, the membrane components (cytochrome <sub>b558</sub>, consisting of gp91<sup>phox</sup> and p22<sup>phox</sup>) are physically separate from the cytosolic components (p67<sup>phox</sup>, p47<sup>phox</sup>, p40<sup>phox</sup>, and the small GTPase Rac) (Fig. 6). Phosphorylation of the cytosolic factor p47<sup>phox</sup> is necessary to change its conformation and unmask the N-terminal PX domain for interaction with the membrane components p22<sup>phox</sup> and phosphatidylinositol (3,4)-bisphosphate (DeLeo and Quinn, 1996). Several studies have demonstrated that fMLF can activate kinases known to phosphorylate p47<sup>phox</sup>, including several PKC isoforms (Dang et al., 2001), Akt (Chen et al., 2003), and the MAPKs ERK and p38 (El Benna et al., 1996; Dewas et al., 2000). In recon-




FIG. 6. The fMLF-induced signaling events leading to the activation of phagocyte NADPH oxidase (Nox2). Depicted schematically are major signaling pathways activated by fMLF that results in NADPH oxidase activation in neutrophils. Upon activation of the  $G_{\alpha_i}$  proteins, the released  $G_{\beta\gamma}$  subunits trigger PI3K activation, resulting in the production PIP3 and its degradation products. The  $G_{\beta\gamma}$  and PIP3-mediated p-Rex1 activation is key to the conversion of GDP-bound Rac small GTPase to GTP-bound Rac, which translocates to membrane and associates with  $gp91^{phox}$ .  $G_{\beta\gamma}$  is also responsible for PLC $\beta$ 2 activation, leading to the production of the second messengers diacyl glycerol and 1,4,5-inositol trisphosphate (IP3), which stimulate PKC activation. Isoforms of PKC (PKC $\delta$ , PKC $\zeta$ , PKC $\beta$ II, and PKC $\alpha$ ), MAPK (p38, ERK), and Akt are known to catalyze the phosphorylation of  $p47^{phox}$  in fMLF-stimulated neutrophils, prompting membrane translocation of the cytosolic factors. The PX domain in  $p47^{phox}$  also facilitates its membrane association. Assembly of a membrane complex of NADPH oxidase is key to its conversion of molecular oxygen to superoxide. Omitted in this drawing is the phospholipase D (PLD) activation pathway, which is reported to contribute to fMLF-induced superoxide generation.

stituted  $COS^{phox}$  cells engineered to express all membrane and cytosolic phox proteins of phagocyte NADPH oxidase (Price et al., 2002), fMLF-induced superoxide generation requires a novel PKC isoform, PKC $\delta$  (He et al., 2004). It has been shown that fMLF stimulates rapid phosphorylation of PKC $\delta$  in its activation loop, and mutating the activation loop Thr507 to an alanine abolishes fMLF-induced, PKC $\delta$ -dependent superoxide production (Cheng et al., 2007). It is noteworthy that PAF, which is a weak agonist for activating NADPH oxidase, also fails to stimulate phosphorylation of PKC $\delta$  at Thr507. Therefore, rapid phosphorylation of PKC $\delta$  may be a mechanism for the superoxide production induced by fMLF. In addition to PKC $\delta$ , several other isoforms of PKC, including PKC $\alpha$ , PKC $\beta$ II, and PKC $\zeta$ , are likely to contribute to fMLF-induced superoxide generation in neutrophils. Genetic deletion of PLC $\beta$ 2, which affects activation of the conventional and novel PKCs, abrogates fMLF-induced superoxide production in mouse neutrophils (Jiang et al., 1997).

The small GTPase Rac is an essential cytosolic component for NADPH oxidase activation (Abo et al., 1991; Knaus et al., 1991). In mouse neutrophils, deletion of Rac abrogates fMLF- and PMA-induced superoxide generation (Roberts et al., 1999). One of the mechanisms for chemoattractant-induced Rac activation involves p-Rex1, a guanine nucleotide exchange factor for Rac that is activated by PtdIns(3,4,5)P $_3$  and  $G_{\beta\gamma}$  from activated heterotrimeric  $G_i$  proteins (Welch et al., 2002). The fMLF-induced Rac activation and superox-

ide generation is significantly reduced in mouse neutrophils lacking p-Rex1 (Dong et al., 2005; Welch et al., 2005). In addition to p-Rex1, other Rac guanine nucleotide exchange factors, such as Vav1, also play a role in fMLF-induced NADPH oxidase activation (Kim et al., 2003).

Studies of the individual kinases and guanine nucleotide exchange factors have shown a prominent role for PI3K in fMLF-induced functions, including superoxide generation. PI3K catalyzes the conversion of phosphatidylinositol (4,5)-bisphosphate to PtdIns(3,4,5)P $_3$ . In addition to activating p-Rex1, PtdIns(3,4,5)P $_3$  is responsible for Akt and PKC $\delta$  activation, either directly or through the action of phosphoinositide-dependent kinase 1 (Dutil et al., 1998). The degradation products of PtdIns(3,4,5)P $_3$ , including phosphatidylinositol (3,4)-bisphosphate and phosphatidylinositol phosphate, serve as membrane binding sites for the PX domain of  $p47^{phox}$  and  $p40^{phox}$ , respectively (Kanai et al., 2001). Genetic deletion of PI3K $\gamma$  significantly reduces chemoattractant-stimulated neutrophil superoxide production (Hirsch et al., 2000; Li et al., 2000; Sasaki et al., 2000). The roles of other type I PI3Ks have also been implicated in fMLF-induced phagocyte NADPH oxidase activation (Condliffe et al., 2005; Boulven et al., 2006).

Human neutrophils in suspension that are pretreated with priming agents such as granulocyte macrophage-colony-stimulating factor, TNF $\alpha$ , PAF, LPS, and certain chemokines produce significantly more superoxide when stimulated with fMLF. These agents do not induce sig-

nificant superoxide production when applied to the neutrophils in concentrations sufficient for priming. There is a large body of literature on the priming mechanisms that range from partial phosphorylation of p47<sup>phox</sup> (Dewas et al., 2003) and p67<sup>phox</sup> (Brown et al., 2004) to enhancement in PI3K activity (Kodama et al., 1999), up-regulation of NADPH oxidase assembly (DeLeo et al., 1998), and mobilization of receptors from intracellular stores (Almkvist et al., 2001; Almkvist et al., 2004), suggesting that there are multiple mechanisms for the potentiation of phagocyte NADPH oxidase activation. It is noteworthy that when used at low concentrations (e.g., 5 nM), fMLF is unable to stimulate neutrophil superoxide production but can prime the cells for a more robust response to PMA (Karnad et al., 1989). Conversely, PMA can prime the response to fMLF, as illustrated by the study performed by Tardif et al. (1998) with a variant clone of the myeloid HL60 cell line. When added to the dibutyryl cyclic AMP-differentiated HL-60 variant, PMA per se is unable to stimulate superoxide production, but it primes the cells for a robust and sustained response to fMLF at concentrations 1000-fold lower than those required for superoxide production when used alone. These findings indicate that the fMLF- and PMA-activated signaling pathways synergize and complement one another. The different studies that deal with the priming issue suggest that the ability of fMLF to stimulate neutrophil superoxide production is a function of its potency (the amount required to produce an effect of given intensity). For most priming agents, their inability to directly stimulate neutrophil superoxide production is an intrinsic property of the ligand and cannot be overcome simply by increasing ligand concentrations. Of particular interest is that IL-8 (CXCL8), known as a priming agent in some studies (e.g., Guichard et al., 2005), has been shown to directly activate neutrophil NADPH oxidase in other studies (Thelen et al., 1988; Fu et al., 2004b). Pretreatment of neutrophils with IL-8 can prime the cells for a more robust response to fMLF, whereas pretreatment of cells with fMLF reduces the IL-8-stimulated superoxide production (Fu et al., 2004b). This pattern is similar to the order of desensitization by these agonists. It remains to be determined whether experimental conditions (e.g., pretreatment of neutrophils with cytochalasin B) or properties of the reagents (e.g., lengths of the chemokine) are contributing factors for the discrepancy in IL-8 priming and activation studies carried out in different laboratories.

**3. Degranulation.** In addition to the induction of superoxide generation, formyl peptides at higher concentrations (usually 10–50 times higher than the optimal concentration for chemotaxis) stimulate release of granule constituents from neutrophils (Showell et al., 1976; Bentwood and Henson, 1980). The fMLF concentrations required for the release of secretory vesicles and gelatinase granules are lower than that required for the re-

lease of other granules (Borregaard et al., 1993). The fMLF-induced mobilization of granules produces several effects, including proteolytic cleavage of membrane-localized adhesion molecules such as L-selectin, cell surface expression of new adhesion molecules, and release of proinflammatory matrix proteins such as the aforementioned FPR2/ALX agonist LL-37, as well as enzymes that can cause tissue degradation and killing of bacteria. For instance, neutrophil myeloperoxidase released from the azurophil granules helps to convert hydrogen peroxide to hypochlorous acid (Nauseef, 2007), a metabolite of importance both for killing invading microbes and for the resolution of inflammation. Most of the membrane-associated NADPH oxidase components, gp91<sup>phox</sup> (Nox2) and p22<sup>phox</sup>, are localized to mobilizable granules and these become more available for the assembly of NADPH oxidase when the granules fuse with the plasma membrane or with the membranes that surround engulfed microorganisms. Last but not the least, there are intracellular pools of FPR1 as well as of FPR2/ALX that are up-regulated to the cell surface when cells are stimulated with inflammatory mediators that mobilize the granules (Sengeløv et al., 1994; Bylund et al., 2002).

Degranulation in fMLF-stimulated neutrophils involves the second messenger diacylglycerol and PKCs that are activated by diacylglycerol and Ca<sup>2+</sup> (Smith et al., 1988). However, fMLF is still able to induce secretory granule release when extracellular and intracellular Ca<sup>2+</sup> is chelated, suggesting the presence of Ca<sup>2+</sup>-independent pathways for degranulation (Sengeløv et al., 1993). It has been reported that adhesion-dependent and fMLF-induced degranulation requires the Src family protein kinase Fgr and Hck, which also involves p38 MAPK (Mócsai et al., 1999, 2000). In addition, the activation of PI3K has been indicated as an important step leading to neutrophil granule release (Arcaro and Wymann, 1993; Thelen et al., 1994). The cGMP-dependent kinase PKG-1 $\alpha$  has been implicated in fMLF-induced degranulation in transfected RBL-2H3 cells expressing FPR1 and in primary neutrophils (Nanamori et al., 2007). Further investigations will be necessary to determine the cross-talk between these kinases and their roles in regulating soluble *N*-ethylmaleimide-sensitive factor attachment protein receptors (SNARE) that mediate vesicular fusion during degranulation.

## B. Other Cellular Functions

**1. Transcriptional Regulation and Anti-Inflammatory Functions.** Although neutrophils are terminally differentiated myeloid cells with special bactericidal functions, these cells retain the ability to synthesize selected proteins, including certain cytokines (Lloyd and Oppenheim, 1992). fMLF has been found to stimulate neutrophil transcriptional regulation and cytokine production (Cassatella et al., 1992). fMLF-induced IL-8 secretion is accompanied by the activation of NF- $\kappa$ B, a nuclear factor



for transcription of a large number of proinflammatory genes (McDonald et al., 1997). NF- $\kappa$ B activation is also mediated by FPR2/ALX, in response to SAA stimulation, which leads to IL-8 secretion (He et al., 2003). Other studies have shown that agonists for FPR2/ALX, such as SAA and WKYMVm, can induce the expression of matrix metalloproteinase in fibroblast-like synoviocytes and monocytes (O'Hara et al., 2004; Sodin-Semrl et al., 2004; Lee et al., 2005). These results are consistent with the ability of certain GPCRs to regulate transcriptional activation that contributes to the proinflammatory activities of the respective ligands (Ye, 2001).

In contrast, FPR2/ALX ligands such as LXA4 and ANXA1 exhibit anti-inflammatory activities (Perretti et al., 2002; Chiang et al., 2006). ANXA1 has been shown to cause detachment of leukocytes and prevent transendothelial migration (diapedesis) (Perretti, 2003). In comparison, the anti-inflammatory effect of LXA4 is shown to involve suppression of proinflammatory gene expression. József et al. (2002) reported that LXA4 analogs attenuated nuclear accumulation of activator protein 1 and NF- $\kappa$ B in both polymorphonuclear and mononuclear leukocytes, resulting in the inhibition of LPS-induced IL-8 mRNA expression and IL-8 release by 50 to 65%. It is not entirely clear whether this action of LXA4 is mediated through FPR2/ALX-dependent negative signaling or blockade of FPR2/ALX binding of and activation by an endogenous, proinflammatory agonist for this receptor. The first possibility was suggested in a study that demonstrated LXA4 induction of NAB1, a transcriptional corepressor induced by glucocorticoids (Qiu et al., 2001). More recently, it was also shown that LXA4 and its analogs induce SOCS-2 expression (Machado et al., 2006) and SOCS-2-dependent proteosomal degradation of TRAF6, a molecule involved in Toll-like receptor signaling (Machado et al., 2008). The utilization of FPR2/ALX in these studies was suggested by the inhibition of LXA4 effects with pertussis toxin treatment (Qiu et al., 2001) and with Boc2 (Machado et al., 2006), although neither is specific for FPR2/ALX. In the latter study, the AhR, which is a receptor for LXA4 (Schaldach et al., 1999), has been shown to also mediate SOCS-2 expression induced by LXA4. The possibility that LXA4 blocks the access of proinflammatory FPR2/ALX ligands is suggested by a more recent study showing that LXA4 can compete off the binding of iodinated WKYMVm in a GPCR heterodimer consisting of BLT1 and a chimeric BLT1 receptor encompassing the third extracellular loop and the seventh transmembrane domain of FPR2/ALX (Damian et al., 2008). The chimeric BLT1 was previously shown to bind LXA4 and share some properties with wild-type FPR2/ALX (Chiang et al., 2000). Therefore, the possibility remains that the anti-inflammatory effect of LXA4 is mediated in part by its blocking effect (i.e., acting as an antagonist of the receptor). This notion is supported by experimental data showing that LXA4 does not have the property of a typical agonist for FPR2/

ALX, despite its ability to stimulate GTPase activity in transfected Chinese hamster ovary cells (Fiore et al., 1994; Takano et al., 1997). In functional assays, most agonists for FPR2/ALX stimulate signaling pathways that lead to calcium mobilization, ERK phosphorylation, and terminal functions of neutrophils. However, LXA4 stimulation does not lead to calcium mobilization in several types of transfected cells or induce neutrophil degranulation and superoxide generation (Su et al., 1999b; Bae et al., 2003; Prat et al., 2006). In contrast, LXA4 has been shown to activate monocytes and the monocytic cell line THP-1 (Maddox et al., 1997). The discrepancy suggests that signaling molecules essential for certain LXA4-induced functions might be missing in neutrophils and epithelial cells. It also reflects that LXA4 lacks full agonistic activities at FPR2/ALX.

**2. Neutrophil Apoptosis.** Neutrophils released to blood circulation have a half-life of 8 to 10 h. If not activated, these cells are destined for apoptosis (Savill, 1997). Stimulation of neutrophils with proinflammatory cytokines such as granulocyte macrophage–colony-stimulating factor, G-CSF, and IL-1 $\beta$ , but not with fMLF, C5a, or IL-8, prolongs the lifespan of neutrophils (Colotta et al., 1992). Other reports have shown that stimulation of neutrophils with fMLF can induce apoptosis, and this process requires superoxide generation (Kettritz et al., 1997). Neutrophil apoptosis and phagocytosis of apoptotic neutrophils are related to resolution of inflammation.

Several ligands for the formyl peptide receptors are found to play different roles in neutrophil apoptosis. SAA suppresses neutrophil apoptosis induced by anti-Fas antibody (Christenson et al., 2008). However, this effect is thought to involve the nucleotide receptor P2X7 and not FPR2/ALX. In comparison, ANXA1 has been shown to accelerate neutrophil apoptosis (Solito et al., 2003), although glucocorticoids that regulate ANXA1 production have antiapoptotic effects (Liles et al., 1995). El Kebir et al. (2007) reported that SAA could stimulate concurrent activation of the ERK and PI3K/Akt signaling pathways, resulting in the phosphorylation of BCL-XL/BCL2-associated death promoter (BAD) protein at Ser112 and Ser136, respectively, thereby preventing collapse of mitochondrial transmembrane potential, cytochrome *c* release, and caspase-3 activation. It is noteworthy that LXA4 treatment could reverse the antiapoptotic effect of SAA. The underlying mechanism has not been fully identified, although ERK and Akt can work together to delay neutrophil apoptosis. These results suggest a potential role of FPR2/ALX in the regulation of neutrophil apoptosis, but other receptors may be involved as well. It will be important to determine how a single class of receptors mediates different functions in cell survival and apoptosis when stimulated with different ligands.

## VII. A New IUPHAR-Approved Nomenclature for the Formyl Peptide Receptor Family

Research conducted in the past 3 decades has shown that members of the human FPR gene family are G protein-coupled receptors expressed primarily in differentiated myeloid cells. All three receptors bind *N*-formylated peptides and respond to these peptides with similar cellular functions. The sequence identity between these receptors (56–69%) is high among G protein-coupled receptors detecting immunomodulatory peptides such as chemokines (Murphy, 1993). Therefore, these receptors are structurally similar and functionally related proteins of one gene family. Recent discovery of multiple agonists for the FPR family of receptors has raised new questions regarding their nomenclature, their physiological functions, and the mechanisms by which one receptor mediates both proinflammatory and anti-inflammatory activities. The pharmacological principle for receptor nomenclature has been based on the specific agonists for the receptor and not sequence homology between the receptor and another protein, as in the case of FPRL1 and FPRL2. This and other considerations for the nomenclature of the FPR family are discussed below.

### A. Nomenclature of FPR1

Most mammalian receptors are known for their abilities to specifically recognize and respond to agonists produced by the host. Although receptor interactions with endogenous agonists are critical for homeostasis and numerous physiological functions, a fundamental basis for innate and acquired immunity is the ability of host cells to recognize exogenous ligands such as bacterial and viral products, known as nonself or foreign antigens. The structural basis for the host to recognize a large number of foreign antigens has been illustrated in both humoral and cytotoxic cell-mediated immune responses. As a first line of host defense against invading microorganisms, cells of the innate immune system serve to detect invading pathogens, and activate their antimicrobial mechanisms, such as phagocytosis, degranulation, and superoxide production. The discovery of FPR1, the first mammalian receptor known to bind *N*-formylated peptides, represents a major milestone in the understanding of how innate immune cells are mobilized when the host is infected with bacteria. The *N*-formylmethionine, characteristic of bacterial protein synthesis, can be considered a PAMP akin to bacterial cell wall peptidoglycans, CpG DNA, and lipopolysaccharides (Medzhitov and Janeway, 2000). Unlike these bacterial products, *N*-formylated peptides may be derived from both bacterial proteins and, in mammalian cells, mitochondrial proteins. Therefore, FPR1 may also serve homeostasis functions. Despite the presence of nonformyl peptides (e.g., acetylated peptides and peptides with no N-terminal modification) as high-affinity li-

gands for FPR1, there is overwhelming evidence for a key role of the *N*-formylmethionine in promoting ligand interaction with the receptor. Studies have shown that addition of an *N*-formylmethionine potentiates binding affinity as well as potency of the peptides tested on FPR1 (Freer et al., 1980, 1982; Chen et al., 1995). A physiological function of FPR1 in host defense is demonstrated in a study using mice lacking *Fpr1*, which displayed increased susceptibility to *L. monocytogenes* infection (Gao et al., 1999). Taken together, numerous studies have shown that FPR1 is a cognate receptor for *N*-formylated peptides of bacterial and mammalian origins.

### B. Nomenclature of FPR2/ALX

The second receptor of the FPR family was initially identified through molecular cloning based on sequence homology to FPR1. It has been referred in the literature as FPR2, FPRL1 (the most frequently used name), FPRH1, HM63, LXA4R, and ALX. Of these, the name FPR2 reflects the ability of this receptor to bind *N*-formyl peptides, albeit at low affinity for fMLF (Ye et al., 1992). The name FPRL1 refers to the sequence homology shared between this receptor and FPR1, which in pharmacological terms does not reflect its ligand binding property. This receptor has been shown to bind a large number of endogenous and exogenous ligands, all peptides except LXA4 and ATLS. Among the characterized endogenous peptides, humanin and CCL23 $\beta$  are highly potent agonists (Elagoz et al., 2004; Harada et al., 2004; Ying et al., 2004; Miao et al., 2007), and SAA is an acute-phase protein with well defined physiological functions in amyloid and inflammatory diseases (Su et al., 1999b; He et al., 2003; O'Hara et al., 2004; Sodin-Semrl et al., 2004; Lee et al., 2006a,b; El Kebir et al., 2007). The primary structures of these peptides have very little in common, and these peptides come from different sources. Therefore, it is difficult to name this receptor after any one of the peptide agonists. Whereas the ALX nomenclature ratified by IUPHAR in 2003 (Brink et al., 2003) precisely indicates that LXA4 and ATLS are endogenous agonists for this receptor, it does not reflect the fact that the majority of the agonists for this receptor are peptides. One prominent feature of this receptor, shared with other members of the FPR family, is the ability to preferentially bind and respond to *N*-formylated peptides compared with nonformylated peptides of the same sequence (Murphy et al., 1992; Ye et al., 1992; Harada et al., 2004; Rabiet et al., 2005). For instance, an *N*-formylated humanin is nearly 300-fold more potent than nonformylated humanin (Harada et al., 2004). Studies conducted by Rabiet et al. (2005) showed that the receptor preferentially binds endogenous *N*-formyl peptides derived from mitochondrial proteins, compared with exogenous *N*-formyl peptides from bacterial sources. This property of the receptor is reflected in the name FPR2, indicating the receptor is the

second member of the FPR family that interacts with *N*-formylated peptides. Based on these considerations as well as the ability of the receptor to bind and mediate the anti-inflammatory effects of LXA4 and ATs, we recommend for this receptor the nomenclature FPR2/ALX (ALX/FPR2 when the lipoxin-binding property is of primary concern).

### C. Nomenclature of FPR3

The third member of the FPR family often appears in the literature as FPR2, a name describing its structural homology to other members of the same family. Compared with FPR1 and FPR2/ALX, fewer ligands have been identified for this receptor. One of its endogenous agonists is F2L, an acetylated N-terminal cleavage product of the mammalian heme-binding protein that is chemotactic to dendritic cells and monocytes (Migeotte et al., 2005). Like FPR2/ALX, this receptor also binds and responds to the mitochondrial peptide fMMY-ALF (Rabiet et al., 2005). Therefore, we recommend the nomenclature of FPR3 for this receptor.

In summary, the presence of an extraordinarily numerous and structurally diverse group of agonistic ligands for the FPR family of receptors has been a challenge to their nomenclature. We recommend the names FPR1, FPR2/ALX, and FPR3 primarily based on a shared property of these receptors to bind *N*-formylated peptides and, for the second member of this family, its ability to bind LXA4 and ATs. Despite the diverse ligand recognition properties, these receptors are genetically clustered and functionally related proteins that play important roles in host defense and inflammation.

## VIII. Unmet Challenges and Future Perspectives

Research on the FPRs conducted in the past 3 decades has generated important information for our understanding of chemoattractant receptors as GPCRs that transduce chemotactic signals in innate immune cells. Despite tremendous progress made in this area of research, a number of issues and questions remain to be addressed in the future.

### A. The Functions of the Formyl Peptide Receptor Family of Receptors in Other Mammals

Most functional studies of the FPRs have been conducted in human neutrophils and, in early studies, rabbit neutrophils. In comparison, much less is known about the functions of FPRs in other species. In mice, deletion of the *Fpr1* gene leads to increased susceptibility to *L. monocytogenes* infection (Gao et al., 1999). This and other studies indicate that mFpr1 is the ortholog of human FPR1 that serves to detect bacterially derived *N*-formylated peptides (Gao and Murphy, 1993; Southgate et al., 2008). However, the number of receptors in the mouse *Fpr* gene family (8) far exceeds the number of receptors in humans (3). The three mouse receptors

found in leukocytes (mFpr1, mFpr2, and mFpr-rs1), which have been studied to some extent, have overlapping structural and functional properties with the three human receptors (Gao and Murphy, 1993; Takano et al., 1997; Gao et al., 1998, 2007; Liang et al., 2000; Vaughn et al., 2002; Wang and Ye, 2002; Southgate et al., 2008). Therefore, although mFpr1 is the ortholog of human FPR1, its putative ligand binding domain contains structural features resembling those of FPR2/ALX. The absence of high-affinity binding for fMLF and the ability to effectively recognize several other *N*-formylated peptides are also features shared between mFpr1 and FPR2/ALX (Rabiet et al., 2005; Southgate et al., 2008). Both mFpr2 and mFpr-rs1 share significant sequence homology with FPR2/ALX and FPR3. It is presently difficult to determine whether the gene product of *Fpr-rs1* or *Fpr-rs2* is the ortholog of FPR2/ALX, because both of the mouse receptors are able to detect and respond to LXA4 (Takano et al., 1997; Vaughn et al., 2002), although LXA4 activation through mFpr2 requires the presence of G $\alpha_{16}$  (Vaughn et al., 2002). Further complicating the nomenclature of the mouse receptors is the recent finding that mFpr2 serves as a receptor for F2L, a highly potent and specific agonist for human FPR3 (Gao et al., 2007). These observations suggest that the human FPRs, especially FPR1 and FPR3, have better defined and more specialized ligand-binding properties than the mouse receptors. Clearly, a great deal of work has yet to be performed to understand the biological functions of these and other distantly related members of the mouse *Fpr* gene family.

### B. Structural Basis for the Diversity of the Formyl Peptide Receptor Family in Ligand Recognition

The human FPRs are able to detect an incredibly diverse group of agonists. More than a dozen ligands of different structures and origins have been found for FPR2/ALX alone, making it one of the most promiscuous GPCRs characterized to date. It is presently unclear how FPR2/ALX is able to bind such a diverse group of agonists that include peptides and the eicosanoid LXA4. Studies employing chimeric receptors between FPR2/ALX and structurally similar receptors have produced useful information suggesting that this receptor has structural properties different from those of the FPR1 (Quehenberger et al., 1993; Chiang et al., 2000). With respect to binding of both peptides and LXA4 by FPR2/ALX, computer-assisted 3-D modeling of the dimensions and spatial volumes for retinal, fMLF, and LXA4 have shown that they are remarkably similar in the three-dimensional spatial volumes (Mills et al., 1999). However, this does not explain why structurally similar receptors such as FPR1 do not interact with LXA4. It is hypothesized that certain features of the FPR2/ALX binding pocket are required for its interaction with a multitude of agonists with different sequence composition and chemical nature. First, the binding pocket must

be sufficiently large to accommodate ligands of various sizes, including longer peptides such as SAA (104 amino acids). Second, the receptor structure involved in ligand binding must be sufficiently flexible for proper contact with a variety of ligands. In this regard, charge-charge interactions such as those found in FPR1 (Mills et al., 1998) may actually restrict ligand access, whereas hydrophobic interactions can facilitate the binding of these ligands. In fact, most agonists for FPR2/ALX are hydrophobic or contain a significant number of hydrophobic residues. Finally, there may be multiple ligand binding sites on FPR2/ALX, thereby permitting its interaction with two or more different classes of agonists. This notion is supported by the diverse structures of the FPR2/ALX ligands, including recently identified synthetic ligands with relatively rigid structures. Moreover, binding studies have shown that the different ligands for FPR2/ALX can only partially compete off radiolabeled WKYMVM or LXA4 (Chiang et al., 2000; Nanamori et al., 2004a). More detailed structural studies will be required to test these and other possibilities and to identify the structural basis for the diverse recognition by FPR2/ALX.

### *C. Mechanisms Used by FPR2/ALX to Respond to Two Different Classes of Ligands for Proinflammatory and Anti-Inflammatory Effects*

LXA4 and its stable analogs have been found to exert potent anti-inflammatory effects in vivo (for review, see Chiang et al., 2006). In comparison, the majority of the peptide agonists for FPR2/ALX stimulate proinflammatory activities in leukocytes through the activation of signaling pathways that are well defined for the G protein-coupled chemoattractant receptors as a class. Studies have shown that LXA4 and its stable analogs induce small G protein-dependent actin reorganization in monocytes but not in neutrophils (Maderna et al., 2002), even though FPR2/ALX is abundant in both types of cells. The mechanism underlying this difference remains undefined. Early characterization of the LXA4 receptor identified it as a high-affinity binding site for the eicosanoid and, when expressed in transfected Chinese hamster ovary cells, could mediate LXA4-stimulated activation of  $G\alpha_i$  proteins (Fiore et al., 1994). However, unlike most peptide agonists for FPR2/ALX, high-affinity binding of LXA4 is not translated into the activation of signaling pathways from the G proteins that couples to the receptor. For instance, LXA4 does not stimulate phosphorylation of ERK1 and ERK2 even when used at micromolar concentrations (Bae et al., 2003). In comparison, several partial agonists for FPR2/ALX such as Quin-C1 are able to induce ERK phosphorylation (Nanamori et al., 2004a). In another study, it was shown that LXA4 could stimulate ERK2 phosphorylation, but through a mechanism independent of FPR2/ALX and FPR3 (Christophe et al., 2002). The same study also demonstrated that LXA4 did not affect neutrophil acti-

vation by WKYMVM, a selective agonist for FPR2/ALX. Apparently, the agonistic activity of LXA4 is not typical compared with most other agonists for FPR2/ALX. This difference may account for the anti-inflammatory effect produced by LXA4 and ATLS. Currently, there is no clear mechanism for the bifurcation of signaling pathways downstream of FPR2/ALX that may lead to proinflammatory and anti-inflammatory functions. It is possible that the anti-inflammatory effects of LXA4 require more than FPR2/ALX. Indeed, published reports indicate that, in addition to ALX, AhR is a receptor for LXA4 (Schaldach et al., 1999). AhR has been shown to be partially responsible for the anti-inflammatory functions of LXA4 in dendritic cells (Machado et al., 2006). Although AhR is structurally different from FPR2/ALX, it has been suggested that both receptors can activate SOCS-2, which is a mechanism for the anti-inflammatory actions (Machado et al., 2006, 2008). The observation that LXB4 does not compete for LXA4 binding in neutrophils but exhibits anti-inflammatory properties similar to that of LXA4 (Fiore et al., 1992; Maddox and Serhan, 1996) suggests the presence of a LXB4 receptor that differs from FPR2/ALX.

The potent anti-inflammatory effects of LXA4 and ATLS have been demonstrated in numerous in vivo studies (for review, see Serhan, 2005; O'Meara et al., 2008). Therefore, it is of great interest to understand how LXA4 and ATLS exert their anti-inflammatory functions through receptor-mediated mechanisms. An ongoing effort is focused on the identification and characterization of peptides and small molecules that serve as ligands for FPR2/ALX and that exhibit anti-inflammatory properties (Bürli et al., 2006; Hecht et al., 2009). Comparative studies of these molecules along with pharmacologically characterized FPR2/ALX ligands, such as WKYMVM, SAA, and LXA4, will likely uncover the different structural and functional changes that the ligands bring to the receptor and the receptor-mediated signaling pathways. It is our hope that future research, including the development of mice genetically deficient in *Fpr-rs1* and *Ahr*, the two reported mouse LXA4 receptors, will lead to a better understanding of the mechanisms whereby GPCRs such as FPR2/ALX transduce multiple signals for different biological functions.

*Acknowledgments.* This work was supported in part by the National Institutes of Health National Institute of Allergy and Infectious Diseases [Grant AI033503]; the National Institutes of Health National Institute of General Medical Sciences [Grant GM066182] (to R.D.Y.); by the Commissariat à l'Energie Atomique, the Centre National de la Recherche Scientifique, and the University Joseph Fourier, France (to F.B.); by the Swedish Medical Research Council [Grant 298-20-3] (to C.D.); and by the Intramural Research Programs of the National Institutes of Health National Cancer Institute (to J.M.W.) and the National Institutes of Health National Institute of Allergy and Infectious Diseases (to P.M.M.). We thank Jing Deng for drawing chemical structures and Benjamin Gantner for critical reading of the manuscript.

## References

- Abo A, Pick E, Hall A, Totty N, Teahan CG, and Segal AW (1991) Activation of the NADPH oxidase involves the small GTP-binding protein p21rac1. *Nature* **353**: 668–670.
- Agarwal S, Reynolds MA, Duckett LD, and Suzuki JB (1989) Altered free cytosolic calcium changes and neutrophil chemotaxis in patients with juvenile periodontitis. *J Periodontol Res* **24**:149–154.
- Ali H, Richardson RM, Haribabu B, and Snyderman R (1999) Chemoattractant receptor cross-desensitization. *J Biol Chem* **274**:6027–6030.
- Ali H, Richardson RM, Tomhave ED, Didsbury JR, and Snyderman R (1993) Differences in phosphorylation of formylpeptide and C5a chemoattractant receptors correlate with differences in desensitization. *J Biol Chem* **268**:24247–24254.
- Aliberti J, Hieng S, Reis e Sousa C, Serhan CN, and Sher A (2002a) Lipoxin-mediated inhibition of IL-12 production by DCs: a mechanism for regulation of microbial immunity. *Nat Immunol* **3**:76–82.
- Aliberti J, Serhan C, and Sher A (2002b) Parasite-induced lipoxin A4 is an endogenous regulator of IL-12 production and immunopathology in *Toxoplasma gondii* infection. *J Exp Med* **196**:1253–1262.
- Allen RA, Tolley JO, and Jesaitis AJ (1986) Preparation and properties of an improved photoaffinity ligand for the N-formyl peptide receptor. *Biochim Biophys Acta* **882**:271–280.
- Almkvist J, Dahlgren C, Leffler H, and Karlsson A (2004) Newcastle disease virus neuraminidase primes neutrophils for stimulation by galectin-3 and formyl-Met-Leu-Phe. *Exp Cell Res* **298**:74–82.
- Almkvist J, Földt J, Dahlgren C, Leffler H, and Karlsson A (2001) Lipopolysaccharide-induced gelatinase granule mobilization primes neutrophils for activation by galectin-3 and formylmethionyl-Leu-Phe. *Infect Immun* **69**:832–837.
- Alvarez V, Coto E, Setién F, González-Roces S, and López-Larrea C (1996) Molecular evolution of the N-formyl peptide and C5a receptors in non-human primates. *Immunogenetics* **44**:446–452.
- Amatruda TT 3rd, Dragas-Graonic S, Holmes R, and Perez HD (1995) Signal transduction by the formyl peptide receptor. Studies using chimeric receptors and site-directed mutagenesis define a novel domain for interaction with G-proteins. *J Biol Chem* **270**:28010–28013.
- Amatruda TT 3rd, Steele DA, Slepak VZ, and Simon MI (1991) G alpha 16, a G protein alpha subunit specifically expressed in hematopoietic cells. *Proc Natl Acad Sci U S A* **88**:5587–5591.
- Arcaro A and Wymann MP (1993) Wortmannin is a potent phosphatidylinositol 3-kinase inhibitor: the role of phosphatidylinositol 3,4,5-trisphosphate in neutrophil responses. *Biochem J* **296**:297–301.
- Aruffo A and Seed B (1987) Molecular cloning of a CD28 cDNA by a high-efficiency COS cell expression system. *Proc Natl Acad Sci U S A* **84**:8573–8577.
- Aswanikumar S, Corcoran B, Schifmann E, Day AR, Freer RJ, Showell HJ, and Becker EL (1977) Demonstration of a receptor on rabbit neutrophils for chemotactic peptides. *Biochem Biophys Res Commun* **74**:810–817.
- Babior BM, Lambeth JD, and Nauseef W (2002) The neutrophil NADPH oxidase. *Arch Biochem Biophys* **397**:342–344.
- Bae YS, Lee HY, Jo EJ, Kim JI, Kang HK, Ye RD, Kwak JY, and Ryu SH (2004) Identification of peptides that antagonize formyl peptide receptor-like 1-mediated signaling. *J Immunol* **173**:607–614.
- Bae YS, Park JC, He R, Ye RD, Kwak JY, Suh PG, and Ho Ryu S (2003) Differential signaling of formyl peptide receptor-like 1 by Trp-Lys-Tyr-Met-Val-Met-CONH<sub>2</sub> or lipoxin A4 in human neutrophils. *Mol Pharmacol* **64**:721–730.
- Baek SH, Seo JK, Chae CB, Suh PG, and Ryu SH (1996) Identification of the peptides that stimulate the phosphoinositide hydrolysis in lymphocyte cell lines from peptide libraries. *J Biol Chem* **271**:8170–8175.
- Bao L, Gerard NP, Eddy RL Jr, Shows TB, and Gerard C (1992) Mapping of genes for the human C5a receptor (C5AR), human FMLP receptor (FPR), and two FMLP receptor homologue orphan receptors (FPRH1, FPRH2) to chromosome 19. *Genomics* **13**:437–440.
- Baranova IN, Vishnyakova TG, Bocharov AV, Kurlander R, Chen Z, Kimelman ML, Remaley AT, Csako G, Thomas F, Eggerman TL, et al. (2005) Serum amyloid A binding to CLA-1 (CD36 and LIMPII analogous-1) mediates serum amyloid A protein-induced activation of ERK1/2 and p38 mitogen-activated protein kinases. *J Biol Chem* **280**:8031–8040.
- Becker EL, Forouhar FA, Grunnet ML, Boulay F, Tardif M, Bormann BJ, Sodja D, Ye RD, Woska JR Jr, and Murphy PM (1998) Broad immunocytochemical localization of the formylpeptide receptor in human organs, tissues, and cells. *Cell Tissue Res* **292**:129–135.
- Bellner L, Thorén F, Nygren E, Liljeqvist JA, Karlsson A, and Eriksson K (2005) A proinflammatory peptide from herpes simplex virus type 2 glycoprotein G affects neutrophil, monocyte, and NK cell functions. *J Immunol* **174**:2235–2241.
- Bennett JP, Hirth KP, Fuchs E, Sarvas M, and Warren GB (1980) The bacterial factors which stimulate neutrophils may be derived from procaryote signal peptides. *FEBS Lett* **116**:57–61.
- Bennett TA, Foutz TD, Gurevich VV, Sklar LA, and Prossnitz ER (2001) Partial phosphorylation of the N-formyl peptide receptor inhibits G protein association independent of arrestin binding. *J Biol Chem* **276**:49195–49203.
- Bentwood BJ and Henson PM (1980) The sequential release of granule constituents from human neutrophils. *J Immunol* **124**:855–862.
- Betten A, Bylund J, Christophe T, Cristophe T, Boulay F, Romero A, Hellstrand K, and Dahlgren C (2001) A proinflammatory peptide from *Helicobacter pylori* activates monocytes to induce lymphocyte dysfunction and apoptosis. *J Clin Invest* **108**:1221–1228.
- Björkman L, Karlsson J, Karlsson A, Rabiet MJ, Boulay F, Fu H, Bylund J, and Dahlgren C (2008) Serum amyloid A mediates human neutrophil production of reactive oxygen species through a receptor independent of formyl peptide receptor like-1. *J Leukoc Biol* **83**:245–253.
- Bokoch GM and Gilman AG (1984) Inhibition of receptor-mediated release of arachidonic acid by pertussis toxin. *Cell* **39**:301–308.
- Bokoch GM, Katada T, Northup JK, Ui M, and Gilman AG (1984) Purification and properties of the inhibitory guanine nucleotide-binding regulatory component of adenylate cyclase. *J Biol Chem* **259**:3560–3567.
- Bommakanti RK, Bokoch GM, Tolley JO, Schreiber RE, Siemsen DW, Klotz KN, and Jesaitis AJ (1992) Reconstitution of a physical complex between the N-formyl chemotactic peptide receptor and G protein. Inhibition by pertussis toxin-catalyzed ADP ribosylation. *J Biol Chem* **267**:7576–7581.
- Bommakanti RK, Dratz EA, Siemsen DW, and Jesaitis AJ (1994) Characterization of complex formation between Gi2 and octyl glucoside solubilized neutrophil N-formyl peptide chemoattractant receptor by sedimentation velocity. *Biochim Biophys Acta* **1209**:69–76.
- Bommakanti RK, Dratz EA, Siemsen DW, and Jesaitis AJ (1995) Extensive contact between Gi2 and N-formyl peptide receptor of human neutrophils: mapping of binding sites using receptor-mimetic peptides. *Biochemistry* **34**:6720–6728.
- Bommakanti RK, Klotz KN, Dratz EA, and Jesaitis AJ (1993) A carboxyl-terminal tail peptide of neutrophil chemotactic receptor disrupts its physical complex with G protein. *J Leukoc Biol* **54**:572–577.
- Bonnans C, Gras D, Chavis C, Mainprice B, Vachier I, Godard P, and Chanez P (2007) Synthesis and anti-inflammatory effect of lipoxins in human airway epithelial cells. *Biomed Pharmacother* **61**:261–267.
- Borregaard N, Lollike K, Kjeldsen L, Sengeløv H, Bastholm L, Nielsen MH, and Bainton DF (1993) Human neutrophil granules and secretory vesicles. *Eur J Haematol* **51**:187–198.
- Boulay F, Tardif M, Brouchon L, and Vignais P (1990a) The human N-formylpeptide receptor. Characterization of two cDNA isolates and evidence for a new subfamily of G-protein-coupled receptors. *Biochemistry* **29**:11123–11133.
- Boulay F, Tardif M, Brouchon L, and Vignais P (1990b) Synthesis and use of a novel N-formyl peptide derivative to isolate a human N-formyl peptide receptor cDNA. *Biochem Biophys Res Commun* **168**:1103–1109.
- Boulven I, Levasseur S, Marois S, Paré G, Rollet-Labelle E, and Naccache PH (2006) Class IA phosphatidylinositol 3-kinases, rather than p110 gamma, regulate formylmethionyl-leucyl-phenylalanine-stimulated chemotaxis and superoxide production in differentiated neutrophil-like PLB-985 cells. *J Immunol* **176**:7621–7627.
- Boxer LA, Yoder M, Bonsib S, Schmidt M, Ho P, Jersild R, and Baehner RL (1979) Effects of a chemotactic factor, N-formylmethionyl peptide, on adherence, superoxide anion generation, phagocytosis, and microtubule assembly of human polymorphonuclear leukocytes. *J Lab Clin Med* **93**:506–514.
- Brazil TJ, Rossi AG, Haslett C, McGorum B, Dixon PM, and Chilvers ER (1998) Priming induces functional coupling of N-formyl-methionyl-leucyl-phenylalanine receptors in equine neutrophils. *J Leukoc Biol* **63**:380–388.
- Brink C, Dahlén SE, Drazen J, Evans JF, Hay DW, Nicotia S, Serhan CN, Shimizu T, and Yokomizo T (2003) International Union of Pharmacology. XXXVII. Nomenclature for leukotriene and lipoxin receptors. *Pharmacol Rev* **55**:195–227.
- Brown GE, Stewart MQ, Bissonnette SA, Elia AE, Wilker E, and Yaffe MB (2004) Distinct ligand-dependent roles for p38 MAPK in priming and activation of the neutrophil NADPH oxidase. *J Biol Chem* **279**:27059–27068.
- Burg M, Raffetseder U, Grove M, Klos A, Köhl J, and Bautsch W (1995) G alpha-16 complements the signal transduction cascade of chemotactic receptors for complement factor C5a (C5a-R) and N-formylated peptides (fMLF-R) in *Xenopus* oocytes: G alpha-16 couples to chemotactic receptors in *Xenopus* oocytes. *FEBS Lett* **377**:426–428.
- Bürli RW, Xu H, Zou X, Muller K, Golden J, Frohn M, Adlam M, Plant MH, Wong M, McElvain M, et al. (2006) Potent hFPR1 (ALXR) agonists as potential anti-inflammatory agents. *Bioorg Med Chem Lett* **16**:3713–3718.
- Bylund J, Karlsson A, Boulay F, and Dahlgren C (2002) Lipopolysaccharide-induced granule mobilization and priming of the neutrophil response to *Helicobacter pylori* peptide Hp(2–20), which activates formyl peptide receptor-like 1. *Infect Immun* **70**:2908–2914.
- Cai H, Song C, Endoh I, Goyette J, Jessup W, Freedman SB, McNeil HP, and Gezey CL (2007) Serum amyloid A induces monocyte tissue factor. *J Immunol* **178**:1852–1860.
- Cai L, de Beer MC, de Beer FC, and van der Westhuyzen DR (2005) Serum amyloid A is a ligand for scavenger receptor class B type I and inhibits high density lipoprotein binding and selective lipid uptake. *J Biol Chem* **280**:2954–2961.
- Campbell JJ, Foxman EF, and Butcher EC (1997) Chemoattractant receptor cross talk as a regulatory mechanism in leukocyte adhesion and migration. *Eur J Immunol* **27**:2571–2578.
- Camps M, Carozzi A, Schnabel P, Scheer A, Parker PJ, and Gierschik P (1992a) Isozyme-selective stimulation of phospholipase C-beta 2 by G protein beta gamma subunits. *Nature* **360**:684–686.
- Camps M, Hou C, Sidiropoulos D, Stock JB, Jakobs KH, and Gierschik P (1992b) Stimulation of phospholipase C by guanine-nucleotide-binding protein beta gamma subunits. *Eur J Biochem* **206**:821–831.
- Carp H (1982) Mitochondrial N-formylmethionyl proteins as chemoattractants for neutrophils. *J Exp Med* **155**:264–275.
- Cassatella MA, Bazzoni F, Ceska M, Ferro I, Baggiolini M, and Berton G (1992) IL-8 production by human polymorphonuclear leukocytes. The chemoattractant formylmethionyl-leucyl-phenylalanine induces the gene expression and release of IL-8 through a pertussis toxin-sensitive pathway. *J Immunol* **148**:3216–3220.
- Chen J, Bernstein HS, Chen M, Wang L, Ishii M, Turck CW, and Coughlin SR (1995) Tethered ligand library for discovery of peptide agonists. *J Biol Chem* **270**:23398–23401.
- Chen K, Iribarren P, Hu J, Chen J, Gong W, Cho EH, Lockett S, Dunlop NM, and Wang JM (2006) Activation of Toll-like receptor 2 on microglia promotes cell uptake of Alzheimer disease-associated amyloid beta peptide. *J Biol Chem* **281**: 3651–3659.
- Chen Q, Powell DW, Rane MJ, Singh S, Butt W, Klein JB, and McLeish KR (2003) Akt phosphorylates p47phox and mediates respiratory burst activity in human neutrophils. *J Immunol* **170**:5302–5308.
- Chen Q, Wade D, Kurosaka K, Wang ZY, Oppenheim JJ, and Yang D (2004) Temporin A and related frog antimicrobial peptides use formyl peptide receptor-like 1 as a receptor to chemoattract phagocytes. *J Immunol* **173**:2652–2659.

- Chen X, Mellon RD, Yang L, Dong H, Oppenheim JJ, and Howard OM (2002) Regulatory effects of deoxycholic acid, a component of the anti-inflammatory traditional Chinese medicine Niu Huang, on human leukocyte response to chemoattractants. *Biochem Pharmacol* **63**:533–541.
- Chen X, Yang D, Shen W, Dong HF, Wang JM, Oppenheim JJ, and Howard MZ (2000) Characterization of chenodeoxycholic acid as an endogenous antagonist of the G-coupled formyl peptide receptors. *Inflamm Res* **49**:744–755.
- Cheng N, He R, Tian J, Dinauer MC, and Ye RD (2007) A critical role of protein kinase C delta activation loop phosphorylation in formyl-methionyl-leucyl-phenylalanine-induced phosphorylation of p47(phox) and rapid activation of nicotinamide adenine dinucleotide phosphate oxidase. *J Immunol* **179**:7720–7728.
- Cheng N, He R, Tian J, Ye PP, and Ye RD (2008) Cutting edge: TLR2 is a functional receptor for acute-phase serum amyloid A. *J Immunol* **181**:22–26.
- Chiang N, Fierro IM, Gronert K, and Serhan CN (2000) Activation of lipoxin A(4) receptors by aspirin-triggered lipoxins and select peptides evokes ligand-specific responses in inflammation. *J Exp Med* **191**:1197–1208.
- Chiang N, Serhan CN, Dahlén SE, Drazen JM, Hay DW, Rovati GE, Shimizu T, Yokomizo T, and Brink C (2006) The lipoxin receptor ALX: potent ligand-specific and stereoselective actions in vivo. *Pharmacol Rev* **58**:463–487.
- Christenson K, Björkman L, Tängemo C, and Bylund J (2008) Serum amyloid A inhibits apoptosis of human neutrophils via a P2X7-sensitive pathway independent of formyl peptide receptor-like 1. *J Leukoc Biol* **83**:139–148.
- Christophe T, Karlsson A, Dugave C, Rabiet MJ, Boulay F, and Dahlgren C (2001) The synthetic peptide Trp-Lys-Tyr-Met-Val-Met-NH<sub>2</sub> specifically activates neutrophils through FPRL1/lipoxin A4 receptors and is an agonist for the orphan monocyte-expressed chemoattractant receptor FPRL2. *J Biol Chem* **276**:21585–21593.
- Christophe T, Karlsson A, Rabiet MJ, Boulay F, and Dahlgren C (2002) Phagocyte activation by Trp-Lys-Tyr-Met-Val-Met, acting through FPRL1/LXA4R, is not affected by lipoxin A4. *Scand J Immunol* **56**:470–476.
- Clark RA, Page RC, and Wilde G (1977) Defective neutrophil chemotaxis in juvenile periodontitis. *Infect Immun* **18**:694–700.
- Colgan SP, Serhan CN, Parkos CA, Delp-Archer C, and Madara JL (1993) Lipoxin A4 modulates transmigration of human neutrophils across intestinal epithelial monolayers. *J Clin Invest* **92**:75–82.
- Colotta F, Re F, Polentarutti N, Sozzani S, and Mantovani A (1992) Modulation of granulocyte survival and programmed cell death by cytokines and bacterial products. *Blood* **80**:2012–2020.
- Condliffe AM, Davidson K, Anderson KE, Ellison CD, Crabbe T, Okkenhaug K, Vanhaesebroeck B, Turner M, Webb L, Wymann MP, et al. (2005) Sequential activation of class IB and class IA PI3K is important for the primed respiratory burst of human but not murine neutrophils. *Blood* **106**:1432–1440.
- Cui Y, Le Y, Yazawa H, Gong W, and Wang JM (2002) Potential role of the formyl peptide receptor-like 1 (FPRL1) in inflammatory aspects of Alzheimer's disease. *J Leukoc Biol* **72**:628–635.
- Czapiga M, Gao JL, Kirk A, and Lekstrom-Himes J (2005) Human platelets exhibit chemotaxis using functional N-formyl peptide receptors. *Exp Hematol* **33**:73–84.
- Damian M, Mary S, Martin A, Pin JP, and Banères JL (2008) G protein activation by the leukotriene B4 receptor dimer. Evidence for an absence of trans-activation. *J Biol Chem* **283**:21084–21092.
- D'Andrea AD, Lodish HF, and Wong GG (1989) Expression cloning of the murine erythropoietin receptor. *Cell* **57**:277–285.
- Dang PM, Fontayne A, Hakim J, El Benna J, and Périannin A (2001) Protein kinase C zeta phosphorylates a subset of selective sites of the NADPH oxidase component p47phox and participates in formyl peptide-mediated neutrophil respiratory burst. *J Immunol* **166**:1206–1213.
- De Lean A, Stadel JM, and Lefkowitz RJ (1980) A ternary complex model explains the agonist-specific binding properties of the adenylate cyclase-coupled  $\beta$ -adrenergic receptor. *J Biol Chem* **255**:7108–7117.
- de Paulis A, Montuori N, Prevede N, Fiorentino I, Rossi FW, Visconte V, Rossi G, Marone G, and Ragno P (2004a) Urokinase induces basophil chemotaxis through a urokinase receptor epitope that is an endogenous ligand for formyl peptide receptor-like 1 and -like 2. *J Immunol* **173**:5739–5748.
- de Paulis A, Prevede N, Fiorentino I, Walls AF, Curto M, Petraroli A, Castaldo V, Ceppa P, Fiocca R, and Marone G (2004b) Basophils infiltrate human gastric mucosa at sites of *Helicobacter pylori* infection, and exhibit chemotaxis in response to *H. pylori*-derived peptide Hp(2–20). *J Immunol* **172**:7734–7743.
- DeLeo FR and Quinn MT (1996) Assembly of the phagocyte NADPH oxidase: molecular interaction of oxidase proteins. *J Leukoc Biol* **60**:677–691.
- DeLeo FR, Renee J, McCormick S, Nakamura M, Apicella M, Weiss JP, and Nauseef WM (1998) Neutrophils exposed to bacterial lipopolysaccharide upregulate NADPH oxidase assembly. *J Clin Invest* **101**:455–463.
- Deng X, Ueda H, Su SB, Gong W, Dunlop NM, Gao JL, Murphy PM, and Wang JM (1999) A synthetic peptide derived from human immunodeficiency virus type 1 gp120 downregulates the expression and function of chemokine receptors CCR5 and CXCR4 in monocytes by activating the 7-transmembrane G-protein-coupled receptor FPRL1/LXA4R. *Blood* **94**:1165–1173.
- Derian CK, Solomon HF, Higgins JD 3rd, Beblavy MJ, Santulli RJ, Bridger GJ, Pike MC, Kroon DJ, and Fischman AJ (1996) Selective inhibition of N-formylpeptide-induced neutrophil activation by carbamate-modified peptide analogues. *Biochemistry* **35**:1265–1269.
- Devchand PR, Arita M, Hong S, Bannenberg G, Moussignac RL, Gronert K, and Serhan CN (2003) Human ALX receptor regulates neutrophil recruitment in transgenic mice: roles in inflammation and host defense. *Faseb J* **17**:652–659.
- Dewas C, Dang PM, Gougerot-Pocidalo MA, and El-Benna J (2003) TNF- $\alpha$  induces phosphorylation of p47(phox) in human neutrophils: partial phosphorylation of p47phox is a common event of priming of human neutrophils by TNF- $\alpha$  and granulocyte-macrophage colony-stimulating factor. *J Immunol* **171**:4392–4398.
- Dewas C, Fay M, Gougerot-Pocidalo MA, and El-Benna J (2000) The mitogen-activated protein kinase extracellular signal-regulated kinase 1/2 pathway is involved in formyl-methionyl-leucyl-phenylalanine-induced p47phox phosphorylation in human neutrophils. *J Immunol* **165**:5238–5244.
- Didsbury JR, Uhing RJ, Tomhave E, Gerard C, Gerard N, and Snyderman R (1991) Receptor class desensitization of leukocyte chemoattractant receptors. *Proc Natl Acad Sci U S A* **88**:11564–11568.
- Didsbury JR, Uhing RJ, Tomhave E, Gerard C, Gerard N, and Snyderman R (1992) Functional high efficiency expression of cloned leukocyte chemoattractant receptor cDNAs. *FEBS Lett* **297**:275–279.
- Dixon RA, Kobilka BK, Strader DJ, Benovic JL, Dohlman HG, Frielle T, Bolanowski MA, Bennett CD, Rands E, Diehl RE, et al. (1986) Cloning of the gene and cDNA for mammalian beta-adrenergic receptor and homology with rhodopsin. *Nature* **321**:75–79.
- Dohlman HG, Thorner J, Caron MG, and Lefkowitz RJ (1991) Model systems for the study of seven-transmembrane-segment receptors. *Annu Rev Biochem* **60**:653–688.
- Dolmatch B and Nield J (1983) Formyl peptide chemotactic receptor. Evidence for an active proteolytic fragment. *J Biol Chem* **258**:7570–7577.
- Donabedian H and Gallin JI (1981) Deactivation of human neutrophil chemotaxis by chemoattractants: effect on receptors for the chemotactic factor f-Met-Leu-Phe. *J Immunol* **127**:839–844.
- Dong X, Mo Z, Bokoch G, Guo C, Li Z, and Wu D (2005) P-Rex1 is a primary Rac2 guanine nucleotide exchange factor in mouse neutrophils. *Curr Biol* **15**:1874–1879.
- Dormann D, Weijer G, Parent CA, Devreotes PN, and Weijer CJ (2002) Visualizing PI3 kinase-mediated cell-cell signaling during Dictyostelium development. *Curr Biol* **12**:1178–1188.
- Dutil EM, Tokar A, and Newton AC (1998) Regulation of conventional protein kinase C isozymes by phosphoinositide-dependent kinase 1 (PKD-1). *Curr Biol* **8**:1366–1375.
- Edwards BS, Bologna C, Young SM, Balakin KV, Prossnitz ER, Savchuck NP, Sklar LA, and Oprea TI (2005) Integration of virtual screening with high-throughput flow cytometry to identify novel small molecule formylpeptide receptor antagonists. *Mol Pharmacol* **68**:1301–1310.
- El Benna J, Han J, Park JW, Schmid E, Ulevitch RJ, and Babior BM (1996) Activation of p38 in stimulated human neutrophils: phosphorylation of the oxidase component p47phox by p38 and ERK but not by JNK. *Arch Biochem Biophys* **334**:395–400.
- El Kebir D, Jözsef L, Khreiss T, Pan W, Petasis NA, Serhan CN, and Filep JG (2007) Aspirin-triggered lipoxins override the apoptosis-delaying action of serum amyloid A in human neutrophils: a novel mechanism for resolution of inflammation. *J Immunol* **179**:616–622.
- Elagöz A, Henderson D, Babu PS, Salter S, Grahames C, Bowers L, Roy MO, Laplante P, Grazzini E, Ahmad S, et al. (2004) A truncated form of CKbeta-1 is a potent agonist for human formyl peptide-receptor-like 1 receptor. *Br J Pharmacol* **141**:37–46.
- Ernst S, Lange C, Wilbers A, Goebeler V, Gerke V, and Rescher U (2004a) An annexin 1 N-terminal peptide activates leukocytes by triggering different members of the formyl peptide receptor family. *J Immunol* **172**:7669–7676.
- Ernst S, Zobiack N, Boecker K, Gerke V, and Rescher U (2004b) Agonist-induced trafficking of the low-affinity formyl peptide receptor FPRL1. *Cell Mol Life Sci* **61**:1684–1692.
- Finney DA and Sklar LA (1983) Ligand/receptor internalization: a kinetic, flow cytometric analysis of the internalization of N-formyl peptides by human neutrophils. *Cytometry* **4**:54–60.
- Fiore S, Maddox JF, Perez HD, and Serhan CN (1994) Identification of a human cDNA encoding a functional high affinity lipoxin A4 receptor. *J Exp Med* **180**:253–260.
- Fiore S, Romano M, Reardon EM, and Serhan CN (1993) Induction of functional lipoxin A4 receptors in HL-60 cells. *Blood* **81**:3395–3403.
- Fiore S, Ryeom SW, Weller PF, and Serhan CN (1992) Lipoxin recognition sites. Specific binding of labeled lipoxin A4 with human neutrophils. *J Biol Chem* **267**:16168–16176.
- Fletcher MP and Gallin JI (1983) Human neutrophils contain an intracellular pool of putative receptors for the chemoattractant N-formyl-methionyl-leucyl-phenylalanine. *Blood* **62**:792–799.
- Foxman EF, Campbell JJ, and Butcher EC (1997) Multistep navigation and the combinatorial control of leukocyte chemotaxis. *J Cell Biol* **139**:1349–1360.
- Foxman EF, Kunkel EJ, and Butcher EC (1999) Integrating conflicting chemotactic signals. The role of memory in leukocyte navigation. *J Cell Biol* **147**:577–588.
- Freer RJ, Day AR, Muthukumaraswamy N, Pinon D, Wu A, Showell HJ, and Becker EL (1982) Formyl peptide chemoattractants: a model of the receptor on rabbit neutrophils. *Biochemistry* **21**:257–263.
- Freer RJ, Day AR, Radding JA, Schiffmann E, Aswanikumar S, Showell HJ, and Becker EL (1980) Further studies on the structural requirements for synthetic peptide chemoattractants. *Biochemistry* **19**:2404–2410.
- Fu H, Björkman L, Janney P, Karlsson A, Karlsson J, Movitz C, and Dahlgren C (2004a) The two neutrophil members of the formylpeptide receptor family activate the NADPH-oxidase through signals that differ in sensitivity to a gelsolin derived phosphoinositide-binding peptide. *BMC Cell Biol* **5**:50.
- Fu H, Bylund J, Karlsson A, Pellmé S, and Dahlgren C (2004b) The mechanism for activation of the neutrophil NADPH-oxidase by the peptides formyl-Met-Leu-Phe and Trp-Lys-Tyr-Met-Val-Met differs from that for interleukin-8. *Immunology* **112**:201–210.
- Fu H, Dahlgren C, and Bylund J (2003) Subinhibitory concentrations of the deformed-lase inhibitor actinonin increase bacterial release of neutrophil-activating peptides: a new approach to antimicrobial chemotherapy. *Antimicrob Agents Chemother* **47**:2545–2550.
- Funamoto S, Meili R, Lee S, Parry L, and Firtel RA (2002) Spatial and temporal regulation of 3-phosphoinositides by PI 3-kinase and PTEN mediates chemotaxis. *Cell* **109**:611–623.
- Furlan F, Orlando S, Laudanna C, Resnati M, Basso V, Blasi F, and Mondino A (2004) The soluble D2D3(88–274) fragment of the urokinase receptor inhibits monocyte chemotaxis and integrin-dependent cell adhesion. *J Cell Sci* **117**:2909–2916.

- Gao JL, Becker EL, Freer RJ, Muthukumaraswamy N, and Murphy PM (1994) A high potency nonformylated peptide agonist for the phagocyte N-formylpeptide chemotactic receptor. *J Exp Med* **180**:2191–2197.
- Gao JL, Chen H, Filie JD, Kozak CA, and Murphy PM (1998) Differential expansion of the N-formylpeptide receptor gene cluster in human and mouse. *Genomics* **51**:270–276.
- Gao JL, Guillaubert A, Hu J, Le Y, Urizar E, Seligman E, Fang KJ, Yuan X, Imbault V, Communi D, et al. (2007) F2L, a peptide derived from heme-binding protein, chemoattracts mouse neutrophils by specifically activating Fpr2, the low-affinity N-formylpeptide receptor. *J Immunol* **178**:1450–1456.
- Gao JL, Lee EJ, and Murphy PM (1999) Impaired antibacterial host defense in mice lacking the N-formylpeptide receptor. *J Exp Med* **189**:657–662.
- Gao JL and Murphy PM (1993) Species and subtype variants of the N-formyl peptide chemotactic receptor reveal multiple important functional domains. *J Biol Chem* **268**:25395–25401.
- Gargiulo L, Longanesi-Cattani I, Bifulco K, Franco P, Raiola R, Campiglia P, Grieco P, Peluso G, Stoppelli MP, and Carriero MV (2005) Cross-talk between fMLP and vitronectin receptors triggered by urokinase receptor-derived SRSRY peptide. *J Biol Chem* **280**:25225–25232.
- Gavins FN, Yona S, Kamal AM, Flower RJ, and Perretti M (2003) Leukocyte antiadhesive actions of annexin 1: ALXR- and FPR-related anti-inflammatory mechanisms. *Blood* **101**:4140–4147.
- Gewirtz AT, Neish AS, and Madara JL (2002) Mechanisms of active intestinal inflammation and potential down-regulation via lipoxins. *Adv Exp Med Biol* **507**:229–236.
- Gierschik P, Sidiropoulos D, and Jakobs KH (1989) Two distinct G<sub>i</sub>-proteins mediate formyl peptide receptor signal transduction in human leukemia (HL-60) cells. *J Biol Chem* **264**:21470–21473.
- Gilbert TL, Bennett TA, Maestas DC, Cimino DF, and Prossnitz ER (2001) Internalization of the human N-formyl peptide and C5a chemoattractant receptors occurs via clathrin-independent mechanisms. *Biochemistry* **40**:3467–3475.
- Godson C, Mitchell S, Harvey K, Petasis NA, Hogg N, and Brady HR (2000) Cutting edge: lipoxins rapidly stimulate nonphlogistic phagocytosis of apoptotic neutrophils by monocyte-derived macrophages. *J Immunol* **164**:1663–1667.
- Goh J, Baird AW, O'Keane C, Watson RW, Cottell D, Bernasconi G, Petasis NA, Godson C, Brady HR, and MacMathuna P (2001) Lipoxin A(4) and aspirin-triggered 15-epi-lipoxin A(4) antagonize TNF-alpha-stimulated neutrophil-enterocyte interactions in vitro and attenuate TNF-alpha-induced chemokine release and colonocyte apoptosis in human intestinal mucosa ex vivo. *J Immunol* **167**:2772–2780.
- Goodman OB Jr, Krupnick JG, Santini F, Gurevich VV, Penn RB, Gagnon AW, Keen JH, and Benovic JL (1996) Beta-arrestin acts as a clathrin adaptor in endocytosis of the beta2-adrenergic receptor. *Nature* **383**:447–450.
- Gripentrog JM, Jesaitis AJ, and Miettinen HM (2000) A single amino acid substitution (N297A) in the conserved NPXXY sequence of the human N-formyl peptide receptor results in inhibition of desensitization and endocytosis, and a dose-dependent shift in p42/44 mitogen-activated protein kinase activation and chemotaxis. *Biochem J* **352**:399–407.
- Gripentrog JM and Miettinen HM (2008) Formyl peptide receptor-mediated ERK1/2 activation occurs through G(i) and is not dependent on beta-arrestin1/2. *Cell Signal* **20**:424–431.
- Gripentrog JM, Mills JS, Saari GJ, and Miettinen HM (2008) Variable responses of formyl peptide receptor haplotypes toward bacterial peptides. *Immunogenetics* **60**:83–93.
- Grumbach Y, Quynh NV, Chiron R, and Urbach V (2009) LXA4 stimulates ZO-1 expression and transepithelial electrical resistance in human airway epithelial (16HBE14o-) cells. *Am J Physiol Lung Cell Mol Physiol* **296**:L101–108.
- Guichard C, Pedruzzi E, Dewas C, Fay M, Pouzet C, Bens M, Vandewalle A, Ogier-Denis E, Gougerot-Pocidallo MA, and Elblim C (2005) Interleukin-8-induced priming of neutrophil oxidative burst requires sequential recruitment of NADPH oxidase components into lipid rafts. *J Biol Chem* **280**:37021–37032.
- Gulden PH, Fischer P 3rd, Sherman NE, Wang W, Engelhard VH, Shabanowitz J, Hunt DF, and Pamer EG (1996) A Listeria monocytogenes pentapeptide is presented to cytolytic T lymphocytes by the H2-M3 MHC class Ib molecule. *Immunology* **5**:73–79.
- Gunji T, Onouchi Y, Nagasawa T, Katagiri S, Watanabe H, Kobayashi H, Arakawa S, Noguchi K, Hata A, Izumi Y, et al. (2007) Functional polymorphisms of the FPR1 gene and aggressive periodontitis in Japanese. *Biochem Biophys Res Commun* **364**:7–13.
- Gwinn MR, Sharma A, and De Nardin E (1999) Single nucleotide polymorphisms of the N-formyl peptide receptor in localized juvenile periodontitis. *J Periodontol* **70**:1194–1201.
- Haas PJ, de Haas CJ, Kleibeuker W, Poppelier MJ, van Kessel KP, Kruijtz JA, Liskamp RM, and van Strijp JA (2004) N-terminal residues of the chemotaxis inhibitory protein of Staphylococcus aureus are essential for blocking formylated peptide receptor but not C5a receptor. *J Immunol* **173**:5704–5711.
- Hachicha M, Pouliot M, Petasis NA, and Serhan CN (1999) Lipoxin (LX)A4 and aspirin-triggered 15-epi-LXA4 inhibit tumor necrosis factor 1alpha-initiated neutrophil responses and trafficking: regulators of a cytokine-chemokine axis. *J Exp Med* **189**:1923–1930.
- Han M, Gurevich VV, Vishnivetskii SA, Sigler PB, and Schubert C (2001) Crystal structure of beta-arrestin at 1.9 Å: possible mechanism of receptor binding and membrane translocation. *Structure* **9**:869–880.
- Harada M, Habata Y, Hosoya M, Nishi K, Fujii R, Kobayashi M, and Hinuma S (2004) N-Formylated humanin activates both formyl peptide receptor-like 1 and 2. *Biochem Biophys Res Commun* **324**:255–261.
- HARRIS H (1954) Role of chemotaxis in inflammation. *Physiol Rev* **34**:529–562.
- Hartt JK, Barish G, Murphy PM, and Gao JL (1999) N-formylpeptides induce two distinct concentration optima for mouse neutrophil chemotaxis by differential interaction with two N-formylpeptide receptor (FPR) subtypes. Molecular characterization of FPR2, a second mouse neutrophil FPR. *J Exp Med* **190**:741–747.
- Hartt JK, Liang T, Sahagun-Ruiz A, Wang JM, Gao JL, and Murphy PM (2000) The HIV-1 cell entry inhibitor T-20 potentially chemoattracts neutrophils by specifically activating the N-formylpeptide receptor. *Biochem Biophys Res Commun* **272**:699–704.
- Hashimoto Y, Niikura T, Ito Y, Sudo H, Hata M, Arakawa E, Abe Y, Kita Y, and Nishimoto I (2001) Detailed characterization of neuroprotection by a rescue factor humanin against various Alzheimer's disease-relevant insults. *J Neurosci* **21**:9235–9245.
- Hayhoe RP, Kamal AM, Solito E, Flower RJ, Cooper D, and Perretti M (2006) Annexin 1 and its bioactive peptide inhibit neutrophil-endothelium interactions under flow: indication of distinct receptor involvement. *Blood* **107**:2123–2130.
- He R, Browning DD, and Ye RD (2001) Differential roles of the NPXXY motif in formyl peptide receptor signaling. *J Immunol* **166**:4099–4105.
- He R, Nanamori M, Sang H, Yin H, Dinauer MC, and Ye RD (2004) Reconstitution of chemotactic peptide-induced nicotinamide adenine dinucleotide phosphate (reduced) oxidase activation in transgenic COS-phox cells. *J Immunol* **173**:7462–7470.
- He R, Sang H, and Ye RD (2003) Serum amyloid A induces IL-8 secretion through a G protein-coupled receptor, FPR1/LXA4R. *Blood* **101**:1572–1581.
- Hecht I, Rong J, Sampaio AL, Hermesh C, Rutledge C, Shemesh R, Toporik A, Beiman M, Dassa L, Niv H, et al. (2009) A novel peptide agonist of formyl-peptide receptor-like 1 (ALX) displays anti-inflammatory and cardioprotective effects. *J Pharmacol Exp Ther* **328**:426–434.
- Heit B, Robbins SM, Downey CM, Guan Z, Colarusso P, Miller BJ, Jirik FR, and Kubus P (2008) PTEN functions to 'prioritize' chemotactic cues and prevent 'distraction' in migrating neutrophils. *Nat Immunol* **9**:743–752.
- Heit B, Tavener S, Rahaarjo E, and Kubus P (2002) An intracellular signaling hierarchy determines direction of migration in opposing chemotactic gradients. *J Cell Biol* **159**:91–102.
- Hepler JR, Kozasa T, Smrcka AV, Simon MI, Rhee SG, Sternweis PC, and Gilman AG (1993) Purification from Sf9 cells and characterization of recombinant G<sub>αq</sub> and G<sub>11α</sub>. Activation of purified phospholipase C isozymes by G<sub>α</sub> subunits. *J Biol Chem* **268**:14367–14375.
- Higgins JD 3rd, Bridger GJ, Derian CK, Beblavy MJ, Hernandez PE, Gaul FE, Abrams MJ, Pike MC, and Solomon HF (1996) N-terminus urea-substituted chemotactic peptides: new potent agonists and antagonists toward the neutrophil fMLF receptor. *J Med Chem* **39**:1013–1015.
- Hirsch E, Katanaev VL, Garlanda C, Azzolino O, Pirolo L, Silengo L, Sozzani S, Mantovani A, Altruda F, and Wymann MP (2000) Central role for G protein-coupled phosphoinositide 3-kinase gamma in inflammation. *Science* **287**:1049–1053.
- Hsu MH, Chiang SC, Ye RD, and Prossnitz ER (1997) Phosphorylation of the N-formyl peptide receptor is required for receptor internalization but not chemotaxis. *J Biol Chem* **272**:29426–29429.
- Hu JY, Le Y, Gong W, Dunlop NM, Gao JL, Murphy PM, and Wang JM (2001) Synthetic peptide MMK-1 is a highly specific chemotactic agonist for leukocyte FPR1. *J Leukoc Biol* **70**:155–161.
- Huet E, Boulay F, Barral S, and Rabiet MJ (2007) The role of beta-arrestins in the formyl peptide receptor-like 1 internalization and signaling. *Cell Signal* **19**:1939–1948.
- Hughes BJ, Hollers JC, Crockett-Torabi E, and Smith CW (1992) Recruitment of CD11b/CD18 to the neutrophil surface and adherence-dependent cell locomotion. *J Clin Invest* **90**:1687–1696.
- Jesaitis AJ, Naemura JR, Sklar LA, Cochrane CG, and Painter RG (1984) Rapid modulation of N-formyl chemotactic peptide receptors on the surface of human granulocytes: formation of high-affinity ligand-receptor complexes in transient association with cytoskeleton. *J Cell Biol* **98**:1378–1387.
- Jesaitis AJ, Tolley JO, Painter RG, Sklar LA, and Cochrane CG (1985) Membrane-cytoskeleton interactions and the regulation of chemotactic peptide-induced activation of human granulocytes: the effects of dihydrocytochalasin B. *J Cell Biochem* **27**:241–253.
- Jiang H, Kuang Y, Wu Y, Xie W, Simon MI, and Wu D (1997) Roles of phospholipase C beta2 in chemoattractant-elicited responses. *Proc Natl Acad Sci U S A* **94**:7971–7975.
- Jones BE, Miettinen HM, Jesaitis AJ, and Mills JS (2003) Mutations of F110 and C126 of the formyl peptide receptor interfere with G-protein coupling and chemotaxis. *J Periodontol* **74**:475–484.
- József L, Zouki C, Petasis NA, Serhan CN, and Filep JG (2002) Lipoxin A4 and aspirin-triggered 15-epi-lipoxin A4 inhibit peroxynitrite formation, NF-kappa B and AP-1 activation, and IL-8 gene expression in human leukocytes. *Proc Natl Acad Sci U S A* **99**:13266–13271.
- Kamal AM, Hayhoe RP, Paramasivam A, Cooper D, Flower RJ, Solito E, and Perretti M (2006) Antiflammin-2 activates the human formyl-peptide receptor like 1. *ScientificWorldJournal* **6**:1375–1384.
- Kanai F, Liu H, Field SJ, Akbary H, Matsuo T, Brown GE, Cantley LC, and Yaffe MB (2001) The PX domains of p47phox and p40phox bind to lipid products of PI(3)K. *Nat Cell Biol* **3**:675–678.
- Kang HK, Lee HY, Kim MK, Park KS, Park YM, Kwak JY, and Bae YS (2005) The synthetic peptide Trp-Lys-Tyr-Met-Val-D-Met inhibits human monocyte-derived dendritic cell maturation via formyl peptide receptor and formyl peptide receptor-like 2. *J Immunol* **175**:685–692.
- Karlsson J, Fu H, Boulay F, Bylund J, and Dahlgren C (2006) The peptide Trp-Lys-Tyr-Met-Val-D-Met activates neutrophils through the formyl peptide receptor only when signaling through the formylpeptide receptor like 1 is blocked. A receptor switch with implications for signal transduction studies with inhibitors and receptor antagonists. *Biochem Pharmacol* **71**:1488–1496.
- Karlsson J, Fu H, Boulay F, Dahlgren C, Hellstrand K, and Movitz C (2005) Neutrophil NADPH-oxidase activation by an annexin AI peptide is transduced by the formyl peptide receptor (FPR), whereas an inhibitory signal is generated independently of the FPR family receptors. *J Leukoc Biol* **78**:762–771.
- Karnad AB, Hartshorn KL, Wright J, Myers JB, Schwartz JH, and Tauber AI (1989)

- Priming of human neutrophils with N-formyl-methionyl-leucyl-phenylalanine by a calcium-independent, pertussis toxin-insensitive pathway. *Blood* **74**:2519–2526.
- Kettritz R, Falk RJ, Jennette JC, and Gaido ML (1997) Neutrophil superoxide release is required for spontaneous and FMLP-mediated but not for TNF alpha-mediated apoptosis. *J Am Soc Nephrol* **8**:1091–1100.
- Key TA, Foutz TD, Gurevich VV, Sklar LA, and Prossnitz ER (2003) N-formyl peptide receptor phosphorylation domains differentially regulate arrestin and agonist affinity. *J Biol Chem* **278**:4041–4047.
- Key TA, Vines CM, Wagener BM, Gurevich VV, Sklar LA, and Prossnitz ER (2005) Inhibition of chemoattractant N-formyl peptide receptor trafficking by active arrestins. *Traffic* **6**:87–99.
- Khan AI, Heit B, Andonegui G, Colarusso P, and Kubers P (2005) Lipopolysaccharide: a p38 MAPK-dependent disrupter of neutrophil chemotaxis. *Microcirculation* **12**: 421–432.
- Kim C, Marchal CC, Penninger J, and Dinarello MC (2003) The hemopoietic Rho/Rac guanine nucleotide exchange factor Vav1 regulates N-formyl-methionyl-leucyl-phenylalanine-activated neutrophil functions. *J Immunol* **171**:4425–4430.
- Kim JW and Cunningham JM (1993) N-linked glycosylation of the receptor for murine ecotropic retroviruses is altered in virus-infected cells. *J Biol Chem* **268**: 16316–16320.
- Kim Y, Lee BD, Kim O, Bae YS, Lee T, Suh PG, and Ryu SH (2006) Pituitary adenylate cyclase-activating polypeptide 27 is a functional ligand for formyl peptide receptor-like 1. *J Immunol* **176**:2969–2975.
- Klein C, Paul JI, Sauvé K, Schmidt MM, Arcangeli L, Ransom J, Trueheart J, Manfredi JP, Broach JR, and Murphy AJ (1998) Identification of surrogate agonists for the human FPR1-1 receptor by autocrine selection in yeast. *Nat Biotechnol* **16**:1334–1337.
- Knaus UG, Heyworth PG, Evans T, Curnutte JT, and Bokoch GM (1991) Regulation of phagocyte oxygen radical production by the GTP-binding protein Rac 2. *Science* **254**:1512–1515.
- Kobilka BK, Matsui H, Kobilka TS, Yang-Feng TL, Francke U, Caron MG, Lefkowitz RJ, and Regan JW (1987) Cloning, sequencing, and expression of the gene coding for the human platelet alpha 2-adrenergic receptor. *Science* **238**:650–656.
- Kocuzilla R, von Degenfeld G, Kupatt C, Krötz F, Zahler S, Gloe T, Issbrücker K, Unterberger P, Zaiou M, Lebherz C, et al. (2003) An angiogenic role for the human peptide antibiotic LL-37/hCAP-18. *J Clin Invest* **111**:1665–1672.
- Kodama T, Hazeki K, Hazeki O, Okada T, and Ui M (1999) Enhancement of chemotactic peptide-induced activation of phosphoinositide 3-kinase by granulocyte-macrophage colony-stimulating factor and its relation to the cytokine-mediated priming of neutrophil superoxide-anion production. *Biochem J* **337**:201–209.
- Kohout TA, Lin FS, Perry SJ, Conner DA, and Lefkowitz RJ (2001) beta-Arrestin 1 and 2 differentially regulate heptahelical receptor signaling and trafficking. *Proc Natl Acad Sci U S A* **98**:1601–1606.
- König B, Arendt A, McDowell JH, Kahlert M, Hargrave PA, and Hofmann KP (1989) Three cytoplasmic loops of rhodopsin interact with transducin. *Proc Natl Acad Sci U S A* **86**:6878–6882.
- Koo C, Lefkowitz RJ, and Snyderman R (1982) The oligopeptide chemotactic factor receptor on human polymorphonuclear leukocyte membranes exists in two affinity states. *Biochem Biophys Res Commun* **106**:442–449.
- Koo C, Lefkowitz RJ, and Snyderman R (1983) Guanine nucleotides modulate the binding affinity of the oligopeptide chemoattractant receptor on human polymorphonuclear leukocytes. *J Clin Invest* **72**:748–753.
- Korchak HM, Wilkenfeld C, Rich AM, Radin AR, Vienne K, and Rutherford LE (1984) Stimulus response coupling in the human neutrophil. Differential requirements for receptor occupancy in neutrophil responses to a chemoattractant. *J Biol Chem* **259**:7439–7445.
- Kozasa T, Hepler JR, Smrcka AV, Simon MI, Rhee SG, Sternweis PC, and Gilman AG (1993) Purification and characterization of recombinant G16 alpha from Sf9 cells: activation of purified phospholipase C isozymes by G-protein alpha subunits. *Proc Natl Acad Sci U S A* **90**:9176–9180.
- Kubo T, Fukuda K, Mikami A, Maeda A, Takahashi H, Mishina M, Haga K, Ichiyama A, and Kangawa K (1986) Cloning, sequencing and expression of complementary DNA encoding the muscarinic acetylcholine receptor. *Nature* **323**:411–416.
- Kucharzik T, Gewirtz AT, Merlin D, Madara JL, and Williams IR (2003) Lateral membrane LXA4 receptors mediate LXA4's anti-inflammatory actions on intestinal epithelium. *Am J Physiol Cell Physiol* **284**:C888–896.
- Kurosaka K, Chen Q, Yarovsky F, Oppenheim JJ, and Yang D (2005) Mouse cathelin-related antimicrobial peptide chemoattracts leukocytes using formyl peptide receptor-like 1/mouse formyl peptide receptor-like 2 as the receptor and acts as an immune adjuvant. *J Immunol* **174**:6257–6265.
- Kushner I and Rzewnicki D (1999) Acute phase response, in *Inflammation: Basic Principles and Clinical Correlates* (Gallin JI and Snyderman R eds) pp 317–329, Lippincott Williams & Wilkins, Philadelphia.
- Lacy M, Jones J, Whittemore SR, Haviland DL, Wetsel RA, and Barnum SR (1995) Expression of the receptors for the C5a anaphylatoxin, interleukin-8 and FMLP by human astrocytes and microglia. *J Neuroimmunol* **61**:71–78.
- Lad PM, Olson CV, and Smiley PA (1985) Association of the N-formyl-Met-Leu-Phe receptor in human neutrophils with a GTP-binding protein sensitive to pertussis toxin. *Proc Natl Acad Sci U S A* **82**:869–873.
- Lad PM, Welton AF, and Rodbell M (1977) Evidence for distinct guanine nucleotide sites in the regulation of the glucagon receptor and of adenylate cyclase activity. *J Biol Chem* **252**:5942–5946.
- Laporte SA, Miller WE, Kim KM, and Caron MG (2002) beta-Arrestin/AP-2 interaction in G protein-coupled receptor internalization: identification of a beta-arrestin binding site in beta2-adaptin. *J Biol Chem* **277**:9247–9254.
- Lavigne MC, Murphy PM, Leto TL, and Gao JL (2002) The N-formylpeptide receptor (FPR) and a second Gi-coupled receptor mediate fMet-Leu-Phe-stimulated activation of NADPH oxidase in murine neutrophils. *Cell Immunol* **218**:7–12.
- Le Y, Gong W, Li B, Dunlop NM, Shen W, Su SB, Ye RD, and Wang JM (1999) Utilization of two seven-transmembrane, G protein-coupled receptors, formyl peptide receptor-like 1 and formyl peptide receptor, by the synthetic hexapeptide WKYMVMV for human phagocyte activation. *J Immunol* **163**:6777–6784.
- Le Y, Gong W, Tiffany HL, Tumanov A, Nedospasov S, Shen W, Dunlop NM, Gao JL, Murphy PM, Oppenheim JJ, et al. (2001a) Amyloid (beta)42 activates a G-protein-coupled chemoattractant receptor, FPR-like-1. *J Neurosci* **21**:RC123.
- Le Y, Hu J, Gong W, Shen W, Li B, Dunlop NM, Halverson DO, Blair DG, and Wang JM (2000a) Expression of functional formyl peptide receptors by human astrocytoma cell lines. *J Neuroimmunol* **111**:102–108.
- Le Y, Jiang S, Hu J, Gong W, Su S, Dunlop NM, Shen W, Li B, and Ming Wang J (2000b) N36, a synthetic N-terminal heptad repeat domain of the HIV-1 envelope protein gp41, is an activator of human phagocytes. *Clin Immunol* **96**:236–242.
- Le Y, Murphy PM, and Wang JM (2002) Formyl-peptide receptors revisited. *Trends Immunol* **23**:541–548.
- Le Y, Oppenheim JJ, and Wang JM (2001b) Pleiotropic roles of formyl peptide receptors. *Cytokine Growth Factor Rev* **12**:91–105.
- Le Y, Yazawa H, Gong W, Yu Z, Ferrans VJ, Murphy PM, and Wang JM (2001c) The neurotoxic prion peptide fragment PrP(106–126) is a chemotactic agonist for the G protein-coupled receptor formyl peptide receptor-like 1. *J Immunol* **166**:1448–1451.
- Le Y, Ye RD, Gong W, Li J, Iribarren P, and Wang JM (2005) Identification of functional domains in the formyl peptide receptor-like 1 for agonist-induced cell chemotaxis. *FEBS J* **272**:769–778.
- Lee HY, Kim MK, Park KS, Bae YH, Yun J, Park JI, Kwak JY, and Bae YS (2005) Serum amyloid A stimulates matrix-metalloproteinase-9 upregulation via formyl peptide receptor like-1-mediated signaling in human monocytic cells. *Biochem Biophys Res Commun* **330**:989–998.
- Lee HY, Kim MK, Park KS, Shin EH, Jo SH, Kim SD, Jo EJ, Lee YN, Lee C, Baek SH, et al. (2006a) Serum amyloid A induces contrary immune responses via formyl peptide receptor-like 1 in human monocytes. *Mol Pharmacol* **70**:241–248.
- Lee MS, Yoo SA, Cho CS, Suh PG, Kim WU, and Ryu SH (2006b) Serum amyloid A binding to formyl peptide receptor-like 1 induces synovial hyperplasia and angiogenesis. *J Immunol* **177**:5585–5594.
- Lefkowitz RJ and Whalen EJ (2004) beta-arrestins: traffic cops of cell signaling. *Curr Opin Cell Biol* **16**:162–168.
- Lehmeyer JE, Snyderman R, and Johnston RB Jr (1979) Stimulation of neutrophil oxidative metabolism by chemotactic peptides: influence of calcium ion concentration and cytochalasin B and comparison with stimulation by phorbol myristate acetate. *Blood* **54**:35–45.
- Li Z, Jiang H, Xie W, Zhang Z, Smrcka AV, and Wu D (2000) Roles of PLC-beta2 and -beta3 and PI3Kgamma in chemoattractant-mediated signal transduction. *Science* **287**:1046–1049.
- Liang TS, Gao JL, Fatemi O, Lavigne M, Leto TL, and Murphy PM (2001) The endogenous opioid spinorphin blocks fMet-Leu-Phe-induced neutrophil chemotaxis by acting as a specific antagonist at the N-formylpeptide receptor subtype FPR. *J Immunol* **167**:6609–6614.
- Liang TS, Wang JM, Murphy PM, and Gao JL (2000) Serum amyloid A is a chemotactic agonist at FPR2, a low-affinity N-formylpeptide receptor on mouse neutrophils. *Biochem Biophys Res Commun* **270**:331–335.
- Liles WC, Dale DC, and Klebanoff SJ (1995) Glucocorticoids inhibit apoptosis of human neutrophils. *Blood* **86**:3181–3188.
- Linnekin D, Bowles CA, Murano G, and MacVittie TJ (1990) Migration of dog polymorphonuclear neutrophilic leukocytes to formylated peptides. *Inflammation* **14**:691–703.
- Lloyd AR and Oppenheim JJ (1992) Poly's lament: the neglected role of the polymorphonuclear neutrophil in the afferent limb of the immune response. *Immunol Today* **13**:169–172.
- Loor F, Tiberghien F, Wenandy T, Didier A, and Traber R (2002) Cyclosporins: structure-activity relationships for the inhibition of the human FPR1 formylpeptide receptor. *J Med Chem* **45**:4613–4628.
- Machado FS, Esper L, Dias A, Madan R, Gu Y, Hildeman D, Serhan CN, Karp CL, and Aliberti J (2008) Native and aspirin-triggered lipoxins control innate immunity by inducing proteasomal degradation of TRAF6. *J Exp Med* **205**:1077–1086.
- Machado FS, Johnrow JE, Esper L, Dias A, Bafica A, Serhan CN, and Aliberti J (2006) Anti-inflammatory actions of lipoxin A4 and aspirin-triggered lipoxin are SOCS-2 dependent. *Nat Med* **12**:330–334.
- Maddox JF, Hachicha M, Takano T, Petasis NA, Fokin VV, and Serhan CN (1997) Lipoxin A4 stable analogs are potent mimetics that stimulate human monocytes and THP-1 cells via a G-protein-linked lipoxin A4 receptor. *J Biol Chem* **272**:6972–6978.
- Maddox JF and Serhan CN (1996) Lipoxin A4 and B4 are potent stimuli for human monocyte migration and adhesion: selective inactivation by dehydrogenation and reduction. *J Exp Med* **183**:137–146.
- Maderna P, Cottell DC, Berlasconi G, Petasis NA, Brady HR, and Godson C (2002) Lipoxins induce actin reorganization in monocytes and macrophages but not in neutrophils: differential involvement of rho GTPases. *Am J Pathol* **160**:2275–2283.
- Maestes DC, Potter RM, and Prossnitz ER (1999) Differential phosphorylation paradigms dictate desensitization and internalization of the N-formyl peptide receptor. *J Biol Chem* **274**:29791–29795.
- Malech HL, Gardner JP, Heiman DF, and Rosenzweig SA (1985) Asparagine-linked oligosaccharides on formyl peptide chemotactic receptors of human phagocytic cells. *J Biol Chem* **260**:2509–2514.
- Marasco WA, Phan SH, Krutzsch H, Showell HJ, Feltner DE, Nairn R, Becker EL, and Ward PA (1984) Purification and identification of formyl-methionyl-leucyl-phenylalanine as the major peptide neutrophil chemotactic factor produced by *Escherichia coli*. *J Biol Chem* **259**:5430–5439.
- Marchese A, Chen C, Kim YM, and Benovic JL (2003) The ins and outs of G protein-coupled receptor trafficking. *Trends Biochem Sci* **28**:369–376.
- McCoy R, Haviland DL, Molmenti EP, Ziambaras T, Wetsel RA, and Perlmutter DH (1995) N-formylpeptide and complement C5a receptors are expressed in liver cells and mediate hepatic acute phase gene regulation. *J Exp Med* **182**:207–217.



- McDonald PP, Bald A, and Cassatella MA (1997) Activation of the NF-kappaB pathway by inflammatory stimuli in human neutrophils. *Blood* **89**:3421–3433.
- Medzhitov R and Janeway C Jr (2000) Innate immune recognition: mechanisms and pathways. *Immunol Rev* **173**:89–97.
- Mery L and Boulay F (1994) The NH<sub>2</sub>-terminal region of C5aR but not that of FPR is critical for both protein transport and ligand binding. *J Biol Chem* **269**:3457–3463.
- Miao Z, Premack BA, Wei Z, Wang Y, Gerard C, Showell H, Howard M, Schall TJ, and Berahovich R (2007) Proinflammatory proteases liberate a discrete high-affinity functional FPRL1 (CCR12) ligand from CCL23. *J Immunol* **178**:7395–7404.
- Miettinen HM, Gripenrot JM, Mason MM, and Jesaitis AJ (1999) Identification of putative sites of interaction between the human formyl peptide receptor and G protein. *J Biol Chem* **274**:27934–27942.
- Miettinen HM, Mills JS, Gripenrot JM, Dratz EA, Granger BL, and Jesaitis AJ (1997) The ligand binding site of the formyl peptide receptor maps in the transmembrane region. *J Immunol* **159**:4045–4054.
- Migeotte I, Communi D, and Parmentier M (2006) Formyl peptide receptors: a promiscuous subfamily of G protein-coupled receptors controlling immune responses. *Cytokine Growth Factor Rev* **17**:501–519.
- Migeotte I, Riboldi E, Franssen JD, Grégoire F, Loison C, Wittamer V, Detheux M, Robberecht P, Costagliola S, Vassart G, et al. (2005) Identification and characterization of an endogenous chemotactic ligand specific for FPRL2. *J Exp Med* **201**:83–93.
- Milano SK, Pace HC, Kim YM, Brenner C, and Benovic JL (2002) Scaffolding functions of arrestin-2 revealed by crystal structure and mutagenesis. *Biochemistry* **41**:3321–3328.
- Mills JS (2006) Peptides derived from HIV-1, HIV-2, Ebola virus, SARS coronavirus and coronavirus 229E exhibit high affinity binding to the formyl peptide receptor. *Biochim Biophys Acta* **1762**:693–703.
- Mills JS (2007) Differential activation of polymorphisms of the formyl peptide receptor by formyl peptides. *Biochim Biophys Acta* **1772**:1085–1092.
- Mills JS, Miettinen HM, Barnidge D, Vlases MJ, Wimer-Mackin S, Dratz EA, Sunner J, and Jesaitis AJ (1998) Identification of a ligand binding site in the human neutrophil formyl peptide receptor using a site-specific fluorescent photoaffinity label and mass spectrometry. *J Biol Chem* **273**:10428–10435.
- Mills JS, Miettinen HM, Cummings D, and Jesaitis AJ (2000) Characterization of the binding site on the formyl peptide receptor using three receptor mutants and analogs of Met-Leu-Phe and Met-Met-Trp-Leu-Leu. *J Biol Chem* **275**:39012–39017.
- Mills JS, Miettinen HM, Vlases MJ, and Jesaitis AJ (1999) The N-formyl peptide receptor, in *Molecular and Cellular Basis of Inflammation* (Serhan CN and Ward PA eds) pp 215–245. Humana, Totowa, NJ.
- Mócsai A, Jakus Z, Vántus T, Berton G, Lowell CA, and Ligeti E (2000) Kinase pathways in chemoattractant-induced degranulation of neutrophils: the role of p38 mitogen-activated protein kinase activated by Src family kinases. *J Immunol* **164**:4321–4331.
- Mócsai A, Ligeti E, Lowell CA, and Berton G (1999) Adhesion-dependent degranulation of neutrophils requires the Src family kinases Fgr and Hck. *J Immunol* **162**:1120–1126.
- Münch G, Dees C, Hekman M, and Palm D (1991) Multisite contacts involved in coupling of the beta-adrenergic receptor with the stimulatory guanine-nucleotide-binding regulatory protein. Structural and functional studies by beta-receptor-site-specific synthetic peptides. *Eur J Biochem* **198**:357–364.
- Murphy PM (1993) Molecular mimicry and the generation of host defense protein diversity. *Cell* **72**:823–826.
- Murphy PM, Gallin EK, Tiffany HL, and Malech HL (1990) The formyl peptide chemoattractant receptor is encoded by a 2 kilobase messenger RNA. Expression in *Xenopus* oocytes. *FEBS Lett* **261**:353–357.
- Murphy PM and McDermott D (1991) Functional expression of the human formyl peptide receptor in *Xenopus* oocytes requires a complementary human factor. *J Biol Chem* **266**:12560–12567.
- Murphy PM, Özçelik T, Kenney RT, Tiffany HL, McDermott D, and Francke U (1992) A structural homologue of the N-formyl peptide receptor. Characterization and chromosome mapping of a peptide chemoattractant receptor family. *J Biol Chem* **267**:7637–7643.
- Nanamori M, Chen J, Du X, and Ye RD (2007) Regulation of leukocyte degranulation by cGMP-dependent protein kinase and phosphoinositide 3-kinase: potential roles in phosphorylation of target membrane SNARE complex proteins in rat mast cells. *J Immunol* **178**:416–427.
- Nanamori M, Cheng X, Mei J, Sang H, Xuan Y, Zhou C, Wang MW, and Ye RD (2004a) A novel nonpeptide ligand for formyl peptide receptor-like 1. *Mol Pharmacol* **66**:1213–1222.
- Nanamori M, He R, Sang H, and Ye RD (2004b) Normal cell surface expression and selective loss of functions resulting from Phe110 to Ser and Cys126 to Trp substitutions in the formyl peptide receptor. *Immunol Invest* **33**:193–212.
- Nascimento-Silva V, Arruda MA, Barja-Fidalgo C, Villela CG, and Fierro IM (2005) Novel lipid mediator aspirin-triggered lipoxin A4 induces heme oxygenase-1 in endothelial cells. *Am J Physiol Cell Physiol* **289**:C557–563.
- Nauseef WM (2007) How human neutrophils kill and degrade microbes: an integrated view. *Immunol Rev* **219**:88–102.
- Nelson RD, Quie PG, and Simmons RL (1975) Chemotaxis under agarose: a new and simple method for measuring chemotaxis and spontaneous migration of human polymorphonuclear leukocytes and monocytes. *J Immunol* **115**:1650–1656.
- Neptune ER and Bourne HR (1997) Receptors induce chemotaxis by releasing the betagamma subunit of Gi, not by activating Gq or Gs. *Proc Natl Acad Sci U S A* **94**:14489–14494.
- Neptune ER, Iiri T, and Bourne HR (1999) G $\alpha_i$  is not required for chemotaxis mediated by G $\beta\gamma$ -coupled receptors. *J Biol Chem* **274**:2824–2828.
- Niedel J, Davis J, and Cuatrecasas P (1980a) Covalent affinity labeling of the formyl peptide chemotactic receptor. *J Biol Chem* **255**:7063–7066.
- Niedel J, Kahane I, Lachman L, and Cuatrecasas P (1980b) A subpopulation of cultured human promyelocytic leukemia cells (HL-60) displays the formyl peptide chemotactic receptor. *Proc Natl Acad Sci U S A* **77**:1000–1004.
- Niedel J, Wilkinson S, and Cuatrecasas P (1979a) Receptor-mediated uptake and degradation of 125-I-chemotactic peptide by human neutrophils. *J Biol Chem* **254**:10700–10706.
- Niedel JE, Kahane I, and Cuatrecasas P (1979b) Receptor-mediated internalization of fluorescent chemotactic peptide by human neutrophils. *Science* **205**:1412–1414.
- Nigam S, Fiore S, Luscinskas FW, and Serhan CN (1990) Lipoxin A4 and lipoxin B4 stimulate the release but not the oxygenation of arachidonic acid in human neutrophils: dissociation between lipid remodeling and adhesion. *J Cell Physiol* **143**:512–523.
- Niggli V and Keller H (1997) The phosphatidylinositol 3-kinase inhibitor wortmannin markedly reduces chemotactic peptide-induced locomotion and increases in cytoskeletal actin in human neutrophils. *Eur J Pharmacol* **335**:43–52.
- Nomura H, Nielsen BW, and Matsushima K (1993) Molecular cloning of cDNAs encoding a LD78 receptor and putative leukocyte chemotactic peptide receptors. *Int Immunol* **5**:1239–1249.
- Offermanns S and Simon MI (1995) G alpha 15 and G alpha 16 couple a wide variety of receptors to phospholipase C. *J Biol Chem* **270**:15175–15180.
- O'Hara R, Murphy EP, Whitehead AS, FitzGerald O, and Bresnihan B (2004) Local expression of the serum amyloid A and formyl peptide receptor-like 1 genes in synovial tissue is associated with matrix metalloproteinase production in patients with inflammatory arthritis. *Arthritis Rheum* **50**:1788–1799.
- Olofsson S, Khanna B, and Lycke E (1980) Altered kinetic properties of sialyl and galactosyl transferases associated with herpes simplex virus infection of GMK and BHK cells. *J Gen Virol* **47**:1–9.
- O'Meara SJ, Rodgers K, and Godson C (2008) Lipoxins: update and impact of endogenous pro-resolution lipid mediators. *Rev Physiol Biochem Pharmacol* **160**:47–70.
- Onnheim K, Bylund J, Boulay F, Dahlgren C, and Forsman H (2008) Tumour necrosis factor (TNF)-alpha primes murine neutrophils when triggered via formyl peptide receptor-related sequence 2, the murine orthologue of human formyl peptide receptor-like 1, through a process involving the type I TNF receptor and subcellular granule mobilization. *Immunology* **125**:591–600.
- Onorato JJ, Palczewski K, Regan JW, Caron MG, Lefkowitz RJ, and Benovic JL (1991) Role of acidic amino acids in peptide substrates of the beta-adrenergic receptor kinase and rhodopsin kinase. *Biochemistry* **30**:5118–5125.
- Oseas RS, Boxer LA, Butterick C, and Baehner RL (1980) Differences in polymorphonuclear leukocyte aggregating responses among several species in response to chemotactic stimulation. *J Lab Clin Med* **96**:213–221.
- Pamer EG, Wang CR, Flaherty L, Lindahl KF, and Bevan MJ (1992) H-2M3 presents a *Listeria monocytogenes* peptide to cytotoxic T lymphocytes. *Cell* **70**:215–223.
- Partida-Sánchez S, Iribarren P, Moreno-García ME, Gao JL, Murphy PM, Oppenheimer N, Wang JM, and Lund FE (2004) Chemotaxis and calcium responses of phagocytes to formyl peptide receptor ligands is differentially regulated by cyclic ADP ribose. *J Immunol* **172**:1896–1906.
- Paul-Clark MJ, Van Cao T, Moradi-Bidhendi N, Cooper D, and Gilroy DW (2004) 15-epi-lipoxin A4-mediated induction of nitric oxide explains how aspirin inhibits acute inflammation. *J Exp Med* **200**:69–78.
- Perez HD, Holmes R, Kelly E, McClary J, and Andrews WH (1992) Cloning of a cDNA encoding a receptor related to the formyl peptide receptor of human neutrophils. *Gene* **118**:303–304.
- Perez HD, Holmes R, Vilander LR, Adams RR, Manzana W, Jolley D, and Andrews WH (1993) Formyl peptide receptor chimeras define domains involved in ligand binding. *J Biol Chem* **268**:2292–2295.
- Perez HD, Kelly E, Elfman F, Armitage G, and Winkler J (1991) Defective polymorphonuclear leukocyte formyl peptide receptor(s) in juvenile periodontitis. *J Clin Invest* **87**:971–976.
- Perretti M (2003) The annexin 1 receptor(s): is the plot unravelling? *Trends Pharmacol Sci* **24**:574–579.
- Perretti M, Chiang N, La M, Fierro IM, Marullo S, Getting SJ, Solito E, and Serhan CN (2002) Endogenous lipid- and peptide-derived anti-inflammatory pathways generated with glucocorticoid and aspirin treatment activate the lipoxin A4 receptor. *Nat Med* **8**:1296–1302.
- Perretti M, Getting SJ, Solito E, Murphy PM, and Gao JL (2001) Involvement of the receptor for formylated peptides in the *in vivo* anti-migratory actions of annexin 1 and its mimetics. *Am J Pathol* **158**:1969–1973.
- Pitcher JA, Freedman NJ, and Lefkowitz RJ (1998) G protein-coupled receptor kinases. *Annu Rev Biochem* **67**:653–692.
- Posner RG, Fay SP, Domalewski MD, and Sklar LA (1994) Continuous spectrofluorometric analysis of formyl peptide receptor ternary complex interactions. *Mol Pharmacol* **45**:65–73.
- Potter RM, Maestas DC, Cimino DF, and Prossnitz ER (2006) Regulation of N-formyl peptide receptor signaling and trafficking by individual carboxyl-terminal serine and threonine residues. *J Immunol* **176**:5418–5425.
- Prat C, Besteiro J, de Haas CJ, van Strijp JA, and van Kessel KP (2006) A new staphylococcal anti-inflammatory protein that antagonizes the formyl peptide receptor-like 1. *J Immunol* **177**:8017–8026.
- Price MO, McPhail LC, Lambeth JD, Han CH, Knaus UG, and Dinauer MC (2002) Creation of a genetic system for analysis of the phagocyte respiratory burst: high-level reconstitution of the NADPH oxidase in a nonhematopoietic system. *Blood* **99**:2653–2661.
- Prossnitz ER, Kim CM, Benovic JL, and Ye RD (1995a) Phosphorylation of the N-formyl peptide receptor carboxyl terminus by the G protein-coupled receptor kinase, GRK2. *J Biol Chem* **270**:1130–1137.
- Prossnitz ER, Quehenberger O, Cochrane CG, and Ye RD (1991) Transmembrane signalling by the N-formyl peptide receptor in stably transfected fibroblasts. *Biochem Biophys Res Commun* **179**:471–476.
- Prossnitz ER, Quehenberger O, Cochrane CG, and Ye RD (1993) Signal transducing

- properties of the N-formyl peptide receptor expressed in undifferentiated HL60 cells. *J Immunol* **151**:5704–5715.
- Prossnitz ER, Schreiber RE, Bokoch GM, and Ye RD (1995b) Binding of low affinity N-formyl peptide receptors to G protein. Characterization of a novel inactive receptor intermediate. *J Biol Chem* **270**:10686–10694.
- Qiu FH, Devchand PR, Wada K, and Serhan CN (2001) Aspirin-triggered lipoxin A4 and lipoxin A4 up-regulate transcriptional corepressor NAB1 in human neutrophils. *Faseb J* **15**:2736–2738.
- Quehenberger O, Pan ZK, Prossnitz ER, Cavanagh SL, Cochrane CG, and Ye RD (1997) Identification of an N-formyl peptide receptor ligand binding domain by a gain-of-function approach. *Biochem Biophys Res Commun* **238**:377–381.
- Quehenberger O, Prossnitz ER, Cavanagh SL, Cochrane CG, and Ye RD (1993) Multiple domains of the N-formyl peptide receptor are required for high-affinity ligand binding. Construction and analysis of chimeric N-formyl peptide receptors. *J Biol Chem* **268**:18167–18175.
- Rabiet MJ, Huet E, and Boulay F (2005) Human mitochondria-derived N-formylated peptides are novel agonists equally active on FPR and FPRL1, while *Listeria monocytogenes*-derived peptides preferentially activate FPR. *Eur J Immunol* **35**:2486–2495.
- Rabiet MJ, Huet E, and Boulay F (2007) The N-formyl peptide receptors and the anaphylatoxin C5a receptors: an overview. *Biochimie* **89**:1089–1106.
- Redl H, Flynn PJ, Lamche H, Schiesser A, Schlag G, and Hammerschmidt DE (1983) Aggregation, chemotaxis, and chemiluminescence of canine granulocytes. Studies utilizing improved cell preparation techniques. *Inflammation* **7**:67–80.
- Resnati M, Pallavicini I, Wang JM, Oppenheim J, Serhan CN, Romano M, and Blasi F (2002) The fibrinolytic receptor for urokinase activates the G protein-coupled chemotactic receptor FPRL1/LXA4R. *Proc Natl Acad Sci U S A* **99**:1359–1364.
- Richardson RM, Ali H, Tomhave ED, Haribabu B, and Snyderman R (1995) Cross-desensitization of chemoattractant receptors occurs at multiple levels. Evidence for a role for inhibition of phospholipase C activity. *J Biol Chem* **270**:27829–27833.
- Roberts AW, Kim C, Zhen L, Lowe JB, Kapur R, Petryniak B, Spaetti A, Pollock JD, Borneo JB, Bradford GB, et al. (1999) Deficiency of the hematopoietic cell-specific Rho family GTPase Rac2 is characterized by abnormalities in neutrophil function and host defense. *Immunity* **10**:183–196.
- Rodgers K, McMahon B, Mitchell D, Sadlier D, and Godson C (2005) Lipoxin A4 modifies platelet-derived growth factor-induced pro-fibrotic gene expression in human renal mesangial cells. *Am J Pathol* **167**:683–694.
- Romano M, Maddox JF, and Serhan CN (1996) Activation of human monocytes and the acute monocytic leukemia cell line (THP-1) by lipoxins involves unique signaling pathways for lipoxin A4 versus lipoxin B4: evidence for differential Ca<sup>2+</sup> mobilization. *J Immunol* **157**:2149–2154.
- Rot A, Henderson LE, Copeland TD, and Leonard EJ (1987) A series of six ligands for the human formyl peptide receptor: tetrapeptides with high chemotactic potency and efficacy. *Proc Natl Acad Sci U S A* **84**:7967–7971.
- Rot A, Henderson LE, Sowder R, and Leonard EJ (1989) Staphylococcus aureus tetrapeptide with high chemotactic potency and efficacy for human leukocytes. *J Leukoc Biol* **45**:114–120.
- Rotrosen D, Malech HL, and Gallin JI (1987) Formyl peptide leukocyte chemoattractant uptake and release by cultured human umbilical vein endothelial cells. *J Immunol* **139**:3034–3040.
- Rovati GE, Capra V, and Neubig RR (2007) The highly conserved DRY motif of class A G protein-coupled receptors: beyond the ground state. *Mol Pharmacol* **71**:959–964.
- Sahagun-Ruiz A, Colla JS, Juhn J, Gao JL, Murphy PM, and McDermott DH (2001) Contrasting evolution of the human leukocyte N-formylpeptide receptor subtypes FPR and FPRL1R. *Genes Immun* **2**:335–342.
- Sandri S, Rodriguez D, Gomes E, Monteiro HP, Russo M, and Campa A (2008) Is serum amyloid A an endogenous TLR4 agonist? *J Leukoc Biol* **83**:1174–1180.
- Sasaki T, Irie-Sasaki J, Jones RG, Oliveira-dos-Santos AJ, Stanford WL, Bolon B, Wakeham A, Itie A, Bouchard D, Koziarzdzki I, et al. (2000) Function of P13Kgamma in thymocyte development, T cell activation, and neutrophil migration. *Science* **287**:1040–1046.
- Savarese TM and Fraser CM (1992) In vitro mutagenesis and the search for structure-function relationships among G protein-coupled receptors. *Biochem J* **283**:1–19.
- Savill J (1997) Apoptosis in resolution of inflammation. *J Leukoc Biol* **61**:375–380.
- Schaldach CM, Riby J, and Bjeldanes LF (1999) Lipoxin A4: a new class of ligand for the Ah receptor. *Biochemistry* **38**:7594–7600.
- Schepetkin IA, Kirpotina LN, Khlebnikov AI, and Quinn MT (2007) High-throughput screening for small-molecule activators of neutrophils: identification of novel N-formyl peptide receptor agonists. *Mol Pharmacol* **71**:1061–1074.
- Schepetkin IA, Kirpotina LN, Tian J, Khlebnikov AI, Ye RD, and Quinn MT (2008) Identification of novel formyl peptide receptor-like 1 agonists that induce macrophage tumor necrosis factor- $\alpha$  production. *Mol Pharmacol* **74**:392–402.
- Schiffmann E, Corcoran BA, and Wahl SM (1975a) N-formylmethionyl peptides as chemoattractants for leukocytes. *Proc Natl Acad Sci U S A* **72**:1059–1062.
- Schiffmann E, Showell HV, Corcoran BA, Ward PA, Smith E, and Becker EL (1975b) The isolation and partial characterization of neutrophil chemotactic factors from *Escherichia coli*. *J Immunol* **114**:1831–1837.
- Schreiber RE, Prossnitz ER, Ye RD, Cochrane CG, and Bokoch GM (1994) Domains of the human neutrophil N-formyl peptide receptor involved in G protein coupling. Mapping with receptor-derived peptides. *J Biol Chem* **269**:326–331.
- Schreiber RE, Prossnitz ER, Ye RD, Cochrane CG, Jesaitis AJ, and Bokoch GM (1993) Reconstitution of recombinant N-formyl chemotactic peptide receptor with G protein. *J Leukocyte Biol* **53**:470–474.
- Schultz P, Stannek P, Voigt M, Jakobs KH, and Gierschik P (1992) Complementation of formyl peptide receptor-mediated signal transduction in *Xenopus laevis* oocytes. *Biochem J* **284**:207–212.
- Seifert R and Wenzel-Seifert K (2001) Defective Gi protein coupling in two formyl peptide receptor mutants associated with localized juvenile periodontitis. *J Biol Chem* **276**:42043–42049.
- Selleri C, Montuori N, Ricci P, Visconte V, Carriero MV, Sidenius N, Serio B, Blasi F, Rotoli B, Rossi G, et al. (2005) Involvement of the urokinase-type plasminogen activator receptor in hematopoietic stem cell mobilization. *Blood* **105**:2198–2205.
- Sengeløv H, Boulay F, Kjeldsen L, and Borregaard N (1994) Subcellular localization and translocation of the receptor for N-formylmethionyl-leucyl-phenylalanine in human neutrophils. *Biochem J* **299**:473–479.
- Sengeløv H, Kjeldsen L, and Borregaard N (1993) Control of exocytosis in early neutrophil activation. *J Immunol* **150**:1535–1543.
- Serhan CN (2005) Lipoxins and aspirin-triggered 15-epi-lipoxins are the first lipid mediators of endogenous anti-inflammation and resolution. *Prostaglandins Leukot Essent Fatty Acids* **73**:141–162.
- Serhan CN (2007) Resolution phase of inflammation: novel endogenous anti-inflammatory and proresolving lipid mediators and pathways. *Annu Rev Immunol* **25**:101–137.
- Serhan CN, Fiore S, Brezinski DA, and Lynch S (1993) Lipoxin A4 metabolism by differentiated HL-60 cells and human monocytes: conversion to novel 15-oxo and dihydro products. *Biochemistry* **32**:6313–6319.
- Serhan CN, Maddox JF, Petasis NA, Akritopoulou-Zanze I, Papiyianni A, Brady HR, Colgan SP, and Madara JL (1995) Design of lipoxin A4 stable analogs that block transmigration and adhesion of human neutrophils. *Biochemistry* **34**:14609–14615.
- Servant G, Weiner OD, Neptune ER, Sedat JW, and Bourne HR (1999) Dynamics of a chemoattractant receptor in living neutrophils during chemotaxis. *Mol Biol Cell* **10**:1163–1178.
- Sha'afi RI, and Molski TF (1988) Activation of the neutrophil. *Prog Allergy* **42**:1–64.
- Shen W, Proost P, Li B, Gong W, Le Y, Sargeant R, Murphy PM, Van Damme J, and Wang JM (2000) Activation of the chemotactic peptide receptor FPRL1 in monocytes phosphorylates the chemokine receptor CCR5 and attenuates cell responses to selected chemokines. *Biochem Biophys Res Commun* **272**:276–283.
- Shenoy SK and Lefkowitz RJ (2003) Multifaceted roles of beta-arrestins in the regulation of seven-membrane-spanning receptor trafficking and signalling. *Biochem J* **375**:503–515.
- Shi G, Partida-Sánchez S, Misra RS, Tighe M, Borchers MT, Lee JJ, Simon MI, and Lund FE (2007) Identification of an alternative G $\alpha_q$ -dependent chemokine receptor signal transduction pathway in dendritic cells and granulocytes. *J Exp Med* **204**:2705–2718.
- Showell HJ, Freer RJ, Zigmund SH, Schiffmann E, Aswanikumar S, Corcoran B, and Becker EL (1976) The structure-activity relations of synthetic peptides as chemotactic factors and inducers of lysosomal secretion for neutrophils. *J Exp Med* **143**:1154–1169.
- Simon MI, Strathmann MP, and Gautam N (1991) Diversity of G proteins in signal transduction. *Science* **252**:802–808.
- Sklar LA, Bokoch GM, Button D, and Smolen JE (1987) Regulation of ligand-receptor dynamics by guanine nucleotides. Real-time analysis of interconverting states for the neutrophil formyl peptide receptor. *J Biol Chem* **262**:135–139.
- Sklar LA, Fay SP, Seligmann BE, Freer RJ, Muthukumaraswamy N, and Mueller H (1990) Fluorescence analysis of the size of a binding pocket of a peptide receptor at natural abundance. *Biochemistry* **29**:313–316.
- Sklar LA and Finney DA (1982) Analysis of ligand-receptor interactions with the fluorescence activated cell sorter. *Cytometry* **3**:161–165.
- Sklar LA, Finney DA, Oades ZG, Jesaitis AJ, Painter RG, and Cochrane CG (1984) The dynamics of ligand-receptor interactions. Real-time analyses of association, dissociation, and internalization of an N-formyl peptide and its receptors on the human neutrophil. *J Biol Chem* **259**:5661–5669.
- Sklar LA, Oades ZG, Jesaitis AJ, Painter RG, and Cochrane CG (1981) Fluorescein-labeled chemotactic peptide and high-affinity anti-fluorescein antibody as a probe of the temporal characteristics of neutrophil stimulation. *Proc Natl Acad Sci U S A* **78**:7540–7544.
- Smith RJ, Sam LM, and Justen JM (1988) Diacylglycerols modulate human polymorphonuclear neutrophil responsiveness: effects on intracellular calcium mobilization, granule exocytosis, and superoxide anion production. *J Leukoc Biol* **43**:411–419.
- Snyderman R and Pike MC (1980) N-Formylmethionyl peptide receptors on equine leukocytes initiate secretion but not chemotaxis. *Science* **209**:493–495.
- Snyderman R and Uhing RJ (1992) Phagocytic cells: stimulus-response coupling mechanisms, in *Inflammation: Basic Principles and Clinical Correlates* (Gallin JI, Goldstein IM, and Snyderman R eds) pp 421–439, Raven Press, New York.
- Sodin-Semrl S, Spagnolo A, Mikus R, Barbaro B, Varga J, and Fiore S (2004) Opposing regulation of interleukin-8 and NF- $\kappa$ B responses by lipoxin A4 and serum amyloid A via the common lipoxin A receptor. *Int J Immunopathol Pharmacol* **17**:145–156.
- Sodin-Semrl S, Taddeo B, Tseng D, Varga J, and Fiore S (2000) Lipoxin A4 inhibits IL-1 beta-induced IL-6, IL-8, and matrix metalloproteinase-3 production in human synovial fibroblasts and enhances synthesis of tissue inhibitors of metalloproteinases. *J Immunol* **164**:2660–2666.
- Solito E, Kamal A, Russo-Marie F, Buckingham JC, Marullo S, and Perretti M (2003) A novel calcium-dependent proapoptotic effect of annexin 1 on human neutrophils. *Faseb J* **17**:1544–1546.
- Solito E, Romero IA, Marullo S, Russo-Marie F, and Wexler BB (2000) Annexin 1 binds to U937 monocytic cells and inhibits their adhesion to microvascular endothelium: involvement of the alpha 4 beta 1 integrin. *J Immunol* **165**:1573–1581.
- Southgate EL, He RL, Gao JL, Murphy PM, Nanamori M, and Ye RD (2008) Identification of formyl peptides from *Listeria monocytogenes* and *Staphylococcus aureus* as potent chemoattractants for mouse neutrophils. *J Immunol* **181**:1429–1437.
- Souza DG, Fagundes CT, Amaral FA, Cisalpino D, Sousa LP, Vieira AT, Pinho V, Nicoli JR, Vieira LQ, Fierro IM, et al. (2007) The required role of endogenously produced lipoxin A4 and annexin-1 for the production of IL-10 and inflammatory hyporesponsiveness in mice. *J Immunol* **179**:8533–8543.
- Stenfeldt AL, Karlsson J, Wennerås C, Bylund J, Fu H, and Dahlgren C (2007) Cyclosporin H, Boc-MLF and Boc-FLFLF are antagonists that preferentially inhibit activity triggered through the formyl peptide receptor. *Inflammation* **30**:224–229.

- Su SB, Gao J, Gong W, Dunlop NM, Murphy PM, Oppenheim JJ, and Wang JM (1999a) T21/DP107, A synthetic leucine zipper-like domain of the HIV-1 envelope gp41, attracts and activates human phagocytes by using G-protein-coupled formyl peptide receptors. *J Immunol* **162**:5924–5930.
- Su SB, Gong W, Gao JL, Shen W, Murphy PM, Oppenheim JJ, and Wang JM (1999b) A seven-transmembrane, G protein-coupled receptor, FPRL1, mediates the chemotactic activity of serum amyloid A for human phagocytic cells. *J Exp Med* **189**:395–402.
- Su SB, Gong WH, Gao JL, Shen WP, Grimm MC, Deng X, Murphy PM, Oppenheim JJ, and Wang JM (1999c) T20/DP178, an ectodomain peptide of human immunodeficiency virus type 1 gp41, is an activator of human phagocyte N-formyl peptide receptor. *Blood* **93**:3885–3892.
- Sun R, Iribarren P, Zhang N, Zhou Y, Gong W, Cho EH, Lockett S, Chertov O, Bednar F, Rogers TJ, et al. (2004) Identification of neutrophil granule protein cathepsin G as a novel chemotactic agonist for the G protein-coupled formyl peptide receptor. *J Immunol* **173**:428–436.
- Svensson CI, Zattoni M, and Serhan CN (2007) Lipoxins and aspirin-triggered lipoxin inhibit inflammatory pain processing. *J Exp Med* **204**:245–252.
- Takano T, Clish CB, Gronert K, Petasis N, and Serhan CN (1998) Neutrophil-mediated changes in vascular permeability are inhibited by topical application of aspirin-triggered 15-epi-lipoxin A4 and novel lipoxin B4 stable analogues. *J Clin Invest* **101**:819–826.
- Takano T, Fiore S, Maddox JF, Brady HR, Petasis NA, and Serhan CN (1997) Aspirin-triggered 15-epi-lipoxin A4 (LXA4) and LXA4 stable analogues are potent inhibitors of acute inflammation: evidence for anti-inflammatory receptors. *J Exp Med* **185**:1693–1704.
- Tardif M, Mery L, Brouchon L, and Boulay F (1993) Agonist-dependent phosphorylation of N-Formylpeptide and activation peptide from the fifth component of C (C5a) chemoattractant receptors in differentiated HL60 cells. *J Immunol* **150**:3534–3545.
- Tardif M, Rabiet MJ, Christophe T, Milcent MD, and Boulay F (1998) Isolation and characterization of a variant HL60 cell line defective in the activation of the NADPH oxidase by phorbol myristate acetate. *J Immunol* **161**:6885–6895.
- Thelen M, Peveri P, Kernen P, von Tscharnner V, Walz A, and Baggiolini M (1988) Mechanism of neutrophil activation by NAF, a novel monocyte-derived peptide agonist. *Faseb J* **2**:2702–2706.
- Thelen M, Wymann MP, and Langen H (1994) Wortmannin binds specifically to 1-phosphatidylinositol 3-kinase while inhibiting guanine nucleotide-binding protein-coupled receptor signaling in neutrophil leukocytes. *Proc Natl Acad Sci U S A* **91**:4960–4964.
- Thomas KM, Pyun HY, and Navarro J (1990) Molecular cloning of the fMet-Leu-Phe receptor from neutrophils. *J Biol Chem* **265**:20061–20064.
- Thomas KM, Taylor L, and Navarro J (1991) The interleukin-8 receptor is encoded by a neutrophil-specific cDNA clone, F3R. *J Biol Chem* **266**:14839–14841.
- Tiffany HL, Lavigne MC, Cui YH, Wang JM, Leto TL, Gao JL, and Murphy PM (2001) Amyloid-beta induces chemotaxis and oxidant stress by acting at formylpeptide receptor 2, a G protein-coupled receptor expressed in phagocytes and brain. *J Biol Chem* **276**:23645–23652.
- Toniolo C, Crisma M, Moretto V, Freer RJ, and Becker EL (1990) N alpha-formylated and tert-butylloxycarbonylated Phe-(Leu-Phe)<sub>n</sub> and (Leu-Phe)<sub>n</sub> peptides as agonists and antagonists of the chemotactic formylpeptide receptor of the rabbit peritoneal neutrophil. *Biochim Biophys Acta* **1034**:67–72.
- Tuluc F, Meshki J, and Kunapuli SP (2003) Membrane lipid microdomains differentially regulate intracellular signaling events in human neutrophils. *Int Immunopharmacol* **3**:1775–1790.
- Van Dyke TE, Zinney W, Winkel K, Taufiq A, Offenbacher S, and Arnold RR (1986) Neutrophil function in localized juvenile periodontitis. Phagocytosis, superoxide production and specific granule release. *J Periodontol* **57**:703–708.
- Vaughn MW, Proske RJ, and Haviland DL (2002) Identification, cloning, and functional characterization of a murine lipoxin A4 receptor homologue gene. *J Immunol* **169**:3363–3369.
- Verghese MW, Charles L, Jakoi L, Dillon SB, and Snyderman R (1987) Role of a guanine nucleotide regulatory protein in the activation of phospholipase C by different chemoattractants. *J Immunol* **138**:4374–4380.
- Villanueva MS, Beckers CJ, and Pamer EG (1994) Infection with *Listeria monocytogenes* impairs sialic acid addition to host cell glycoproteins. *J Exp Med* **180**:2137–2145.
- Vines CM, Revankar CM, Maestas DC, LaRusch LL, Cimino DF, Kohout TA, Lefkowitz RJ, and Prossnitz ER (2003) N-formyl peptide receptors internalize but do not recycle in the absence of arrestins. *J Biol Chem* **278**:41581–41584.
- Vitkuskas G, Showell HJ, and Becker EL (1980) Specific binding of synthetic chemotactic peptides to rabbit peritoneal neutrophils: effects on dissociability of bound peptide, receptor activity and subsequent biologic responsiveness (deactivation). *Mol Immunol* **17**:171–180.
- von der Weid PY, Hollenberg MD, Fiorucci S, and Wallace JL (2004) Aspirin-triggered, cyclooxygenase-2-dependent lipoxin synthesis modulates vascular tone. *Circulation* **110**:1320–1325.
- von Wartburg A and Traber R (1986) Chemistry of the natural cyclosporin metabolites. *Prog Allergy* **38**:28–45.
- Walther A, Riehemann K, and Gerke V (2000) A novel ligand of the formyl peptide receptor: annexin I regulates neutrophil extravasation by interacting with the FPR. *Mol Cell* **5**:831–840.
- Wang F, Herzmark P, Weiner OD, Srinivasan S, Servant G, and Bourne HR (2002) Lipid products of PI(3)Ks maintain persistent cell polarity and directed motility in neutrophils. *Nat Cell Biol* **4**:513–518.
- Wang ZG and Ye RD (2002) Characterization of two new members of the formyl peptide receptor gene family from 129S6 mice. *Gene* **299**:57–63.
- Ward PA, Lepow IH, and Newman LJ (1968) Bacterial factors chemotactic for polymorphonuclear leukocytes. *Am J Pathol* **52**:725–736.
- Ward PA, Sulavik MC, and Johnson KJ (1984) Rat neutrophil activation and effects of lipoxigenase and cyclooxygenase inhibitors. *Am J Pathol* **116**:223–233.
- Weingarten R, Ransnas L, Mueller H, Sklar LA, and Bokoch GM (1990) Mastoparan interacts with the carboxyl terminus of the alpha subunit of Gi. *J Biol Chem* **265**:11044–11049.
- Welch HC, Coadwell WJ, Ellson CD, Ferguson GJ, Andrews SR, Erdjument-Bromage H, Tempst P, Hawkins PT, and Stephens LR (2002) P-Rex1, a PtdIns(3,4,5)P<sub>3</sub>- and Gbetagamma-regulated guanine-nucleotide exchange factor for Rac. *Cell* **108**:809–821.
- Welch HC, Condliffe AM, Milne LJ, Ferguson GJ, Hill K, Webb LM, Okkenhaug K, Coadwell WJ, Andrews SR, Thelen M, et al. (2005) P-Rex1 regulates neutrophil function. *Curr Biol* **15**:1867–1873.
- Wenzel-Seifert K, Arthur JM, Liu HY, and Seifert R (1999) Quantitative analysis of formyl peptide receptor coupling to G<sub>α1</sub>, G<sub>α2</sub>, and G<sub>α3</sub>. *J Biol Chem* **274**:33259–33266.
- Wenzel-Seifert K, Grünbaum L, and Seifert R (1991) Differential inhibition of human neutrophil activation by cyclosporins A, D, and H. Cyclosporin H is a potent and effective inhibitor of formyl peptide-induced superoxide formation. *J Immunol* **147**:1940–1946.
- Wenzel-Seifert K, Hurt CM, and Seifert R (1998) High constitutive activity of the human formyl peptide receptor. *J Biol Chem* **273**:24181–24189.
- Wenzel-Seifert K and Seifert R (1993) Cyclosporin H is a potent and selective formyl peptide receptor antagonist. Comparison with N-t-butoxycarbonyl-L-phenylalanyl-L-leucyl-L-phenylalanyl-L-leucyl-L-phenylalanine and cyclosporins A, B, C, D, and E. *J Immunol* **150**:4591–4599.
- Wilkinson PC (1979) Synthetic peptide chemotactic factors for neutrophils: the range of active peptides, their efficacy and inhibitory activity, and susceptibility of the cellular response to enzymes and bacterial toxins. *Immunology* **36**:579–588.
- Williams LT, Snyderman R, Pike MC, and Lefkowitz RJ (1977) Specific receptor sites for chemotactic peptides on human polymorphonuclear leukocytes. *Proc Natl Acad Sci U S A* **74**:1204–1208.
- Wu D, Jiang H, and Simon MI (1995) Different alpha 1-adrenergic receptor sequences required for activating different G alpha subunits of Gq class of G proteins. *J Biol Chem* **270**:9828–9832.
- Wu SH, Liao PY, Dong L, and Chen ZQ (2008) Signal pathway involved in inhibition by lipoxin A(4) of production of interleukins induced in endothelial cells by lipopolysaccharide. *Inflamm Res* **57**:430–437.
- Wu SH, Lu C, Dong L, Zhou GP, He ZG, and Chen ZQ (2005) Lipoxin A4 inhibits TNF-alpha-induced production of interleukins and proliferation of rat mesangial cells. *Kidney Int* **68**:35–46.
- Xu J, Wang F, Van Keymeulen A, Herzmark P, Straight A, Kelly K, Takuwa Y, Sugimoto N, Mitchison T, and Bourne HR (2003) Divergent signals and cytoskeletal assemblies regulate self-organizing polarity in neutrophils. *Cell* **114**:201–214.
- Xue M, Hsieh G, Raymond-Stintz MA, Pfeiffer J, Roberts D, Steinberg SL, Oliver JM, Prossnitz ER, Lidke DS, and Wilson BS (2007) Activated N-formyl peptide receptor and high-affinity IgE receptor occupy common domains for signaling and internalization. *Mol Biol Cell* **18**:1410–1420.
- Yamamoto Y, Kanazawa T, Shimamura M, Ueki M, and Hazato T (1997) Inhibitory effects of spinorphin, a novel endogenous regulator, on chemotaxis, O<sub>2</sub>-generation, and exocytosis by N-formylmethionyl-leucyl-phenylalanine (FMLP)-stimulated neutrophils. *Biochem Pharmacol* **54**:695–701.
- Yan P, Nanamori M, Sun M, Zhou C, Cheng N, Li N, Zheng W, Xiao L, Xie X, Ye RD, et al. (2006) The immunosuppressant cyclosporin A antagonizes human formyl peptide receptor through inhibition of cognate ligand binding. *J Immunol* **177**:7050–7058.
- Yang D, Chen Q, Gertz B, He R, Phulsuksombati M, Ye RD, and Oppenheim JJ (2002) Human dendritic cells express functional formyl peptide receptor-like-2 (FPRL2) throughout maturation. *J Leukoc Biol* **72**:598–607.
- Yang D, Chen Q, Le Y, Wang JM, and Oppenheim JJ (2001a) Differential regulation of formyl peptide receptor-like 1 expression during the differentiation of monocytes to dendritic cells and macrophages. *J Immunol* **166**:4092–4098.
- Yang D, Chen Q, Schmidt AP, Anderson GM, Wang JM, Wooters J, Oppenheim JJ, and Chertov O (2000) LL-37, the neutrophil granule- and epithelial cell-derived cathelicidin, utilizes formyl peptide receptor-like 1 (FPRL1) as a receptor to chemoattract human peripheral blood neutrophils, monocytes, and T cells. *J Exp Med* **192**:1069–1074.
- Yang M, Sang H, Rahman A, Wu D, Malik AB, and Ye RD (2001b) G alpha 16 couples chemoattractant receptors to NF-kappa B activation. *J Immunol* **166**:6885–6892.
- Yazawa H, Yu ZX, Takeda, Le Y, Gong W, Ferrans VJ, Oppenheim JJ, Li CC, and Wang JM (2001) Beta amyloid peptide (Aβ42) is internalized via the G-protein-coupled receptor FPRL1 and forms fibrillar aggregates in macrophages. *Faseb J* **15**:2454–2462.
- Ye RD (2001) Regulation of nuclear factor kappaB activation by G-protein-coupled receptors. *J Leukoc Biol* **70**:839–848.
- Ye RD, Cavanagh SL, Quehenberger O, Prossnitz ER, and Cochrane CG (1992) Isolation of a cDNA that encodes a novel granulocyte N-formyl peptide receptor. *Biochem Biophys Res Commun* **184**:582–589.
- Ye RD, Quehenberger O, Thomas KM, Navarro J, Cavanagh SL, Prossnitz ER, and Cochrane CG (1993) The rabbit neutrophil N-formyl peptide receptor. cDNA cloning, expression, and structure/function implications. *J Immunol* **150**:1383–1394.
- Ying G, Iribarren P, Zhou Y, Gong W, Zhang N, Yu ZX, Le Y, Cui Y, and Wang JM (2004) Humanin, a newly identified neuroprotective factor, uses the G protein-coupled formylpeptide receptor-like-1 as a functional receptor. *J Immunol* **172**:7078–7085.
- Zhang Y, Syed R, Uygur C, Pallos D, Gorry MC, Firatli E, Cortelli JR, VanDyke TE, Hart PS, Feingold E, et al. (2003) Evaluation of human leukocyte N-formylpeptide receptor (FPR1) SNPs in aggressive periodontitis patients. *Genes Immun* **4**:22–29.
- Zhou C, Zhang S, Nanamori M, Zhang Y, Liu Q, Li N, Sun M, Tian J, Ye PP, Cheng N, et al. (2007) Pharmacological characterization of a novel nonpeptide antagonist for formyl peptide receptor-like 1. *Mol Pharmacol* **72**:976–983.
- Zigmond SH (1977) Ability of polymorphonuclear leukocytes to orient in gradients of chemotactic factors. *J Cell Biol* **75**:606–616.