ASSOCIATE EDITOR: ALAN V. SMRCKA

Genetic Analysis of Rare Human Variants of Regulators of G Protein Signaling Proteins and Their Role in Human Physiology and Disease^[S]

Katherine E. Squires, Carolina Montañez-Miranda, Rushika R. Pandya, Matthew P. Torres, and John R. Hepler

Department of Pharmacology, Emory University School of Medicine, Atlanta, Georgia (K.E.S., C.M.-M., J.R.H.); and School of Biological Sciences, Georgia Institute of Technology, Atlanta, Georgia (R.R.P., M.P.T.)

Abstract.....	446
I. Introduction.....	447
A. G Protein Signaling.....	447
B. A (Very) Brief History of Regulators of G Protein Signaling.....	447
C. RGS Protein Rare Human Variants in Complex Diseases.....	448
II. RGS Proteins in Physiology and Human Disease.....	449
A. The R4 Family.....	449
1. R4 Family Overview.....	449
2. R4 Family Proteins in Human Physiology and Disease.....	450
B. The R7 Family.....	454
1. R7 Family Overview.....	454
2. R7 Family Proteins in Human Physiology and Disease.....	454
C. The R12 Family.....	456
1. R12 Family Overview.....	456
2. R12 Family Proteins in Human Physiology and Disease.....	456
D. The RZ Family.....	457
1. RZ Family Overview.....	457
2. RZ Family Proteins in Human Physiology and Disease.....	457
III. Analysis of Rare Human Variants of RGS Proteins.....	458
A. R4 Family: RGS4 Rare Variants.....	461
B. R7 Family: RGS9 Rare Variants.....	462
C. R12 Family: RGS10 Rare Variants.....	464
D. RZ Family: RGS17 Rare Variants.....	465
IV. Pharmacological Impact of Human Variants in RGS Proteins.....	466
Acknowledgments.....	467
References.....	467

Abstract—Regulators of G protein signaling (RGS) proteins modulate the physiologic actions of many neurotransmitters, hormones, and other signaling molecules. Human RGS proteins comprise a family of 20 canonical proteins that bind directly to G protein-coupled receptors/G protein complexes to limit the lifetime of their signaling events, which regulate all

aspects of cell and organ physiology. Genetic variations account for diverse human traits and individual predispositions to disease. RGS proteins contribute to many complex polygenic human traits and pathologies such as hypertension, atherosclerosis, schizophrenia, depression, addiction, cancers, and many others. Recent analysis indicates that most human diseases are

Address correspondence to: Dr. John R. Hepler, Emory University School of Medicine, G206 Rollins Research Center, 1510 Clifton Road G205, Atlanta, GA 30329-3090. E-mail: jhepler@emory.edu

This work was supported, in whole or in part, by National Institutes of Health [Grants 5R01NS037112 and 1R21 NS087488, both awarded to J.R.H., and Grant R01-GM117400 to M.P.T.]. R.R.P. was supported by the GT Bioinformatics Faculty Research Award, and K.E.S. was supported by National Institutes of Health [Training Grant T32 GM008602].

<https://doi.org/10.1124/pr.117.015354>.

^[S]This article has supplemental material available at pharmrev.aspetjournals.org.

due to extremely rare genetic variants. In this study, we summarize physiologic roles for RGS proteins and links to human diseases/traits and report rare variants found within each human RGS protein exome sequence derived from global population studies. Each RGS sequence is analyzed using recently described bioinformatics and proteomic tools for measures of missense tolerance ratio paired with combined annotation-dependent depletion scores, and protein post-translational modification (PTM) alignment cluster analysis. We highlight selected variants

within the well-studied RGS domain that likely disrupt RGS protein functions and provide comprehensive variant and PTM data for each RGS protein for future study. We propose that rare variants in functionally sensitive regions of RGS proteins confer profound change-of-function phenotypes that may contribute, in newly appreciated ways, to complex human diseases and/or traits. This information provides investigators with a valuable database to explore variation in RGS protein function, and for targeting RGS proteins as future therapeutic targets.

I. Introduction

A. *G Protein Signaling*

Cells communicate by releasing chemical messengers that dictate cell and organ physiology. These messengers include many natural ligands such as hormones, neurotransmitters, cytokines, nutrients, sensory input, and other natural molecules. Because most of these messengers cannot cross the outer cell (plasma) membrane, membrane-bound receptors and their cognate heterotrimeric G proteins ($G\alpha\beta\gamma$) act as cellular transducers. At rest, G protein-coupled receptors (GPCRs) are in close proximity with an inactive $G\alpha\beta\gamma$ protein complex, with GDP bound to $G\alpha$. Upon receptor activation by ligand binding, the chemical message is transduced across the plasma membrane through the receptor to promote the release of GDP from the associated $G\alpha$. Due to the high concentration of GTP in the cytosol, $G\alpha$ quickly binds a GTP to form $G\alpha$ -GTP, which has reduced affinity for both receptor and $G\beta\gamma$, and thus dissociates. The free $G\alpha$ -GTP and $G\beta\gamma$ are then able to relate the chemical message to an intracellular cascade of effectors and second messengers that dictate all aspects of cell and organ physiology. The duration of signaling by $G\alpha$ -GTP and $G\beta\gamma$ is governed by the intrinsic GTPase activity of the $G\alpha$ subunit (Gilman, 1987; Bourne et al., 1990; Simon et al., 1991; Hepler and Gilman, 1992; Hamm, 1998). Upon hydrolysis of the GTP to GDP, $G\alpha$ -GDP reassociates with $G\beta\gamma$ to terminate signaling. Under normal physiologic conditions, a $G\alpha$ acts as a molecular switch capable of turning itself off by GTP hydrolysis. However, researchers noted early on that the rate of GTP hydrolysis by purified $G\alpha$ proteins in vitro was much slower than that observed by $G\alpha$ in cells, hinting at a missing piece of the puzzle: cellular factor(s) that governs the off rate of G protein signaling (Wagner et al., 1988; Vuong and Chabre, 1990).

B. A (Very) Brief History of Regulators of G Protein Signaling

The missing piece was identified based on foundational observations dating back to the early 1980s. SST2, a yeast protein important for mating, was found

to regulate sensitivity to pheromones (Chan and Otte, 1982). Later it was shown that SST2 does so via regulation of Dohlman et al. (1995) and physical association with (Dohlman et al., 1996) yeast $G\alpha$ proteins, which share homology to mammalian systems. These findings followed the discovery of a class of mammalian proteins that could enhance the GTPase activity of Ras and Ras-like small G proteins (Trahey and McCormick, 1987), speeding up the turnover of GTP and accordingly the termination of signaling [termed GTPase-accelerating proteins (GAPs)]. The postulated missing piece, a mammalian GAP for heterotrimeric $G\alpha$ proteins, was discovered soon thereafter as a family of novel proteins very similar to the yeast SST2 (De Vries et al., 1995; Druey et al., 1996; Koelle and Horvitz, 1996). These specialized $G\alpha$ GAPs were named regulators of G protein signaling (RGS), and thus the RGS field was born.

Since their initial discovery, 20 canonical RGS (Fig. 1) and 19 RGS-like proteins have been identified, and extensive characterization of these proteins has revealed multifunctional roles (Ross and Wilkie, 2000; Hollinger and Hepler, 2002; Willars, 2006; Evans et al., 2015). All canonical RGS proteins share a conserved, approximately 120 amino acid RGS domain, which binds active $G\alpha$ -GTP and catalyzes the transition state of GTP hydrolysis by $G\alpha$, demonstrated by the fact that the RGS domains bind preferentially to $G\alpha$ -GDP activated with the transition state mimetic, AlF_4^- (Tesmer et al., 1997). Although RGS proteins are classified according to the presence of a RGS domain, we now understand that RGS proteins encompass a wide diversity of multidomain signaling and scaffolding proteins that are categorized by sequence similarity (Fig. 1). Furthermore, we have come to appreciate that regulation of G protein signaling is crucial for normal cellular function, and improperly regulated G protein signaling underlies many disease states (Gerber et al., 2016; Sjögren, 2017), signifying the potential for RGS proteins as therapeutic targets. In this review, we limit and focus our discussion to the 20 canonical RGS proteins. Due to their sequence heterogeneity, we have grouped the discussion and analysis of each RGS within its respective RGS subfamily.

ABBREVIATIONS: CADD, combined annotation-dependent depletion; CoF, change of function; DEP, disheveled EGL10-Pleckstrin; ERK, extracellular signal-regulated kinase; GAP, GTPase-accelerating protein; GGL, G protein γ subunit-like; GoF, gain of function; GPCR, G protein-coupled receptor; GPR, G protein regulatory; LoF, loss of function; MAP, modified alignment position; MTR, missense tolerance ratio; PTM, post-translational modification; RGS, regulators of G protein signaling; SNP, single nucleotide polymorphism; SNV, single nucleotide variant.

Fig. 1. Phylogenetic tree of the human RGS protein family. RGS proteins (human sequence) were phylogenically mapped using Clustal W, and each subfamily was highlighted by color. The R7 subfamily is labeled in blue, followed by the R12 subfamily in gold, the RZ subfamily in red, and finally the R4 family in green. Protein domains are indicated by color: blue RGS are regulator of G protein signaling domains; green DEP are disheveled EGL10-Pleckstrin homology domains; yellow GGL are G protein γ subunit-like domains; pink R1 and R2 are Ras/Rap-binding domains; orange G are G protein regulator motifs; and lavender PTB is phosphotyrosine binding domain.

C. RGS Protein Rare Human Variants in Complex Diseases

Recent genetic analysis of human exomes indicates that an explosion of variants within protein-coding regions arose between 5000 and 10,000 years ago, leading to diverse human traits and a broad range of potential disease determinants (Fu et al., 2013). Population and clinical genome sequencing has generated new information regarding the etiology of disease, and the recent release of large-scale genome and exome sequencing data [Genome Aggregation Database, GnomAD, of the Broad Institute (Lek et al., 2016)] has allowed for the analysis and possible discovery of new disease-linked rare variants. Although some proteins are functionally well-positioned to participate in monogenic diseases (Yuan et al., 2015a; Ogden et al., 2017), others such as RGS proteins are likely involved in more complex pathways that contribute to disease progression (Bansal et al., 2007; Gerber et al., 2016; Xie et al., 2016). Whereas genetic deletion of the ubiquitously expressed heterotrimeric G proteins can result in severe defects or embryonic lethality (Offermanns et al., 1997, 1998; Yu et al., 1998; Wettschureck and Offermanns, 2005; Okabe and Iwakura, 2010; Plummer et al., 2012; Moon et al., 2014), genetic loss of their more discretely expressed modulators, the RGS proteins, results (broadly speaking) in subtle and less detrimental phenotypes with only a few exceptions (see below). RGS proteins therefore participate in complicated multifactorial physiologic processes and disease states (Williams et al., 2004; Zhang and Mende, 2014; Evans et al., 2015; Ganss, 2015; Ahlers et al., 2016), and large-scale genomic association studies may not be able to detect disease-associated polymorphisms (Bomba et al., 2017). Due to the purifying drive of

natural selection, very rare variants (1%–2% or less) in fact seem to play a far greater role as genetic determinants of disease compared with the more common and less deleterious polymorphisms (Nelson et al., 2012; Tennessen et al., 2012), an idea supported by the observed inverse relationship between minor allele frequency and disease risk (Park et al., 2011). The rarity of these variants means that most carriers are heterozygous, as the likelihood of inheriting the same single nucleotide variant (SNV) from each parent is incredibly low. However, both diseases and traits can arise from heterozygosity. For instance, a single variant allele can cause a dominant-negative phenotype (Kenakin and Miller, 2010), an intermediate phenotype (as compared with wild type or knockout) (Okamoto et al., 2017), mislocalization of proteins to cellular compartments leading to aberrant signaling (Lo Bello et al., 2017), or a change of function (CoF) in other ways.

Given this context, in this work we describe a genomics approach to identify likely deleterious rare human variants (defined in this study as less than 2% prevalence) in functionally sensitive regions of RGS proteins, with an emphasis on the well-defined RGS domain. Due to the likely involvement of RGS proteins in complex disease states (Sjögren, 2017), these variants may contribute to a first hit, leaving the carrier more vulnerable to disease given a secondary insult (e.g., environmental, subsequent mutation of a protein in related signaling pathway, or otherwise). Furthermore, both a loss-of-function (LoF) as well as a gain-of-function (GoF) variant could equally disrupt cellular systems in delicate equilibrium. Thus, we suggest an unbiased approach that measures a CoF, which encompasses both variant forms. Finally, it is important to consider that, due to cultural practices, variants may be

expressed more commonly or exclusively within a single ethnic group, and may give rise to unique human traits, rather than the more obvious and deleterious disease states.

In this review, we examine the 20 canonical human RGS proteins (Fig. 1) and their known links to disease or traits, and then present human variants for these gene exomes, extracted from the publically available GnomAD project of the Broad Institute (Lek et al., 2016). In our analysis, we use recently developed bioinformatics, proteomic and structural tools, including combined annotation-dependent depletion (CADD) (Kircher et al., 2014), missense tolerance ratio (MTR) (Traynelis et al., 2017), and post-translational modification (PTM) cluster analysis (Dewhurst et al., 2015; Torres et al., 2016).

With the availability of large-scale databases for human exome sequences comes the major challenge in medical genomics of determining the significance of (or lack thereof) any particular genetic variant. At present, variants of uncertain significance represent the vast majority of those described in genetic reports. MTR offers a new method for summarizing available human variation data within genes to capture population level genetic variation and measures the ratio of tolerance within exome sequences to genetic mutations (Traynelis et al., 2017). This analysis uses publicly available human mutation data to make predictions about domains and motifs that are resistant to, and atypically low in number of, missense mutations. From these ratios, we can predict which exome regions are likely to be functionally sensitive to mutation. Integrating MTR with other selected bioinformatic tools (e.g., CADD and PTM cluster analysis) for any particular exome sequence provides a way of predicting pathogenic missense variants from background missense variation in disease genes. CADD is a complementary bioinformatic measure of mutation severity, but takes into account multiple measures of deleteriousness. Unlike MTR, CADD analysis tells us not where, but how a mutation might affect protein function. CADD independently integrates various diverse annotations into a single measure (C score) for each variant. The score takes into account measures of sequence conservation and amino acid side chain properties and prioritizes functional, deleterious, and pathogenic variants across many functional categories, effect sizes, and genetic architectures to provide researchers with a valuable tool for selecting variants for study. PTMs play critical roles in regulating and determining protein function, the disruption of which can cause disease (Jensen et al., 2002; Hornbeck et al., 2012; Lothrop et al., 2013; Dewhurst et al., 2015; Torres et al., 2016). Inherently, PTMs alter the structure of proteins and therefore have the potential to alter their function as well. In addition, PTMs can have pronounced effects on protein-protein interactions, serving in many cases as handles for PTM-specific binding domains (Walsh et al., 2005). A handful of

experimentally verified PTMs is reported in human RGS proteins (Alqinyah and Hooks, 2018), including phosphorylation, ubiquitination, acetylation, methylation, and palmitoylation. The function of most PTMs (in RGS or other proteins) remains undetermined due in large part to the time and challenge requirements of conducting biochemical experiments as well as a dearth of suitable methods for functional prioritization of existing knowledge.

Due to their important roles in protein structure and function, we identified and reported a comprehensive list of experimentally verified and publically curated PTMs found on human RGS proteins (Li et al., 2014; Huang et al., 2016b) and mapped them with respect to each RGS protein. For this study, we use a recently described protein family-specific PTM alignment analysis method that has proven utility for revealing functional PTM hotspots in protein families (Dewhurst et al., 2015; Torres et al., 2016), and we report those PTMs in RGS proteins that overlap with human variants. Using all of this information (human variants, CADD, and PTMs) in combination, we have prioritized a narrow list of select variants that we predict will disrupt human RGS protein function. As a proof-of-principle to validate the approach, we test one of these selected variants to demonstrate a profound CoF, and highlight others for future study. Although we focus in this work on the RGS domain, we recognize that other RGS protein regions and domains are also essential for RGS protein function. As such, the comprehensive dataset examining these measures for the entire exome sequence for each canonical RGS protein is also provided for any investigator to use (Supplemental Material). We reason that if rare variants occur in functionally sensitive regions of RGS proteins (e.g., the RGS domain) such that they confer a profound CoF phenotype, then these variants likely make important (and previously unappreciated) contributions to complex human disease states and/or unique human traits. As such, we believe that computationally identified rare variants, combined with experiments that validate a CoF for such variants, can provide a deeper understanding of the etiology of complex disease states and the evolution of human traits, while also providing investigators and clinicians a valuable wealth of information toward personalized medicines.

II. RGS Proteins in Physiology and Human Disease

A. The R4 Family

1. R4 Family Overview. Members of the R4 subfamily compose the largest and best characterized of the RGS proteins due to their early discovery and simplicity in structure. R4 family RGS proteins include the following: RGS1, RGS2, RGS3, RGS4, RGS5, RGS8, RGS13, RGS16, RGS18, and RGS21 (Fig. 1). All R4 family members exhibit capacity to bind and act as GAPs for both *Gai/o* and *Gαq* (Hollinger and Hepler,

2002) proteins, although with varying specificity (Heximer et al., 1997, 1999; Huang et al., 1997; Heximer, 2004) based, in some cases, on subtle structural differences (Nance et al., 2013). For example, RGS2 demonstrates much greater selectivity for G α_q (Heximer et al., 1997), whereas RGS4 demonstrates greater selectivity for G α_i (Hepler et al., 1997; Heximer et al., 1999). In addition to the canonical RGS domain, these small RGS proteins also share an N-terminal amphipathic α helix, which, in coordination with N-terminal palmitoylation (Tu et al., 2001), facilitates plasma membrane localization (Bernstein et al., 2000; Heximer et al., 2001; Gu et al., 2007) and consequent actions on G α proteins. Outside of this, the N termini of certain R4 RGS proteins show considerable diversity that determines their specificity for receptor coupling (Bernstein et al., 2004; Neitzel and Hepler, 2006) and regulation by cellular protein degradation pathways (Davydov and Varshavsky, 2000; Bodenstein et al., 2007). Thus, although this review focuses primarily on the conserved RGS domain, readers should note that human variants in these other protein regions can cause a profound CoF phenotype. Table 1 contains a brief overview of each protein within the R4 family, its tissue distribution, and its reported links to physiology and disease. Figure 1 shows a phylogenetic map of all RGS family proteins, with R4 family proteins highlighted in green.

2. R4 Family Proteins in Human Physiology and Disease. Although not initially recognized as a RGS protein, RGS1 was cloned from activated B lymphocytes as an early activation gene, and designated BL34 (Hong et al., 1993) shortly before the RGS domain was characterized. Since its discovery, much has been learned about the role of RGS1 in immune physiology and pathology (Xie et al., 2016). In B and T lymphocytes, RGS1 controls G α_i -mediated chemotaxis and migration (Hwang et al., 2010; Gibbons et al., 2011). RGS1 knockout mice exhibited abnormal B cell migration, exaggerated germinal center formation, and atypical spleens (Moratz et al., 2004). Unsurprisingly, RGS1 has been linked with multiple T- and B-cell-related diseases such as inflammatory bowel disease, multiple sclerosis (Johnson et al., 2010), type 1 diabetes (Smyth et al., 2008), and celiac disease (Hunt et al., 2008; Smyth et al., 2008). RGS1 has also been implicated in atherosclerosis and is upregulated in atherosclerotic plaques and aortic aneurysms, where it regulates monocyte and macrophage chemotaxis (Patel et al., 2015). A recent report demonstrated that elevated RGS1 expression in diffuse large B-cell lymphoma was associated with poor overall survival (Carreras et al., 2017). In this light, LoF variants may play a role in RGS1-mediated immune disorders, or susceptibility to these disorders.

RGS2 is a widely expressed (Kehrl and Sinnarajah, 2002) immediate early gene that is induced by various stimuli (Song et al., 2001), although it is best understood for its roles in the vasculature and the brain. In

the hippocampus, a brain region related to learning and memory, high frequency stimulation (a method used to produce synaptic plasticity and long-term potentiation) induces RGS2 expression (Ingi et al., 1998), and RGS2 knockout mice exhibit impaired basal neuronal activity (Oliveira-Dos-Santos et al., 2000), supporting a role for RGS2 in synaptic plasticity and memory (Gerber et al., 2016). Due to its widespread expression throughout the brain, RGS2 has been linked to many neurologic diseases and affective disorders, including anxiety, post-traumatic stress disorder, and suicide (Amstadter et al., 2009a,b; Koenen et al., 2009; Lifschytz et al., 2012; Okimoto et al., 2012; Hohoff et al., 2015). In particular, a variant in the 3' untranslated region of RGS2, rs4606, is associated with reduced mRNA expression of RGS2 (Semplicini et al., 2006), and is linked to anxiety (Leygraf et al., 2006; Koenen et al., 2009) and suicide (Cui et al., 2008; Amstadter et al., 2009b). In the periphery, RGS2 is expressed in vascular smooth muscle cells, where it is regulated by nitric oxide and controls vascular relaxation (Tang et al., 2003; Sun et al., 2005). RGS2 knockout mice are hypertensive (Heximer et al., 2003), and hypertensive human patients have reduced RGS2 mRNA compared with controls (Semplicini et al., 2006). Furthermore, hypertensive patients were more likely to have a SNV in the 3' untranslated region (rs4606), which correlated with RGS2 expression (Semplicini et al., 2006). In a Japanese cohort, several N-terminal coding mutations (Q2L, M5V, and R44H) were associated with hypertension (Yang et al., 2005). In a subsequent study, it was found that the Q2L mutation destabilized RGS2 protein (but was reversed by proteasomal inhibition), whereas Q2R (another SNV found in both cases and controls) did not (Bodenstein et al., 2007). A more recent study demonstrated that four human variants, including Q2L, R188H, R44H, and D40Y, showed reduced capacity to inhibit Ca²⁺ release by the angiotensin II receptor (Phan et al., 2017). For further review of RGS2 function in physiology, see Bansal et al. (2007). Altogether, RGS2 expression has been found to regulate multiple aspects of normal and pathophysiology, and LoF variants would likely generate similar phenotypes to loss of protein. Finally, RGS2 also has been shown to negatively regulate protein translation by binding directly to the eukaryotic initiation factor 2B ϵ (eIF2B ϵ) subunit via a 37-amino-acid segment within the RGS domain (Nguyen et al., 2009), suggesting variants in this region could impact protein translation.

In contrast to RGS2, less is known about RGS3. RGS3 is found in the heart (Zhang et al., 1998), and mouse overexpression studies have shown that it protects against cardiac hypertrophy (Liu et al., 2014) and also regulates the survival/differentiation responses in multiple cell types (Nishiura et al., 2009; Qiu et al., 2010). In humans, the longest splice variant (and canonical PDZ-containing isoform) RGS3-1 may promote epithelial–mesenchymal

TABLE 1

Summary of RGS distribution, function, disease links, and predicted loss of function effects on human physiology

In this table, we summarize the broad tissue distribution and characterized function on physiology for each RGS family member. Note that this is not an all-encompassing list, but rather where each RGS is best defined. Disease links are sourced from human studies when available and animal studies otherwise, usually correlative data between disease phenotype and protein or transcript expression, and are described in the text.

Overview of RGS Protein Distribution and Function				
RGS Family	RGS	Major Tissue Distribution	Function	Knockout Phenotype
R4 subfamily	RGS1	Immune system (B and T lymphocytes, monocytes, natural killer cells, etc.)	Suppress chemotaxis	Abnormal B cell migration, exaggerated germinal center formation, and atypical spleens
	RGS2	Cardiovascular system and brain	Vascular relaxation	Impaired basal neuronal activity and hypertension
	RGS3	Heart	Still under investigation	Overexpression protects against cardiac hypertrophy
	RGS4	Heart and brain	Respond to cardiovascular stress and modulate neuronal signaling	Susceptible to atrial fibrillation
R5	RGS5	Vasculature	Control vascular remodeling	Reduced blood pressure, improved immune-mediated destruction of solid tumors and survival
	RGS8	Brain (cerebellum)	Still under investigation	None noted
	RGS13	Immune system (B and T lymphocytes, mast cells, etc.)	Suppress immune response	Enhanced B cell responses and anaphylaxis
	RGS16	Retina, heart, brain, liver, immune system	Regulate inflammatory response and circadian rhythms	Enhanced inflammatory response in the lung and dysregulated circadian signaling
R6	RGS18	Platelets	Regulation of hemostasis	Reduced bleeding and hyper-responsive platelet activation
	RGS21	Bitter and sweet taste cells	Gustatory sensation	Unknown
	RGS6	Brain and heart	Regulate drug reward, anxiety, and depression	Protection from alcohol-seeking behaviors, anxiety and depression-like behavior; enhanced dopamine neurodegeneration
	RGS7	Brain	Regulates anxiety and drug reward	Enhanced reward, withdrawal tolerance to morphine
R7 subfamily	RGS9	Retina; brain striatum	Regulation of photoreceptor and striatal signaling	Disrupted striatal dopamine signaling related to

Overview of RGS Protein Distribution and Function

References

Moratz et al., 2004; Hunt et al., 2008; Smyth et al., 2008; Johnson et al., 2010; Patel et al., 2015

Oliveira-Dos-Santos et al., 2000; Heximer et al., 2003; Yang et al., 2005; Leygraf et al., 2006; Semplicini et al., 2006; Amstadter et al., 2009a,b; Koenen et al., 2009; Lifschytz et al., 2012; Okimoto et al., 2012; Hohoff et al., 2015

Ooe et al., 2007; Liu et al., 2014; Chen et al., 2016; Wang et al., 2017a

Owen et al., 2001; Chowdari et al., 2002, 2008; Mittmann et al., 2002; Chen et al., 2004; Williams et al., 2004; Prasad et al., 2005; Emilsson et al., 2006; Guo et al., 2006; Campbell et al., 2008a; So et al., 2008; Ding and Hegde, 2009; Ho et al., 2010; Felkin et al., 2011; Lee et al., 2012; Jaba et al., 2013; Opel et al., 2015

Chang et al., 2007; Cho et al., 2008; Hamzah et al., 2008; Nisancioglu et al., 2008; Holobovskyy et al., 2013; Cheng et al., 2015; Daniel et al., 2016

Gold et al., 2002

Bansal et al., 2008a,b; Pise-Masison et al., 2009; Hwang et al., 2013; Sethakorn and Dulin, 2013; Raedler et al., 2015

Doi et al., 2011; Hayasaka et al., 2011; Shankar et al., 2012; Hu et al., 2016; Jones et al., 2016

Schossner et al., 2011; Haggmark et al., 2014; Mao et al., 2014; Alshbool et al., 2015

Sharma et al., 2016

Bifsha et al., 2014; Schizophrenia Working Group of the Psychiatric Genomics, 2014; Stewart et al., 2014; Moon et al., 2015; Stewart et al., 2015; Ahlers et al., 2016

Crovetto et al., 2001; Gelernter et al., 2001; Hohoff et al., 2009; McCauley et al., 2009

Nishiguchi et al., 2004; Cheng et al., 2007; Hartong et al., 2007; Stockman et al., 2008; Michaelides et al., 2010

(continued)

TABLE 1—Continued

Overview of RGS Protein Distribution and Function					References
RGS Family	RGS	Major Tissue Distribution	Function	Knockout Phenotype	Proposed Disease Links
R12 subfamily	RGS11	Retina	Regulation of photoreceptor signaling	stimulants and motor deficits	Retinal dysfunction
	RGS10	Brain and immune system	Suppression of neuroinflammation	Unknown	Parkinson's disease, possible link to schizophrenia and chemoresistant ovarian cancer
	RGS12	Brain, lung, testis, heart, bone, and spleen	Regulates osteoclast differentiation	Growth retardation and increased bone mass	Potential role for osteoporosis
	RGS14	Brain, spleen, and heart	Regulation of blood calcium homeostasis, long-term potentiation regulation	Improved spatial learning	Kidney disease, possible link to social memory
RZ subfamily	RGS17	Brain	Regulation of drug dependence	Decreased tumor growth and proliferation	Smoking and substance dependence
	RGS19	Heart, lung, brain, and liver	Potential links to cardiac development and regulation of cell proliferation	Enhanced analgesia and heart defects	Multiple sclerosis, ovarian cancers
	RGS20	Brain and eye	Still under investigation	Enhanced analgesia and tolerance to morphine	Hypertension and breast cancer

transition via WNT signaling (Shi et al., 2012), and have a role in docetaxel-resistant breast cancer (Ooe et al., 2007). Other reports link RGS3 expression to various types of cancers, mediated by microRNAs, which bind the 3' untranslated region to regulate RGS3 expression (Chen et al., 2016; Wang et al., 2017a).

RGS4 is selectively expressed in the heart and the brain (Zhang et al., 1998; Ingi and Aoki, 2002), where its expression and protein stability is tightly controlled by its N terminus (Bodenstein et al., 2007; Bastin et al., 2012). Although RGS4 protein in the heart is found at very low levels under normal cardiac physiology (Stewart et al., 2012), it can be upregulated during pathophysiology and cardiac remodeling (Owen et al., 2001; Mittmann et al., 2002; Felkin et al., 2011; Lee et al., 2012; Jaba et al., 2013). Additionally, RGS4 knockout mice are susceptible to atrial fibrillation (Opel et al., 2015). A great deal is known about RGS4 in the brain. First, RGS4 mRNA is decreased in the frontal cortex of schizophrenic patients, whereas other R4 family transcripts were not (Mirnics et al., 2001). Furthermore, several noncoding SNVs [single nucleotide polymorphism (SNP)1, SNP4, SNP7, and SNP18] are found to be associated with RGS4 expression and schizophrenia diagnosis in multiple populations (Chowdari et al., 2002, 2008; Chen et al., 2004; Williams et al., 2004; Prasad et al., 2005; Guo et al., 2006; Campbell et al., 2008a; So et al., 2008). Aside from schizophrenia, RGS4 mRNA levels and SNVs have been linked with Alzheimer's disease (Emilsson et al., 2006) and alcoholism (Ho et al., 2010). RGS4 is associated with Parkinson's disease development, via decreased regulation of dopamine receptors and/or muscarinic acetylcholine receptors (Ding et al., 2006; Lerner and Kreitzer, 2012; Min et al., 2012), although results are varied (Ashrafi et al., 2017). In dopamine-depleted mice, a mouse model for Parkinson's disease, RGS4 is upregulated and inhibits M4 muscarinic autoreceptors, an effect that is mimicked by RGS4 infusion onto untreated cells (Ding et al., 2006). Remarkably, in that study, infusion of a RGS4 construct that lacks the N terminus (and is less potent at M4 receptors) reversed native RGS4-mediated attenuation of M4 signaling in dopamine-depleted mice. That is to say, this dysfunctional RGS4 demonstrated a dominant-negative effect on RGS4-mediated cell signaling, suggesting that fully functional knockouts may arise from only one LoF allele. This supports the prediction that, whereas rare variants overwhelmingly occur in a heterozygous manner (as discussed above), diseases and traits can still arise from one minor allele, in this case as a dominant-negative effect.

RGS5 is found broadly in tissues, including but not limited to the cardiovascular system and brain (Seki et al., 1998). Although there have been several genome-wide association studies suggesting a link between

RGS5 variants and neurologic disorders, such as schizophrenia and bipolar disorder (Campbell et al., 2008b; Smith et al., 2009), its role is best defined in the vasculature. In this work, RGS5 is expressed in arterial smooth muscle cells, and its expression is correlated with protection from hypertension and atherosclerosis (Holobotovskyy et al., 2013; Cheng et al., 2015; Daniel et al., 2016). Specifically, RGS5 is downregulated in various hypertensive animal models (Kirsch et al., 2001; Grayson et al., 2007) as well as atherosclerotic plaques in nonhuman primates (Li et al., 2004) and humans (Adams et al., 2006). Furthermore, RGS5 seems to be an important regulator of vascular remodeling, under both normal and pathophysiological conditions (Armulik et al., 2005; Berger et al., 2005; Wang et al., 2016). For example, vascularization of tumors is normalized in RGS5 knockout mice, and immune-mediated destruction of solid tumors was more effective, resulting in greatly improved survival (Hamzah et al., 2008). This may be due to the role of RGS5 in pericytes (Cho et al., 2003), which critically support nascent vascularization (Mitchell et al., 2008). In this study, RGS5 inhibits multiple signaling pathways induced by angiotensin II, endothelin I, and others (Cho et al., 2003). RGS5-expressing pericytes have a demonstrated role in tumor angiogenesis (Bergers and Song, 2005; Ribeiro and Okamoto, 2015) and, together with previous work (Hamzah et al., 2008), strongly support a role for RGS5 in this disease model. Mouse models also show that blood pressure is reduced in RGS5-null mice (Cho et al., 2008; Nisancioglu et al., 2008). Accordingly, a human genome-wide screen found RGS5 expression was linked to blood pressure regulation (Chang et al., 2007). The role of RGS5 in vascular physiology and pathophysiology was recently reviewed (Ganss, 2015).

RGS8 appears to be enriched throughout the brain (Larminie et al., 2004), especially in Purkinje cells of the cerebellum (Gold et al., 1997; Saitoh and Odagiri, 2003; Saitoh et al., 2003). Interestingly, when RGS8 cDNA is expressed in non-neuronal cells, it accumulates in the nucleus, whereas in Purkinje neurons RGS8 is found within the soma and dendrites (Itoh et al., 2001). Coexpression with constitutively active G α o protein, or expression of RGS8 lacking the N terminus, reversed the nuclear localization (Saitoh et al., 2001, 2003), suggesting either robust nuclear export in Purkinje neurons of RGS8, or a cytosolic-localizing binding partner of RGS8 that is specific to Purkinje neurons versus non-neuronal cells. Furthermore, unlike canonical RGS function on G protein-gated potassium channels that accelerate channel desensitization (Chen et al., 2014; Ostrovskaya et al., 2014; Wydeven et al., 2014), RGS8 speeds up both the on-rate as well as the off-rate channel kinetics (Saitoh et al., 1997). RGS8 knockout mice have been generated, but no overt phenotype or histologic abnormality was found (Kuwata et al., 2008). Thus, RGS8's role in physiology remains uncertain

although likely important for key regulatory processes. Although not much is known regarding RGS8 links to human disease, one study found that electroconvulsive seizures in rats caused an increase in RGS8 mRNA in the prefrontal cortex 2 hours following acute shock and significantly reduced RGS8 mRNA in hippocampus 24 hours following both acute and chronic shock (Gold et al., 2002), suggesting a potential role for RGS8 in seizures.

RGS13 is another immune-specific modulator that is expressed in B and T lymphocytes (Shi et al., 2002; Estes et al., 2004), as well as mast cells (Bansal et al., 2008a,b). In B and T cells, RGS13 acts to desensitize chemokine receptor signaling (Shi et al., 2002; Estes et al., 2004; Han et al., 2006) similar to RGS1, and RGS13 knockout mice have enhanced B cell responses (Hwang et al., 2013). In mast cells, RGS13 constrains allergic responses generated by IgE (Bansal et al., 2008a,b), and RGS13 knockout mice exhibit enhanced IgE-mediated anaphylaxis. RGS13 transcript is greatly increased in adult T cell leukemia/lymphoma (Pisemason et al., 2009; Sethakorn and Dulin, 2013) and asthma (Raedler et al., 2015), underscoring both its role in immune cells and the importance of homeostatic balance of RGS13 signaling.

Originally cloned from the retina (Chen et al., 1996; Natochin et al., 1997), RGS16 has a relatively broad expression pattern, including the heart (Patten et al., 2002), brain (Grafstein-Dunn et al., 2001), liver (Kurraasch et al., 2004), and immune system (Beadling et al., 1999; Shi et al., 2004; Kveberg et al., 2005; Kim et al., 2006). RGS16 localization and GAP activity are regulated by addition of a palmitate at multiple cysteine residues (Hiol et al., 2003; Osterhout et al., 2003). As with many R4 family members, one of the most well-defined functions for RGS16 has been its role in adaptive immunity. RGS16 is involved in trafficking and migration of T lymphocytes (Estes et al., 2004) in response to an allergen challenge (Lippert et al., 2003), and RGS16 knockout mice have an enhanced inflammatory response in the lung (Shankar et al., 2012). Apart from its defined roles in allergic responses, RGS16 also has an intriguing role in regulating circadian systems (Goto et al., 2017). Within the brain, RGS16 is expressed in both the suprachiasmatic nucleus (SCN) and the thalamus (Grafstein-Dunn et al., 2001; Ueda et al., 2002). The SCN sets the global circadian clocks through cyclical signaling pathways, where RGS16 expression is cyclical. Loss of RGS16 causes dysregulation in circadian signaling in the SCN as well as delayed, shorter circadian behavioral activity in mice (Doi et al., 2011; Hayasaka et al., 2011). Another component of SCN circadian regulation is feeding behavior, which is synchronized with circadian rhythms coregulated by the liver. In mice, food anticipatory activity was found to be attenuated in RGS16 knockdown mice during a restricted feeding schedule (Hayasaka et al., 2011). Complementary studies in the

liver showed that RGS16 knockout mice had higher rates of fatty acid oxidation, whereas mice that over-express RGS16 had lower rates of fatty acid oxidation and higher blood triglyceride levels (Pashkov et al., 2011). Interestingly, in humans, noncoding variants in RGS16 have been linked with self-reported “morning people” (Hu et al., 2016; Jones et al., 2016). Finally, RGS16 expression has been linked with various cancers (Liang et al., 2009; Miyoshi et al., 2009; Carper et al., 2014).

RGS18 expression is mostly confined to bone marrow-derived cells (Nagata et al., 2001; Park et al., 2001; Yowe et al., 2001), more specifically in platelets (Gagnon et al., 2002). Although very little is known about RGS18 beyond its expression, mechanistically it seems to be important for regulating platelet activation (Gegenbauer et al., 2012; Ma et al., 2012). RGS18 knockout mice have reduced bleeding compared with wild-type mice, hyper-responsive platelet activation (Alshbool et al., 2015), and reduced platelet recovery following acute thrombocytopenia (Delesque-Touchard et al., 2014). In humans, RGS18 mRNA is elevated in aspirin-resistant platelets (Mao et al., 2014), and RGS18 protein was elevated in amyotrophic lateral sclerosis patients (Hagmark et al., 2014). Beyond its role in platelets, human SNVs near the RGS18 gene were associated with suicide attempts (Schosser et al., 2011), although no clear mechanism for this is known.

Among the R4 family, RGS21 is the smallest and most recently cloned RGS protein (von Buchholtz et al., 2004). It was originally cloned from bitter and sweet taste cells, although the protein may be expressed much more broadly (Li et al., 2005). In bitter taste cells, RGS21 was found to inhibit bitter taste signaling to cAMP, suggesting a role of RGS21 in the gustatory system (Cohen et al., 2012). Since then, human SNVs have linked RGS21 with celiac disease (Sharma et al., 2016). Nonetheless, roles for RGS21 in physiology and disease remain largely unexplored.

B. The R7 Family

1. R7 Family Overview. The R7 family of RGS proteins is composed of RGS6, RGS7, RGS9, and RGS11 (Fig. 1). These are highly homologous proteins mostly expressed in the nervous system, where they have a role in neuronal G protein signaling controlling nociception, reward behavior, motor control, and vision (Gold et al., 1997; Anderson et al., 2009; Gerber et al., 2016). The R7 RGS proteins contain distinctive domains that form stable stoichiometric heterotrimeric complexes with accessory binding partners that control protein-protein interaction, subcellular localization, and protein stability (Anderson et al., 2009; Sjögren, 2011). Besides the canonical RGS domain, other domains include the disheveled EGL10-Pleckstrin (DEP) homology domain, a R7 homology domain, and a G protein γ subunit-like

(GGL) domain (Gold et al., 1997; Sjögren, 2011; Ahlers et al., 2016; Gerber et al., 2016). The RGS domain is located at the C terminus, where it stimulates GTP hydrolysis on Gai/o protein subunits (Snow et al., 1998b; Posner et al., 1999; He et al., 2000; Hooks et al., 2003; Martemyanov and Arshavsky, 2004; Anderson et al., 2009; Masuho et al., 2013; Stewart et al., 2015). The GGL domain, located upstream from the RGS domain, is structurally homologous to conventional γ subunits of G proteins (Posner et al., 1999; Anderson et al., 2009) and binds G β 5 (type 5 G protein β subunit) as an obligatory partner (Anderson et al., 2009), which is crucial for protein stability (Snow et al., 1998b; Anderson et al., 2009; Sjögren, 2011; Gerber et al., 2016). Consistent with the brain expression patterns of R7 family members, various neurologic conditions such as anxiety, schizophrenia, drug dependence, and visual complications have been linked with the function of these proteins. Table 1 contains a brief overview of each protein within the R7 family, its tissue distribution, and its reported links to physiology and disease. Figure 1 shows a phylogenetic map of all RGS family proteins, with R7 family proteins highlighted in blue.

2. R7 Family Proteins in Human Physiology and Disease. RGS6 is highly expressed, at both the mRNA level and protein level, in brain tissue and in the heart (Gold et al., 1997; Ahlers et al., 2016). Within heart, RGS6 functions as an essential modulator of parasympathetic activation to prevent parasympathetic override and severe bradycardia (Yang et al., 2010). Studies relating RGS6 to human diseases are limited, although literature suggests that RGS6-specific modulation of G α may be involved in regulating several central nervous system diseases such as alcoholism (Stewart et al., 2015), anxiety and depression (Stewart et al., 2014), Parkinson's disease (Bifsha et al., 2014), Alzheimer's disease (Moon et al., 2015), schizophrenia (Schizophrenia Working Group of the Psychiatric Genomics, 2014), and vision (Chograni et al., 2014). Prolonged exposure to alcohol upregulates RGS6 protein in a brain region known as the ventral tegmental area of wild-type mice. RGS6 knockout mice have reduced striatal dopamine, ameliorated alcohol-seeking behavior, and a reduction in alcohol-conditioned reward and withdrawal (Stewart et al., 2015). RGS6 knockout mice also showed protection from pathologic effects of chronic alcohol consumption on peripheral tissues, believed to be due to direct or indirect regulation by RGS6 of reactive oxygen species (Stewart et al., 2015; Ahlers et al., 2016). RGS6 is also enriched in dopaminergic neurons of the substantia nigra pars compacta, which are characteristically lost in Parkinson's disease (Ahlers et al., 2016). Studies have shown that RGS6 knockout mice, but not wild-type mice, suffered from age-onset neurodegeneration of these neurons by the first year. These results also correlate with a decrease in

gene products associated with differentiation and maintenance of dopamine neurons during development (Li et al., 2009; Ahlers et al., 2016), and whose expression is dysregulated in RGS6 knockout mice (Ahlers et al., 2016). Within mouse brain, RGS6 is also expressed in cortical and hippocampal neurons, where it mediates anxiety and depression (Stewart et al., 2015). RGS6 knockout mice displayed spontaneous anxiolytic and antidepressant behaviors that are sensitive to 5-HT_{1A} receptor antagonism (Ahlers et al., 2016). Several studies have suggested RGS6 SNVs are significantly associated with multiple central diseases, including Alzheimer's disease (rs4899412) (Moon et al., 2015; Ahlers et al., 2016) and schizophrenia (rs2332700) (Schizophrenia Working Group of the Psychiatric Genomics, 2014; Ahlers et al., 2016). Accordingly, the RGS6 gene can influence the pathophysiological processes underlying Alzheimer's disease (Moon et al., 2015). Outside of the brain, atypical RGS6 protein expression is linked to several forms of cancer. A C→T SNV located in the 3' untranslated region of the RGS6 gene was associated with a 34% reduction in bladder cancer and an increase in RGS6 protein (rs2074647) (Berman et al., 2004; Ahlers et al., 2016). RGS6 expression was found to be negatively correlated with human pancreatic cancer (Ahlers et al., 2016), human breast cancer progression (Maity et al., 2011, 2013; Ahlers et al., 2016), and resistance to chemotherapies (Maity et al., 2013). Finally, in roles unrelated to cancer, a splice mutation in RGS6 was identified as a genetic cause of autosomal recessive congenital cataract, mental retardation, and microcephaly in two Tunisian siblings (Chograni et al., 2014).

RGS7 is highly expressed in brain, in particular regions linked to anxiety such as the amygdala, hippocampus, brain stem, and hypothalamus (Larminie et al., 2004; Hohoff et al., 2009). RGS7 mRNA is found in abundance in neurons of ventral tegmental area and nucleus accumbens (Sutton et al., 2016), which have roles in drug reward and reinforcement. Within this circuit, the euphoric and analgesic effects of morphine are mediated by the μ -opioid receptor. RGS7 knockout mice showed an enhancement in reward behavior, increased analgesia, delayed tolerance, and heightened withdrawal in response to morphine administration (Sutton et al., 2016). Furthermore, chromosome 1q43, which contains the RGS7 gene, has been reported as a risk loci for panic disorder (Hohoff et al., 2009), and RGS7 SNV rs11805657 is associated with panic disorder (Crowe et al., 2001; Gelernter et al., 2001; Hohoff et al., 2009). Outside of the brain, a Genome-wide Association Study shows modest evidence for the involvement of RGS7 intron variants (rs4660010 and rs261809) in multiple sclerosis. Accordingly, it has been suggested that alteration in RGS7 function could potentially impair the normal dampening of the inflammatory response, leading to multiple sclerosis (McCauley

et al., 2009). Overall, less is known about RGS7's link to physiology and pathophysiology relative to other RGS, and the human variant information provided in this review may prove helpful in defining RGS7's involvement in potential traits or disease.

The RGS9 gene forms two products: a short retina-specific transcript variant (RGS9-1) where it acts as a GAP on mammalian photoreceptor-linked G proteins (He et al., 1998), and a long brain-specific transcript variant (RGS9-2) enriched in the striatum (Zhang et al., 1999). The full-length human RGS9-1 consists of 484 amino acids, whereas RGS9-2 contains almost 200 extra amino acids at its C terminus (Zhang et al., 1999). Both RGS9 variants have a RGS domain and a DEP domain that binds to adaptor proteins such as R9AP (RGS9-1 anchor protein) in the retina, and R7BP (R7 binding protein) in the brain (Traynor et al., 2009). RGS9-1 accelerates the GTPase activity of transducin and colocalizes with other members of the phototransduction cascade (He et al., 1998; Zhang et al., 1999). During the recovery phase of visual transduction, RGS9 is anchored to a photoreceptor outer segment by R9AP, and mutations in both proteins have been associated with stationary retinal dysfunction syndrome, including RGS9 W299R (Nishiguchi et al., 2004; Cheng et al., 2007; Hartong et al., 2007; Stockman et al., 2008; Michaelides et al., 2010). RGS9-1 loss of function in the retina leads to bradyopsia, an inability to see moving objects during sudden changes in light intensity (Nishiguchi et al., 2004). Importantly, variants such as R128X (a nonsense mutation) in RGS9-1 create a truncated gene product lacking important domains crucial for a functional protein (Michaelides et al., 2010). RGS9-2 is expressed in the striatum of rat (Gold et al., 1997; Zhang et al., 1999) and human brain (Thomas et al., 1998; Zhang et al., 1999; Rahman et al., 2003; Liou et al., 2009), a region highly involved in antipsychotic-induced tardive dyskinesia. Genetic differences in RGS9-2 may play a role in patients developing tardive dyskinesia after antipsychotic treatment, because of RGS9-2 regulation of the D2 dopamine receptor (Rahman et al., 2003; Cabrera-Vera et al., 2004; Kovoov et al., 2005; Cerver et al., 2010; Waugh et al., 2011). Intronic SNVs rs8077696, rs8070231, and rs2292593 were reported to likely alter binding efficiency of RGS9-2 to D2DR and play an important role in the development of tardive dyskinesia (Liou et al., 2009). RGS9-2 also regulates the μ -opioid receptor (Zachariou et al., 2003; Psifogeorgou et al., 2007, 2011; Waugh et al., 2011) and modulates reward responses through both opioid and dopamine receptors (Hooks et al., 2008; Waugh et al., 2011). These neurotransmitter systems regulate feeding behavior and body weight (Bodnar, 2004; Gainetdinov, 2007; Waugh et al., 2011) in addition to reward response (Le Merrer et al., 2009; Johnson and Kenny, 2010; Waugh et al., 2011). A study showed an association between rs3215227

(an intronic variant) and significant higher body mass index in East Asian subjects (Waugh et al., 2011). RGS9-2 involvement in the dopamine reward pathway has suggested its involvement in addiction behavior. Cocaine self-administration in rats shows decreased RGS9-2 levels in the striatum compared with controls (Rahman et al., 2003; Traynor et al., 2009). RGS9 knockout mice have distinct locomotor-activating actions of dopaminergic or opioidergic agents such as cocaine, amphetamine, or morphine when compared with wild-type mice (Rahman et al., 2003; Blundell et al., 2008; Traynor et al., 2009). The absence of RGS9 also shows accelerated locomotor sensitization and increased reward sensitivity (Psifogeorgou et al., 2007; Blundell et al., 2008; Traynor et al., 2009).

RGS11 is highly expressed in the brain, especially in retinal bipolar and nerve cells (Rao et al., 2007; Cao et al., 2012; Shim et al., 2012), as well as outside the brain (Yang et al., 2016b). RGS11 interacts with mGluR6 at the dendritic tips of ON bipolar cells to regulate light-evoked responses (Cao et al., 2012). *Gβ5* knockout mice have greatly reduced RGS11 expression in the retina, resulting in dysfunctional photoreceptor signaling (Rao et al., 2007; Cao et al., 2012; Shim et al., 2012). Beyond the brain, recent studies have shown that upregulation of RGS11 might play a role in cancer. RGS11 is highly overexpressed in multiple tumors and associated with increased primary tumor status, nodal metastasis, and disease stage (Yang et al., 2016b). In colorectal cancer, RGS11 is upregulated and involved in chemotherapy resistance (Martinez-Cardus et al., 2009; Yang et al., 2016b). Although RGS11's role in retinal bipolar and nerve cells has been described, its role in cancer and other diseases remains yet to be fully defined (Yang et al., 2016b).

C. The R12 Family

1. R12 Family Overview. The R12 family is a diverse group of RGS proteins, consisting of three members: RGS10, RGS12, and RGS14. Each has its own unique structure and function, but share a conserved RGS sequence and dynamic nuclear shuttling (Burgon et al., 2001; Chatterjee and Fisher, 2002; Cho et al., 2005; Waugh et al., 2005; Shu et al., 2007). Whereas RGS10 is a small, simple RGS protein that resembles the R4 family members, RGS12 and RGS14 have larger and more complex structures that share homology. Both RGS12 and RGS14 contain accessory domains, including two tandem Ras/Rap-binding domains (R1 and R2) and a G protein regulatory (GPR) motif. The R1 domains of RGS12 and RGS14 each interact with small G proteins such as Rap2 and H-Ras to regulate mitogen-activated protein kinase signaling (Traver et al., 2000; Willard et al., 2009; Shu et al., 2010; Vellano et al., 2013). The GPR motif binds inactive (as opposed to active GTP-bound) $G\alpha$ proteins and serves as an inhibitor of GDP release (Kimple et al., 2001, 2002, 2004; Mittal and

Linder, 2004) and also a regulator of RGS protein subcellular localization and membrane attachment (Shu et al., 2007; Brown et al., 2015b). RGS12 is expressed in humans as multiple splice variants (Chatterjee and Fisher, 2000), the longest of which (called trans-spliced, RGS12-TS) contains two additional domains: a PDZ domain and a PTB domain. PDZ domains are important regulators of localization and interaction with binding partners (Dunn and Ferguson, 2015). For example, RGS12-TS binds to CXCR2 via its PDZ domain (Snow et al., 1998a) as a means of directing to its target signaling partners. The PTB domain binds phosphotyrosines, and one report demonstrated that the PTB domain of RGS12 can attenuate platelet-derived growth factor-induced phosphorylated extracellular signal-regulated kinase (ERK) (Sambi et al., 2006). The demonstrated roles of these accessory domains are important to consider in RGS protein function/regulation beyond the canonical RGS domains highlighted in our review in this work. Table 1 contains a brief overview of each protein within the R12 family, its tissue distribution, and its reported links to physiology and disease. Figure 1 shows a phylogenetic map of all RGS family proteins, with R12 family proteins highlighted in gold.

2. R12 Family Proteins in Human Physiology and Disease. RGS10, at 20 kDa, is one of the smallest RGS family proteins, and is highly expressed in the brain and immune system (Gold et al., 1997; Haller et al., 2002). In humans, there are three splice variants of RGS10, differing by only a few amino acids at the N terminus. However, these small differences can have a substantial effect on RGS10 function, as the shortest splice variant (lacking only 14 amino acids) has impaired GAP activity (Ajit and Young, 2005). RGS10 is also dynamically regulated within the cell. Palmitoylation of an N-terminal cysteine targets RGS10 to the plasma membrane and enhances its GAP activity (Tu et al., 1999), whereas phosphorylation of a C-terminal serine targets RGS10 to the nucleus and impedes its GAP activity (Burgon et al., 2001). RGS10 has been documented in the nuclei of microglia and neurons (Waugh et al., 2005), where it may serve to regulate neuroinflammation. Indeed, RGS10 has been shown to promote survival of dopaminergic neurons via regulation of neuroinflammatory pathways in nigrostriatal circuits (Lee et al., 2008, 2011), implicating a neuroprotective role for RGS10 in dopaminergic disorders such as Parkinson's disease (Tansey and Goldberg, 2010). Interestingly, a polymorphism (V38M or V44M in canonical sequence) in RGS10 was found in Japanese patients with schizophrenia, but it was not found to be significantly associated with disease due to sample size (Hishimoto et al., 2004). In peripheral immune cells, RGS10 regulates macrophage activation (Lee et al., 2013) and platelet activation (Hensch et al., 2016) and T lymphocytes (Lee et al., 2016), with potential roles in

clotting or autoimmune diseases. Additionally, loss of RGS10 in aged mice is linked with dysregulated peripheral immune cells and inflammatory cytokines (Kannarkat et al., 2015). Last, there is a curious link between RGS10 and chemoresistant ovarian cancer (Hooks et al., 2010; Ali et al., 2013; Cacan et al., 2014; Hooks and Murph, 2015), potentially via a Rheb-GTP/mTOR pathway (Altman et al., 2015). A comprehensive review of the roles of RGS10 in neurons and immune cells was recently published (Lee and Tansey, 2015).

RGS12, in contrast to RGS10, is the largest RGS protein family member, with multiple splice variants ranging in size from 55 to 155 kDa. As outlined above, RGS12 contains additional signaling domains other than a RGS domain (Snow et al., 1997, 1998a; Ponting, 1999) that interact with various proteins. Beyond this, RGS12 also has been shown to interact with calcium channels in neurons (Schiff et al., 2000; Richman et al., 2005), and de novo mutations have been linked with schizophrenia (Xu et al., 2011). RGS12 expression has been reported throughout the body, including the brain, lung, testis, heart, and spleen (Snow et al., 1997; Doupnik et al., 2001). Like RGS10, RGS12 also shuttles in and out of the nucleus, where it has been shown to repress transcription (Chatterjee and Fisher, 2002; Lopez-Aranda et al., 2006). RGS12 is also expressed in osteoclasts and regulates differentiation (Yang and Li, 2007). Accordingly, RGS12 knockout mice have aberrant bone mass (Yang et al., 2013; Yuan et al., 2015b), suggesting a potential role in osteoporosis. Finally, RGS12 has been linked with cardiac hypertrophy (Huang et al., 2016a) and various cancers (Dai et al., 2011; Wang et al., 2017b).

RGS14 is a ~60-kDa protein within the R12 family that is expressed in brain, heart, and spleen of rodents (Snow et al., 1997; Hollinger et al., 2001; Li et al., 2016). Although its brain expression pattern in adult rodents is largely limited to hippocampal area CA2, RGS14 has a wider brain distribution pattern in monkey and human brain (Squires et al., 2018), including multiple nuclei of the basal ganglia. Of note, within striatum of monkey brain, RGS14 appears to express several shorter splice variants not observed in rodents (Squires et al., 2018). Although RGS14 has not been conclusively linked with any specific diseases in the brain, a Genome-wide Association Study identified RGS14 as a risk factor for multiple sclerosis (Ryu et al., 2014), and a follow-up mouse study confirmed differential expression in a mouse model of multiple sclerosis (Sevastou et al., 2016). RGS14 also suppresses hippocampal-based synaptic plasticity and learning in the CA2 region of the hippocampus in mice (Lee et al., 2010), which has been linked to social and contextual memory (Hitti and Siegelbaum, 2014; Alexander et al., 2016; Dudek et al., 2016; Piskorowski et al., 2016). Although RGS14's role in these behaviors has not yet been fully elucidated

(Evans et al., 2015), its high level of expression in area CA2 suggests it may be a key regulator of hippocampal-based learning and memory. Outside of the hippocampus, RGS14's role within the basal ganglia suggests a link to movement disorders, such as Parkinson's disease, and transcriptional studies of Parkinson's patients showing decreases in RGS14 mRNA support a possible role in this process (Vogt et al., 2006). In the periphery, RGS14 expression is downregulated in failing human hearts, which suppresses cardiac remodeling through regulation of the mitogen-activated protein kinase kinase/ERK pathway (Li et al., 2016). RGS14 interacts with active H-Ras-GTP and Raf-1 and to block ERK signaling (Willard et al., 2009; Shu et al., 2010; Vellano et al., 2013), and RGS14 actions on cardiac remodeling presumably are mediated through one or both Ras/Rap-binding domains on RGS14 (Li et al., 2016), highlighting the importance of accessory domains on RGS protein functions independent of the canonical RGS domain. Finally, multiple genetic studies have found variants in the proximity of the RGS14 gene that are associated with kidney disease (Urabe et al., 2012; Yasui et al., 2013; Mahajan et al., 2016) and altered serum concentrations of both parathyroid hormone (Robinson-Cohen et al., 2017) and phosphorous (Kestenbaum et al., 2010), implicating a potential role for RGS14 in regulating the homeostasis of serum phosphate and other ions.

D. The RZ Family

1. RZ Family Overview. The RZ family is composed of RGS17, RGS19, and RGS20. These all are small, simple RGS proteins similar to the R4 family members. However, unique to the RZ family members is a conserved string of cysteine residues found near their N termini that is palmitoylated and regulates both their membrane localization and interaction with binding partners (De Vries et al., 1996; Nunn et al., 2006). RZ proteins also function as adapter proteins for $G\alpha$ subunit degradation and play important roles in the regulation of signaling and cytoskeletal events in the brain (Mao et al., 2004). They are also highly conserved in metazoans and most closely related to the R4 RGS family (Sierra et al., 2002; Nunn et al., 2006). All members of this family can bind to certain members of the $G\alpha_i$ and $G\alpha_q$ subfamily, but with some selectivity (Tu et al., 1997; Glick et al., 1998; Mao et al., 2004). Table 1 contains a brief overview of each protein within the RZ family, its tissue distribution, and its reported links to physiology and disease. Figure 1 shows a phylogenetic map of all RGS family proteins, with RZ family proteins highlighted in red.

2. RZ Family Proteins in Human Physiology and Disease. RGS17, also known as RGSZ2, demonstrates GAP activity for $G\alpha_i/o$, $G\alpha_z$, and $G\alpha_q$ (Mao et al., 2004). In humans, RGS17 is expressed in the nucleus accumbens, hippocampus, and putamen, with highest

expression found in the cerebellum (Mao et al., 2004; Hayes and Roman, 2016). However, outside the brain, RGS17 has been reported to be overexpressed in human lung adenocarcinomas and prostate cancer (Mao et al., 2004; James et al., 2009; You et al., 2009). In lung, colon, and prostate tumor cell lines, knocking down RGS17 results in decreased tumor growth and tumor cell proliferation; conversely, overexpression of RGS17 in these cell lines resulted in increased tumor growth (James et al., 2009; You et al., 2009). Underscoring this, SNVs in the first intron of the RGS17 gene (rs6901126, rs4083914, and rs9479510) are associated with lung cancer (You et al., 2009). RGS17 also is overexpressed in human liver cancer (Hayes and Roman, 2016). Furthermore, when ovarian cancer cells are treated with chemotherapeutic agents, there is a loss of RGS17 expression, suggesting a role for RGS17 in chemoresistance, perhaps by promoting cell survival via phosphatidylinositol 3-kinase/AKT signaling in these cells (Hooks et al., 2010; Hayes and Roman, 2016). Outside of RGS17 links to cancer, postmortem brain samples from patients with clinical depression show a decrease in RGS17 expression (Shelton et al., 2011; Hayes and Roman, 2016). Furthermore, SNVs in RGS17 found in the promoter region (rs596359) and introns (rs6931160, rs9397585, rs1933258, rs9371276, rs516557, and rs545323) are associated with substance dependence (Zhang et al., 2012; Hayes and Roman, 2016). Finally, two intronic RGS17 SNVs are associated with smoking initiation, rs7747583 and rs2349433 (Yoon et al., 2012; Hayes and Roman, 2016). Whereas possible mechanisms are unknown, expression of RGS17 in brain regions known to be linked to substance dependence supports the relationship between SNVs and these diseases.

Comparatively less is known about RGS19, which is highly expressed (by mRNA) in the heart, lung, and liver, but very low in brain, and seems to regulate proliferation of embryonic stem cells (De Vries et al., 1995; Ji et al., 2015). Unlike the other two RZ family members, RGS19 preferentially interacts with Gai3 (De Vries et al., 1995). One report showed loss of RGS19 slightly enhances opioid-induced analgesia at μ -opioid receptors (Garzon et al., 2004), a brain-specific function that suggests even low expression can have a functional impact. RGS19 expression levels are reportedly upregulated in several disease states, including multiple sclerosis (Igci et al., 2016) and ovarian cancers (Tso et al., 2011). RGS19 is also highly expressed in human neuroblastoma SH-SY5Y cells (Wang and Traynor, 2013). However, in other instances, RGS19 has been reported to inhibit Ras activation by upregulating Nm23, a tumor metastasis suppressor (Wang et al., 2013). Transgenic mice overexpressing RGS19 exhibited multiple heart defects during development and increased expression of heart failure-related biomarkers, including B-type natriuretic peptide and β -major

histocompatibility complex (Ji et al., 2010). Although there are few reports to date defining RGS19 in human disease, the overexpression studies in mice highlight the potential heart disease contribution of GoF variants within the coding region.

RGS20, also known as RGSZ1 and Ret RGS, selectively interacts with G α z and Gai2 subunits (Wang et al., 1998, 2002). The RGS20 transcript is highly expressed in human caudate nucleus and temporal lobe (Wang et al., 1998), and RGS20 splice variants are detectable in the eye (Barker et al., 2001; Yang et al., 2016a). Loss of RGS20 in mice leads to enhanced μ -opioid-induced analgesia and tolerance to morphine (Garzon et al., 2004). Outside the brain, RGS20 is found at significantly high levels in melanoma and metastatic breast cancer cells (Yang et al., 2016a). Furthermore, expression of RGS20 in HeLa, breast adenocarcinoma MDA-MB-231, non-small cell lung carcinoma H1299, and A549 cells results in enhanced cell aggregation, migration, invasion, and adhesion, suggesting a role for RGS20 in tumor metastasis (Yang et al., 2016a). A recent study on triple-negative breast cancer reported RGS20 was overexpressed in those tissues, and that protein expression correlated with disease progression/prognosis, suggesting a novel target for therapy (Li et al., 2017). Finally, RGS20 is reported to be significantly associated with hypertension, where it may synergistically interact with other genes to predispose patients to hypertension (Kohara et al., 2008).

III. Analysis of Rare Human Variants of RGS Proteins

As outlined above, RGS proteins play key roles in human physiology and disease, and rare human variants are thought to underlie many complex human diseases and traits. Therefore, we took advantage of recently available human exome sequencing databases and newly described bioinformatics/proteomic analytical tools to identify rare human variants in canonical RGS proteins that we predict will have a marked CoF phenotype. Using knowledge (published structural Protein Data Bank files) of the binding interface between the RGS domain and partner G α gained by crystallography and sequence conservation, we focus on individual variants of interest derived from these vast datasets that are likely to disrupt RGS-G α interactions. Whereas a LoF or GoF may lead to disease states, a GoF in one pathway may in fact lead to a LoF in a completely separate pathway. Thus, as described above, we consider a CoF that accounts for either LoF or GoF. We postulate that these variants may be missed by genome association studies due to their rarity (1%–2% or less), but nonetheless confer the same phenotype (i.e., multiple genotypes may have the same phenotype), or may redirect the RGS function (e.g., by mislocalization) to affect atypical pathways. Given these caveats, in this

TABLE 2

Human rare variants in each representative family member's RGS domain are predicted to disrupt function due to structural changes or inability to bind/GAP Gα

Residues were selected based on our criteria (outlined in the text) of CADD >20, residues that overlap with conserved sites important for Gα interaction (determined by structural insights), and PTM alignment analysis. Each residue mutation (native, residue position, mutation) has a description as to why it is predicted to disrupt function ("LoF Rationale"), its associated combined annotation-dependent depletion score ("CADD"), associated genic intolerance score ("MTR"), number of reported PTMs, number of neighboring PTMs (where a PTM is found within seven amino acids before/after), and prevalence in the population ("Prevalence"). Note that if the human variants are reported in the noncanonical transcript, the analogous residue position for the canonical sequence is coded in gray.

RGS Protein	Selected Human RGS Protein Rare Variants						
	Residue Mutation	LoF Rationale	CADD	MTR	PTM	Neighboring PTMs	Prevalence
RGS4	L170(73)P	Proline in α helix	23	1.01	0	0	0.0043% (South Asian)
	E214(117)K	Salt bridge partner	27.3	0.50	0	0	0.0063% (Latino)
	D227(130)G	100% conserved, salt bridge partner	29.6	0.90	0	0	0.0065% (African)
	R231(134)W	Highly conserved, stabilizes switch III	29.8	1.04	0	0	0.0065% (African)
	D260(163)N	Highly conserved, salt bridge partner, very high CADD	32	1.19	0	0	0.0065% (African)
	D260(163)G	Highly conserved, salt bridge partner	29.4	1.19	0	0	0.0009% (European)
	R263(166)C	Contact with α helical domain of Gαi1	28.4	1.07	0	0	0.0116% (East Asian)
	R263(166)H	Contact with α helical domain of Gαi1	24.5	1.07	0	0	0.0032% (Global)
	R264(167)C	Highly conserved, salt bridge partner, very high CADD	35	1.06	0	0	0.0009% (European)
RGS9	W299R	Link to disease, participates in electrostatic interaction	29.1	0.71	0	1	0.0229% (European)
	R364C	Stabilizes α5-α6 loop of RGS domain	29.5	0.78	0	0	0.0018% (European)
	K400Q	100% conserved, contact with α helical domain	28.1	0.97	0	0	0.0018% (European)
	R406C	100% conserved, salt bridge partner, very high CADD	35	0.98	0	1	0.0030% (Latino)
	R406H	100% conserved, salt bridge partner, very high CADD	34	0.98	0	1	0.0009% (European)
RGS10	Y413C	Phosphorylation	27.2	0.94	1	1	0.0032% (South Asian)
	L46(38)P	Proline in α helix	29.6	0.78	0	2	0.0032% (South Asian)
	V52(44)M	Possible link to schizophrenia, hydrophobic core partner	34	0.78	0	1	1.9239% (East Asian)
	D141(133)N	Highly conserved, ionic stabilization, very high CADD	35	0.90	0	2	0.0183% (Other)
	R145(137)C	Highly conserved, ionic stabilization, very high CADD	35	0.86	0	2	0.0617% (East Asian)
RGS17	K148(140)R	Ubiquitination	25	0.81	1	1	0.0009% (European)
	E148G	100% conserved, likely ionic interaction	28.2	0.87	0	0	0.0010% (European)
	P166L	100% conserved within and across family, high CADD	28.3	0.73	0	1	0.0009% (European)
	R189T	100% conserved, likely ionic interaction	28.1	1.03	0	0	0.0009% (European)

study we analyze the entire coding sequence for all 20 canonical RGS proteins (Supplemental Material) and provide a detailed case study of a representative RGS protein from each subfamily that we highlight in the main body of this review (RGS4 from R4, RGS9 from R7, RGS10 from R12, and RGS17 from RZ) (Figs. 2–6).

Selection of rare variants that have potential to produce CoF phenotypes was based on several factors described in this work. First and foremost, we identified rare variants as those for which prevalence is well below 2% in a given population (for this dataset: minimum = 0.0009%, maximum = 1.92%, median = 0.0032%). Second, we used combined annotation-dependent depletion (CADD) analysis, presented as a C score (Supplemental Data; Table 2), to estimate the potential deleteriousness of each variant—a feature that strongly correlates with both molecular functionality and pathogenicity (Kircher et al., 2014). A CADD C score above 20 (top 1%), which we used as a hard filter, indicates a very high likelihood of deleteriousness (for this dataset: minimum = 23, maximum = 35, median = 29.4). Third, we considered structural data and sequence conservation when available (Figs. 2A, 4A, 5A, and 6A). Variants

that overlap with identified contact points between the RGS domain and Gα provide a strong case for CoF and were selected. Similarly, variants that overlap with highly conserved residues (particularly those that reside in critical regions of the RGS domain as determined by high-resolution structural data) were selected. This criterion was particularly important for eliminating variants that fall on highly variable residues within the family (i.e., if the amino acid properties are not conserved within the family, it is likely a noncritical residue). Fourth, we used a newly described bioinformatics tool to measure the tolerance of each position in each RGS protein for mutation, expressed as a MTR (Figs. 2B, 4B, 5B, and 6B; Supplemental Material; Table 2), which estimates the functional sensitivity of a given residue to mutation (Traynelis et al., 2017). MTR represents a novel measure of purifying selection acting on missense variants in a 31-codon sliding window across the sequence of a gene. Neutrality, or a MTR of 1.0, represents the point at which the observed number of missense variants is equivalent to the expected number in the sliding window. MTR is a useful tool for interpreting missense protein variants in the

Fig. 2. The R4 subfamily of RGS proteins: analysis of RGS4 human variants. (A) Clustal W was used to align sequences of the RGS domains within the R4 subfamily of human RGS proteins. The RGS4:Gai1-ALF₄ crystal structure (Protein Data Bank: 1AGR) was used to extract residues on RGS4 critical for interaction with Gai1 (highlighted in yellow). We then added pale yellow-filled boxes to extend this information to all R4 subfamily members. (B) MTR and PTM cluster analysis plot for RGS4-3 (UniProt ID P49798-3; see text for details). The RGS domain of RGS4 (highlighted in blue) is under selective pressure, indicated by a low MTR (median, red line; 25%, gray dashed line; 5%, green dashed line). RGS4 is palmitoylated on three residues, and although one is found to be mutated in the human population (GnomAD version 2.0), it is not located within the RGS domain and therefore not highlighted in this analysis (but see Supplemental Material; Supplemental Table 1) (C) Rare human variants of RGS4 plotted along the sequence using Lollipops (<https://github.com/pbnjay/loollipop>). Missense (amino acid change) variants are displayed on top in teal, whereas silent (no amino acid change) variants are displayed on the bottom in purple. (D) From our analyses, we selected nine variants in the RGS domain predicted to have a CoF phenotype and mapped them onto the RGS4 sequence using Lollipops (<https://github.com/pbnjay/loollipop>). (E) These selected variants are mapped onto the reported crystal structure (Protein Data Bank: 1AGR) of RGS4 (gray, selected residues in red) in complex with Gai1-ALF₄ (light blue).

context of monogenic diseases (Traynelis et al., 2017) and, although may be less predictive for proteins linked to polygenic diseases, is nonetheless reported in this work (for this dataset: minimum = 0.5, maximum = 1.19, median = 0.94) (Supplemental Material; Table 2).

As a further level of analysis, we consider sites of PTM found in human RGS proteins that, in many cases, overlap with rare variants that exhibit high C scores (Figs. 2B, 4B, 5B, and 6B). As outlined above, PTMs serve critical regulatory roles in protein function and are often overlooked or missed in disease assessment (Torres et al., 2016). A recent advance in the analysis of PTMs (Dewhurst et al., 2015; Torres et al., 2016; Dewhurst and Torres, 2017) offers a powerful bioinformatics/proteomics tool for characterizing experimentally verified PTMs in the context of their alignment within protein or domain families [i.e., modified alignment positions (MAPs)] (see Supplemental Information and Hunter et al. [2009], Edgar [2004], and Hornbeck et al. [2015]) and provides a functionally impactful analysis that is complementary to CADD and MTR data. This approach has shown further promise in the discovery of PTMs associated with disease-linked mutations (Torres et al., 2016). Therefore, we surveyed the coincidence of PTMs/MAPs and mutations within each RGS protein, with the goal of highlighting positions where CoF might be attributable to a change in

PTM status. For this purpose, PTMs unique to each human RGS protein are superimposed onto the MTR plots where the specific type of modification (Ph = phosphorylation, Ub = ubiquitination, Ac = acetylation, Pm = palmitoylation) is noted above the graph, and the median, 25%, and 5% MTR values are indicated by horizontal red, gray, and green lines, respectively, for representative family members (Figs. 2B, 4B, 5B, and 6B; Supplemental Material; Supplemental Table 1). In each case, only positions found to be modified in the given human RGS protein are shown (Figs. 2–6, red circles). Within the RGS domain of each protein, the size of each PTM site reflects the total number of PTMs observed within the domain MAP (see Supplemental Material), whereas positions outside the RGS domain simply indicate the position of a single PTM for the given protein. Each PTM site is further annotated to indicate when there is evidence of function for the PTM in the given protein (green outer ring) or evidence for function within the MAP for the domain family (red outer ring). Disease-linked variant positions are also indicated (orange circles). Although not included in our primary analysis, we also report neighboring PTM count (Table 2), which is a summation of reported PTMs found within a ± 7 residue window surrounding the human variant. Variants that fall within this window may in fact conflict with the ability of a modifying enzyme to dock on the target protein. We

therefore indicate in our analysis below when this occurs and propose this may be one mechanism for a CoF phenotype. A comprehensive table of each experimentally observed PTM for all human RGS proteins is also included in Supplemental Table 1. Notably, in each RGS protein case study we present, the RGS domain is both under mutational selective pressure and a hotbed for PTM activity, supporting the idea that disruption of these domains by mutation is likely to alter critical cellular functions.

Taking all of this information into account, we plot the genic distribution of both missense (top of Figs. 2C, 4C, 5C, and 6C) and silent (bottom of Figs. 2C, 4C, 5C, and 6C) variants for each RGS protein, along with a mapping of the selected rare variants of interest onto the protein sequence (Figs. 2D, 4D, 5D, and 6D). (Note: Human RGS protein exome sequences selected for analysis and presentation in this review are identified by UniProt IDs (Bairoch et al., 2005) listed in Supplemental Material and in the figure legends.) These selected variants, identified from the predictive criteria, are finally placed onto the reported crystal structure of the RGS protein (Figs. 2E, 4E, 5E, and 6E) and listed in Table 2. Importantly, whereas our focus and discussion are centered on the RGS domain of each protein subfamily, we present the comprehensive set of missense variants, MTR, CADD C score, and PTMs for the entire sequence of all 20 canonical RGS proteins (see Supplemental Material). Lastly, we validate this overall approach by testing one of the selected variants for RGS4 to show that this variant, as predicted, results in a profound CoF phenotype (Fig. 3). We suggest that this overall approach and dataset can be used and expanded to other individual RGS proteins across each family (and any protein family in general) to prioritize rare variants and PTMs for CoF studies and improved understanding of human pathophysiology.

A. R4 Family: RGS4 Rare Variants

Among the R4 subfamily of RGS proteins, RGS4 is perhaps the best characterized with abundant functional information, including the first solved RGS protein crystal structure (Tesmer et al., 1997). Therefore, RGS4 is well suited as a case study for analyzing and predicting CoF human variants within the R4 family. Consistent with this idea, a human mutation in RGS4, S30C, has previously been reported to display a GoF phenotype (enhanced GAP activity) that translated to RGS16 when an analogous mutation was made (Hill et al., 2008). This demonstrates the idea that insights gained from mutational analysis in one RGS protein may extend to other close family members, although in this study we emphasize variants unique to each individual RGS protein (Supplemental Material).

Human RGS4 has five reported splice variants with predicted sizes ranging from 93 to 304 amino acids (Ding et al., 2007). The RGS4 variants in the GnomAD

database (Lek et al., 2016) are reported with respect to the longest splice variant (RGS4-3; Uniprot ID: P49798-3), which adds almost 100 amino acids to the N terminus of the commonly used canonical human RGS4 reference sequence (RGS4-1; 205 amino acids; Uniprot ID: P49798-1) (Ding et al., 2007). Therefore, for our purposes in this work, we list first the reported variant amino acid location within the longer RGS4-3 sequence, followed by the canonical residue position in parentheses (Table 2). In our analysis of RGS4, we first compare an alignment of all R4 family members and identify residues that are contact points for $G\alpha$ highlighted in yellow, as determined by X-ray crystallography (Tesmer et al., 1997) (Fig. 2A). Conserved RGS: $G\alpha$ interface contact regions across R4 family members are highlighted by boxes. Next, we examine MTR for our case study protein, RGS4 (Fig. 2B), with reported PTMs unique to human RGS4 plotted onto the MTR data and identified above. From this analysis, it is clear that the RGS domain of RGS4 is under selective pressure, suggesting that human variants in this region are likely to change the function of the protein. Thus, we take advantage of the reported crystal structure for RGS4 in complex with $G\alpha i1$ (Tesmer et al., 1997), extracting information about contact points between the two proteins. From these structural, PTM, and CADD analyses, we identified nine variants, including the following: E214(117)K, D227(130)G, D260(163)N, D260(163)G, R264(167)C, L170(73)P, R231(134)W, R263(166)C, and R263(166)H (Fig. 2, C and D; Table 2). Of these, E214(117)K (C score 27.3), D227(130)G (C score 29.6), D260(163)N (C score 32), D260(163)G (C score 29.4), and R264(167)C (C score 35) are all highly or completely conserved residues among the R4 family (Fig. 2A) and participate in salt bridge interactions. Mutation of these residues to a noncharged [as with variants D227(130)G, D260(163)N, and D260(163)G], or opposite charge [as with E214(117)K] residue is predicted to decrease the stability of a RGS4: $G\alpha i1$ complex. L170(73)P (C score 23) is located within an α helix and is, therefore, predicted to disrupt secondary structure of the RGS domain. Amino acid R231(134) is another highly conserved residue within the family that participates in stabilizing switch III on the $G\alpha$ subunit, an important interaction for promoting GTP hydrolysis. As such, the R231(134)W (C score 29.8) variant very likely disrupts RGS4 capacity to interact with $G\alpha$ (see below). Finally, R263(166)C (C score 28.4) and R263(166)H (C score 24.5) are predicted to disrupt contact between the RGS domain and the all helical domain of $G\alpha$. In these cases, disruption of binding to $G\alpha$, or disruption of RGS4 capacity to stabilize switch regions on $G\alpha$, should similarly affect downstream G protein signaling events and disease states. Table 2 lists human variants in RGS4 that are predicted to disrupt RGS4 interaction with $G\alpha$, and/or GAP activity.

Further analysis of the RGS4 sequence using PTM alignment analysis identified three residues in human

Fig. 3. RGS4 human variant R231(134)W disrupts RGS domain interaction with activated *Gai1*. To validate our methods, we selected one variant from the list to generate a mutation and test binding between RGS4 (wild type or mut) and *Gai1*-AIF₄⁻. We chose R231(134)W (henceforth RW) and highlighted this residue on the structure of RGS4:*Gai1*-AIF₄⁻ in red (A). This arginine residue is positively charged and interacts with the carbonyl oxygen of glutamate 236 on *Gai1* in the switch III region. Disruption of this interaction by mutation to a tryptophan (uncharged) is predicted to reduce RGS4 binding to *Gai1*. (B) We generated this mutation and performed an immunoprecipitation of RGS4 (wild type or RW) and found that the mutation greatly reduced *Gai1*-AIF₄⁻ coimmunoprecipitation. (C) We next examined this interaction in live cells and found that, again, the RW mutant greatly reduced Net bioluminescence resonance energy transfer signal (Brown et al., 2015a) between RGS4 RW and *Gai1*-AIF₄⁻, compared with wild-type RGS4, indicating that the mutant is not interacting with *Gai1*-AIF₄⁻.

RGS4 that undergo PTM. C99 and C109, both located outside the RGS domain, are palmitoylated. Of note, C109 corresponds to reported human variant C109(12) R and is an important regulator of RGS4 subcellular localization and GAP activity (Bastin et al., 2012). Therefore, this human variant, although not within our focus in this study on the RGS domain, is predicted to disrupt the function of RGS4 and thus may mimic loss of RGS4, as is the case for multiple diseases described above. In addition, C192(95), located within the RGS domain of RGS4, is palmitoylated and serves an important role in regulating RGS4 membrane localization and its GAP activity (Tu et al., 1999). However, this cysteine residue does not correspond with a reported human variant from this dataset. Thus, we mapped our selected rare variants for RGS4 onto the sequence (Fig. 2D) and structure (Fig. 2E) of RGS4.

In all, we identified nine variants (corresponding to seven residues) within the RGS domain of RGS4 that are predicted to exhibit CoF phenotypes. To test this prediction and validate our approach, we selected one human variant in RGS4 from our list, R231(134)W, to test for a CoF phenotype. Compared with wild-type RGS4, R231(134)W (Fig. 3A) exhibited a loss of direct *Gai1*-AIF₄⁻ binding (Fig. 3B), as well as a dramatic decrease in *Gα*-YFP binding in live cells by bioluminescence resonance energy transfer analysis (Brown et al., 2015a), indicating a profound LoF phenotype (Fig. 3C). These data suggest that the capacity of the RGS domain of RGS4 to bind *Gα*, and thus serve as a GAP, is disrupted by this naturally-occurring human variant, and may therefore be a determinant in RGS4-linked diseases found in carriers. For example, multiple neurologic diseases, including schizophrenia and alcoholism, correlate with low RGS4 expression (Mirnics et al., 2001; Ho et al., 2010), a phenotype that would also be expected for a LoF variant in the RGS domain of RGS4. Therefore, one possible hypothesis is that LoF

variant R231(134)W is a risk factor for schizophrenia—a classification that could be difficult to make through unbiased genome-wide studies due to the extremely low prevalence of the variant in the population (0.0065% of the African population). Indeed, we propose that hypotheses generated through an integrated bioinformatics analysis such as this one could reveal several such cases wherein disease states can be more confidently linked to discrete changes in protein structure and function. Other LoF variants listed in Table 2 are predicted to share the same phenotype and outcome. Below, we take a similar approach to extend our analyses to other RGS subfamilies.

B. R7 Family: RGS9 Rare Variants

Unlike the R4 family, the R7 family of RGS proteins is all larger multidomain proteins (Fig. 1). Among the R7 subfamily of RGS proteins, RGS9 was chosen as a representative family member to analyze because of its solved protein crystal structure in complex with *Gα* and known RGS9 links to disease (Slep et al., 2001; Nishiguchi et al., 2004; Cheng et al., 2007; Hartong et al., 2007; Stockman et al., 2008; Michaelides et al., 2010). Structural insight for the RGS9 RGS domain cocrystallized in complex with a *Gai1/t* chimera reveals a mostly charged interface (Slep et al., 2001), similar to other RGS:*Gα* interfaces, and provides information about residues that are both highly conserved and crucial for interaction (highlighted in Fig. 4A). The RGS9 MTR plot (Fig. 4B) demonstrates that the RGS domain, particularly the C-terminal half of the domain, is under selective pressure. This information indicates that the latter half of the domain is predicted to be sensitive to missense variation.

We next examined whether reported human PTMs aligned with human variants (Fig. 4B). Although there are many PTMs outside of the RGS domain, two rare-variant phosphorylation sites were found within the

RGS domain, S304R and Y413C. Whereas Y413C has a CADD C score of 27.2, S304R has a CADD C score of 3.7, which did not meet our cutoff. We mapped the rare human variants to the sequence of RGS9 (Fig. 4C) and noticed a large degree of variation across the gene, suggesting a great deal of sequence diversity. Based on the PTM, CADD, and structural analyses, we selected six variants of interest (W299R, R364C, K400Q, R406C, R406H, and Y413C; Table 2) and mapped them onto the RGS9 sequence (Fig. 4D) and structure (Fig. 4E). Most variants map to the C-terminal half of the RGS domain (Fig. 4B), with the exception of W299R, which was found in the literature to underlie some cases of retinal dysfunction (Nishiguchi et al., 2004; Cheng et al., 2007; Hartong et al., 2007; Stockman et al., 2008; Michaelides et al., 2010). Based on the crystal structure of RGS9 (Slep et al., 2001), W299 participates in an electrostatic interaction with a nearby lysine (K311) to stabilize the tertiary structure of the RGS domain, and mutation to a positively charged arginine in the W299R variant (C score 29.1) could disrupt this tertiary structure. As noted above, we report human variants that fall within a ± 7 residue window of PTM. As such, W299R has one neighboring PTM, a phosphorylation on S304. Thus, one possibility is that the W299R variant leads to retinal dysfunction by disrupting phosphorylation of S304, a prospect that must be further explored. Y413C,

although modified itself, also falls within a neighboring PTM window, an acetylation on K419. R364C (C score 29.5) makes a water-mediated contact with Gai1/t and stabilizes the $\alpha 5$ - $\alpha 6$ loop, which undergoes the greatest conformational change when serving as a GAP for G α . K400Q (C score 28.1) is perfectly conserved within the R7 family and makes a contact with the α helical domain of G α , which is important for RGS:G α specificity (Skiba et al., 1999). Finally, R406C and R406H, also perfectly conserved within the family, form a cis salt bridge with a nearby aspartate (D402) on RGS9 to help stabilize G α switch I, and these variants have very high CADD C scores (35 and 34, respectively). R406C/H also falls within a neighboring PTM, that of Y413C.

Due to the known link between W299R and retinal dysfunction, we propose that these as-yet undefined variants listed in this study may have a similar role in retinal or other RGS9-linked diseases. Elsewhere within the CNS, RGS9-2 has a role in the striatum (Rahman et al., 1999) regulating dopamine signaling, which has consequences for addiction, Parkinson's disease, and schizophrenia (Rahman et al., 2003; Kovoov et al., 2005; Seeman et al., 2007). Notably, these phenotypes, such as enhanced sensitivity to stimulants, arise from a loss of RGS9-2 protein. Thus, LoF variants described in this dataset may contribute to susceptibility of these diseases in carriers. Furthermore, as the R7

Fig. 4. The R7 subfamily of RGS proteins: analysis of RGS9 human variants. (A) Clustal W was used to align sequences of the RGS domains within the R7 subfamily of human RGS proteins. The RGS9:Gai1/t- AlF_4^- crystal structure (Protein Data Bank: 1FQK) was used to extract residues on RGS9 critical for interaction with Gai1/t (highlighted in yellow). We then added pale yellow-filled boxes to extend this information to all R7 subfamily members. (B) MTR and PTM cluster analysis plot for RGS9 (UniProt ID O75916-1; see text for details). The RGS domain of RGS9 (highlighted in blue) is under selective pressure, indicated by a low MTR (median, red line; 25%, gray dashed line; 5%, green dashed line). Human RGS9 contains many reported post-translational modifications, including multiple acetylations and phosphorylations, one of which (phosphorylation) is found to be mutated in the RGS domain (GnomAD version 2.0) and met our criteria for inclusion (Supplemental Table 1; Table 2). (C) Rare human variants of RGS9 plotted along the sequence using Lollipops (<https://github.com/pbnjay/loollipop>). Missense (amino acid change) variants are displayed on top in teal, whereas silent (no amino acid change) variants are displayed on the bottom in purple. (D) From our analyses, six variants in the RGS domain of RGS9 were predicted to have a CoF phenotype and mapped onto the RGS9 sequence using Lollipops (<https://github.com/pbnjay/loollipop>). (E) Selected variants were mapped onto the crystal structure (Protein Data Bank: 1FQK) of RGS9 (gray, selected residues in red) and Gai1/t- AlF_4^- (light blue).

subfamily proteins contain additional signaling domains that are critical for protein stability and proper subcellular localization (Anderson et al., 2007, 2009), LoF variants in those domains could also cause LoF phenotypes that should be explored further. Table 2 lists human variants in the RGS domain of RGS9 that we predict disrupt RGS9 interaction with $G\alpha$ and/or GAP activity.

C. R12 Family: RGS10 Rare Variants

Among the R12 subfamily of RGS proteins, RGS10 is the smallest and simplest, and is the only family member that has been crystallized in complex with a $G\alpha$, $G\alpha i3$ (Soundararajan et al., 2008), providing an excellent case study for the R12 family. We first compared amino sequence conservation and highlighted contact points between RGS10 and $G\alpha i3$ (Fig. 5A). As with RGS4, human RGS10 has three splice variants that range in size from 167 to 181 amino acids (Rivero et al., 2010; Ali et al., 2013; Lee and Tansey, 2015). The RGS10 variants in the GnomAD database (Lek et al., 2016) are reported with respect to the longest splice variant (RGS10-3; UniProt ID O43665-3) and mapped onto this sequence in Fig. 5C, which adds only

eight amino acids to the N terminus of the commonly used canonical human RGS10 reference sequence (RGS10-1; Uniprot ID O43665-1; 173 amino acids).

Compared with other solved RGS domain- $G\alpha$ crystal structures (Tesmer et al., 1997; Slep et al., 2001), the RGS domain of RGS10 makes fewer contact points with $G\alpha$ (compare with highlighted regions in Fig. 4A; Fig. 2A), yet five variants were found in our predictive CADD/PTM/structural analysis: L46(38)P, V52(44)M, D141(133)N, R145(137)C, and K148(140)R (mapped onto the RGS10 sequence in Fig. 5D and RGS10 structure in Fig. 5E). The RGS10 MTR plot shows a high degree of selective pressure across the entire RGS domain (Fig. 5B). Furthermore, there are several reported human PTMs within the RGS domain, including K53 and K148 (ubiquitination), K78 (acetylation), and Y94 and Y143 (phosphorylation). Of these, only K148 is included in the GnomAD dataset as K148(140)R (Fig. 5B; Supplemental Material). Of note, K148(140)R has a CADD C score of 25, which met our criteria for a CoF candidate.

In addition to the highlighted variant residues above, D141(133)N, R145(137)C, and L46(38)P met our structural and sequence analysis criteria. Residues D141(133)N and R145(137)C both have very high

Fig. 5. The R12 subfamily of RGS proteins: analysis of RGS10 human variants. (A) Clustal W was used to align sequences of the RGS domains within the R12 subfamily of human RGS proteins. The RGS10: $G\alpha i3$ -ALF₄ crystal structure (Protein Data Bank: 2IHB) was used to extract residues on RGS10 critical for interaction with $G\alpha i3$ (highlighted in yellow). We then added pale yellow-filled boxes to extend this information to all R12 subfamily members. (B) MTR and PTM cluster analysis plot for RGS10-3 (UniProt ID O43665-3). The RGS domain of RGS10 (highlighted in blue) is under selective pressure, indicated by a low MTR (median, red line; 25%, gray dashed line; 5%, green dashed line). Human RGS10 contains many reported post-translational modifications, including multiple ubiquitinations, acetylations, and phosphorylations, one of which is found to be mutated in the RGS domain (GnomAD version 2.0) and met our criteria for predicted CoF phenotype (Supplemental Table 1; Table 2). (C) Rare human variants of RGS10 plotted along the sequence using Lollipop (https://github.com/pbnjay/lollipop). Missense (amino acid change) variants are displayed on top in teal, whereas silent (no amino acid change) variants are displayed on the bottom in purple. (D) From our analyses, we selected six variants in the RGS domain with predicted CoF phenotype and mapped them onto the sequence of RGS10 using Lollipop (https://github.com/pbnjay/lollipop). (E) Finally, we mapped these selected variants onto the crystal structure (Protein Data Bank: 2IHB) of RGS10 (gray, selected residues in red) and $G\alpha i3$ -ALF₄ (light blue).

CADD C scores of 35 and are close to two reported PTMs, a phosphorylation on Y143(135) and a ubiquitination on K148(140), indicating that these variants are likely to be deleterious to protein function. Consistent with this idea, both variants occur at highly conserved residues that are important for ionic stabilization. Changing the charge of the side chain from negative to uncharged, as is the case for D141(133)N and R145(137)C, is likely to disrupt efficient binding of the RGS domain to the $G\alpha$ protein. L46(38)P (C score 29.6) inserts a proline into an α helix, which we predict will cause disruption of the secondary structure. Furthermore, L46(38)P falls near two neighboring PTMs, a phosphorylation on S41(33) and a ubiquitination on K53(45). We postulate that disrupting secondary structure via insertion of a proline into an α helix would greatly inhibit the ability of a modifying enzyme to dock.

Finally, V52(44)M has a high CADD C score of 34 and is next to a modified residue, K53(45), suggesting that this mutation could lead to LoF. In support of this evidence, we observed that V52(44)M is putatively linked to schizophrenia (Hishimoto et al., 2004). V52(44) appears to participate in a hydrophobic core of the RGS domain; thus, a mutational change to methionine will likely disrupt this interaction. Interestingly, V52(44)M is found in nearly 2% of the East Asian population, and as such, we speculate that it could be an attractive target for deeper study. Furthermore, as loss of RGS10 protein in mice is linked with an increase in neuroinflammation (Lee et al., 2008, 2011, 2016; Lee and Tansey, 2015), LoF variants highlighted in this work may lead to susceptibility for neurodegenerative diseases such as Parkinson's disease. Table 2 lists human variants in RGS10 that are predicted to disrupt RGS10 interaction with $G\alpha$ and/or GAP activity.

D. RZ Family: RGS17 Rare Variants

Unlike the other RGS subfamilies, no member of the RZ subfamily (RGS17, RGS19, or RG20) has been crystalized in complex with a $G\alpha$ partner. However, a crystal structure of RGS17 alone has been reported (Soundararajan et al., 2008), allowing us to use this structure and conserved sequence alignments with other RGS family members to determine predicted CoF variants. RGS17 also has been linked to a number of diseases (Mao et al., 2004; James et al., 2009; You et al., 2009; Hayes and Roman, 2016), and therefore provides an attractive case study example for the RZ subfamily. For this, we first aligned the RGS domain of RGS17 with each of the case study proteins (RGS4, RGS9, RGS10; Figs. 2, 4, and 5) and highlight in yellow the residues that are indicated as contact points with $G\alpha$, based on their respective crystal structures (Supplemental Fig. 1). We noted a great deal of conservation for residues that participate in $G\alpha$ interaction. For example, residues 145–155 correspond to the $\alpha 5$ – $\alpha 6$ loop on each of the comparison family members (Tesmer et al., 1997;

Slep et al., 2001; Soundararajan et al., 2008), a region found within RGS domains that is critical for stabilization of switch II in $G\alpha$ and GAP activity. We then used this information to predict contact points between RGS17 and $G\alpha$, and highlighted in yellow those predicted contact points in Fig. 6A for all members of the RZ family. Each of those predicted contact points is either highly or completely conserved (highly conserved denoted by “:,” completely conserved denoted by “*,” yellow boxes). Again, MTR analysis for RGS17 (Fig. 6B) indicates that the RGS domain is under selective pressure, with the exception of residues 134–143, which immediately precede the highly conserved $\alpha 5$ – $\alpha 6$ loop. Using PTM alignment analysis, we identified two phosphorylation sites reported in the RGS domain of human RGS17, Y137 and Y171. Whether either phosphosite is functional for the protein is yet unknown. Although Y171 is a known variant site (Y171F), its low PTM observation frequency and CADD C score (9.1) exclude this variant from our selected dataset. Amino acid Y137 contributes to a MAP that is one of the most frequently observed phosphorylation sites in the RGS domain, but whose function is currently unknown. Thus, although not aligned with a genetic variant, PTM at this position appears to be very common in the RGS domain of several RGS proteins, a highly predictive feature for functional significance (Dewhurst et al., 2015; Torres et al., 2016; Dewhurst and Torres, 2017). Indeed, the lack of genetic variance at or near this position suggests that it is of critical importance, and any future variant at this position would be a prime target for functional analysis.

Based on our combined structural, conservation, CADD, and PTM analysis, three variants met our predictive criteria: E148G, P166L, and R189T (Fig. 6D). Amino acid E148 in RGS17 is a charged residue found in the highly conserved $\alpha 5$ – $\alpha 6$ loop that serves as a key $G\alpha$ contact point for each of the representative RGS (Fig. 6A; Supplemental Fig. 1). E148 in RGS17 is also 100% conserved within the RZ family. The charged residues analogous to E148 in other RGS protein crystal structures participate in ionic bonding with opposite charge residues to stabilize this $\alpha 5$ – $\alpha 6$ loop; thus, this negatively charged glutamate in RGS17 is a likely ionic binding partner serving the same role. As such, the E148G variant (C score 28.2) of RGS17 could disrupt RGS17 interactions with target $G\alpha$ subunits. We also identified P166L (C score 28.3), which is 100% conserved both within the RZ family (Fig. 6A) and across the other RGS subfamilies (Supplemental Fig. 1). This conserved proline appears to mediate the critical turn in the $\alpha 6$ – $\alpha 7$ loop that is present in all RGS domains. Furthermore, it is located near a neighboring PTM, a phosphorylation on Y171. Due to the drastically different side chain properties of leucine versus proline, we speculate that P166L could result in CoF. Residue R189T (C score 28.1), which was also identified as a potential CoF candidate, like P166, is 100% conserved within the RZ subfamily and across all other RGS

Fig. 6. The RZ subfamily of RGS proteins: analysis of RGS17 human variants. (A) Clustal W was used to align sequences of the RGS domains within the RZ subfamily of human RGS proteins. A crystal structure for RGS17 in complex with a $G\alpha$ is not yet available, although a crystal structure of RGS17 alone has been reported (Protein Data Bank: 1ZV4). We therefore compared critical residues across all RGS subfamilies and highlighted residues in RGS17 that are, based on conservation, predicted to be contact points for $G\alpha$ interaction. We then added pale yellow-filled boxes to extend this information to all RZ subfamily members. (B) MTR and PTM cluster analysis plot for RGS17 (UniProt ID Q9UGC6-1). The RGS domain of RGS17 (highlighted in blue) is under selective pressure, indicated by a low MTR (red line is median MTR for gene). RGS17 contains two phosphorylations, both within the RGS domain (Supplemental Table 1; Table 2). Although one is found to be mutated in humans (GnomAD version 2.0), it did not meet our criteria for inclusion in our CoF table, but nonetheless should be noted for study. (C) Rare human variants of RGS17 plotted along the sequence using Lollipop (https://github.com/pbnjay/lollipops). Missense (amino acid change) variants are displayed on top in teal, whereas silent (no amino acid change) variants are displayed on the bottom in purple. (D) From our analyses, we selected three variants in the RGS domain with predicted CoF phenotype and mapped them onto the RGS17 sequence using Lollipop (https://github.com/pbnjay/lollipops). (E) Finally, we structurally aligned the apo RGS17 structure (Protein Data Bank: 1ZV4) with the RGS4:Gai1- AlF_4^- structure (Protein Data Bank: 1AGR) and mapped these selected variants onto the model of RGS17 (gray, selected residues in red) and Gai1 (light blue).

subfamilies. We predict this variant could disrupt RGS17 ionic interactions with target $G\alpha$ subunits, as shown in a structural homology model based on the solved crystal structure of the RGS4:Gai1- AlF_4^- complex (Tesmer et al., 1997) (Fig. 6E).

IV. Pharmacological Impact of Human Variants in RGS Proteins

GPCRs have dominated the field of drug discovery for decades, and resulting new therapeutics have revolutionized modern medicine (Lundstrom, 2009). However, despite their success in ameliorating many diseases, drugs that target GPCRs often have off-target effects, making their therapeutic utility less than ideal. In more recent years, drug discovery efforts have turned to new ways of modulating GPCR signaling, including but not limited to such novel approaches as biased agonism (Bologna et al., 2017), positive and negative allosteric modulators (i.e., PAMs and NAMs) (Kenakin and Miller, 2010), and targeting the regulation of G protein signaling via RGS proteins (Sjögren et al., 2010). The rationale (Neubig and Siderovski, 2002) for targeting RGS proteins would be to do the following: 1) potentiate the effects of “dirty” GPCR agonists, thereby lowering

the requisite therapeutic dose and off-target actions, and 2) enhance specific tissue and/or receptor signaling output when confronted with reduced natural agonist (e.g., depression, neurodegenerative diseases, others). Currently, available small-molecule modulators that target RGS proteins do so by binding to and blocking RGS domain interactions with $G\alpha$ (Hayes et al., 2018). The emergence of personal genomics raises the realistic prospect of personalized medicine (Cardon and Harris, 2016). Understanding the specific drug and target mechanism of action, and how that is intertwined with genetic variation (pharmacogenetics), could lead to better predictions of therapeutic efficacy.

The nascent field of pharmacogenetics aims to develop new medications that are tailored to an individual’s genetic makeup. As we move closer to this new world, the benefits of understanding the effects of genetic variation on target protein GoF versus LoF would be of great value, as demonstrated by recent reports of the profound impact of human variants on prescription drug effects on GPCR (Hauser et al., 2018) and NMDAR actions (Ogden et al., 2017). If a RGS protein is implicated in a complex disease, particularly by enhancement of the RGS domain/ GAP activity, a RGS inhibitor may prove to be a superior target compared

with a drug that targets a GPCR upstream of the RGS. For example, RGS5 mediates angiogenesis, and, remarkably, vascularization of tumors is normalized in RGS5 knockout mice (Hamzah et al., 2008). Thus, normal or GoF variants of RGS5 would promote tumor angiogenesis, and, in this case, a selective targeted inhibitor of RGS5 may make immune-mediated destruction of solid tumors more effective, greatly improving survival. Similarly, if a GoF mutation in the RGS domain of RGS4, for instance (Hill et al., 2008), is found to underlie an individual's cardiovascular disease, a RGS inhibitor that selectively blocks RGS4 actions such as CCG-203769 (Blazer et al., 2015) may offer therapeutic benefit. By contrast, if a LoF in RGS4 or other RGS protein is found to contribute to disease progression, such a RGS inhibitor may exacerbate the disease and should thus be avoided. A prime example of the power of this analysis is a recent report on RGS2 human variants (Phan et al., 2017) that are linked to hypertension (Yang et al., 2005). The authors found that these RGS2 human variants, located in the N terminus, reduced RGS2-mediated inhibition of Ca^{2+} signaling via protein degradation and mislocalization and provided a mechanism for their association with hypertension. However, simply knowing a link between a variant and a disease does not account for functional directionality. Instead, one must determine whether the variant enhances or reduces protein function. Knowing this information becomes important in the case of RGS proteins because they regulate the timing of G protein on/off rates; thus, a drug that stimulates a GPCR may exacerbate the disease state in a patient with a LoF or GoF variant in a downstream regulatory RGS. Just as mutations in a metabolic enzyme dysregulate intended bioavailability or half-life of a drug, dysregulated RGS proteins may lead to extended or exaggerated drug actions. Therefore, investigators and clinicians will want to understand how natural genetic variation may affect druggable targets, both current and future.

Although we have focused this review on the effects of rare human variants within the canonical RGS domain, we must note that other domains and regions on RGS proteins are, in many ways, equally important for RGS protein function. Other RGS protein domains/regions dictate subcellular localization, modify GAP activity, stabilize the protein, and engage additional signaling pathways independent of G protein signaling. For example, RGS4 is robustly degraded by an N-terminal cysteine degradation signal (Bodenstein et al., 2007), and genetic modification (C→S) of this N-terminal cysteine increases RGS4 protein expression by nearly 50-fold. This could have major implications in diseases linked with RGS4 expression, such as schizophrenia and alcoholism (Mirnics et al., 2001; Ho et al., 2010). The subcellular localization of RGS14 is regulated by a C-terminal GPR motif (Shu et al., 2007; Brown et al., 2015b), which binds an inactive $\text{G}\alpha\text{-GDP}$. Ablation of

this interaction may have consequences for the role of RGS14 as a regulator of hippocampal-based learning and synaptic long-term potentiation (Lee et al., 2010). RGS7 family members all contain DEP domains (binds R7BP) that are important for plasma membrane localization (Drenan et al., 2006) and GGL domains (binds $\text{G}\beta 5$) that are essential for protein stability (Chen et al., 2003). Loss of interaction with either of these critical binding partners would disrupt the function of RGS7 completely independent of the RGS domain, and the DEP domain could, in theory, serve as a beneficial drug target. Indeed, information gained from LoF or GoF genetic variation throughout each of the functional domains of these heterogeneous RGS proteins may provide key insight into the etiology of disease and, should drug modulators become available, may offer the best course of treatment. The widespread accessibility of genome sequencing and parallel improvements in *in silico* predictions of protein functions brings precision medicine and pharmacogenomics much closer to reality. Within this context, modern drug development and usage should take into account whether, and how, natural genetic variation may affect patient responses to therapies targeting complex disease states.

Acknowledgments

We thank Joshua L. Traynelis, who kindly provided MTR analysis and data and constructive suggestions on how to improve the manuscript. We thank Suneela Ramineni for technical support and Kyle Gerber for motivation to work faster. We also thank Jeff Squires, who provided endless guidance on using Excel for complex data management.

Authorship Contributions

Contributed experiments: Squires, Montañez-Miranda, Hepler.

Contributed new reagents or analytic tools: Squires, Pandya, Torres.

Wrote or contributed to the writing of the manuscript: Squires, Montañez-Miranda, Pandya, Torres, Hepler.

References

- Adams LD, Geary RL, Li J, Rossini A, and Schwartz SM (2006) Expression profiling identifies smooth muscle cell diversity within human intima and plaque fibrous cap: loss of RGS5 distinguishes the cap. *Arterioscler Thromb Vasc Biol* **26**:319–325.
- Ahlers KE, Chakravarti B, and Fisher RA (2016) RGS6 as a novel therapeutic target in CNS diseases and cancer. *AAPS J* **18**:560–572.
- Ajit SK and Young KH (2005) Analysis of chimeric RGS proteins in yeast for the functional evaluation of protein domains and their potential use in drug target validation. *Cell Signal* **17**:817–825.
- Alexander GM, Farris S, Pirone JR, Zheng C, Colgin LL, and Dudek SM (2016) Social and novel contexts modify hippocampal CA2 representations of space. *Nat Commun* **7**:10300.
- Ali MW, Cacan E, Liu Y, Pierce JY, Creasman WT, Murph MM, Govindarajan R, Eblen ST, Greer SF, and Hooks SB (2013) Transcriptional suppression, DNA methylation, and histone deacetylation of the regulator of G-protein signaling 10 (RGS10) gene in ovarian cancer cells. *PLoS One* **8**:e60185.
- Alqinyah M and Hooks SB (2018) Regulating the regulators: epigenetic, transcriptional, and post-translational regulation of RGS proteins. *Cell Signal* **42**:77–87.
- Alshbool FZ, Karim ZA, Vemana HP, Conlon C, Lin OA, and Khasawneh FT (2015) The regulator of G-protein signaling 18 regulates platelet aggregation, hemostasis and thrombosis. *Biochem Biophys Res Commun* **462**:378–382.
- Altman MK, Alshamrani AA, Jia W, Nguyen HT, Fambrough JM, Tran SK, Patel MB, Hoseinzadeh P, Beedle AM, and Murph MM (2015) Suppression of the GTPase-activating protein RGS10 increases Rheb-GTP and mTOR signaling in ovarian cancer cells. *Cancer Lett* **369**:175–183.
- Amstadter AB, Koenen KC, Ruggiero KJ, Acierio R, Galea S, Kilpatrick DG, and Gelernter J (2009a) Variant in RGS2 moderates posttraumatic stress symptoms following potentially traumatic event exposure. *J Anxiety Disord* **23**:369–373.

- Amstadter AB, Koenen KC, Ruggiero KJ, Acierno R, Galea S, Kilpatrick DG, and Gelernter J (2009b) Variation in RGS2 is associated with suicidal ideation in an epidemiological study of adults exposed to the 2004 Florida hurricanes. *Arch Suicide Res* 13:349–357.
- Anderson GR, Posokhova E, and Martemyanov KA (2009) The R7 RGS protein family: multi-subunit regulators of neuronal G protein signaling. *Cell Biochem Biophys* 54:33–46.
- Anderson GR, Semenov A, Song JH, and Martemyanov KA (2007) The membrane anchor R7BP controls the proteolytic stability of the striatal specific RGS protein, RGS9-2. *J Biol Chem* 282:4772–4781.
- Armulik A, Abramsson A, and Betsholtz C (2005) Endothelial/pericyte interactions. *Circ Res* 97:512–523.
- Ashrafi A, Garcia P, Kollmus H, Schughart K, Del Sol A, Buttini M, and Glaab E (2017) Absence of regulator of G-protein signaling 4 does not protect against dopamine neuron dysfunction and injury in the mouse 6-hydroxydopamine lesion model of Parkinson's disease. *Neurobiol Aging* 58:30–33.
- Bairoch A, Apweiler R, Wu CH, Barker WC, Boeckmann B, Ferro S, Gasteiger E, Huang H, Lopez R, Magrane M, et al. (2005) The universal protein resource (UniProt). *Nucleic Acids Res* 33:D154–D159.
- Bansal G, DiVietro JA, Kuehn HS, Rao S, Nocka KH, Gilfillan AM, and Druey KM (2008a) RGS13 controls G protein-coupled receptor-evoked responses of human mast cells. *J Immunol* 181:7882–7890.
- Bansal G, Druey KM, and Xie Z (2007) R4 RGS proteins: regulation of G-protein signaling and beyond. *Pharmacol Ther* 116:473–495.
- Bansal G, Xie Z, Rao S, Nocka KH, and Druey KM (2008b) Suppression of immunoglobulin E-mediated allergic responses by regulator of G protein signaling 13. *Nat Immunol* 9:73–80.
- Barker SA, Wang J, Sierra DA, and Ross EM (2001) RGSZ1 and Ret RGS: two of several splice variants from the gene RGS20. *Genomics* 78:223–229.
- Bastin G, Singh K, Dissanayake K, Mighiu AS, Nurmoheamed A, and Heximer SP (2012) Amino-terminal cysteine residues differentially influence RGS4 protein plasma membrane targeting, intracellular trafficking, and function. *J Biol Chem* 287:28966–28974.
- Beadling C, Druey KM, Richter G, Kehrl JH, and Smith KA (1999) Regulators of G protein signaling exhibit distinct patterns of gene expression and target G protein specificity in human lymphocytes. *J Immunol* 162:2677–2682.
- Berger M, Bergers G, Arnold B, Hammerling GJ, and Ganss R (2005) Regulator of G-protein signaling-5 induction in pericytes coincides with active vessel remodeling during neovascularization. *Blood* 105:1094–1101.
- Bergers G and Song S (2005) The role of pericytes in blood-vessel formation and maintenance. *Neuro-oncol* 7:452–464.
- Berman DM, Wang Y, Liu Z, Dong Q, Burke LA, Liotta LA, Fisher R, and Wu X (2004) A functional polymorphism in RGS6 modulates the risk of bladder cancer. *Cancer Res* 64:6820–6826.
- Bernstein LS, Grillo AA, Loranger SS, and Linder ME (2000) RGS4 binds to membranes through an amphipathic alpha-helix. *J Biol Chem* 275:18520–18526.
- Bernstein LS, Ramineni S, Hague C, Cladman W, Chidiac P, Levey AI, and Hepler JR (2004) RGS2 binds directly and selectively to the M1 muscarinic acetylcholine receptor third intracellular loop to modulate Gq/11alpha signaling. *J Biol Chem* 279:21248–21256.
- Bifisha P, Yang J, Fisher RA, and Drouin J (2014) Rgs6 is required for adult maintenance of dopaminergic neurons in the ventral substantia nigra. *PLoS Genet* 10:e1004863.
- Blazer LL, Storaska AJ, Jutkiewicz EM, Turner EM, Calcagno M, Wade SM, Wang Q, Huang XP, Traynor JR, Husbands SM, et al. (2015) Selectivity and anti-Parkinson's potential of thiadiazolidinone RGS4 inhibitors. *ACS Chem Neurosci* 6:911–919.
- Blundell J, Hoang CV, Potts B, Gold SJ, and Powell CM (2008) Motor coordination deficits in mice lacking RGS9. *Brain Res* 1190:78–85.
- Bodenstein J, Sunahara RK, and Neubig RR (2007) N-terminal residues control proteasomal degradation of RGS2, RGS4, and RGS5 in human embryonic kidney 293 cells. *Mol Pharmacol* 71:1040–1050.
- Bodnar RJ (2004) Endogenous opioids and feeding behavior: a 30-year historical perspective. *Peptides* 25:697–725.
- Bologna Z, Teoh JP, Bayoumi AS, Tang Y, and Kim IM (2017) Biased G protein-coupled receptor signaling: new player in modulating physiology and pathology. *Biomol Ther (Seoul)* 25:12–25.
- Bomba L, Walter K, and Soranzo N (2017) The impact of rare and low-frequency genetic variants in common disease. *Genome Biol* 18:77.
- Bourne HR, Sanders DA, and McCormick F (1990) The GTPase superfamily: a conserved switch for diverse cell functions. *Nature* 348:125–132.
- Brown NE, Blumer JB, and Hepler JR (2015a) Bioluminescence resonance energy transfer to detect protein-protein interactions in live cells. *Methods Mol Biol* 1278:457–465.
- Brown NE, Goswami D, Branch MR, Ramineni S, Ortlund EA, Griffin PR, and Hepler JR (2015b) Integration of G protein α ($G\alpha$) signaling by the regulator of G protein signaling 14 (RGS14). *J Biol Chem* 290:9037–9049.
- Burgen PG, Lee WL, Nixon AB, Peralta EG, and Casey PJ (2001) Phosphorylation and nuclear translocation of a regulator of G protein signaling (RGS10). *J Biol Chem* 276:32828–32834.
- Cabrera-Vera TM, Hernandez S, Earls LR, Medkova M, Sundgren-Andersson AK, Surmeier DJ, and Hamm HE (2004) RGS9-2 modulates D2 dopamine receptor-mediated Ca^{2+} channel inhibition in rat striatal cholinergic interneurons. *Proc Natl Acad Sci USA* 101:16339–16344.
- Cacan E, Ali MW, Boyd NH, Hooks SB, and Greer SF (2014) Inhibition of HDAC1 and DNMT1 modulate RGS10 expression and decrease ovarian cancer chemoresistance. *PLoS One* 9:e87455.
- Campbell DB, Ebert PJ, Skelly T, Stroup TS, Lieberman J, Levitt P, and Sullivan PF (2008a) Ethnic stratification of the association of RGS4 variants with antipsychotic treatment response in schizophrenia. *Biol Psychiatry* 63:32–41.
- Campbell DB, Lange LA, Skelly T, Lieberman J, Levitt P, and Sullivan PF (2008b) Association of RGS2 and RGS5 variants with schizophrenia symptom severity. *Schizophr Res* 101:67–75.
- Cao Y, Pahlberg J, Sarria I, Kamasawa N, Sampath AP, and Martemyanov KA (2012) Regulators of G protein signaling RGS7 and RGS11 determine the onset of the light response in ON bipolar neurons. *Proc Natl Acad Sci USA* 109:7905–7910.
- Cardon LR and Harris T (2016) Precision medicine, genomics and drug discovery. *Hum Mol Genet* 25:R166–R172.
- Carper MB, Denvir J, Boskovic G, Primerano DA, and Claudio PP (2014) RGS16, a novel p53 and pRb cross-talk candidate inhibits migration and invasion of pancreatic cancer cells. *Genes Cancer* 5:420–435.
- Carreras J, Kikuti YY, Beà S, Miyaoka M, Hiraiwa S, Ikoma H, Nagao R, Tomita S, Martin-Garcia D, Salaverria I, et al. (2017) Clinicopathological characteristics and genomic profile of primary sinonasal tract diffuse large B cell lymphoma (DLBCL) reveals gain at 1q31 and RGS1 encoding protein; high RGS1 immunohistochemical expression associates with poor overall survival in DLBCL not otherwise specified (NOS). *Histopathology* 70:595–621.
- Celver J, Sharma M, and Kooroor A (2010) RGS9-2 mediates specific inhibition of agonist-induced internalization of D2-dopamine receptors. *J Neurochem* 114:739–749.
- Chan RK and Otte CA (1982) Isolation and genetic analysis of *Saccharomyces cerevisiae* mutants supersensitive to G1 arrest by a factor and alpha factor pheromones. *Mol Cell Biol* 2:11–20.
- Chang YP, Liu X, Kim JD, Ikeda MA, Layton MR, Weder AB, Cooper RS, Kardias SL, Rao DC, Hunt SC, et al. (2007) Multiple genes for essential-hypertension susceptibility on chromosome 1q. *Am J Hum Genet* 80:253–264.
- Chatterjee TK and Fisher RA (2000) Novel alternative splicing and nuclear localization of human RGS12 gene products. *J Biol Chem* 275:29660–29671.
- Chatterjee TK and Fisher RA (2002) RGS12TS-S localizes at nuclear matrix-associated subnuclear structures and represses transcription: structural requirements for subnuclear targeting and transcriptional repression. *Mol Cell Biol* 22:4334–4345.
- Chen CK, Eversole-Cire P, Zhang H, Mancino V, Chen YJ, He W, Wensel TG, and Simon MI (2003) Instability of GGL domain-containing RGS proteins in mice lacking the G protein beta-subunit Gbeta5. *Proc Natl Acad Sci USA* 100:6604–6609.
- Chen CK, Wieland T, and Simon MI (1996) RGS-r, a retinal specific RGS protein, binds an intermediate conformation of transducin and enhances recycling. *Proc Natl Acad Sci USA* 93:12885–12889.
- Chen IS, Furutani K, Inanobe A, and Kurachi Y (2014) RGS4 regulates partial agonism of the M2 muscarinic receptor-activated K^{+} currents. *J Physiol* 592:1237–1248.
- Chen X, Dunham C, Kendler S, Wang X, O'Neill FA, Walsh D, and Kendler KS (2004) Regulator of G-protein signaling 4 (RGS4) gene is associated with schizophrenia in Irish high density families. *Am J Med Genet B Neuropsychiatr Genet* 129B:23–26.
- Chen Z, Wu Y, Meng Q, and Xia Z (2016) Elevated microRNA-25 inhibits cell apoptosis in lung cancer by targeting RGS3. *In Vitro Cell Dev Biol Anim* 52:62–67.
- Cheng JY, Luu CD, Yong VH, Mathur R, Aung T, and Vithana EN (2007) Bradyopsia in an Asian man. *Arch Ophthalmol* 125:1138–1140.
- Cheng WL, Wang PX, Wang T, Zhang Y, Du C, Li H, and Ji Y (2015) Regulator of G-protein signalling 5 protects against atherosclerosis in apolipoprotein E-deficient mice. *Br J Pharmacol* 172:5676–5689.
- Cho H, Kim DU, and Kehrl JH (2005) RGS14 is a centrosomal and nuclear cytoplasmic shuttling protein that traffics to promyelocytic leukemia nuclear bodies following heat shock. *J Biol Chem* 280:805–814.
- Cho H, Kozasa T, Bondjers C, Betsholtz C, and Kehrl JH (2003) Pericyte-specific expression of Rgs5: implications for PDGF and EDG receptor signaling during vascular maturation. *FASEB J* 17:440–442.
- Cho H, Park C, Hwang IY, Han SB, Schimel D, Despres D, and Kehrl JH (2008) Rgs5 targeting leads to chronic low blood pressure and a lean body habitus. *Mol Cell Biol* 28:2590–2597.
- Chograni M, Alkuraya FS, Maazoul F, Lariani I, and Chaabouni-Bouhamed H (2014) RGS6: a novel gene associated with congenital cataract, mental retardation, and microcephaly in a Tunisian family. *Invest Ophthalmol Vis Sci* 55:1261–1266.
- Chowdari KV, Bamne M, Wood J, Talkowski ME, Mirnics K, Levitt P, Lewis DA, and Nimgaonkar VL (2008) Linkage disequilibrium patterns and functional analysis of RGS4 polymorphisms in relation to schizophrenia. *Schizophr Bull* 34:118–126.
- Chowdari KV, Mirnics K, Semwal P, Wood J, Lawrence E, Bhatia T, Deshpande SN, B K T, Ferrell RE, Middleton FA, et al. (2002) Association and linkage analyses of RGS4 polymorphisms in schizophrenia [published correction appears in *Hum Mol Genet* (2003) 12:1781]. *Hum Mol Genet* 11:1373–1380.
- Cohen SP, Buckley BK, Kosloff M, Garland AL, Bosch DE, Cheng G Jr, Radhakrishna H, Brown MD, Willard FS, Arshavsky VY, et al. (2012) Regulator of G-protein signaling-21 (RGS21) is an inhibitor of bitter gustatory signaling found in lingual and airway epithelia. *J Biol Chem* 287:41706–41719.
- Crowe RR, Goedken R, Samuelson S, Wilson R, Nelson J, and Noyes R Jr (2001) Genomewide survey of panic disorder. *Am J Med Genet* 105:105–109.
- Cui H, Nishiguchi N, Ivleva E, Yanagi M, Fukutake M, Nushida H, Ueno Y, Kitamura N, Maeda K, and Shirakawa O (2008) Association of RGS2 gene polymorphisms with suicide and increased RGS2 immunoreactivity in the postmortem brain of suicide victims. *Neuropsychopharmacology* 33:1537–1544.
- Dai J, Gu J, Lu C, Lin J, Stewart D, Chang D, Roth JA, and Wu X (2011) Genetic variations in the regulator of G-protein signaling genes are associated with survival in late-stage non-small cell lung cancer. *PLoS One* 6:e21120.
- Daniel JM, Prock A, Dutzmann J, Sonnenschein K, Thum T, Bauersachs J, and Sedding DG (2016) Regulator of G-protein signaling 5 prevents smooth muscle cell proliferation and attenuates neointima formation. *Arterioscler Thromb Vasc Biol* 36:317–327.

- Davydov IV and Varshavsky A (2000) RGS4 is arginylated and degraded by the N-end rule pathway in vitro. *J Biol Chem* **275**:22931–22941.
- Delesque-Touchard N, Pendaries C, Volle-Challier C, Millet L, Salel V, Hervé C, Plieger AM, Berthou-Soulie L, Prades C, Sorg T, et al. (2014) Regulator of G-protein signaling 18 controls both platelet generation and function. *PLoS One* **9**: e113215.
- De Vries L, Elenko E, Hubler L, Jones TL, and Farquhar MG (1996) GAIP is membrane-anchored by palmitoylation and interacts with the activated (GTP-bound) form of G alpha i subunits. *Proc Natl Acad Sci USA* **93**:15203–15208.
- De Vries L, Mousli M, Wurmser A, and Farquhar MG (1995) GAIP, a protein that specifically interacts with the trimeric G protein G alpha i3, is a member of a protein family with a highly conserved core domain. *Proc Natl Acad Sci USA* **92**: 11916–11920.
- Dewhurst HM, Choudhury S, and Torres MP (2015) Structural analysis of PTM hotspots (SAPH-ire)—a quantitative informatics method enabling the discovery of novel regulatory elements in protein families. *Mol Cell Proteomics* **14**:2285–2297.
- Dewhurst HM and Torres MP (2017) Systematic analysis of non-structural protein features for the prediction of PTM function potential by artificial neural networks. *PLoS One* **12**:e0172572.
- Ding J, Guzman JN, Tkatch T, Chen S, Goldberg JA, Ebert PJ, Levitt P, Wilson CJ, Hamm HE, and Surmeier DJ (2006) RGS4-dependent attenuation of M4 autoreceptor function in striatal cholinergic interneurons following dopamine depletion. *Nat Neurosci* **9**:832–842.
- Ding L and Hegde AN (2009) Expression of RGS4 splice variants in dorsolateral prefrontal cortex of schizophrenic and bipolar disorder patients. *Biol Psychiatry* **65**: 541–545.
- Ding L, Mychaleckyj JC, and Hegde AN (2007) Full length cloning and expression analysis of splice variants of regulator of G-protein signaling RGS4 in human and murine brain. *Gene* **401**:46–60.
- Dohlman HG, Apianeski D, Chen Y, Song J, and Nusskern D (1995) Inhibition of G-protein signaling by dominant gain-of-function mutations in Sst2p, a pheromone desensitization factor in *Saccharomyces cerevisiae*. *Mol Cell Biol* **15**:3635–3643.
- Dohlman HG, Song J, Ma D, Courchesne WE, and Thorne J (1996) Sst2, a negative regulator of pheromone signaling in the yeast *Saccharomyces cerevisiae*: expression, localization, and genetic interaction and physical association with Gpa1 (the G-protein alpha subunit). *Mol Cell Biol* **16**:5194–5209.
- Doi M, Ishida A, Miyake A, Sato M, Komatsu R, Yamazaki F, Kimura I, Tsuchiya S, Kori H, Seo K, et al. (2011) Circadian regulation of intracellular G-protein signalling mediates intercellular synchrony and rhythmicity in the suprachiasmatic nucleus. *Nat Commun* **2**:327.
- Doupnik CA, Xu T, and Shinaman JM (2001) Profile of RGS expression in single rat atrial myocytes. *Biochim Biophys Acta* **1522**:97–107.
- Drenan RM, Doupnik CA, Jayaraman M, Buchwalter AL, Kaltenbronn KM, Huettner JE, Linder ME, and Blumer KJ (2006) R7BP augments the function of RGS7*Gbeta5 complexes by a plasma membrane-targeting mechanism. *J Biol Chem* **281**:28222–28231.
- Druey KM, Blumer KJ, Kang VH, and Kehrl JH (1996) Inhibition of G-protein-mediated MAP kinase activation by a new mammalian gene family. *Nature* **379**: 742–746.
- Dudek SM, Alexander GM, and Farris S (2016) Rediscovering area CA2: unique properties and functions. *Nat Rev Neurosci* **17**:89–102.
- Dunn HA and Ferguson SS (2015) PDZ protein regulation of G protein-coupled receptor trafficking and signaling pathways. *Mol Pharmacol* **88**:624–639.
- Edgar RC (2004) MUSCLE: a multiple sequence alignment method with reduced time and space complexity. *BMC Bioinformatics* **5**:113.
- Emilsson L, Sætre P, and Jazin E (2006) Low mRNA levels of RGS4 splice variants in Alzheimer's disease: association between a rare haplotype and decreased mRNA expression. *Synapse* **59**:173–176.
- Estes JD, Thacker TC, Hampton DL, Kell SA, Keele BF, Palenski EA, Druey KM, and Burton GF (2004) Follicular dendritic cell regulation of CXCR4-mediated germinal center CD4 T cell migration. *J Immunol* **173**:6169–6178.
- Evans PR, Dudek SM, and Hepler JR (2015) Regulator of G protein signaling 14: a molecular brake on synaptic plasticity linked to learning and memory. *Prog Mol Biol Transl Sci* **133**:169–206.
- Felkin LE, Lara-Pezzi EA, Hall JL, Birks EJ, and Barton PJ (2011) Reverse remodelling and recovery from heart failure are associated with complex patterns of gene expression. *J Cardiovasc Transl Res* **4**:321–331.
- Fu W, O'Connor TD, Jun G, Kang HM, Abecasis G, Leal SM, Gabriel S, Rieder MJ, Altshuler D, Shendure J, et al.; NHLBI Exome Sequencing Project (2013) Analysis of 6,515 exomes reveals the recent origin of most human protein-coding variants [published correction appears in *Nature* (2013) 495:270]. *Nature* **493**: 216–220.
- Gagnon AW, Murray DL, and Leadley RJ (2002) Cloning and characterization of a novel regulator of G protein signalling in human platelets. *Cell Signal* **14**:595–606.
- Gainetdinov RR (2007) Mesolimbic dopamine in obesity and diabetes. *Am J Physiol Regul Integr Comp Physiol* **293**:R601–R602.
- Ganess R (2015) Keeping the balance right: regulator of G protein signaling 5 in vascular physiology and pathology. *Prog Mol Biol Transl Sci* **133**:93–121.
- Garzón J, Rodríguez-Muñoz M, López-Fando A, García-España A, and Sánchez-Blázquez P (2004) RGSZ1 and GAIP regulate mu- but not delta-opioid receptors in mouse CNS: role in tachyphylaxis and acute tolerance. *Neuropsychopharmacology* **29**:1091–1104.
- Gegenbauer K, Elia G, Blanco-Fernandez A, and Smolenski A (2012) Regulator of G-protein signaling 18 integrates activating and inhibitory signaling in platelets. *Blood* **119**:3799–3807.
- Gelernter J, Bonvicini K, Page G, Woods SW, Goddard AW, Kruger S, Pauls DL, and Goodson S (2001) Linkage genome scan for loci predisposing to panic disorder or agoraphobia. *Am J Med Genet* **105**:548–557.
- Gerber KJ, Squires KE, and Hepler JR (2016) Roles for regulator of G protein signaling proteins in synaptic signaling and plasticity. *Mol Pharmacol* **89**:273–286.
- Gibbons DL, Abeler-Dörner L, Raine T, Hwang IY, Jandke A, Wencker M, Deban L, Rudd CE, Irving PM, Kehrl JH, et al. (2011) Cutting edge: regulator of G protein signaling-1 selectively regulates gut T cell trafficking and colitic potential. *J Immunol* **187**:2067–2071.
- Gilman AG (1987) G proteins: transducers of receptor-generated signals. *Annu Rev Biochem* **56**:615–649.
- Glick JL, Meigs TE, Miron A, and Casey PJ (1998) RGSZ1, a Gz-selective regulator of G protein signaling whose action is sensitive to the phosphorylation state of Galpha. *J Biol Chem* **273**:26008–26013.
- Gold SJ, Heifets BD, Pudiak CM, Potts BW, and Nestler EJ (2002) Regulation of regulators of G protein signaling mRNA expression in rat brain by acute and chronic electroconvulsive seizures. *J Neurochem* **82**:828–838.
- Gold SJ, Ni YG, Dohlman HG, and Nestler EJ (1997) Regulators of G-protein signaling (RGS) proteins: region-specific expression of nine subtypes in rat brain. *J Neurosci* **17**:8024–8037.
- Goto K, Doi M, Wang T, Kunisue S, Murai I, and Okamura H (2017) G-protein-coupled receptor signaling through Gpr176, Gz, and RGS16 tunes time in the center of the circadian clock [Review]. *Endocr J* **64**:571–579.
- Grafstein-Dunn E, Young KH, Cockett MI, and Khawaja XZ (2001) Regional distribution of regulators of G-protein signaling (RGS) 1, 2, 13, 14, 16, and GAIP messenger ribonucleic acids by in situ hybridization in rat brain. *Brain Res Mol Brain Res* **88**:113–123.
- Grayson TH, Ohms SJ, Brackenbury TD, Meaney KR, Peng K, Pittelkow YE, Wilson SR, Sandow SL, and Hill CE (2007) Vascular microarray profiling in two models of hypertension identifies caveolin-1, Rgs2 and Rgs5 as antihypertensive targets. *BMC Genomics* **8**:404.
- Gu S, He J, Ho WT, Ramineni S, Thal DM, Natesh R, Tesmer JJ, Hepler JR, and Heximer SP (2007) Unique hydrophobic extension of the RGS2 amphipathic helix domain imparts increased plasma membrane binding and function relative to other RGS R4/B subfamily members. *J Biol Chem* **282**:33064–33075.
- Guo S, Tang W, Shi Y, Huang K, Xi Z, Xu Y, Feng G, and He L (2006) RGS4 polymorphisms and risk of schizophrenia: an association study in Han Chinese plus meta-analysis. *Neurosci Lett* **406**:122–127.
- Häggmark A, Mikus M, Mohsenian A, Hong MG, Forsström B, Gajewska B, Barančzyk-Kuźma A, Uhlén M, Schwenk JM, Kuźma-Kozakiewicz M, et al. (2014) Plasma profiling reveals three proteins associated to amyotrophic lateral sclerosis. *Ann Clin Transl Neurol* **1**:544–553.
- Haller C, Fillatreau S, Hoffmann R, and Agenes F (2002) Structure, chromosomal localization and expression of the mouse regulator of G-protein signaling10 gene (mRGS10). *Gene* **297**:39–49.
- Hamm HE (1998) The many faces of G protein signaling. *J Biol Chem* **273**:669–672.
- Hamzah J, Jugold M, Kiessling F, Rigby P, Manzur M, Marti HH, Rabie T, Kaden S, Gröne HJ, Hämmelring GJ, et al. (2008) Vascular normalization in Rgs5-deficient tumours promotes immune destruction. *Nature* **453**:410–414.
- Han JJ, Huang NN, Kim DU, and Kehrl JH (2006) RGS1 and RGS13 mRNA silencing in a human B lymphoma line enhances responsiveness to chemoattractants and impairs desensitization. *J Leukoc Biol* **79**:1357–1368.
- Hartong DT, Pott JW, and Kooijman AC (2007) Six patients with bradyopsia (slow vision): clinical features and course of the disease. *Ophthalmology* **114**:2323–2331.
- Hauser AS, Chavali S, Masuho I, Jahn LJ, Martemyanov KA, Gloriam DE, and Babu MM (2018) Pharmacogenomics of GPCR drug targets. *Cell* **172**:41–54.e19.
- Hayasaka N, Aoki K, Kinoshita S, Yamaguchi S, Wakefield JK, Tsuji-Kawahara S, Horikawa K, Ikegami H, Wakana S, Murakami T, et al. (2011) Attenuated food anticipatory activity and abnormal circadian locomotor rhythms in Rgs16 knock-down mice. *PLoS One* **6**:e17655.
- Hayes MP, Bodle CR, and Roman DL (2018) Evaluation of the selectivity and cysteine dependence of inhibitors across the regulator of G protein-signaling family. *Mol Pharmacol* **93**:25–35.
- Hayes MP and Roman DL (2016) Regulator of G protein signaling 17 as a negative modulator of GPCR signaling in multiple human cancers. *AAPS J* **18**:550–559.
- He W, Cowan CW, and Wensel TG (1998) RGS9, a GTPase accelerator for phototransduction. *Neuron* **20**:95–102.
- He W, Lu L, Zhang X, El-Hodiri HM, Chen CK, Slep KC, Simon MI, Jamrich M, and Wensel TG (2000) Modules in the photoreceptor RGS9-1.Gbeta 5L GTPase-accelerating protein complex control effector coupling, GTPase acceleration, protein folding, and stability. *J Biol Chem* **275**:37093–37100.
- Hensch NR, Karim ZA, Druey KM, Tansey MG, and Khasawneh FT (2016) RGS10 negatively regulates platelet activation and thrombogenesis. *PLoS One* **11**: e0165984.
- Hepler JR, Berman DM, Gilman AG, and Kozasa T (1997) RGS4 and GAIP are GTPase-activating proteins for Gq alpha and block activation of phospholipase C beta by gamma-thio-GTP-Gq alpha. *Proc Natl Acad Sci USA* **94**:428–432.
- Hepler JR and Gilman AG (1992) G proteins. *Trends Biochem Sci* **17**:383–387.
- Heximer SP (2004) RGS2-mediated regulation of Gqalpha. *Methods Enzymol* **390**: 65–82.
- Heximer SP, Knutsen RH, Sun X, Kaltenbronn KM, Rhee MH, Peng N, Oliveira-dos-Santos A, Penninger JM, Muslin AJ, Steinberg TH, et al. (2003) Hypertension and prolonged vasoconstrictor signaling in RGS2-deficient mice. *J Clin Invest* **111**: 445–452.
- Heximer SP, Lim H, Bernard JL, and Blumer KJ (2001) Mechanisms governing subcellular localization and function of human RGS2. *J Biol Chem* **276**: 14195–14203.
- Heximer SP, Srinivasa SP, Bernstein LS, Bernard JL, Linder ME, Hepler JR, and Blumer KJ (1999) G protein selectivity is a determinant of RGS2 function. *J Biol Chem* **274**:34253–34259.
- Heximer SP, Watson N, Linder ME, Blumer KJ, and Hepler JR (1997) RGS2/G0S8 is a selective inhibitor of Gqalpha function. *Proc Natl Acad Sci USA* **94**:14389–14393.
- Hill C, Brownlie Z, Davey J, Milligan G, and Ladds G (2008) Isolation and characterization of a novel human RGS mutant displaying gain-of-function activity. *Cell Signal* **20**:323–336.

- Hiol A, Davey PC, Osterhout JL, Waheed AA, Fischer ER, Chen CK, Milligan G, Druey KM, and Jones TL (2003) Palmitoylation regulates regulators of G-protein signaling (RGS) 16 function. I. Mutation of amino-terminal cysteine residues on RGS16 prevents its targeting to lipid rafts and palmitoylation of an internal cysteine residue. *J Biol Chem* **278**:19301–19308.
- Hishimoto A, Shirakawa O, Nishiguchi N, Aoyama S, Ono H, Hashimoto T, and Maeda K (2004) Novel missense polymorphism in the regulator of G-protein signaling 10 gene: analysis of association with schizophrenia. *Psychiatry Clin Neurosci* **58**:579–581.
- Hitti FL and Siegelbaum SA (2014) The hippocampal CA2 region is essential for social memory. *Nature* **508**:88–92.
- Ho AM, MacKay RK, Dodd PR, and Lewohl JM (2010) Association of polymorphisms in RGS4 and expression of RGS transcripts in the brains of human alcoholics. *Brain Res* **1340**:1–9.
- Hohoff C, Neumann A, Domschke K, Jacob C, Maier W, Fritze J, Bandelow B, Krakowitzky P, Rothermundt M, Arolt V, et al. (2009) Association analysis of Rgs7 variants with panic disorder. *J Neural Transm (Vienna)* **116**:1523–1528.
- Hohoff C, Weber H, Richter J, Domschke K, Zwanzger PM, Ohrmann P, Bauer J, Suslow T, Kugel H, Baumann C, et al. (2015) RGS2 genetic variation: association analysis with panic disorder and dimensional as well as intermediate phenotypes of anxiety. *Am J Med Genet B Neuropsychiatr Genet* **168B**:211–222.
- Hollinger S and Hepler JR (2002) Cellular regulation of RGS proteins: modulators and integrators of G protein signaling. *Pharmacol Rev* **54**:527–559.
- Hollinger S, Taylor JB, Goldman EH, and Hepler JR (2001) RGS14 is a bifunctional regulator of Galphai/o activity that exists in multiple populations in brain. *J Neurochem* **79**:941–949.
- Holobovskyy V, Manzur M, Tare M, Burchell J, Bolitho E, Viola H, Hool LC, Arnolda LF, McKittrick DJ, and Ganss R (2013) Regulator of G-protein signaling 5 controls blood pressure homeostasis and vessel wall remodeling. *Circ Res* **112**:781–791.
- Hong JX, Wilson GL, Fox CH, and Kehrl JH (1993) Isolation and characterization of a novel B cell activation gene. *J Immunol* **150**:3895–3904.
- Hooks SB, Callihan P, Altman MK, Hurst JH, Ali MW, and Murph MM (2010) Regulators of G-protein signaling RGS10 and RGS17 regulate chemoresistance in ovarian cancer cells. *Mol Cancer* **9**:289.
- Hooks SB, Martemyanov K, and Zachariou V (2008) A role of RGS proteins in drug addiction. *Biochem Pharmacol* **75**:76–84.
- Hooks SB and Murph MM (2015) Cellular deficiency in the RGS10 protein facilitates chemoresistant ovarian cancer. *Future Med Chem* **7**:1483–1489.
- Hooks SB, Waldo GL, Corbitt J, Bodor ET, Krumins AM, and Harden TK (2003) RGS6, RGS7, RGS9, and RGS11 stimulate GTPase activity of Gi family G-proteins with differential selectivity and maximal activity. *J Biol Chem* **278**:10087–10093.
- Hornbeck PV, Kornhauser JM, Tkachev S, Zhang B, Skrzypek E, Murray B, Latham V, and Sullivan M (2012) PhosphoSitePlus: a comprehensive resource for investigating the structure and function of experimentally determined post-translational modifications in man and mouse. *Nucleic Acids Res* **40**:D261–D270.
- Hornbeck PV, Zhang B, Murray B, Kornhauser JM, Latham V, and Skrzypek E (2015) PhosphoSitePlus, 2014: mutations, PTMs and recalibrations. *Nucleic Acids Res* **43**:D512–D520.
- Hu Y, Shmygelska A, Tran D, Eriksson N, Tung JY, and Hinds DA (2016) GWAS of 89,283 individuals identifies genetic variants associated with self-reporting of being a morning person. *Nat Commun* **7**:10448.
- Huang C, Hepler JR, Gilman AG, and Mumby SM (1997) Attenuation of Gi- and Gq-mediated signaling by expression of RGS4 or GAI1 in mammalian cells. *Proc Natl Acad Sci USA* **94**:6159–6163.
- Huang J, Chen L, Yao Y, Tang C, Ding J, Fu C, Li H, and Ma G (2016a) Pivotal role of regulator of G-protein signaling 12 in pathological cardiac hypertrophy. *Hypertension* **67**:1228–1236.
- Huang KY, Su MG, Kao HJ, Hsieh YC, Jhong JH, Cheng KH, Huang HD, and Lee TY (2016b) dbPTM 2016: 10-year anniversary of a resource for post-translational modification of proteins. *Nucleic Acids Res* **44**:D435–D446.
- Hunt KA, Zhernakova A, Turner G, Heap GA, Franke L, Bruinenberg M, Romanos J, Dinesen LC, Ryan AW, Panesar D, et al. (2008) Newly identified genetic risk variants for celiac disease related to the immune response. *Nat Genet* **40**:395–402.
- Hunter S, Appweiler R, Attwood TK, Bairoch A, Bateman A, Binns D, Bork P, Das U, Daugherty L, Duquenne L, et al. (2009) InterPro: the integrative protein signature database. *Nucleic Acids Res* **37**:D211–D215.
- Hwang IY, Hwang KS, Park C, Harrison KA, and Kehrl JH (2013) Rgs13 constrains early B cell responses and limits germinal center sizes. *PLoS One* **8**:e60139.
- Hwang IY, Park C, Harrison KA, Huang NN, and Kehrl JH (2010) Variations in Gnai2 and Rgs1 expression affect chemokine receptor signaling and the organization of secondary lymphoid organs. *Genes Immun* **11**:384–396.
- Igci M, Baysan M, Yigiter R, Ulasli M, Geyik S, Bayraktar R, Bozgeyik I, Bozgeyik E, Bayram A, and Cakmak EA (2016) Gene expression profiles of autophagy-related genes in multiple sclerosis. *Gene* **588**:38–46.
- Ingi T and Aoki Y (2002) Expression of RGS2, RGS4 and RGS7 in the developing postnatal brain. *Eur J Neurosci* **15**:929–936.
- Ingi T, Krumins AM, Chidiac P, Brothers GM, Chung S, Snow BE, Barnes CA, Lanahan AA, Siderovski DP, Ross EM, et al. (1998) Dynamic regulation of RGS2 suggests a novel mechanism in G-protein signaling and neuronal plasticity. *J Neurosci* **18**:7178–7188.
- Itoh M, Odagiri M, Abe H, and Saitoh O (2001) RGS8 protein is distributed in dendrites and cell body of cerebellar Purkinje cell. *Biochem Biophys Res Commun* **287**:223–228.
- Jaba IM, Zhuang ZW, Li N, Jiang Y, Martin KA, Sinusas AJ, Papademetris X, Simons M, Sessa WC, Young LH, et al. (2013) NO triggers RGS4 degradation to coordinate angiogenesis and cardiomyocyte growth. *J Clin Invest* **123**:1718–1731.
- James MA, Lu Y, Liu Y, Vikis HG, and You M (2009) RGS17, an overexpressed gene in human lung and prostate cancer, induces tumor cell proliferation through the cyclic AMP-PKA-CREB pathway. *Cancer Res* **69**:2108–2116.
- Jensen LJ, Gupta R, Blom N, Devos D, Tamames J, Kesmir C, Nielsen H, Staerfeldt HH, Rapacki K, Workman C, et al. (2002) Prediction of human protein function from post-translational modifications and localization features. *J Mol Biol* **319**:1257–1265.
- Ji YR, Kim HJ, Park SJ, Bae KB, Park SJ, Jang WY, Kang MC, Jeong J, Sung YH, Choi M, et al. (2015) Critical role of Rgs19 in mouse embryonic stem cell proliferation and differentiation. *Differentiation* **89**:42–50.
- Ji YR, Kim MO, Kim SH, Yu DH, Shin MJ, Kim HJ, Yuh HS, Bae KB, Kim JY, Park HD, et al. (2010) Effects of regulator of G protein signaling 19 (RGS19) on heart development and function. *J Biol Chem* **285**:28627–28634.
- Johnson BA, Wang J, Taylor EM, Caillier SJ, Herbert J, Khan OA, Cross AH, De Jager PL, Gourraud PA, Cree BC, et al. (2010) Multiple sclerosis susceptibility alleles in African Americans. *Genes Immun* **11**:343–350.
- Johnson PM and Kenny PJ (2010) Dopamine D2 receptors in addiction-like reward dysfunction and compulsive eating in obese rats. *Nat Neurosci* **13**:635–641.
- Jones SE, Tyrrell J, Wood AR, Beaumont RN, Ruth KS, Tuke MA, Yaghootkar H, Hu Y, Teder-Laving M, Hayward C, et al. (2016) Genome-wide association analyses in 128,266 individuals identifies new morningness and sleep duration loci. *PLoS Genet* **12**:e1006125.
- Kannarkat GT, Lee JK, Ramsey CP, Chung J, Chang J, Porter I, Oliver D, Shepherd K, and Tansey MG (2015) Age-related changes in regulator of G-protein signaling (RGS)-10 expression in peripheral and central immune cells may influence the risk for age-related degeneration. *Neurobiol Aging* **36**:1982–1993.
- Kehrl JH and Sinnarajah S (2002) RGS2: a multifunctional regulator of G-protein signaling. *Int J Biochem Cell Biol* **34**:432–438.
- Kenakin T and Miller LJ (2010) Seven transmembrane receptors as shapeshifting proteins: the impact of allosteric modulation and functional selectivity on new drug discovery. *Pharmacol Rev* **62**:265–304.
- Kestenbaum B, Glazer NL, Köttgen A, Felix JF, Hwang SJ, Liu Y, Lohman K, Kritchevsky SB, Hausman DB, Petersen AK, et al. (2010) Common genetic variants associate with serum phosphorus concentration. *J Am Soc Nephrol* **21**:1223–1232.
- Kim SD, Sung HJ, Park SK, Kim TW, Park SC, Kim SK, Cho JY, and Rhee MH (2006) The expression patterns of RGS transcripts in platelets. *Platelets* **17**:493–497.
- Kimple RJ, De Vries L, Tronchère H, Behe CI, Morris RA, Gist Farquhar M, and Siderovski DP (2001) RGS12 and RGS14 GoLoco motifs are G alpha(i) interaction sites with guanine nucleotide dissociation inhibitor activity. *J Biol Chem* **276**:29275–29281.
- Kimple RJ, Kimple ME, Betts L, Sondek J, and Siderovski DP (2002) Structural determinants for GoLoco-induced inhibition of nucleotide release by Galph subunits. *Nature* **416**:878–881.
- Kimple RJ, Willard FS, and Siderovski DP (2004) Purification and in vitro functional analyses of RGS12 and RGS14 GoLoco motif peptides. *Methods Enzymol* **390**:416–436.
- Kircher M, Witten DM, Jain P, O’Roak BJ, Cooper GM, and Shendure J (2014) A general framework for estimating the relative pathogenicity of human genetic variants. *Nat Genet* **46**:310–315.
- Kirsch T, Wellner M, Luft FC, Haller H, and Lippoldt A (2001) Altered gene expression in cerebral capillaries of stroke-prone spontaneously hypertensive rats. *Brain Res* **910**:106–115.
- Koelle MR and Horvitz HR (1996) EGL-10 regulates G protein signaling in the *C. elegans* nervous system and shares a conserved domain with many mammalian proteins. *Cell* **84**:115–125.
- Koenen KC, Amstadter AB, Ruggiero KJ, Acierio R, Galea S, Kilpatrick DG, and Gelernter J (2009) RGS2 and generalized anxiety disorder in an epidemiologic sample of hurricane-exposed adults. *Depress Anxiety* **26**:309–315.
- Kohara K, Tabara Y, Nakura J, Imai Y, Okubo T, Hata A, Soma M, Nakayama T, Umemura S, Hirawa N, et al. (2008) Identification of hypertension-susceptibility genes and pathways by a systemic multiple candidate gene approach: the millennium genome project for hypertension. *Hypertens Res* **31**:203–212.
- Kovoor A, Seyffarth P, Ebert J, Barghshoon S, Chen CK, Schwarz S, Axelrod JD, Cheyette BN, Simon MI, Lester HA, et al. (2005) D2 dopamine receptors colocalize regulator of G-protein signaling 9-2 (RGS9-2) via the RGS9 DEP domain, and RGS9 knock-out mice develop dyskinesias associated with dopamine pathways. *J Neurosci* **25**:2157–2165.
- Kurash DM, Huang J, Wilkie TM, and Repa JJ (2004) Quantitative real-time polymerase chain reaction measurement of regulators of G-protein signaling mRNA levels in mouse tissues. *Methods Enzymol* **389**:3–15.
- Kuwata H, Nakao K, Harada T, Matsuda I, and Aiba A (2008) Generation of RGS8 null mutant mice by Cre/loxP system. *Kobe J Med Sci* **53**:275–281.
- Kveberg L, Ryan JC, Rolstad B, and Innegjerd M (2005) Expression of regulator of G protein signalling proteins in natural killer cells, and their modulation by Ly49A and Ly49D. *Immunology* **115**:358–365.
- Larminie C, Murdock P, Walhin JP, Duckworth M, Blumer KJ, Scheideler MA, and Garnier M (2004) Selective expression of regulators of G-protein signaling (RGS) in the human central nervous system. *Brain Res Mol Brain Res* **122**:24–34.
- Lee JK, Chung J, Kannarkat GT, and Tansey MG (2013) Critical role of regulator G-protein signaling 10 (RGS10) in modulating macrophage M1/M2 activation. *PLoS One* **8**:e81785.
- Lee JK, Chung J, McAlpine FE, and Tansey MG (2011) Regulator of G-protein signaling-10 negatively regulates NF- κ B in microglia and neuroprotects dopaminergic neurons in hemiparkinsonian rats. *J Neurosci* **31**:11879–11888.
- Lee JK, Kannarkat GT, Chung J, Joon Lee H, Graham KL, and Tansey MG (2016) RGS10 deficiency ameliorates the severity of disease in experimental autoimmune encephalomyelitis. *J Neuroinflammation* **13**:24.
- Lee JK, McCoy MK, Harms AS, Ruhn KA, Gold SJ, and Tansey MG (2008) Regulator of G-protein signaling 10 promotes dopaminergic neuron survival via regulation of the microglial inflammatory response. *J Neurosci* **28**:8517–8528.

- Lee JK and Tansey MG (2015) Physiology of RGS10 in neurons and immune cells. *Prog Mol Biol Transl Sci* **133**:153–167.
- Lee MJ, Kim DE, Zakrzewska A, Yoo YD, Kim SH, Kim ST, Seo JW, Lee YS, Dorn GW II, Oh U, et al. (2012) Characterization of arginylation branch of N-end rule pathway in G-protein-mediated proliferation and signaling of cardiomyocytes. *J Biol Chem* **287**:24043–24052.
- Lee SE, Simons SB, Heldt SA, Zhao M, Schroeder JP, Vellano CP, Cowan DP, Ramineni S, Yates CK, Feng Y, et al. (2010) RGS14 is a natural suppressor of both synaptic plasticity in CA2 neurons and hippocampal-based learning and memory. *Proc Natl Acad Sci USA* **107**:16994–16998.
- Lek M, Karczewski KJ, Minikel EV, Samocha KE, Banks E, Fennell T, O'Donnell-Luria AH, Ware JS, Hill AJ, Cummings BB, et al.; Exome Aggregation Consortium (2016) Analysis of protein-coding genetic variation in 60,706 humans. *Nature* **536**:285–291.
- Le Merrier J, Becker JA, Befort K, and Kieffer BL (2009) Reward processing by the opioid system in the brain. *Physiol Rev* **89**:1379–1412.
- Lerner TN and Kreitzer AC (2012) RGS4 is required for dopaminergic control of striatal LTD and susceptibility to parkinsonian motor deficits. *Neuron* **73**:347–359.
- Leygraf A, Hohoff C, Freitag C, Willis-Owen SA, Krakowicz P, Fritze J, Franke P, Bandelow B, Fimmers R, Flint J, et al. (2006) Rgs 2 gene polymorphisms as modulators of anxiety in humans? *J Neural Transm (Vienna)* **113**:1921–1925.
- Li J, Adams LD, Wang X, Pabon L, Schwartz SM, Sane DC, and Geary RL (2004) Regulator of G protein signaling 5 marks peripheral arterial smooth muscle cells and is downregulated in atherosclerotic plaque. *J Vasc Surg* **40**:519–528.
- Li J, Dani JA, and Le W (2009) The role of transcription factor Pitx3 in dopamine neuron development and Parkinson's disease. *Curr Top Med Chem* **9**:855–859.
- Li J, Jia J, Li H, Yu J, Sun H, He Y, Lv D, Yang X, Glocker MO, Ma L, et al. (2014) SysPTM 2.0: an updated systematic resource for post-translational modification. *Database (Oxford)* **2014**:bau025.
- Li Q, Jin W, Cai Y, Yang F, Chen E, Ye D, Wang Q, and Guan X (2017) Regulator of G protein signaling 20 correlates with clinicopathological features and prognosis in triple-negative breast cancer. *Biochem Biophys Res Commun* **485**:693–697.
- Li X, Chen L, Ji C, Liu B, Gu J, Xu J, Zou X, Gu S, and Mao Y (2005) Isolation and expression pattern of RGS21 gene, a novel RGS member. *Acta Biochim Pol* **52**:943–946.
- Li Y, Tang XH, Li XH, Dai HJ, Miao RJ, Cai JJ, Huang ZJ, Chen AF, Xing XW, Lu Y, et al. (2016) Regulator of G protein signalling 14 attenuates cardiac remodelling through the MEK-ERK1/2 signalling pathway. *Basic Res Cardiol* **111**:47.
- Liang B, Bansal G, Xie Z, and Druey KM (2009) RGS16 inhibits breast cancer cell growth by mitigating phosphatidylinositol 3-kinase signaling. *J Biol Chem* **284**:21719–21727.
- Lifschytz T, Broner EC, Zozulinsky P, Slonimsky A, Eitan R, Greenbaum L, and Lerer B (2012) Relationship between Rgs2 gene expression level and anxiety and depression-like behaviour in a mutant mouse model: serotonergic involvement. *Int J Neuropsychopharmacol* **15**:1307–1318.
- Liou YJ, Chen ML, Wang YC, Chen JY, Liao DL, Bai YM, Lin CC, Chen TT, Mo GH, and Lai IC (2009) Analysis of genetic variations in the RGS9 gene and antipsychotic-induced tardive dyskinesia in schizophrenia. *Am J Med Genet B Neuropsychiatr Genet* **150B**:239–242.
- Lippert E, Yowe DL, Gonzalo JA, Justice JP, Webster JM, Fedyk ER, Hodge M, Miller C, Gutierrez-Ramos JC, Borrego F, et al. (2003) Role of regulator of G protein signaling 16 in inflammation-induced T lymphocyte migration and activation. *J Immunol* **171**:1542–1555.
- Liu Y, Huang H, Zhang Y, Zhu XY, Zhang R, Guan LH, Tang Q, Jiang H, and Huang C (2014) Regulator of G protein signaling 3 protects against cardiac hypertrophy in mice. *J Cell Biochem* **115**:977–986.
- Lo Bello M, Di Fini F, Notaro A, Spataro R, Conforti FL, and La Bella V (2017) ALS-related mutant FUS protein is mislocalized to cytoplasm and is recruited into stress granules of fibroblasts from asymptomatic FUS P525L mutation carriers. *Neurodegener Dis* **17**:292–303.
- López-Aranda MF, Acevedo MJ, Carballo FJ, Gutiérrez A, and Khan ZU (2006) Localization of the GoLoco motif carrier regulator of G-protein signaling 12 and 14 proteins in monkey and rat brain. *Eur J Neurosci* **23**:2971–2982.
- Lothrop AP, Torres MP, and Fuchs SM (2013) Deciphering post-translational modification codes. *FEBS Lett* **587**:1247–1257.
- Lundstrom K (2009) An overview on GPCRs and drug discovery: structure-based drug design and structural biology on GPCRs. *Methods Mol Biol* **552**:51–66.
- Ma P, Cierniewska A, Signarvic R, Cieslak M, Kong H, Sinnamon AJ, Neubig RR, Newman DK, Stalker TJ, and Brass LF (2012) A newly identified complex of spinophilin and the tyrosine phosphatase, SHP-1, modulates platelet activation by regulating G protein-dependent signaling. *Blood* **119**:1935–1945.
- Mahajan A, Rodan AR, Le TH, Gaulton KJ, Haessler J, Stilp AM, Kamatani Y, Zhu G, Sofer T, Puri S, et al.; SUMMIT Consortium; BioBank Japan Project (2016) Trans-ethnic fine mapping highlights kidney-function genes linked to salt sensitivity. *Am J Hum Genet* **99**:636–646.
- Maity B, Stewart A, O'Malley Y, Askeland RW, Sugg SL, and Fisher RA (2013) Regulator of G protein signaling 6 is a novel suppressor of breast tumor initiation and progression. *Carcinogenesis* **34**:1747–1755.
- Maity B, Yang J, Huang J, Askeland RW, Bera S, and Fisher RA (2011) Regulator of G protein signaling 6 (RGS6) induces apoptosis via a mitochondrial-dependent pathway not involving its GTPase-activating protein activity. *J Biol Chem* **286**:1409–1419.
- Mao H, Zhao Q, Daigle M, Ghahremani MH, Chidiac P, and Albert PR (2004) RGS17/RGSZ2, a novel regulator of Gi/o, Gz, and Gq signaling. *J Biol Chem* **279**:26314–26322.
- Mao Y, Lei L, Su J, Yu Y, Liu Z, and Huo Y (2014) Regulators of G protein signaling are up-regulated in aspirin-resistant platelets from patients with metabolic syndrome. *Pharmazie* **69**:371–373.
- Martemyanov KA and Arshavsky VY (2004) Kinetic approaches to study the function of RGS9 isoforms. *Methods Enzymol* **390**:196–209.
- Martinez-Cardús A, Martinez-Balibrea E, Bandrés E, Malumbres R, Ginés A, Manzanao JL, Taron M, Garcia-Foncillas J, and Abad A (2009) Pharmacogenomic approach for the identification of novel determinants of acquired resistance to oxaliplatin in colorectal cancer. *Mol Cancer Ther* **8**:194–202.
- Masuho I, Xie K, and Martemyanov KA (2013) Macromolecular composition dictates receptor and G protein selectivity of regulator of G protein signaling (RGS) 7 and 9-2 protein complexes in living cells. *J Biol Chem* **288**:25129–25142.
- McCauley JL, Zuvich RL, Bradford Y, Kenealy SJ, Schnetz-Boutaud N, Gregory SG, Hauser SL, Oksenberg JR, Mortlock DP, Pericak-Vance MA, et al. (2009) Follow-up examination of linkage and association to chromosome 1q43 in multiple sclerosis. *Genes Immun* **10**:624–630.
- Michaelides M, Li Z, Rana NA, Richardson EC, Hykin PG, Moore AT, Holder GE, and Webster AR (2010) Novel mutations and electrophysiologic findings in RGS9- and R9AP-associated retinal dysfunction (Bradyopsia). *Ophthalmology* **117**:120–127 e121.
- Min C, Cheong SY, Cheong SJ, Kim M, Cho DI, and Kim KM (2012) RGS4 exerts inhibitory activities on the signaling of dopamine D2 receptor and D3 receptor through the N-terminal region. *Pharmacol Res* **65**:213–220.
- Mirnes K, Middleton FA, Stanwood GD, Lewis DA, and Levitt P (2001) Disease-specific changes in regulator of G-protein signaling 4 (RGS4) expression in schizophrenia. *Mol Psychiatry* **6**:293–301.
- Mitchell TS, Bradley J, Robinson GS, Shima DT, and Ng YS (2008) RGS5 expression is a quantitative measure of pericyte coverage of blood vessels. *Angiogenesis* **11**:141–151.
- Mittal V and Linder ME (2004) The RGS14 GoLoco domain discriminates among Galphai isoforms. *J Biol Chem* **279**:46772–46778.
- Mittmann C, Chung CH, Höppner G, Michalek C, Nose M, Schuler C, Schuh A, Eschenhagen T, Weil J, Pieske B, et al. (2002) Expression of ten RGS proteins in human myocardium: functional characterization of an upregulation of RGS4 in heart failure. *Cardiovasc Res* **55**:778–786.
- Miyoshi N, Ishii H, Sekimoto M, Doki Y, and Mori M (2009) RGS16 is a marker for prognosis in colorectal cancer. *Ann Surg Oncol* **16**:3507–3514.
- Moon AM, Stauffer AM, Schwindinger WF, Sheridan K, Firmant A, and Robishaw JD (2014) Disruption of G-protein $\gamma 5$ subtype causes embryonic lethality in mice. *PLoS One* **9**:e90970.
- Moon SW, Dinov ID, Kim J, Zamanyan A, Hobel S, Thompson PM, and Toga AW (2015) Structural neuroimaging genetics interactions in Alzheimer's disease. *J Alzheimers Dis* **48**:1051–1063.
- Moratz C, Hayman JR, Gu H, and Kehrl JH (2004) Abnormal B-cell responses to chemokines, disturbed plasma cell localization, and distorted immune tissue architecture in Rgs1-/- mice. *Mol Cell Biol* **24**:5767–5775.
- Nagata Y, Oda M, Nakata H, Shozaki Y, Kozasa T, and Todokoro K (2001) A novel regulator of G-protein signaling bearing GAP activity for Galphai and Galphaq in megakaryocytes. *Blood* **97**:3051–3060.
- Nance MR, Kreutz B, Tesmer VM, Sterne-Marr R, Kozasa T, and Tesmer JJ (2013) Structural and functional analysis of the regulator of G protein signaling 2-gq complex. *Structure* **21**:438–448.
- Natochin M, Granovsky AE, and Artemyev NO (1997) Regulation of transducin GTPase activity by human retinal RGS. *J Biol Chem* **272**:17444–17449.
- Neitzel KL and Hepler JR (2006) Cellular mechanisms that determine selective RGS protein regulation of G protein-coupled receptor signaling. *Semin Cell Dev Biol* **17**:383–389.
- Nelson MR, Wegmann D, Ehm MG, Kessner D, St Jean P, Verzilli C, Shen J, Tang Z, Bacanu SA, Fraser D, et al. (2012) An abundance of rare functional variants in 202 drug target genes sequenced in 14,002 people. *Science* **337**:100–104.
- Neubig RR and Siderovski DP (2002) Regulators of G-protein signalling as new central nervous system drug targets. *Nat Rev Drug Discov* **1**:187–197.
- Nguyen CH, Ming H, Zhao P, Hugendubler L, Gros R, Kimball SR, and Chidiac P (2009) Translational control by RGS2. *J Cell Biol* **186**:755–765.
- Nisancioğlu MH, Mahoney WM Jr, Kimmel DD, Schwartz SM, Betsholtz C, and Genové G (2008) Generation and characterization of rgs5 mutant mice. *Mol Cell Biol* **28**:2324–2331.
- Nishiguchi KM, Sandberg MA, Kooijman AC, Martemyanov KA, Pott JW, Hagstrom SA, Arshavsky VY, Berson EL, and Dryja TP (2004) Defects in RGS9 or its anchor protein R9AP in patients with slow photoreceptor deactivation. *Nature* **427**:75–78.
- Nishiura H, Nonaka H, Revollo IS, Semba U, Li Y, Ota Y, Irie A, Harada K, Kehrl JH, and Yamamoto T (2009) Pro- and anti-apoptotic dual functions of the C5a receptor: involvement of regulator of G protein signaling 3 and extracellular signal-regulated kinase. *Lab Invest* **89**:676–694.
- Nunn C, Mao H, Chidiac P, and Albert PR (2006) RGS17/RGSZ2 and the RZ/A family of regulators of G-protein signaling. *Semin Cell Dev Biol* **17**:390–399.
- Offermanns S, Mancino V, Revel JP, and Simon MI (1997) Vascular system defects and impaired cell chemokinesis as a result of Galphal3 deficiency. *Science* **275**:533–536.
- Offermanns S, Zhao LP, Gohla A, Sarosi I, Simon MI, and Wilkie TM (1998) Embryonic cardiomyocyte hypoplasia and craniofacial defects in G alpha q/G alpha 11-mutant mice. *EMBO J* **17**:4304–4312.
- Ogden KK, Chen W, Swanger SA, McDaniel MJ, Fan LZ, Hu C, Tankovic A, Kusumoto H, Kosobucki GJ, Schulien AJ, et al. (2017) Molecular mechanism of disease-associated mutations in the pre-M1 helix of NMDA receptors and potential rescue pharmacology. *PLoS Genet* **13**:e1006536.
- Okae H and Iwakura Y (2010) Neural tube defects and impaired neural progenitor cell proliferation in Gbeta1-deficient mice. *Dev Dyn* **239**:1089–1101.
- Okamoto N, Kohmoto T, Naruto T, Masuda K, Komori T, and Imoto I (2017) Novel *CLCN7* compound heterozygous mutations in intermediate autosomal recessive osteopetrosis. *Hum Genome Var* **4**:17036.
- Okimoto N, Bosch OJ, Slattery DA, Pflaum K, Matsushita H, Wei FY, Ohmori M, Nishiki T, Ohmori I, Hiramatsu Y, et al. (2012) RGS2 mediates the anxiolytic effect of oxytocin. *Brain Res* **1453**:26–33.

- Oliveira-Dos-Santos AJ, Matsumoto G, Snow BE, Bai D, Houston FP, Whishaw IQ, Mariathasan S, Sasaki T, Wakeham A, Ohashi PS, et al. (2000) Regulation of T cell activation, anxiety, and male aggression by RGS2. *Proc Natl Acad Sci USA* **97**: 12272–12277.
- Ooe A, Kato K, and Noguchi S (2007) Possible involvement of CCT5, RGS3, and YKT6 genes up-regulated in p53-mutated tumors in resistance to docetaxel in human breast cancers. *Breast Cancer Res Treat* **101**:305–315.
- Opel A, Nobles M, Montaigne D, Finlay M, Anderson N, Breckenridge R, and Tinker A (2015) Absence of the regulator of G-protein signaling, RGS4, predisposes to atrial fibrillation and is associated with abnormal calcium handling. *J Biol Chem* **290**:19233–19244.
- Osterhout JL, Waheed AA, Hiol A, Ward RJ, Davey PC, Nini L, Wang J, Milligan G, Jones TL, and Druey KM (2003) Palmitoylation regulates regulator of G-protein signaling (RGS) 16 function. II. Palmitoylation of a cysteine residue in the RGS box is critical for RGS16 GTPase accelerating activity and regulation of Gi-coupled signalling. *J Biol Chem* **278**:19309–19316.
- Ostrovskaya O, Xie K, Masuho I, Fajardo-Serrano A, Lujan R, Wickman K, and Martemyanov KA (2014) RGS7/Gβ5/R7BP complex regulates synaptic plasticity and memory by modulating hippocampal GABABR-GIRK signaling. *eLife* **3**: e02053.
- Owen VJ, Burton PB, Mullen AJ, Birks EJ, Barton P, and Yacoub MH (2001) Expression of RGS3, RGS4 and Gi alpha 2 in acutely failing donor hearts and end-stage heart failure. *Eur Heart J* **22**:1015–1020.
- Park IK, Klug CA, Li K, Jerabek L, Li L, Nanamori M, Neubig RR, Hood L, Weissman IL, and Clarke MF (2001) Molecular cloning and characterization of a novel regulator of G-protein signaling from mouse hematopoietic stem cells. *J Biol Chem* **276**:915–923.
- Park JH, Gail MH, Weinberg CR, Carroll RJ, Chung CC, Wang Z, Chanock SJ, Fraumeni JF Jr, and Chatterjee N (2011) Distribution of allele frequencies and effect sizes and their interrelationships for common genetic susceptibility variants. *Proc Natl Acad Sci USA* **108**:18026–18031.
- Pashkov V, Huang J, Parameswara VK, Kedzierski W, Kurrasch DM, Tall GG, Esser V, Gerard RD, Uyeda K, Towle HC, et al. (2011) Regulator of G protein signaling (RGS16) inhibits hepatic fatty acid oxidation in a carbohydrate response element-binding protein (ChREBP)-dependent manner. *J Biol Chem* **286**: 15116–15125.
- Patel J, McNeill E, Douglas G, Hale AB, de Bono J, Lee R, Iqbal AJ, Regan-Komito D, Stylianou E, Greaves DR, et al. (2015) RGS1 regulates myeloid cell accumulation in atherosclerosis and aortic aneurysm rupture through altered chemokine signaling. *Nat Commun* **6**:6614.
- Patten M, Bünemann J, Thoma B, Krämer E, Thoenes M, Stübe S, Mittmann C, and Wieland T (2002) Endotoxin induces desensitization of cardiac endothelin-1 receptor signaling by increased expression of RGS4 and RGS16. *Cardiovasc Res* **53**: 156–164.
- Phan HTN, Sjögren B, and Neubig RR (2017) Human missense mutations in regulator of G protein signaling 2 affect the protein function through multiple mechanisms. *Mol Pharmacol* **92**:451–458.
- Pise-Masison CA, Radonovich M, Dohoney K, Morris JC, O'Mahony D, Lee MJ, Trepel J, Waldmann TA, Janik JE, and Brady JN (2009) Gene expression profiling of ATL patients: compilation of disease-related genes and evidence for TCF4 involvement in BIRC5 gene expression and cell viability. *Blood* **113**:4016–4026.
- Piskorowski RA, Nasrallah K, Diamantopoulou A, Mukai J, Hassan SI, Siegelbaum SA, Gogos JA, and Chevalleyre V (2016) Age-dependent specific changes in area CA2 of the hippocampus and social memory deficit in a mouse model of the 22q11.2 deletion syndrome. *Neuron* **89**:163–176.
- Plummer NW, Spicher K, Malphurs J, Akiyama H, Abramowitz J, Nürnberg B, and Birnbaumer L (2012) Development of the mammalian axial skeleton requires signaling through the Gα(i) subfamily of heterotrimeric G proteins. *Proc Natl Acad Sci USA* **109**:21366–21371.
- Ponting CP (1999) Raf-like Ras/Rap-binding domains in RGS12- and still-life-like signalling proteins. *J Mol Med (Berl)* **77**:695–698.
- Posner BA, Gilman AG, and Harris BA (1999) Regulators of G protein signaling 6 and 7: purification of complexes with Gβ25 and assessment of their effects on G protein-mediated signaling pathways. *J Biol Chem* **274**:31087–31093.
- Prasad KM, Chowdari KV, Nimgaonkar VL, Talkowski ME, Lewis DA, and Keshavan MS (2005) Genetic polymorphisms of the RGS4 and dorsolateral prefrontal cortex morphometry among first episode schizophrenia patients. *Mol Psychiatry* **10**:213–219.
- Psifogeorgou K, Papakosta P, Russo SJ, Neve RL, Kardassios D, Gold SJ, and Zachariou V (2007) RGS9-2 is a negative modulator of mu-opioid receptor function. *J Neurochem* **103**:617–625.
- Psifogeorgou K, Terzi D, Papachatzaki MM, Varidaki A, Ferguson D, Gold SJ, and Zachariou V (2011) A unique role of RGS9-2 in the striatum as a positive or negative regulator of opiate analgesia [published correction appears in *J Neurosci* (2011) 31:7578]. *J Neurosci* **31**:5617–5624.
- Qiu R, Wang J, Tsark W, and Lu Q (2010) Essential role of PDZ-RGS3 in the maintenance of neural progenitor cells. *Stem Cells* **28**:1602–1610.
- Raedler D, Ballenberger N, Klucker E, Böck A, Otto R, Prazeres da Costa O, Holst O, Illig T, Buch T, von Mutius E, et al. (2015) Identification of novel immune phenotypes for allergic and nonallergic childhood asthma. *J Allergy Clin Immunol* **135**: 81–91.
- Rahman Z, Gold SJ, Potenza MN, Cowan CW, Ni YG, He W, Wensel TG, and Nestler EJ (1999) Cloning and characterization of RGS9-2: a striatal-enriched alternatively spliced product of the RGS9 gene. *J Neurosci* **19**:2016–2026.
- Rahman Z, Schwarz J, Gold SJ, Zachariou V, Wein MN, Choi KH, Kovoor A, Chen CK, DiLeone RJ, Schwarz SC, et al. (2003) RGS9 modulates dopamine signaling in the basal ganglia. *Neuron* **38**:941–952.
- Rao A, Dallman R, Henderson S, and Chen CK (2007) Gbeta5 is required for normal light responses and morphology of retinal ON-bipolar cells. *J Neurosci* **27**: 14199–14204.
- Ribeiro AL and Okamoto OK (2015) Combined effects of pericytes in the tumor microenvironment. *Stem Cells Int* **2015**:868475.
- Richman RW, Strock J, Hains MD, Cabanilla NJ, Lau KK, Siderovski DP, and Diversé-Pierluissi M (2005) RGS12 interacts with the SNARE-binding region of the Cav2.2 calcium channel. *J Biol Chem* **280**:1521–1528.
- Rivero G, Gabilondo AM, García-Sevilla JA, La Harpe R, Morentin B, and Javier Meana J (2010) Characterization of regulators of G-protein signaling RGS4 and RGS10 proteins in the postmortem human brain. *Neurochem Int* **57**:722–729.
- Robinson-Cohen C, Lutsey PL, Kleber ME, Nielson CM, Mitchell BD, Bis JC, Eny KM, Portas L, Eriksson J, Lorentzon M, et al. (2017) Genetic variants associated with circulating parathyroid hormone. *J Am Soc Nephrol* **28**:1553–1565.
- Ross EM and Wilkie TM (2000) GTPase-activating proteins for heterotrimeric G proteins: regulators of G protein signaling (RGS) and RGS-like proteins. *Annu Rev Biochem* **69**:795–827.
- Ryu J, Woo J, Shin J, Ryoo H, Kim Y, and Lee C (2014) Profile of differential promoter activity by nucleotide substitution at GWAS signals for multiple sclerosis. *Medicine (Baltimore)* **93**:e281.
- Saitoh O, Kubo Y, Miyatani Y, Asano T, and Nakata H (1997) RGS8 accelerates G-protein-mediated modulation of K⁺ currents. *Nature* **390**:525–529.
- Saitoh O, Masuho I, Itoh M, Abe H, Komori K, and Odagiri M (2003) Distribution of regulator of G protein signaling 8 (RGS8) protein in the cerebellum. *Cerebellum* **2**: 154–160.
- Saitoh O, Masuho I, Terakawa I, Nomoto S, Asano T, and Kubo Y (2001) Regulator of G protein signaling 8 (RGS8) requires its NH2 terminus for subcellular localization and acute desensitization of G protein-gated K⁺ channels. *J Biol Chem* **276**: 5052–5058.
- Saitoh O and Odagiri M (2003) RGS8 expression in developing cerebellar Purkinje cells. *Biochem Biophys Res Commun* **309**:836–842.
- Sambi BS, Hains MD, Waters CM, Connell MC, Willard FS, Kimple AJ, Pyne S, Siderovski DP, and Pyne NJ (2006) The effect of RGS12 on PDGFβ receptor signalling to p42/p44 mitogen activated protein kinase in mammalian cells. *Cell Signal* **18**:971–981.
- Schiff ML, Siderovski DP, Jordan JD, Brothers G, Snow B, De Vries L, Ortiz DF, and Diversé-Pierluissi M (2000) Tyrosine-kinase-dependent recruitment of RGS12 to the N-type calcium channel. *Nature* **408**:723–727.
- Schizophrenia Working Group of the Psychiatric Genomics Consortium (2014) Biological insights from 108 schizophrenia-associated genetic loci. *Nature* **511**: 421–427.
- Schossner A, Butler AW, Ising M, Perroud N, Uher R, Ng MY, Cohen-Woods S, Craddock N, Owen MJ, Korszun A, et al. (2011) Genomewide association scan of suicidal thoughts and behaviour in major depression. *PLoS One* **6**:e20690.
- Seeman P, Ko F, Jack E, Greenstein R, and Dean B (2007) Consistent with dopamine supersensitivity, RGS9 expression is diminished in the amphetamine-treated animal model of schizophrenia and in postmortem schizophrenia brain. *Synapse* **61**: 303–309.
- Seki N, Sugano S, Suzuki Y, Nakagawara A, Ohira M, Muramatsu M, Saito T, and Hori T (1998) Isolation, tissue expression, and chromosomal assignment of human RGS5, a novel G-protein signaling regulator gene. *J Hum Genet* **43**: 202–205.
- Semplicini A, Lenzi L, Sartori M, Papparella I, Calò LA, Pagnin E, Strapazzon G, Benna C, Costa R, Avogaro A, et al. (2006) Reduced expression of regulator of G-protein signaling 2 (RGS2) in hypertensive patients increases calcium mobilization and ERK1/2 phosphorylation induced by angiotensin II. *J Hypertens* **24**: 1115–1124.
- Sethakorn N and Dulin NO (2013) RGS expression in cancer: oncoming the cancer microarray data. *J Recept Signal Transduct Res* **33**:166–171.
- Sevastou I, Pryce G, Baker D, and Selwood DL (2016) Characterisation of transcriptional changes in the spinal cord of the progressive experimental autoimmune encephalomyelitis biozzi ABH mouse model by RNA sequencing. *PLoS One* **11**: e0157754.
- Shankar SP, Wilson MS, DiVietro JA, Mentink-Kane MM, Xie Z, Wynn TA, and Druey KM (2012) RGS16 attenuates pulmonary Th2/Th17 inflammatory responses. *J Immunol* **188**:6347–6356.
- Sharma A, Liu X, Hadley D, Hagopian W, Liu E, Chen WM, Onengut-Gumuscu S, Simell V, Rewers M, Ziegler AG, et al.; TEDDY Study Group (2016) Identification of non-HLA genes associated with celiac disease and country-specific differences in a large, international pediatric cohort. *PLoS One* **11**:e0152476.
- Shelton RC, Claiborne J, Sidoryk-Wegrzynowicz M, Reddy R, Aschner M, Lewis DA, and Mirmics K (2011) Altered expression of genes involved in inflammation and apoptosis in frontal cortex in major depression. *Mol Psychiatry* **16**:751–762.
- Shi CS, Huang NN, and Kehrl JH (2012) Regulator of G-protein signaling 3 isoform 1 (PDZ-RGS3) enhances canonical Wnt signaling and promotes epithelial mesenchymal transition. *J Biol Chem* **287**:33480–33487.
- Shi GX, Harrison K, Han SB, Moratz C, and Kehrl JH (2004) Toll-like receptor signaling alters the expression of regulator of G protein signaling proteins in dendritic cells: implications for G protein-coupled receptor signaling. *J Immunol* **172**:5175–5184.
- Shi GX, Harrison K, Wilson GL, Moratz C, and Kehrl JH (2002) RGS13 regulates germinal center B lymphocytes responsiveness to CXCL12 and CXCL13. *J Immunol* **169**:2507–2515.
- Shim H, Wang CT, Chen YL, Chau VQ, Fu KG, Yang J, McQuiston AR, Fisher RA, and Chen CK (2012) Defective retinal depolarizing bipolar cells in regulators of G protein signaling (RGS) 7 and 11 double null mice. *J Biol Chem* **287**: 14873–14879.
- Shu FJ, Ramineni S, Amyot W, and Hepler JR (2007) Selective interactions between Gi alpha1 and Gi alpha3 and the GoLoco/GPR domain of RGS14 influence its dynamic subcellular localization. *Cell Signal* **19**:163–176.
- Shu FJ, Ramineni S, and Hepler JR (2010) RGS14 is a multifunctional scaffold that integrates G protein and Ras/Raf MAPkinase signalling pathways. *Cell Signal* **22**: 366–376.

- Sierra DA, Gilbert DJ, Householder D, Grishin NV, Yu K, Ukidwe P, Barker SA, He W, Wensel TG, Otero G, et al. (2002) Evolution of the regulators of G-protein signaling multigene family in mouse and human. *Genomics* **79**:177–185.
- Simon MI, Strathmann MP, and Gautam N (1991) Diversity of G proteins in signal transduction. *Science* **252**:802–808.
- Sjögren B (2011) Regulator of G protein signaling proteins as drug targets: current state and future possibilities. *Adv Pharmacol* **62**:315–347.
- Sjögren B (2017) The evolution of regulators of G protein signalling proteins as drug targets - 20 years in the making: IUPHAR review 21. *Br J Pharmacol* **174**:427–437.
- Sjögren B, Blazer LL, and Neubig RR (2010) Regulators of G protein signaling proteins as targets for drug discovery. *Prog Mol Biol Transl Sci* **91**:81–119.
- Skiba NP, Yang CS, Huang T, Bae H, and Hamm HE (1999) The alpha-helical domain of Galphat determines specific interaction with regulator of G protein signaling 9. *J Biol Chem* **274**:8770–8778.
- Slep KC, Kercher MA, He W, Cowan CW, Wensel TG, and Sigler PB (2001) Structural determinants for regulation of phosphodiesterase by a G protein at 2.0 Å. *Nature* **409**:1071–1077.
- Smith EN, Bloss CS, Badner JA, Barrett T, Belmonte PL, Berrettini W, Byerley W, Coryell W, Craig D, Edenberg HJ, et al. (2009) Genome-wide association study of bipolar disorder in European American and African American individuals. *Mol Psychiatry* **14**:755–763.
- Smyth DJ, Plagnol V, Walker NM, Cooper JD, Downes K, Yang JH, Howson JM, Stevens H, McManus R, Wijmenga C, et al. (2008) Shared and distinct genetic variants in type 1 diabetes and celiac disease. *N Engl J Med* **359**:2767–2777.
- Snow BE, Antonio L, Suggs S, Gutstein HB, and Siderovski DP (1997) Molecular cloning and expression analysis of rat Rgs12 and Rgs14. *Biochem Biophys Res Commun* **233**:770–777.
- Snow BE, Hall RA, Krumins AM, Brothers GM, Bouchard D, Brothers CA, Chung S, Mangion J, Gilman AG, Lefkowitz RJ, et al. (1998a) GTPase activating specificity of RGS12 and binding specificity of an alternatively spliced PDZ (PSD-95/Dlg/ZO-1) domain. *J Biol Chem* **273**:17749–17755.
- Snow BE, Krumins AM, Brothers GM, Lee SF, Wall MA, Chung S, Mangion J, Arya S, Gilman AG, and Siderovski DP (1998b) A G protein gamma subunit-like domain shared between RGS11 and other RGS proteins specifies binding to Gbetas5 subunits. *Proc Natl Acad Sci USA* **95**:13307–13312.
- So HC, Chen RY, Chen EY, Cheung EF, Li T, and Sham PC (2008) An association study of RGS4 polymorphisms with clinical phenotypes of schizophrenia in a Chinese population. *Am J Med Genet B Neuropsychiatr Genet* **147B**:77–85.
- Song L, Zmijewski JW, and Jope RS (2001) RGS2: regulation of expression and nuclear localization. *Biochem Biophys Res Commun* **283**:102–106.
- Soundararajan M, Willard FS, Kimple AJ, Turnbull AP, Ball LJ, Schoch GA, Gileadi C, Fedorov OY, Dowler EF, Higman VA, et al. (2008) Structural diversity in the RGS domain and its interaction with heterotrimeric G protein alpha-subunits. *Proc Natl Acad Sci USA* **105**:6457–6462.
- Squires KE, Gerber KJ, Pare JF, Branch MR, Smith Y, and Hepler JR (2018) Regulator of G protein signaling 14 (RGS14) is expressed pre- and postsynaptically in neurons of hippocampus, basal ganglia, and amygdala of monkey and human brain. *Brain Struct Funct* **223**:233–253.
- Stewart A, Huang J, and Fisher RA (2012) RGS proteins in heart: brakes on the vagus. *Front Physiol* **3**:95.
- Stewart A, Maity B, Anderegg SP, Allamargot C, Yang J, and Fisher RA (2015) Regulator of G protein signaling 6 is a critical mediator of both reward-related behavioral and pathological responses to alcohol. *Proc Natl Acad Sci USA* **112**:E786–E795.
- Stewart A, Maity B, Wunsch AM, Meng F, Wu Q, Wemmie JA, and Fisher RA (2014) Regulator of G-protein signaling 6 (RGS6) promotes anxiety and depression by attenuating serotonin-mediated activation of the 5-HT(1A) receptor-adenylyl cyclase axis. *FASEB J* **28**:1735–1744.
- Stockman A, Smithson HE, Webster AR, Holder GE, Rana NA, Ripamonti C, and Sharpe LT (2008) The loss of the PDE6 deactivating enzyme, RGS9, results in precocious light adaptation at low light levels. *J Vis* **8**:10.1–10.
- Sun X, Kaltenbronn KM, Steinberg TH, and Blumer KJ (2005) RGS2 is a mediator of nitric oxide action on blood pressure and vasoconstrictor signaling. *Mol Pharmacol* **67**:631–639.
- Sutton LP, Ostrovskaya O, Dao M, Xie K, Orlandi C, Smith R, Wee S, and Martemyanov KA (2016) Regulator of G-protein signaling 7 regulates reward behavior by controlling opioid signaling in the striatum. *Biol Psychiatry* **80**:235–245.
- Tang KM, Wang GR, Lu P, Karas RH, Aronovitz M, Heximer SP, Kaltenbronn KM, Blumer KJ, Siderovski DP, Zhu Y, et al. (2003) Regulator of G-protein signaling-2 mediates vascular smooth muscle relaxation and blood pressure [published correction appears in *Nat Med* (2004) 10:105]. *Nat Med* **9**:1506–1512.
- Tansey MG and Goldberg MS (2010) Neuroinflammation in Parkinson's disease: its role in neuronal death and implications for therapeutic intervention. *Neurobiol Dis* **37**:510–518.
- Tennessen JA, Bigham AW, O'Connor TD, Fu W, Kenny EE, Gravel S, McGee S, Do R, Liu X, Jun G, et al.; Broad GO; Seattle GO; NHLBI Exome Sequencing Project (2012) Evolution and functional impact of rare coding variation from deep sequencing of human exomes. *Science* **337**:64–69.
- Tesmer JJ, Berman DM, Gilman AG, and Sprang SR (1997) Structure of RGS4 bound to AlF4-activated G(i alpha1): stabilization of the transition state for GTP hydrolysis. *Cell* **89**:251–261.
- Thomas EA, Danielson PE, and Sutcliffe JG (1998) RGS9: a regulator of G-protein signalling with specific expression in rat and mouse striatum. *J Neurosci Res* **52**:118–124.
- Torres MP, Dewhurst H, and Sundararaman N (2016) Proteome-wide structural analysis of PTM hotspots reveals regulatory elements predicted to impact biological function and disease. *Mol Cell Proteomics* **15**:3513–3528.
- Trahey M and McCormick F (1987) A cytoplasmic protein stimulates normal N-ras p21 GTPase, but does not affect oncogenic mutants. *Science* **238**:542–545.
- Traver S, Bidot C, Spassky N, Baltauss T, De Tand MF, Thomas JL, Zalc B, Janoueix-Lerosey I, and Gunzburg JD (2000) RGS14 is a novel Rap effector that preferentially regulates the GTPase activity of galphao. *Biochem J* **350**:19–29.
- Traynelis J, Silk M, Wang Q, Berkovic SF, Liu L, Ascher DB, Balding DJ, and Petrovski S (2017) Optimizing genomic medicine in epilepsy through a gene-customized approach to missense variant interpretation. *Genome Res* **27**:1715–1729.
- Traynor JR, Terzi D, Caldaroni BJ, and Zachariou V (2009) RGS9-2: probing an intracellular modulator of behavior as a drug target. *Trends Pharmacol Sci* **30**:105–111.
- Tso PH, Yung LY, Wang Y, and Wong YH (2011) RGS19 stimulates cell proliferation by deregulating cell cycle control and enhancing Akt signaling. *Cancer Lett* **309**:199–208.
- Tu Y, Popov S, Slaughter C, and Ross EM (1999) Palmitoylation of a conserved cysteine in the regulator of G protein signaling (RGS) domain modulates the GTPase-activating activity of RGS4 and RGS10. *J Biol Chem* **274**:38260–38267.
- Tu Y, Wang J, and Ross EM (1997) Inhibition of brain Gz GAP and other RGS proteins by palmitoylation of G protein alpha subunits. *Science* **278**:1132–1135.
- Tu Y, Woodson J, and Ross EM (2001) Binding of regulator of G protein signaling (RGS) proteins to phospholipid bilayers: contribution of location and/or orientation to Gtpase-activating protein activity. *J Biol Chem* **276**:20160–20166.
- Ueda HR, Chen W, Adachi A, Wakamatsu H, Hayashi S, Takasugi T, Nagano M, Nakahama K, Suzuki Y, Sugano S, et al. (2002) A transcription factor response element for gene expression during circadian night. *Nature* **418**:534–539.
- Urabe Y, Tanikawa C, Takahashi A, Okada Y, Morizono T, Tsunoda T, Kamatani N, Kohri K, Chayama K, Kubo M, et al. (2012) A genome-wide association study of nephrolithiasis in the Japanese population identifies novel susceptible loci at 5q35.3, 7p14.3, and 13q14.1. *PLoS Genet* **8**:e1002541.
- Vellano CP, Brown NE, Blumer JB, and Hepler JR (2013) Assembly and function of the regulator of G protein signaling 14 (RGS14)-H-Ras signaling complex in live cells are regulated by Gai1 and Gai-linked G protein-coupled receptors. *J Biol Chem* **288**:3620–3631.
- Vogt IR, Lees AJ, Evert BO, Klockgether T, Bonin M, and Wüllner U (2006) Transcriptional changes in multiple system atrophy and Parkinson's disease putamen. *Exp Neurol* **199**:465–478.
- von Buchholtz L, Elischer A, Tareilus E, Gouka R, Kaiser C, Breer H, and Conzelmann S (2004) RGS21 is a novel regulator of G protein signalling selectively expressed in subpopulations of taste bud cells. *Eur J Neurosci* **19**:1535–1544.
- Vuong TM and Chabre M (1990) Subsecond deactivation of transducin by endogenous GTP hydrolysis. *Nature* **346**:71–74.
- Vural A, McQuiston TJ, Blumer JB, Park C, Hwang IY, Williams-Bey Y, Shi CS, Ma DZ, and Kehrl JH (2013) Normal autophagic activity in macrophages from mice lacking Gai3, AGS3, or RGS19. *PLoS One* **8**:e81886.
- Wagner R, Ryba N, and Uhl R (1988) Rapid transducin deactivation in intact stacks of bovine rod outer segment disks as studied by light scattering techniques: arrestin requires additional soluble proteins for rapid quenching of rhodopsin catalytic activity. *FEBS Lett* **235**:103–108.
- Walsh CT, Garneau-Tsodikova S, and Gatto GJ Jr (2005) Protein posttranslational modifications: the chemistry of proteome diversifications. *Angew Chem Int Ed Engl* **44**:7342–7372.
- Wang J, Ducret A, Tu Y, Kozasa T, Aebersold R, and Ross EM (1998) RGSZ1, a Gz-selective RGS protein in brain: structure, membrane association, regulation by Galphaz phosphorylation, and relationship to a Gz gtpase-activating protein subfamily. *J Biol Chem* **273**:26014–26025.
- Wang J, Zhou Y, Fei X, Chen X, and Zhu Z (2017a) Regulator of G-protein signaling 3 targeted by miR-126 correlates with poor prognosis in gastric cancer patients. *Anticancer Drugs* **28**:161–169.
- Wang Q and Traynor JR (2013) Modulation of μ -opioid receptor signaling by RGS19 in SH-SY5Y cells. *Mol Pharmacol* **83**:512–520.
- Wang Y, Ho G, Zhang JJ, Nieuwenhuisen B, Edris W, Chanda PK, and Young KH (2002) Regulator of G protein signaling Z1 (RGSZ1) interacts with Galphai subunits and regulates Galphai-mediated cell signaling. *J Biol Chem* **277**:48325–48332.
- Wang Y, Tong Y, Tso PH, and Wong YH (2013) Regulator of G protein signaling 19 suppresses Ras-induced neoplastic transformation and tumorigenesis. *Cancer Lett* **339**:33–41.
- Wang Y, Wang J, Zhang L, Karatas OF, Shao L, Zhang Y, Castro P, Creighton CJ, and Ittmann M (2017b) RGS12 is a novel tumor-suppressor gene in African American prostate cancer that represses AKT and MNX1 expression. *Cancer Res* **77**:4247–4257.
- Wang Z, Huang H, He W, Kong B, Hu H, Fan Y, Liao J, Wang L, Mei Y, Liu W, et al. (2016) Regulator of G-protein signaling 5 protects cardiomyocytes against apoptosis during in vitro cardiac ischemia-reperfusion in mice by inhibiting both JNK1/2 and P38 signaling pathways. *Biochem Biophys Res Commun* **473**:551–557.
- Waugh JL, Cerver J, Sharma M, Dufresne RL, Terzi D, Risch SC, Fairbrother WG, Neve RL, Kane JP, Malloy MJ, et al. (2011) Association between regulator of G protein signaling 9-2 and body weight. *PLoS One* **6**:e27984.
- Waugh JL, Lou AC, Eisch AJ, Monteggia LM, Muly EC, and Gold SJ (2005) Regional, cellular, and subcellular localization of RGS10 in rodent brain. *J Comp Neurol* **481**:299–313.
- Wetschurek N and Offermanns S (2005) Mammalian G proteins and their cell type specific functions. *Physiol Rev* **85**:1159–1204.
- Willard FS, Willard MD, Kimple AJ, Soundararajan M, Oestreich EA, Li X, Sowa NA, Kimple RJ, Doyle DA, Der CJ, et al. (2009) Regulator of G-protein signaling 14 (RGS14) is a selective H-Ras effector. *PLoS One* **4**:e4884.
- Willars GB (2006) Mammalian RGS proteins: multifunctional regulators of cellular signalling. *Semin Cell Dev Biol* **17**:363–376.

- Williams NM, Preece A, Spurlock G, Norton N, Williams HJ, McCreddie RG, Buckland P, Sharkey V, Chowdari KV, Zammit S, et al. (2004) Support for RGS4 as a susceptibility gene for schizophrenia. *Biol Psychiatry* **55**:192–195.
- Wydeven N, Posokhova E, Xia Z, Martemyanov KA, and Wickman K (2014) RGS6, but not RGS4, is the dominant regulator of G protein signaling (RGS) modulator of the parasympathetic regulation of mouse heart rate. *J Biol Chem* **289**:2440–2449.
- Xie Z, Chan EC, and Druey KM (2016) R4 regulator of G protein signaling (RGS) proteins in inflammation and immunity. *AAPS J* **18**:294–304.
- Xu B, Roos JL, Dexheimer P, Boone B, Plummer B, Levy S, Gogos JA, and Karayiorgou M (2011) Exome sequencing supports a de novo mutational paradigm for schizophrenia. *Nat Genet* **43**:864–868.
- Yang J, Huang J, Maity B, Gao Z, Lorca RA, Gudmundsson H, Li J, Stewart A, Swaminathan PD, Ibeawuchi SR, et al. (2010) RGS6, a modulator of parasympathetic activation in heart. *Circ Res* **107**:1345–1349.
- Yang J, Kamide K, Kokubo Y, Takiuchi S, Tanaka C, Banno M, Miwa Y, Yoshii M, Horio T, Okayama A, et al. (2005) Genetic variations of regulator of G-protein signaling 2 in hypertensive patients and in the general population. *J Hypertens* **23**:1497–1505.
- Yang L, Lee MM, Leung MM, and Wong YH (2016a) Regulator of G protein signaling 20 enhances cancer cell aggregation, migration, invasion and adhesion. *Cell Signal* **28**:1663–1672.
- Yang S and Li YP (2007) RGS12 is essential for RANKL-evoked signaling for terminal differentiation of osteoclasts in vitro. *J Bone Miner Res* **22**:45–54.
- Yang S, Li YP, Liu T, He X, Yuan X, Li C, Cao J, and Kim Y (2013) Mx1-cre mediated Rgs12 conditional knockout mice exhibit increased bone mass phenotype. *Genesis* **51**:201–209.
- Yang SH, Li CF, Chu PY, Ko HH, Chen LT, Chen WW, Han CH, Lung JH, and Shih NY (2016b) Overexpression of regulator of G protein signaling 11 promotes cell migration and associates with advanced stages and aggressiveness of lung adenocarcinoma. *Oncotarget* **7**:31122–31136.
- Yasui T, Okada A, Urabe Y, Usami M, Mizuno K, Kubota Y, Tozawa K, Sasaki S, Higashi Y, Sato Y, et al. (2013) A replication study for three nephrolithiasis loci at 5q35.3, 7p14.3 and 13q14.1 in the Japanese population. *J Hum Genet* **58**:588–593.
- Yoon D, Kim YJ, Cui WY, Van der Vaart A, Cho YS, Lee JY, Ma JZ, Payne TJ, Li MD, and Park T (2012) Large-scale genome-wide association study of Asian population reveals genetic factors in FRMD4A and other loci influencing smoking initiation and nicotine dependence. *Hum Genet* **131**:1009–1021.
- You M, Wang D, Liu P, Vikis H, James M, Lu Y, Wang Y, Wang M, Chen Q, Jia D, et al. (2009) Fine mapping of chromosome 6q23-25 region in familial lung cancer families reveals RGS17 as a likely candidate gene. *Clin Cancer Res* **15**:2666–2674.
- Yowe D, Weich N, Prabhudas M, Poisson L, Errada P, Kapeller R, Yu K, Faron L, Shen M, Cleary J, et al. (2001) RGS18 is a myeloerythroid lineage-specific regulator of G-protein-signalling molecule highly expressed in megakaryocytes. *Biochem J* **359**:109–118.
- Yu S, Yu D, Lee E, Eckhaus M, Lee R, Corria Z, Accili D, Westphal H, and Weinstein LS (1998) Variable and tissue-specific hormone resistance in heterotrimeric Gs protein alpha-subunit (G α) knockout mice is due to tissue-specific imprinting of the g α gene. *Proc Natl Acad Sci USA* **95**:8715–8720.
- Yuan H, Low CM, Moody OA, Jenkins A, and Traynelis SF (2015a) Ionotropic GABA and glutamate receptor mutations and human neurologic diseases. *Mol Pharmacol* **88**:203–217.
- Yuan X, Cao J, Liu T, Li YP, Scannapieco F, He X, Oursler MJ, Zhang X, Vacher J, Li C, et al. (2015b) Regulators of G protein signaling 12 promotes osteoclastogenesis in bone remodeling and pathological bone loss. *Cell Death Differ* **22**:2046–2057.
- Zachariou V, Georgescu D, Sanchez N, Rahman Z, DiLeone R, Berton O, Neve RL, Sim-Selley LJ, Selley DE, Gold SJ, et al. (2003) Essential role for RGS9 in opiate action. *Proc Natl Acad Sci USA* **100**:13656–13661.
- Zhang H, Wang F, Kranzler HR, Anton RF, and Gelernter J (2012) Variation in regulator of G-protein signaling 17 gene (RGS17) is associated with multiple substance dependence diagnoses. *Behav Brain Funct* **8**:23.
- Zhang K, Howes KA, He W, Bronson JD, Pettenati MJ, Chen C, Palczewski K, Wensel TG, and Baehr W (1999) Structure, alternative splicing, and expression of the human RGS9 gene. *Gene* **240**:23–34.
- Zhang P and Mende U (2014) Functional role, mechanisms of regulation, and therapeutic potential of regulator of G protein signaling 2 in the heart. *Trends Cardiovasc Med* **24**:85–93.
- Zhang S, Watson N, Zahner J, Rottman JN, Blumer KJ, and Muslin AJ (1998) RGS3 and RGS4 are GTPase activating proteins in the heart. *J Mol Cell Cardiol* **30**:269–276.