ASSOCIATE EDITOR: LYNETTE C. DAWS

Modulating Dopamine Signaling and Behavior with Chemogenetics: Concepts, Progress, and Challenges

Annika Højrup Runegaard, Ciarán Martin Fitzpatrick, David Paul Drucker Woldbye, Jesper Tobias Andreasen, Andreas Toft Sørensen, and Ulrik Gether

Molecular Neuropharmacology and Genetics Laboratory, Department of Neuroscience (A.H.R., D.P.D.W., A.T.S., U.G.) and Department of Drug Design and Pharmacology (C.M.F., J.T.A.), Faculty of Health and Medical Sciences, University of Copenhagen, Copenhagen, Denmark

Abstract.....	124
I. Introduction.....	124
A. A Portrait of the Dopamine System	125
1. Dopamine Synthesis and Neurotransmission	125
2. Dopamine Pathways and Circuits.....	126
B. Chemogenetics—Pharmacological Remote Control of Neuronal Activity	126
1. Specific Spatial Resolution of Designer Receptors Exclusively Activated by Designer Drugs Activation	127
2. Specific Temporal Resolution of Designer Receptors Exclusively Activated by Designer Drugs Activation.....	129
II. Chemogenetic Approaches for Dissecting Dopamine-Mediated Behaviors.....	130
A. Locomotor Activity.....	140
1. Dopamine Activity Correlates with Locomotor Activity—Proof-of-Concept Studies.....	140
2. The Relative Output-Activity from Direct Medium Spiny Neurons and Indirect Medium Spiny Neurons Determines Locomotor Activity	140
3. Locomotor Activity—A Simple Behavior with Multiple Levels of Regulation.....	141
4. Dysfunctional Motor Control in Parkinson's Disease.....	142
B. Motivation and Reward-Guided Behaviors.....	143
1. Substantiating Dopamine as a Key Regulator of Motivation	143
2. The Role of Direct Medium Spiny Neurons and Indirect Medium Spiny Neurons in Regulating Reward-Motivated Behaviors	143
3. Feeding	144
4. Mating.....	144
5. Sleep-Wake Patterns and Arousal	145
6. Immunity	145
C. Addiction.....	146
1. Locomotor Sensitization	146
2. Reward Perception, Drug-Cue Associations, and Relapse Behaviors.....	146
3. Compulsive Drug Taking	148
D. Mood, Nociception, and Cognition	148
III. Discussion and Future Directions.....	149
A. Designer Receptors Exclusively Activated By Designer Drugs Effects in Theory and in Reality	149

Address correspondence to: Ulrik Gether, Molecular Neuropharmacology and Genetics Laboratory, Department of Neuroscience, University of Copenhagen, Mærsk Tower 5th floor, 3 Blegdamsvej, DK-2200, Copenhagen, Denmark. E-mail: gether@sund.ku.dk; or Andreas Toft Sørensen, Molecular Neuropharmacology and Genetics Laboratory, Department of Neuroscience, University of Copenhagen, Mærsk Tower 5th floor, 3 Blegdamsvej, DK-2200, Copenhagen, Denmark. E-mail: andreass@sund.ku.dk

A.H.R. and C.M.F. contributed equally as first authors.

A.T.S. and U.G. contributed equally as corresponding authors.

This work was supported by the UCPH 2016 Program of Excellence and the Lundbeck Foundation.

<https://doi.org/10.1124/pr.117.013995>.

1. Designer Receptors Exclusively Activated By Designer Drugs Expression—The Meaning of Inertness	149
2. Interpretation of Activating Designer Receptors Exclusively Activated By Designer Drugs with Clozapine N-Oxide.....	150
3. Considerations for Repeated Activation of Designer Receptors Exclusively Activated By Designer Drugs	151
4. Can Selective Dopamine Expression Be Achieved?	151
5. Can Designer Receptors Exclusively Activated By Designer Drugs Help Dissect the Dopaminergic Pathways?	151
B. Have Chemogenetic Studies Advanced Our Understanding of Dopamine-Related Behaviors?	152
C. Future Directions.....	152
IV. Conclusion	152
References	153

Abstract—For more than 60 years, dopamine (DA) has been known as a critical modulatory neurotransmitter regulating locomotion, reward-based motivation, and endocrine functions. Disturbances in DA signaling have been linked to an array of different neurologic and psychiatric disorders, including Parkinson's disease, schizophrenia, and addiction, but the underlying pathologic mechanisms have never been fully elucidated. One major obstacle limiting interpretation of standard pharmacological and transgenic interventions is the complexity of the DA system, which only appears to widen as research progresses. Nonetheless, development of new genetic tools, such as chemogenetics, has led to an entirely new era for functional studies of neuronal signaling. By exploiting receptors that are engineered to respond selectively to an otherwise inert ligand, so-called Designer Receptors Exclusively Activated by

Designer Drugs (DREADDs), chemogenetics enables pharmacological remote control of neuronal activity. Here we review the recent, extensive application of this technique to the DA field and how its use has advanced the study of the DA system and contributed to our general understanding of DA signaling and related behaviors. Moreover, we discuss the challenges and pitfalls associated with the chemogenetic technology, such as the metabolism of the DREADD ligand clozapine N-oxide (CNO) to the D2 receptor antagonist clozapine. We conclude that despite the recent concerns regarding CNO, the chemogenetic toolbox provides an exceptional approach to study neuronal function. The huge potential should promote continued investigations and additional refinements to further expound key mechanisms of DA signaling and circuitries in normal as well as maladaptive behaviors.

I. Introduction

Pioneering studies showed more than 60 years ago that dopamine (DA) is not just a metabolic intermediate in the synthesis of norepinephrine but a neurotransmitter in itself (Carlsson and Waldeck, 1958). The DA system has since been the subject of numerous studies, and it is now fully established that DA is an essential and dominant neurotransmitter within the central nervous system (CNS), modulating a wide array of physiologic functions, such as motor control, reward-based motivation, memory, learning, decision making, and neuroendocrine secretion (Björklund and Dunnett, 2007; Berke, 2018). The DA field has been further fostered by the realization that imbalances in DA homeostasis are critical components of both neurologic

and psychiatric diseases such as Parkinson's disease (PD), schizophrenia, bipolar disorder, Tourette's syndrome, attention-deficit/hyperactivity disorder (ADHD), and addiction (Hornykiewicz, 1966; Beaulieu and Gainetdinov, 2011; Tritsch and Sabatini, 2012).

A major challenge has been to dissect the molecular and cellular mechanisms underlying the ability of DA to regulate a broad spectrum of brain circuits and thereby influence multiple physiologic and pathophysiological processes in the CNS. One main reason is the complexity and heterogeneity of the DA system, which has become even more apparent as research has progressed over the years. As a consequence, multiple aspects of DA function still appear remarkably poorly understood. This situation has called for the implementation of

ABBREVIATIONS: AAV, adeno-associated virus; AC, adenylyl cyclase; ADHD, attention-deficit/hyperactivity disorder; BNST, bed nucleus of the stria terminalis; CAV2, canine adenovirus type 2; CNO, clozapine N-oxide; CNS, central nervous system; CPP, conditioned place preference; D₁₋₅R, dopamine receptors 1–5; DA, dopamine; DAT, dopamine transporter; DIO, double-floxed inverse open reading frame; dMSN, direct medium spiny neuron; DREADDs, designer receptors exclusively activated by designer drugs; ERK, extracellular signal-regulated kinase; Flp, flippase; GIRK, G-protein coupled inward-rectifying potassium channels; Gi/q/s, regulatory G-protein subunit; GPCR, G protein-coupled receptor; hM3Dq, Gq DREADD; hM4Di, Gi DREADD; 5-HT, 5-hydroxytryptamine; iMSN, indirect medium spiny neuron; KO, knockout; KORD, kappa-opioid receptor-DREADD; L-DOPA, L-3,4-dihydroxyphenylalanine; MSN, medium spiny neuron; NAc, nucleus accumbens; 6-OHDA, 6-hydroxy-dopamine; PD, Parkinson's disease; PFC, prefrontal cortex; PKA, protein kinase A; rM3Darr, beta-arrestin based DREADD; rM3Ds, Gs DREADD; SalB, salvinorin B; SN, substantia nigra; TH, tyrosine hydroxylase; VTA, ventral tegmental area; WT, wild type.

novel techniques in the field encompassing particular tools that can harness specific manipulations of neuronal circuits to elucidate mechanisms of homeostatic and pathologic phenotypes. The advancements of transformative molecular-genetic manipulation technologies, such as chemogenetics and optogenetics, are altering the course of behavioral neuroscience with the potential to unprecedentedly advance our knowledge about how DAergic signaling and its complex organization mediates behavior. It is therefore of no surprise that these technologies already have been widely used as strategies to study and dissect DA-related functions. Despite their many obvious advantages, such techniques inevitably also embrace a substantial number of pitfalls and limitations that must be considered when interpreting experimental results.

In this review, we carry out a comprehensive review of studies over the last few years of the DA system in which chemogenetic approaches have been employed; that is, studies, that have taken advantage of DREADDs (“Designer Receptors Exclusively Activated by Designer Drugs”) and thereby used the possibility of activating discrete G protein-coupled signaling cascades in selected target cells to manipulate neuronal functions. We believe that the technique possesses huge potential and can be a powerful approach if used in the appropriate manner. We will provide a short overview of the DA system before introducing the chemogenetic approach and critically reviewing areas of DA research in which chemogenetic approaches have been applied. Next, we will specifically describe pitfalls and limitations, as well as discuss future directions for refining the technology. We hope thereby to generate a detailed overview of what we have learned from the use of chemogenetics and a framework for how chemogenetics can fruitfully be applied to the DA system to continue the exploration of this fascinating neurotransmitter.

A. A Portrait of the Dopamine System

1. Dopamine Synthesis and Neurotransmission. DA neurotransmission depends on DA synthesis that takes place within a restricted set of neurons expressing the enzymes involved in the biosynthetic pathway of DA from dietary tyrosine sources. Tyrosine hydroxylase (TH), the rate-limiting enzyme in DA production, converts tyrosine to L-3,4-dihydroxyphenylalanine (L-DOPA), which in turn is converted by aromatic amino acids decarboxylase to DA. Cytosolic DA is sequestered into synaptic vesicles by the action of the vesicular monoamine transporter 2, acting as a proton antiporter that takes advantage of the low intravesicular pH to transport DA into the vesicles against a large concentration gradient (German et al., 2015). After fusion of these vesicles with the plasma membrane and release from presynaptic terminals, DA exerts its actions via five distinct, but highly homologous, G protein-coupled receptors (GPCRs). The DA receptors are divided into

two subclasses based on their pharmacological profile and coupling: the D1-like receptors (D₁R and D₅R) and the D2-like receptors (D₂R, D₃R, and D₄R) (Beaulieu and Gainetdinov, 2011). The D₂R exists in a short or long variant (D₂S and D₂L) of which D₂S constitutes the primary presynaptic autoreceptor in the DAergic neurons (Lindgren et al., 2003). D1-like receptors are generally characterized by displaying one to two orders of magnitude lower affinity to DA compared with the D2-like receptors, meaning that the latter are much more likely to be saturated with DA during tonic release conditions (Richfield et al., 1987). The coupling of the D1-like and D2-like receptors to intracellular signaling pathways are distinct and generally oppositely directed. D1-like receptors couple to G_{αs} and G_{αolf}, leading to stimulation of adenylyl cyclase (AC), which results in increased levels of cAMP and activation of protein kinase A (PKA). In contrast, D2-like receptors couple to G_{αi} and G_{αo}, leading to inhibition of AC, lower cAMP levels, and less PKA activation. Via the G_{βγ} subunits of the heterotrimeric G protein, the D2-like receptors can also regulate ion channel function, including G protein-coupled inward-rectifying potassium channels (GIRK) (Kuzhikandathil et al., 1998; Lavine et al., 2002) and voltage-gated Ca²⁺-channels (Yan et al., 1997). DA receptors, and in particular D2-like receptors, also signal through G protein-independent pathways. The D2-like receptors can, for example, operate through β-arrestin-dependent pathways involving activation of Akt (also called protein kinase B) and glycogen synthase kinase 3 (Beaulieu et al., 2004, 2005, 2007). For further details on DA receptors, including a more in-depth description of their functional coupling and distribution in the brain, we kindly refer to other reviews (Neve et al., 2004; Beaulieu and Gainetdinov, 2011; Tritsch and Sabatini, 2012; Beaulieu et al., 2015).

To understand the complex regulation of DA signaling, it is essential to highlight the DA transporter (DAT). DAT is expressed in DA neurons and mediates rapid reuptake of released DA. In doing so, DAT plays an essential role in controlling extracellular levels of DA and thereby DA signaling (Jones et al., 1998). DAT belongs to the large family of neurotransmitter:sodium symporters, and thus the transporter utilizes the transmembrane sodium gradient to translocate effectively DA across the membrane against its concentration gradient. Importantly, DAT is the prime target for the stimulatory action of psychostimulants such as cocaine and amphetamine. Whereas cocaine acts as a high-affinity, competitive inhibitor, amphetamine is a substrate that upon entry through the transporter promotes the release of DA via DAT, a process known to involve activation of protein kinases, such as Ca²⁺/calmodulin-dependent protein kinase II and phosphorylation of the transporter itself (Kristensen et al., 2011). DAT is also a target for commonly used medication used against ADHD such as methylphenidate and modafinil. For further

details on DAT, we kindly refer to other excellent reviews (Torres and Amara, 2007; Kristensen et al., 2011; Sulzer, 2011).

2. Dopamine Pathways and Circuits. The majority of the brain's DAergic transmission is organized into four neural pathways that we now know partake in distinct, but also sometimes seemingly overlapping behaviors. 1) The *nigrostriatal pathway* projects from the midbrain substantia nigra (SN) pars compacta to the dorsal striatum and exerts both facilitatory and inhibitory regulation of movements and is involved in habitual and compulsive responses. 2) The *mesolimbic pathway* projects from the ventral tegmental area (VTA) to limbic brain regions, including the amygdala and the nucleus accumbens (NAc) of the ventral striatum. This pathway is involved in motivation, goal-directed behaviors, attributing incentive values of reward-related environmental stimuli and labeling pleasurable emotions with motivational salience. 3) The *mesocortical pathway* projects from the midbrain VTA to the prefrontal cortex (PFC) and chiefly mediates executive function such as selective and sustained attention, flexible responding to stimuli, planning and goal-directed behavior, and monitoring and inhibitory control. 4) The *tuberoinfundibular pathway* projects from the hypothalamus to the pituitary gland, where DA inhibits the secretion of the hormone prolactin (Volkow et al., 2007; Arias-Carrión and Pöppel, 2007; Björklund and Dunnett, 2007).

Although the separation of the DA system into the above-mentioned pathways is still relevant, it is important to emphasize that recent investigations have led to a significantly more complicated picture of the DAergic system with heterogeneous populations of neurons that express and release DA via different mechanisms in different brain regions (Morales and Margolis, 2017). For example, labeling studies have revealed a complex heterogeneity of the midbrain DAergic system in terms of projection patterns, firing and release probability, and intracellular molecular characteristics and properties (Lammel et al., 2008, 2014; Apuschkin et al., 2015; Poulin et al., 2018). Furthermore, it has been demonstrated that some midbrain DA neurons are capable of releasing other neurotransmitters, such as 5-hydroxytryptamine (5-HT) (Zhou et al., 2005), glutamate (Chuhma et al., 2004; Nair-Roberts et al., 2008; Hnasko et al., 2010; Stuber et al., 2010), and GABA (Tritsch et al., 2014; Kim et al., 2015; Berrios et al., 2016). Finally, complex input networks projecting onto DAergic neurons, as well as output target areas (e.g., Beier et al., 2015; Poulin et al., 2018), emphasize a major role of DA as a modulator rather than a regulator of neuronal excitability in target areas. Once DA is released in target areas, DA receptors expressed on postsynaptic membranes convey and determine the further processing of the DAergic signal, which is dependent on the DA receptor profile. While target

areas such as the dorsal striatum, comprised mainly of GABAergic medium spiny neurons (MSNs), have been found to be rather ordered in terms of expressing either D1- or D2-like receptors and with distinct projection patterns, other target areas such as the NAc and PFC lack such architecture or remain to be equally characterized (Beaulieu and Gainetdinov, 2011; Kupchik et al., 2015). Altogether, the DA system displays an anatomic, as well as molecular, complexity, which precludes standard pharmacologic or genetic approaches for deconstructing DA signaling in normal and maladaptive behavioral states. However, these obstacles can be circumvented if DAergic pathways are precisely defined and/or DAergic signaling can be both spatially and temporally controlled. Furthermore, if maladaptive DA states are reversible, a technique that would allow for bidirectional regulation would be a huge advantage for exploring new therapeutic avenues.

B. Chemogenetics—Pharmacological Remote Control of Neuronal Activity

The DREADD technology was developed by Bryan Roth and colleagues at the University of North Carolina and was published for the first time in 2007 (Armbruster et al., 2007). This methodological approach extends from other chemogenetic platforms described in the field of GPCRs, such as allele-specific activation of genetically encoded receptors (Strader et al., 1991), receptors activated solely by synthetic ligands (Conklin et al., 2008) and neoreceptors (Jacobson et al., 2001). The development of DREADDs enabled new study designs as it allowed neuroscientists to bidirectionally manipulate intracellular G protein-coupled signaling of neuronal populations of interest and at sufficient strength—acting like a device for remote control of neuronal signaling. We will briefly describe the rationale behind the DREADDs technique and elaborate on its suitability for studying the DA system. For more information on DREADDs development and structures, see Roth (2016) and Armbruster et al. (2007).

The original DREADDs are mutated muscarinic acetylcholine receptors, which are insensitive to their natural ligand, acetylcholine, and instead respond to an otherwise inert compound, clozapine *N*-oxide (CNO), a metabolite of the atypical antipsychotic clozapine (Roth, 2016). The basic idea was to genetically introduce and express the various DREADD variants within cells of interest and thereby allow targeted manipulation of specific intracellular signaling cascades through the coupling of the distinct receptors to either excitatory (Gs, Gq) or inhibitory (Gi) signaling cascades upon CNO binding (see Fig. 1). Specifically, the Gq-coupled hM3Dq DREADD leads to activation of the phospholipase C cascade and increases intracellular calcium to promote neuronal burst firing (Armbruster et al., 2007), while the Gs-coupled rM3Ds DREADD increases cAMP-mediated depolarization and thus mimics the signaling

Fig. 1. The chemogenetic toolbox contains a variety of engineered receptors (i.e., DREADDs and the KORD) that upon activation, via CNO or SalB, initiate endogenous signaling cascades and regulate canonical signaling and excitability (Roth 2016; Armbruster et al., 2007; Vardy et al., 2015). Four of the engineered receptors are scaffolded on the muscarinic acetylcholine receptor, respond to CNO, and couple to Gq (hM3Dq), Gs (rM3Ds), Gi (hM4Di) or β -arrestin (rM3Darr). KORD is scaffolded on the kappa-opioid receptor and responds to SalB, but signal as hM4Di via the Gi-coupled pathway. However, the kinetic profile of the KORD-SalB is different from hM4Di-CNO and provides a faster onset of action. While hM3Dq and rM3Ds increase neuronal excitability by activating the PKC and PKA pathways, respectively, hM4Di and KORD decrease neuronal excitability by activating GIRK channels and inhibiting the PKA pathway. The beta-arrestin biased rM3Darr initiates beta-arrestin signaling, which distinguishes the late-stage signaling pathway from Gi-coupled receptors signaling via mitogen-activated protein kinase/ERK.

pathway activated by D1-like receptors (Farrell et al., 2013). The Gi-coupled hM4Di DREADD causes membrane hyperpolarization through activation of GIRK and decreases cAMP signaling (Armbruster et al., 2007) analogous to D_2 Rs. Like other GPCRs, activated DREADDs may recruit arrestins, which are able to convey delayed G protein-independent receptor signaling (Roth, 2016). A beta-arrestin based DREADD (rM3Darr) has been generated to explore the physiologic relevance of these arrestin-dependent signaling pathways, but overall this receptor has been used much less than the other aforementioned variants (Nakajima and Wess, 2012; Roth, 2016). Finally, a new DREADD variant, the kappa-opioid receptor DREADD (KORD), constitutes a mutated kappa-opioid receptor that exerts inhibitory actions via a Gi-coupled signaling pathway. This receptor is activated by salvinorin B (SalB), an inert metabolite of the psychotropic salvinorin A (Vardy et al., 2015). This new variant can be used alone, as well as opening up the possibility for combinatorial DREADD studies, where KORD can be applied together with CNO-sensitive DREADDs. Of further interest, the pharmacokinetics of KORD activation by SalB differs greatly from that of CNO-sensitive DREADDs, as it possesses a much more rapid onset of action and response extinction timeframe within 1 hour in contrast to several hours for the effects of CNO (Vardy et al., 2015).

1. Specific Spatial Resolution of Designer Receptors Exclusively Activated by Designer Drugs Activation. A major advantage of chemogenetics, compared with classic pharmacological manipulations, is the specificity by which activation of the receptors can be achieved. The spatial resolution of DREADD-induced manipulations is determined by the ectopic expression of DREADDs in desired neurons, and this can in principle be controlled by the researcher's choice of approach: either directly by placing the DREADD transgene downstream of a specific promoter in transgenic animals or virus vectors (Farrell et al., 2013; Ferguson et al., 2013) or indirectly through Cre-lox-mediated recombination using double-floxed inverse open reading frame (DIO) DREADD-encoding viruses in combination with rodent Cre-driver lines or a second Cre-expressing virus (Bock et al., 2013; Carter et al., 2013; Schwarz et al., 2015). Figure 2 demonstrates how various viral-transgenic approaches can be applied to target mid-brain DA neurons and here exemplified how to obtain gradually better selective expression toward only NAc-projecting neurons. While viral delivery of a DREADD transgene may limit expression to the target area e.g., VTA (Fig. 2A), the DREADD transgene could be introduced downstream of promoters, regulating genes that are selectively expressed within the DA system to obtain more desirable expression. Such an approach has been applied selectively to target DREADDs in dorsostriatal MSNs belonging to either the striatonigral direct

Fig. 2. Viral approaches to obtain increased specificity of chemogenetic manipulations within the DA system. (A–D) Illustrates different viral-transgenic approaches to obtain chemogenetic control of midbrain DA neurons with increasing specificity of DREADD expression toward NAc-projecting neurons (shown for hM4Di-mCherry). (A) The viral approach that limits DREADD expression primarily toward the injection site; a viral vector carrying the DREADD transgene is injected into the target area of a WT mouse. Increased selectivity can be obtained by placing the DREADD transgene downstream of a neuron-specific promoter (e.g., Ferguson et al., 2011; Farrell et al., 2013; Ferguson et al., 2013). (B) The viral Cre-dependent approach is by far the most commonly used to obtain chemogenetic control of a neuronal population of interest; a single Cre-dependent DREADD vector is injected into the target area of a Cre-driver mouse and restricts the expression toward Cre-expressing neurons within the injection site (e.g., Lindeberg et al., 2004; Savitt et al., 2005; Bäckman et al., 2006; Runegaard et al., 2018). (C) The dual viral Cre-dependent approach has been applied to

pathway (i.e., D₁R expressing) or the striatopallidal indirect pathway (i.e., D₂R expressing) by using herpes simplex virus carrying dynorphin and enkephalin promoters, respectively (Ferguson et al., 2011). Targeting of midbrain DA neurons has been successful using a combinatorial approach, with the most commonly applied approach taking advantage of transgenic Cre-driver lines (Fig. 2B). More specifically, injection of a viral vector (e.g., AAV), carrying a DIO-DREADD (the expression of which relies on the presence of Cre), into the target area (e.g., VTA) of Cre-driver lines exploiting classic genetic markers of DA neurons, confines DREADD expression to putative DA neurons in the midbrain. Classic markers include the DA transporter (DAT) and tyrosine hydroxylase (TH), and both DAT-Cre and TH-Cre driver lines have been used to confine DREADD expression to midbrain DA neurons (Lindeberg et al., 2004; Savitt et al., 2005; Bäckman et al., 2006; Runegaard et al., 2018) (Fig. 2B). Alternatively, the Cre-recombinase can be delivered by viral means in a dual viral approach in wild-type (WT) mice (Fig. 2C), as recently demonstrated by use of retrogradely transported recombinant canine adenovirus (CAV2) (Hnasko et al., 2006; Boender et al., 2014; Boekhoudt et al., 2016) or retrograde adeno-associated virus (AAV2-retro) (Tervo et al., 2016). The dual viral approach can even provide extra target specificity toward individual DA projections in Cre-driver lines. In such a setup, combinatorial viral expression determined by a retrograde vector encoding Cre-dependent recombinase flippase (Flp) and a second viral vector encoding Flp-dependent DREADD can confine the expression further (Fig. 2D) (Boender et al., 2014; Runegaard et al., 2018). In addition, although less well characterized for achieving high DA neuron specificity, an AAV vector delivering Cre-recombinase under the control of a fragment of the TH promoter is another option (Stauffer et al., 2016). With a new delivery platform allowing for intravenous AAV delivery, it appears that DA neurons can even be selectively targeted by systemic application (Chan et al., 2017). These two latter viral approaches are interesting, as they technically permit DREADD targeting toward DA neurons in species other than rodents (Stauffer et al., 2016). Taken together, the molecular-genetic toolbox for chemogenetics is large and ideally suited for targeting and studying the DA system (Whissell et al., 2016). Since the signaling cascade of

DREADDs and DA receptors are conveyed through the same downstream molecules, it is also noteworthy to mention that activation of DREADDs possesses the potential ability to mimic, mitigate, or override endogenous DA signaling.

2. Specific Temporal Resolution of Designer Receptors Exclusively Activated by Designer Drugs Activation. The temporal resolution of DREADDs activation represents another important variable. Indeed, as mentioned above, the DREADD and KORD agonists, CNO and SalB, respectively, display distinct temporal dynamics (Roth, 2016). Furthermore, the route of administration can strongly influence the pharmacokinetics and thereby the temporal dynamics of the manipulation. Systemic administration of CNO by intraperitoneal injection causes plasma levels to peak within 30 min and decline over the subsequent 2 h (Guettier et al., 2009). Although plasma levels of CNO decline quickly, behavioral effects may be longer lasting for up to 6–10 h (Alexander et al., 2009; Wang et al., 2013). It is likely that this is linked to the recently reported possible conversion of CNO to clozapine following in vivo administration in rodents (Maclaren et al., 2016; Gomez et al., 2017). Strikingly, it appears that clozapine, but not CNO, crosses the blood-brain barrier, and clozapine binds to and activates DREADD receptors with higher affinity and potency than CNO (Gomez et al., 2017). As clozapine is a D₂R antagonist, this underscores the importance of including CNO-treated non-DREADD-expressing animals as controls, particularly in studies focusing on the DA system. Along these lines, it should be mentioned that the resulting concentrations of clozapine, however, are predicted to be too low to affect D₂R function at the most commonly used concentrations of CNO in behavioral studies (Gomez et al., 2017; Mahler and Aston-Jones, 2018). Nonetheless, these findings support the use of subthreshold doses of clozapine as the DREADD-activating agent instead of high doses of CNO (Gomez et al., 2017). In contrast to CNO, the pharmacokinetics of SalB have not been equally well studied, but behavioral effects after intraperitoneal administration of SalB indicates a 10–15 min onset of action time and roughly 1 h duration depending on the behavior investigated, thus providing both a

obtain selective expression within neurons projecting to a certain area. In WT mice, a Cre-dependent DREADD vector is injected in the area containing the cell nuclei and a retrogradely transported vector, such as CAV2-Cre, is injected in the terminal area of the neurons of interest (e.g., Hnasko et al., 2006; Boender et al., 2014; Boekhoudt et al., 2016; Tervo et al., 2016). One limitation here is that the DREADD may be expressed in any neuron projecting from the target area (e.g., VTA) and traveling along or in proximity to the target area of the neurons of interest where the retrograde vector is injected (e.g., NAc)—even if they do not terminate here. Such an approach does not discriminate between various populations of neurons projecting here unless specified by the promoter driving either Cre or DREADD. (D) The dual viral Flp- and Cre-dependent approach is a way to obtain chemogenetic control within a specific projection pathway of a neuronal population of interest (e.g., Boender et al., 2014; Runegaard et al., 2018; Stauffer et al., 2016). In a Cre-driver line, a Flp-dependent DREADD vector is injected in the area containing the nuclei of the neurons of interest, and in addition, a retrogradely transported and Cre-dependent vector, such as CAV2-CMV-DIO-Flp, is injected in the terminal area of the projection of interest. By this method, DREADD will be expressed only in Flp-transduced neurons, which again depends on the neurons of interest, defined by the Cre-driver line as well as the projection area of interest where the CAV-2 vector is injected.

faster onset and elimination of its effect compared with CNO (Vardy et al., 2015; Marchant, et al., 2016b).

II. Chemogenetic Approaches for Dissecting Dopamine-Mediated Behaviors

In the following sections, we will review how the DREADD technology has been applied to investigate DA circuitries and signaling pathways in an array of different behaviors and pathologic states, including locomotion, PD, reward-motivated behaviors, addiction, and mood disorders. Figure 3 shows an anatomic overview of selected DA-related studies where the

DREADD technology has been employed and which approaches those that have been used to target selective regions, neurons, or projections of the DA system. A more extensive and elaborate overview of all DA-related chemogenetic studies published so far (77 papers in total), covering the period from 2011 till June 2018, is provided in Table 1, including information regarding area of research, test model, chemogenetic vector, target area/cell, strain, and major findings for easy comparison between studies. Considering that DREADDs have demonstrated their usefulness for studying neurotransmitter systems other than DA, we refer to excellent recent reviews covering a broader use

Fig. 3. Sagittal view of a rodent brain presenting an anatomic overview of selected DA-related studies where the DREADD technology has been employed, illustrating which approaches that have been used to target selective regions, neurons, or projections of the DA system. Studies that have revealed DAergic influence on behavior by chemogenetic manipulations within specified regions are indicated by a dashed link. Arrows indicate increases (↑) or decreases (↓) in the respective behaviors. The key provides information on the application of DREADDs in the specific study, including the DREADD variant (Gq, Gi, or Gs), how the expression is specified either directly or by means of a Cre-dependency (i.e., DIO), as well as injection site of DREADDs (which is also indicated by the link to the schematic presentation of the rodent brain). CNO was administered intraperitoneally except in one study (Mahler et al., 2014), where CNO was applied by local microinfusions into VTA, denoted by “#VTA.” For further information, see Table 1.

TABLE 1

Elaborate overview of DA-related chemogenetic behavioral studies categorized by area of research and test model applied and sorted by year of publication

For each study, the chemogenetic vector, target area/cell, rodent species/strain utilized is listed, together with the behavioral effect mediated by each chemogenetic intervention to allow comparison between studies. The respective DREADDs were activated by injection of CNO/SalB, unless stated otherwise. Codes in the index are abbreviated as in the manuscript. The various promoters used in the chemogenetic vectors are denoted by their commonly applied abbreviation. AAV^{CA}: AAV serotype was not reported by authors in the original manuscript.

Area of Research	Test Model	Reference	Chemogenetic Vector *DIO: double-floxed inverted open reading frame (Cre-dependent)	Target Area (Cells)	Rodent Species/Strain	Effect Mediated by Chemogenetic Intervention
Locomotor activity	Novelty induced locomotor activity and habituation	Farrell et al. (2013)	BAC: Adora2A-rM3Ds-IRES- mCherry	Global expression (iMSN)	Mouse: transgenic (Adora2A-rM3Ds)	↓ locomotor activity
		Wang et al. (2013)	AAV2/9-CAG-DIO-hM3Dq-2A- mCherry	VTA/SN (DAergic)	Mouse: DAT-Cre	↑ locomotor activity
		Vardy et al. (2015)	AAV8-hSyn-DIO-KORD-IRES- mCitrine	VTA/SN (GABAergic)	Mouse: VGAT-Cre	↑ locomotor activity
		Parker et al. (2016)	AAV5-CaMKII-hM4Di- mCherry	Thalamus (glutamatergic thalamostriatal neurons)	Mouse: Dorsolateral striatum, 6- OHDA- lesioned	Locomotor recovery of parkinsonian mice; ↑ ambulatory time and average velocity (both by intrapertoneally and intra- dorsolateral striatum CNO), ↓ freezing (intrapertoneally only)
		Bellocchio et al. (2016)	AAV1/2-CAG-DIO-hM3Dq- mCherry	Dorsal striatum (dMSN or iMSN)	Mouse: D ₁ R-Cre, D ₂ R- Cre	Acute dMSN/iMSN Dq-stimulation: ↑/↓ locomotor activity, Chronic iMSN Dq-stimulation: ↑ locomotor activity
		Lemos et al. (2016)	AAV2/5-hSyn-DIO-hM4Di- mCherry, AAV8-hSyn-DIO- KORD-IRES-mCitrine	Dorsal striatum/NAc (iMSN)	Mouse: iMSN- D ₂ R KO (Adora2A-Cre background)	Locomotor recovery of D ₂ R-KO mice mimicking parkinsonian mice (↑ locomotor activity) ↑ locomotor activity
		Carvalho Poyraz et al. (2016)	AAV5-hSyn-DIO-hM4Di- mCherry	Dorsomedial striatum or NAc (iMSN)	Mouse: D ₂ R- (D2R-Cre background), D ₂ R-Cre *overexpressing D ₂ R in iMSN	↑ locomotor activity
		Dobbs et al. (2016)	AAV5-hSyn-DIO-hM4Di- mCherry	NAc (iMSN)	Mouse: Adora2ACre	↑ locomotor activity
		Marchant et al. (2016b)	AAV9-CaMKIIα-HA-KORD- IRES-mCitrine, AAV9- CaMKIIα-HA-hM4Di-IRES- mCitrine	VTA/SN (DAergic)	Rat: WT	↓ locomotor activity
		Chu et al. (2017)	BAC: Adora2A-rM3Ds-IRES- mCherry	Global expression (iMSN)	Mouse: transgenic (Adora2A-rM3Ds)	Acute: ↓ locomotor activity Repeated (2 to 3 days): ↓ reduced vesicular glutamate transporter expressing cortico- subthalamic nucleus axon terminals (i.e. parkinsonian phenotype) Inhibits D ₁ R-induced increase in miniature inhibitory postsynaptic current frequency – supports important role of muscarinic receptor type
		Moehle et al. (2017)	AAV5-hSyn-DIO-hM4Di- mCherry	Dorsal striatum (dMSN)	Mouse: D ₁ R-M4 KO (D ₁ R-Cre background)	Inhibits D ₁ R-induced increase in miniature inhibitory postsynaptic current frequency – supports important role of muscarinic receptor type

(continued)

TABLE 1—Continued

Area of Research	Test Model	Reference	Chemogenetic Vector *DIO: double-floxed inverted open reading frame (Cre-dependent)	Target Area (Cells)	Rodent Species/Strain	Effect Mediated by Chemogenetic Intervention
	Habituated/home- cage locomotor activity	Runegaard et al. (2018)	AAV8-hSyn-DIO-rM3Ds- mCherry, AAV8-hSyn-DIO- hM4Di-mCherry	VTA (DAergic)	Mouse: TH-Cre	4 Gi-signaling in dMSN neurons by regulating locomotor activity via control of AC/cAMP. N.B. no behavioral experiments performed
		Sandhu et al. (2018)	AAV1/2-Eifla-DIO-hM3Dq- mCherry-WPRE	VTA (DAergic)	Mouse: DAT-Cre	Di-/Ds-stimulation: ↓/↑ locomotor activity during habituation
		Bouabid and Zhou (2018)	BAC: Adora2A-rM3Ds-IRES- mCherry	Global expression (iMSN)	Mouse: transgenic (Adora2A-rM3Ds)	↑ locomotor activity
		Blum et al. (2014)	AAV8-DIO-hM3Dq-mCherry	VTA/SN (DAergic)	Mouse: DAT-Cre	↓ locomotor activity (associate with excitation of iMSN and inhibition of globus pallidum neuronal firing)
		Boender et al. (2014)	AAV ^{NA} -hSyn-DIO-hM3Dq- mCherry (VTA), CAV2-Cre	VTA→NAc (DAergic)	Rat: WT	↑ locomotor activity, ↑ ultradian day rhythms by increased active-phase (CNO applied in drinking water)
		Patterson et al. (2015)	AAV ^{NA} -hSyn-DIO-hM3Dq- mCherry	Lateral hypothalamus (neurotensin releasing)	Mouse: neurotensin-Cre	↑ locomotor activity
		Boekhoudt et al. (2016)	AAV5-DIO-hSyn-hM3Dq- mCherry (VTA/SN), CAV2- Cre (NAc/dorsomedial striatum)	VTA, SN, (DAergic) VTA→NAc, SN→Dorsomedial striatum	Rat: TH-Cre, WT with CAV2Cre	VTA and VTA→NAc Dq-stimulation: ↑ long-lasting locomotor activity, SN and SN→dorsomedial striatum stimulation: modest ↑ locomotor activity
		Zhu et al. (2016)	AAV2-hSyn-DIO-hM3Dq- mCherry, AAV2-hSyn-DIO- hM4Di-mCherry	NAc (dMSN or iMSN)	Mouse: D ₁ R-Cre, D ₂ R- Cre	NAc dMSN/iMSN Dq-stimulation: 1/4 locomotor activity and running, NAc iMSN Di-stimulation: ↑ locomotor activity and running
		Qualls- Creekmore et al. (2017)	AAV5-hSyn-DIO-hM3Dq- mCherry	Lateral hypothalamus (galaninergic, GABAergic)	Mouse: GalaninCre, VGAT-Cre	Galanin Dq-stimulation: ↑ locomotor activity GABA Dq-stimulation: ↑ locomotor activity and compulsive-like behaviors
		Dell'Anno et al. (2014)	Lentivirus-hM3Dq-IRES- mCherry	Dorsal striatal transplanted DA neurons induced from fibroblast	Rat: SN 6- OHDA- lesioned	Dq-stimulation: ↑ DA release, excitability and motor recovery of hemi-parkinsonian mice
	Motor coordination and performance assays (Rotarod, cylinder, rotational locomotion, corridor tests, etc.)					

(continued)

TABLE 1—Continued

Area of Research	Test Model	Reference	Chemogenetic Vector *DIO: double-flxed inverted open reading frame (Cre-dependent)	Target Area (Cells)	Rodent Species/Strain	Effect Mediated by Chemogenetic Intervention
		Chen et al. (2016)	AAV1-CAG-DIO-hM3Dq- mCherry, AAV1-CAG-DIO- hM4Di-mCherry	Dorsal striatum transplanted DA neurons differentiated from human embryonic stem cells	Mouse: SN 6- OHDA- lesioned	Di-/Dq-stimulation: ↓/↑ graft cell DA release, excitability and motor recovery of hemi- parkinsonian mice
		Aldrin-Kirk et al. (2016)	AAV8-hSyn-DIO-rM3Ds- mCherry-WPRE, AAV8- hSyn-DIO-hM3Dq-HA, AAV8-hSyn-DIO-KORD- IRES-mCitrine	Dorsal striatum transplanted DA neurons from TH-Cre rat embryos	Rat: striatal 6- OHDA- lesioned	Dq- and Ds-combinatory stimulation: ↑ DA release, excitability and motor performance but also ↑ graft-induced dyskinesia KORD- stimulation: ↓ graft- stimulation: induced recovery
		Alcacer et al. (2017)	AAV5-hSyn-DIO-hM3Dq- mCherry, AAV5-hSyn-DIO- rM3Ds-mCherry	Unilateral dorsolateral striatum (dMSN or iMSN)	Mouse: Adora2aCre, D ₁ R-Cre	iMSN/dMSN Dq-stimulation: ↓/↑ locomotor activity, ipsi/contralateral rotations enhanced iMSN/dMSN Dq-stimulation in hemi- parkinsonian mice: prevent/mimic L-DOPA's effect on contralateral forelimb use: ↓/↑ L-DOPA induced dyskinesia dMSN Dq/Ds-stimulation induces abnormal involuntary movements lower than/similar to L-DOPA
		Lopez-Huerta et al. (2016)	AAV10 ⁺ -hSyn-hM4Di- mCherry, # selective towards neurons of matrix compartment in striatum	Dorsolateral striatum (matrix neurons)	Mouse: WT	↓ accuracy in learned task assaying skilled motor performance
Motivation and reward guided behaviors	Progressive ratio schedule	Aldrin-Kirk et al. (2018)	AAV8-hSyn-DIO-rM3Ds- mCherry-WPRE, AAV8- hSyn-DIO-hM3Dq-HA, AAV8-hSyn-DIO-KORD- IRES-mCitrine	Unilateral dorsal striatum (cholinergic interneurons)	Rat: striatal 6- OHDA- lesioned Choline acetyltransferase Cre	Dq/Ds-stimulation: potentiated low dose L-DOPA to reverse parkinsonian phenotype in stepping task (not cylinder task) and increased abnormal involuntary movement following mid and high dose of L-DOPA (via iMSN output) ↑ effort exertion in progressive ratio task
		Boender et al. (2014)	AAV ^{NA} -hSyn-DIO-hM3Dq- mCherry (VTA), CAV2-Cre (NAc)	VTA→NAc (DAergic)	Rat: WT	
		Boekhoudt et al. (2017a)	AAV2/5-hSyn-DIO-hM3Dq- mCherry	VTA, SN pars compacta (DAergic)	Rat: TH-Cre	↑ sucrose responding through ↑ initiations of reward-seeking actions
						(continued)

TABLE 1—Continued

Area of Research	Test Model	Reference	Chemogenetic Vector *DIO, double-fluxed inverted open reading frame (Cre-dependent)	Target Area (Cells)	Rodent Species/Strain	Effect Mediated by Chemogenetic Intervention
Feeding	Action outcome learning task	Bobadilla et al. (2017)	AAV2-hSyn-DIO-hM3Dq- mCherry	NAc (neuronal nitric oxide synthase (NOS) interneurons)	Mouse: NOS1- Cre	↑ sucrose seeking after extinction ↑ cue-induced reinstatement of sucrose seeking
		Ferguson and Neumaier (2012)	HSV-pDYN-hM4Di-GFP	Dorsal striatum (dMSN)	Rat: WT	↓ acquisition of lever pressing for a reward reinforcer
		Runegaard et al. (2018)	AAV8-hSyn-DIO-hM4Di- mCherry	VTA (DAergic)	Mouse: TH-Cre	↓ motivation to work for reward in a simple must touch test, no effect on reward preference
	High vs. low reward discrimination task	Ferguson et al. (2013)	HSV-pDYN-HA-rM3Ds, HSV- pDYN-HA-hM4Di	Dorsomedial striatum (dMSN)	Rat: WT	Di-stimulation: ↓ acquisition of lever pressing for a reward reinforcer Di-/Ds- stimulation 1/1 stable retention of high-reward optimized strategies
		Chang et al. (2015)	AAV8-hSyn-hM4Di-mCitrine	Ventral pallidum	Rat: WT	↓ acquisition of sign- tracking behaviors
	Progressive ratio schedule, progressive hold- down, outcome devaluation	Carvalho Poyraz et al. (2016)	AAV5-hSyn-DIO-hM4Di- mCherry	Dorsomedial striatum or NAc (iMSN)	Mouse: D ₂ R- ^{ORdev*} (D2R-Cre background), D ₂ R-Cre ^{*overexpressing D₂R in iMSN}	↑ motivation in D ₂ R- ^{ORdev} mice and control littermates in PR and hold-down sessions
		Gallo et al. (2018)	AAV ^{NA} , *DIO-hM4Di- mCherry (Serotype, and promotor information is lacking)	NAc (iMSN)	Mouse: Drd2-Cre	↑ motivation to work for a salient reward by Di stimulation of iMSNs projecting to the ventral pallidum
	Home-cage food intake measurement	Zhu et al. (2016)	AAV2-hSyn-DIO-hM3Dq- mCherry, AAV2-hSyn-DIO- hM4Di-mCherry	NAc (dMSN or iMSN)	Mouse: D ₁ R-Cre, D ₂ R- Cre	dMSN Dq/Di stimulation in home cage under high energy expenditure conditions: 1/↓ food intake
		Boekhoudt, Roelofs et al. (2017)	AAV5-hSyn-DIO-hM3Dq- mCherry (VTA, SN pars compacta), CAV2-Cre (NAc, PFC or amygdala)	VTA, SN, (DAergic), VTA→NAc, VTA→PFC, VTA→amygdala	Rat: TH-Cre, WT with CAV2Cre	VTA Dq- stimulation: ↓ meal size, ↑ locomotor activity, VTA→NAc Dq-stimulation: ↓ mean meal size and eating duration, ↑ mean meal frequency, ↑ locomotor activity
	High fat vs. regular chow diet choice model Intermittent high fat diet exposure or	Foldi et al. (2017)	AAV ^{NA} -hSyn-DIO-hM3Dq- mCherry (VTA), CAV2-Cre (NAc)	VTA→NAc (DAergic)	Rat: WT	↑ food intake, food anticipatory activity, prevented activity-based anorexia weight loss
		Wang et al. (2015)	AAV ^{NA} -hSyn-DIO-hM4Di- mCherry, AAV ^{NA} -hSyn-DIO- hM3Dq-mCherry	Nucleus tractus solitarius (glucagon-like peptide 1 expressing neurons)	Mouse: Phox2b- Cre	Dq/Di-stimulation 1/↑ high- fat food intake (also by intra-VTA CNO)
		Xu et al. (2017)	AAV ^{NA} , *hM3Dq-mCherry, AAV ^{NA} , *hM4Di-mCherry,	VTA/SN (DAergic)	Mouse: DATCreER* ^{*Tamoxifen inducible}	VTA Dq-stimulation: ↓ binge-like eating

(continued)

TABLE 1—Continued

Area of Research	Test Model	Reference	Chemogenetic Vector *DIO: double-flxed inverted open reading frame (Cre-dependent)	Target Area (Cells)	Rodent Species/Strain	Effect Mediated by Chemogenetic Intervention
	continuous chow/high fat diet exposure		*Assumed DIO, but poor description of viral vectors. Serotype, and promotor information is lacking			behaviors, VTA Di-stimulation blocked anti-binge-like eating properties of 5-HT _{2C} receptor agonists
	Progressive ratio schedule	Qualls- Creekmore et al. (2017)	AAV5-hSyn-DIO-hM3Dq- mCherry	Lateral hypothalamus (GABAergic, GABA subpopulation with galanin)	Mouse: VGAT- Cre, Galanin-Cre	Dq-VGAT/Galanin stimulation: ↑ operant food-seeking behaviors, Dq-VGAT stimulation: ↑ food consumption No effect on salt intake
	Salt-depletion model	Sandhu et al. (2018)	AAV1/2-EFlα -DIO-hM3Dq- mCherry-WPRE	VTA (DAergic)	DAT-Cre	
Mating	Behavioral observations during mating, CPP test	Goto et al. (2015)	AAV ^{NA} -hSyn-DIO-rM3Ds- mCherry, AAV ^{NA} -hSyn-DIO- hM4Di-mCherry	Dorsal striatum (dMSN or iMSN)	Mouse: transgenic biosensor (PKA or ERK) in D ₁ R- or D ₂ R- Cre	dMSN/iMSN Di-stimulation: ↓/↑ mating behaviors, dMSN/iMSN Ds-stimulation: ↑/↓ mating behaviors ↓ mating-induced cFos in VTA but no effect on initiation, expression and facilitation of sexual behavior, ↓ subsequent amphetamine CPP following withdrawal in sex experienced rats (cross-sensitization of reward)
		Beloate et al. (2016)	AAV5-hSyn-DIO-hM4Di- mCherry	VTA (DAergic)	Rat: TH-Cre	
Sleep-wake patterns and arousal	Polysomnography recordings	Bellocchio et al. (2016)	AAV ^{NA} -CAG-DIO-hM3Dq- mCherry	Dorsal striatum (dMSN or iMSN)	Mouse: D ₁ R-Cre, D ₂ R- Cre	Acute iMSN-Dq stimulation: ↑ slow-wave phase sleep, Chronic iMSN-Dq stimulation: ↑ wakefulness
		Eban-Rothschild et al. (2016)	AAV5-EF1α-DIO-hM4Di- mCherry	VTA (DAergic)	Mouse: TH-Cre	↓ wakefulness, ↑ pre-sleep- related behaviors, ↓ wakefulness even in the presence of relevant salient-stimuli cues ↓ wakefulness, ↑ sleep duration
		Cho et al. (2017)	AAV2-hSyn-DIO-hM4Di- mCherry	Dorsal raphe nuclei (DAergic)	Mouse: TH-Cre	A2a-Dq/Di indirect pathway stimulation in NAc core: slow-wave sleep induction/inhibition of induction, Ventral pallidum-Dq stimulation: ↑ slow-wave sleep, neuronal disinhibition
		Oishi et al. (2017b)	AAV-hSyn-DIO-hM3Dq- mCherry, AAV-hSyn-DIO- hM4Di-mCherry	NAc core (iMSN), Ventral pallidum (GABAergic)	Mouse: A2aR- Cre, GAD2-Cre	VTA Dq-stimulation: ↑ wakefulness duration, ↓ number of wakefulness episodes
		Oishi et al. (2017a)	AAV10-hSyn-DIO-hM3Dq- mCherry, AAV10-hSyn-DIO- hM4Di-mCherry	VTA, SN (DAergic)	Mouse: DAT-Cre	

(continued)

TABLE 1—Continued

Area of Research	Test Model	Reference	Chemogenetic Vector *DIO: double-flxed inverted open reading frame (Cre-dependent)	Target Area (Cells)	Rodent Species/Strain	Effect Mediated by Chemogenetic Intervention
Drug response and addiction	CPP, immune response assays	Sun et al. (2017)	AAV10-hSyn-hM3Dq- mCherry, AAV10-hSyn- hM4Di-mCherry	VTA	Mouse: WT	Dq-stimulation: ↑ wakefulness duration
		Luo et al. (2018)	AAV ^{NA} -hSyn-DIO-hM3Dq- mCherry, AAV ^{NA} -hSyn-DIO- hM4Di-mCherry	NAc	D ₁ R-Cre	Dq-stimulation: ↑ arousal/wakefulness, ↓ food intake, nesting behaviors, Di-stimulation: ↑ non- REM sleep, nesting ↑ innate and adaptive immunity responses against bacterial challenges, mediates CNO-induced CPP
		Ben-Shaanan et al. (2016)	AAV8-hSyn-DIO-hM3Dq- mCherry	VTA (DAergic)	Mouse: TH-Cre	dMSN/iMSN Di-stimulation: ↓/↑ behavioral sensitization to amphetamine, no effect on acute response
		Ferguson et al. (2011)	HSV-pDyn-HA-hM4Di, HSV- pEnk-HA-hM4Di	Dorsomedial striatum (dMSN or iMSN)	Rat: WT	↓ amphetamine-induced locomotor activity and sensitization ↓ amphetamine-induced locomotion during initiation of sensitization, but no effect on expression, ↑ conditioned locomotor response in amphetamine pre- treated rats
	Acute stimulant induced locomotor activity and behavioral sensitization	Farrell et al. (2013)	BAC: Adora2a-rM3Ds-IRES- mCherry	Global expression (iMSN)	Mouse: transgenic (Adora2a rM3Ds)	Rescues cocaine-induced locomotor response of mice with no D ₂ Rs in iMSN neurons ↓ cocaine-induced hyperlocomotion
		Kerstetter et al. (2016)	AAV5-hSyn-DIO-hM4Di- mCherry (medial PFC), CAV2-Cre (NAc)	Medial PFC→NAc	Rat: WT	
		Dobbs et al. (2016)	AAV5-hSyn-DIO-hM4Di- mCherry	NAc (iMSN)	Mouse: iMSN- D ₂ R KO (Adora2a-Cre background)	
		Marchant et al. (2016b)	AAV9-CaMKIIa-HA-KORD- IRES-mCitrine, AAV9- CaMKIIa-HA-hM4Di-IRES- mCitrine	VTA/SN (DAergic)	Rat: WT	
	CPP test	Wasserman et al. (2016)	AAV ^{NA} -CAG-DIO-hM3Dq- mCherry, AAV ^{NA} -CAG-DIO- hM4Di-mCherry	Rostromedial tegmental area (GABAergic)	Mouse: GAD2Cre	Dq/Di stimulation: ↓/↑ morphine-induced locomotor activity ↓ cocaine-induced hyperlocomotion (acute), but no effect on cocaine- induced locomotion in sensitized mice and repeated Gi stimulation prior to cocaine do not prevent behavioral sensitization
		Runegaard et al. (2018)	AAV8-hSyn-DIO-hM4Di- mCherry	VTA (DAergic)	Mouse: TH-Cre	↓ expression of ethanol place preference dMSN Di-stimulation: ↓ cocaine-induced place
	CPP test	Pina et al. (2015)	AAV5-hSyn-HA-hM4Di-IRES- mCitrine	BNST	Mouse: WT	
		Calipari et al. (2016)	AAV5-hSyn-DIO-hM4Di- mCherry	NAc core (dMSN or iMSN)	Mouse: D ₁ R-Cre, D ₂ R- Cre	

(continued)

TABLE 1—Continued

Area of Research	Test Model	Reference	Chemogenetic Vector *DIO; double-flxed inverted open reading frame (Cre-dependent)	Target Area (Cells)	Rodent Species/Strain	Effect Mediated by Chemogenetic Intervention
		Calipari et al. (2017)	AAV2-hSyn-DIO-hM3Dq- mCherry	VTA (DAergic)	Mouse: TH-Cre	preference (sustained effect) ↑ cocaine CPP in male and diestrus female mice (mimicking increased reward processing in estrus female mice) Disrupt expression of ethanol place preference
		Pina and Cunningham (2017)	AAV8-hSyn-DIO-hM4Di- mCherry (BNST), HSV-Cre (VTA)	BNST→VTA	Mouse: WT	Cocaine CPP unaffected but extinction prolonged IL→NAc shell Dq-stimulation: disrupts expression of toluene place preference (reverse Toluene-induced reduced excitability of these projections) IL→NAc core Di-stimulation: no effect on expression of toluene place preference
		Runegaard et al. (2018)	AAV8-hSyn-DIO-hM4Di- mCherry	VTA (DAergic)	Mouse: TH-Cre	↑ motivation to obtain cocaine during progressive ratio testing; no effect on cocaine intake during training Di-stimulation: ↓ ethanol consumption (limited access model)
		Wayman and Woodward (2018)	AAV8-EF1a-DIO-hM3Dq- mCherry or AAV8-EF1a- DIO-hM4Di-mCherry (IL), CAV2-Cre (NAc core or NAc shell)	IL→NAc core IL→NAc shell	Mouse: TH-Cre Rat: WT	facilitates intracranial self- stimulation ↓ ethanol self-administration after 3 wk abstinence Rostral/caudal ventral pallidum Di-stimulation: ↓ cue-prime induced cocaine reinstatement respectively Di in rostral ventral pallidum (unilateral) + VTA (contralateral): ↓ cue- induced cocaine reinstatement
Self-administration drug/ethanol paradigm		Bock et al. (2013)	AAV1/2-hSyn-DIO-hM4Di- mCherry	NAc core (iMSN)	Mouse: Adora2a-Cre	↓ cue-induced cocaine reinstatement (via mGluR2/3 signaling), no effect on cue-induced reinstatement of sucrose seeking
		Cassatano et al. (2014)	AAV2/5-hSyn-DIO-hM4Di- mCherry, AAV2/5-hSyn- DIO-hM3Dq-mCherry, AAV2/5-hSyn-Cre	NAc	Mouse: WT	↓ compulsivity and prevented persevered
		Bull et al. (2014)	AAV5-GFAP-HA-hM3Dq- IRES-mCitrine	NAc core (astrocytes)	Rat: WT	
		Mahler et al. (2014)	lentivirus-hSyn-HA-hM4Di- GFP, lentivirus-DIO-hSyn- hM4Di-mCherry	Rostral and caudal ventral pallidum, VTA/SN (DAergic)	Rat: WT, TH-Cre	
		Scofield et al. (2015)	AAV5-GFAP-HA-hM3Dq- IRES-mCitrine	NAc core (astrocytes)	Rat: WT	
		Pascoli et al. (2015)	AAV1-CamKIIα-hM4Di- mCherry	Orbitofrontal cortex (DAergic)	Mouse: DAT-Cre	

(continued)

TABLE 1—Continued

Area of Research	Test Model	Reference	Chemogenetic Vector *DIO: double-flxed inverted open reading frame (Cre-dependent)	Target Area (Cells)	Rodent Species/Strain	Effect Mediated by Chemogenetic Intervention
						self-stimulation despire punishment
		Augur et al. (2016)	AAV2-hSyn-HA-hM3Dq-IRES- mCitrine (ventromedial PFC), CAV-2-cre (NAc shell),	Ventromedial PFC → NAc shell	Rat: WT	↓ cue-induced cocaine reinstatement after extinction, no effect on prime-dose-induced reinstatement
		Cheng et al. (2017)	AAV8-hSyn-DIO-hM3Dq- mCherry, AAV8-hSyn-DIO- hM4Di-mCherry	Dorsomedial striatum (dMSN or iMSN)	Mouse: D ₁ R-Cre, D ₂ R- Cre	dMSN Dq/Di stimulation: 1/4 ethanol consumption and preference, iMSN Dq/Di stimulation: 1/1 ethanol consumption and preference
		Kerstetter et al. (2016)	AAV5-hSyn-DIO-hM4Di- mCherry (medial PFC), CAV-2-Cre (NAc)	Medial PFC→NAc	Rat: WT	Di-stimulation during progressive ratio testing: no effect on progressive ratio responding for cocaine but ↑ motivation to seek cocaine during extinction and cocaine- primed reinstatement. Di-stimulation prior to reinstatement: ↓ cocaine- primed reinstatement - only in rats that had Di-stimulation during progressive ratio (normalize) ↓ context-induced ethanol reinstatement
		Marchant et al. (2016a)	AAV1-hSyn-DIO-HA-KORD- IRES-mCitrine (ventral Subiculum), AAV5-Cre (NAc shell)	Ventral subiculum →NAc	Rat: WT	
		Prasad and McNally (2016)	AAV5-hSyn-HA-hM3Dq-IRES- mCitrine, AAV5-hSyn-HA- hM4Di-IRES-mCitrine	Ventral palladium → STN, VTA	Rat: WT	Ventral pallidum Di-/Ds- stimulation: 1/1 cue- and prime-induced reinstatement of extinguished alcohol seeking Di in ventral pallidum (unilateral) + VTA or STN (contralateral): ↓ cue- and prime-induced alcohol reinstatement
		Heinsbroek et al. (2017)	AAV2-hSyn-DIO-hM4Di- mCherry, AAV2-hSyn-DIO- rM3Ds-mCherry, AAV2- hSyn-HA-hM4Di-IRES- mCitrine	NAc for Cre-dependent vectors (dMSN or iMSN) Ventral pallidum for the non-Cre dependent vector)	Mouse: D ₁ R-Cre, D ₂ R- Cre	dMSN/iMSN Ds-stimulation in NAc: 1*/no effect on cue- induced reinstatement in iMSN Di-stimulation in NAc: 1* cue-induced reinstatement *reversed by simultaneous Di-stimulation in ventral pallidum

(continued)

TABLE 1—Continued

Area of Research	Test Model	Reference	Chemogenetic Vector *DIO: double-flxed inverted open reading frame (Cre-dependent)	Target Area (Cells)	Rodent Species/Strain	Effect Mediated by Chemogenetic Intervention
Mood, nociception, and cognition		Wunsch et al. (2017)	AAV5-hSyn-hM4Di-mCherry, AAV5-hSyn-DIO-hM4Di- mCherry (Midline thalamic nuclei (MTN)), CAV-2-Cre (NAc)	MTN, MTN→NAc	Rat: WT	MTN Di-stimulation: ↓ cue- induced and drug-primed reinstatement of cocaine seeking MTN→NAc Di-stimulation: ↓ cocaine- primed reinstatement, ↑ cue-induced reinstatement NAc core and shell Dq/Di- stimulation: ↓/no effect on ethanol consumption NAc core Dq/Di- stimulation: ↓/↑ ethanol consumption NAc shell D Dq/Di-stimulation: no effect on ethanol consumption ↓ anxiogenic-like and depressive-like phenotype of Cdk5 KO mice ↓ immobility time in forced swim test also in control mice (DATCre) ↓ exercise-induced hypoaesthesia (following partial sciatic nerve ligation)
		Purohit et al. (2018)	AAV2-hSyn-HA-hM3Dq-IRES- mCherry, AAV2-hSyn-HA- hM4Di-IRES-mCherry	NAc core + shell, NAc core, NAc shell,	Mouse: WT	
		Zhong et al. (2014)	AAV8-hSyn-DIO-rM3Ds- mCherry	VTA (DAergic)	Mouse: Cdk5 KO (DAT- Cre background)	
	Open-field, elevated- plus maze, novelty suppressed feeding, sucrose preference test, forced swim test Neuropathic pain model, von Frey and thermal stimulation pain assays	Wakaizumi et al. (2016)	AAV10-hSyn-hM4Di-mCherry (VTA), NeuRet-Cre* (NAc) *lentiviral vector for neuron- specific retrograde gene transfer	VTA (DAergic), VTA→NAc	Mouse: DAT-Cre, WT	
	Hot plate, tail-flick assays	Li et al. (2016)	AAV8-Eflα-DIO-hM3Dq- mCherry	Ventral periaqueductal gray/dorsal raphe region (DAergic)	Mouse: TH-Cre	↓ supraspinal nociceptive effect in hot-plate test, no effect on locomotion and tail-flick test ↓ conditioned place aversion to a systemic inflammation agent VTA or SN pars compacta Dq-stimulation: ↑ trial omissions, SN pars compacta Dq-stimulation: ↓ task accuracy
	CPP test	Fritz et al. (2016)	AAV8-hSyn-DIO-hM3Dq- mCherry	VTA (DAergic)	Mouse: DATCreER* *Tamoxifeninducible	
	5-choice serial reaction time task	Boekhoudt, Voets et al. (2017)	AAV5-hSyn-DIO-hM3Dq- mCherry	VTA/SN pars compacta (DAergic)	Rat: TH-Cre	
	Barnes maze, Acoustic startle reflex, Social approach test Electrophysiology	Locke et al. (2018)	AAV ^{NA} -DIO-hM4Di-YFP	Lateral cerebellar nucleus (DlR expressing neurons)	Mouse: D ₁ R-Cre	↓ spatial navigation, working memories, ↓ prepulse inhibition of acoustic startle response Ds-stimulation: mimicked D ₂ R-induced activation- elicits afterdepolarizations Inhibition of relief learning
	Fear conditioning paradigm	Robinson and Sohal (2017)	AAV5-hSyn-DIO-hM3Dq- mCherry, AAV5-hSyn-DIO- rM3Ds-mCherry, AAV5- hSyn-DIO-hM4Di-mCherry AAV8-CaMKIIa-hM4Di- mCherry	Medial PFC (D2R expressing neurons)	Mouse: D ₂ R-Cre	
		Mayer et al. (2018)	AAV8-CaMKIIa-hM4Di- mCherry	Posterior medial VTA → NAc shell (DAergic)	Rat: WT	
		Bouchet et al. (2018)	AAV5-hSyn-DIO-hM3Dq- mCherry	SN pars compacta (DAergic)	Rat: TH-Cre	No effect on fear extinction acquisition, but enhanced fear extinction memory and blocked the renewal of fear in a novel context

of chemogenetics (Smith et al., 2016; Whissell et al., 2016).

A. Locomotor Activity

1. Dopamine Activity Correlates with Locomotor Activity—Proof-of-Concept Studies. Classic behavioral pharmacology experiments have indicated that locomotor hyperactivity can be attributed to increased striatal DA levels (Pijnenburg et al., 1976). Therefore, initial DREADD investigations were carefully applied to behavioral neuroscience with conceptual proof-of-concept studies in which researchers applied DREADDs to confirm pharmacologic manipulation of general neuronal populations with immediate effects on behaviors such as locomotor activity. A direct relationship between the brain DA system and regulation of locomotor activity in rodents was supported by expression of hM3Dq within midbrain (VTA/SN) DAergic neurons of DAT-Cre mice (Wang et al., 2013). CNO application increased the firing frequency of DA neurons in midbrain slices, and this finding was paralleled by strong dose-dependent hyperactivity in mice challenged with CNO (Wang et al., 2013). Likewise, in DAT-Cre mice also with hM3Dq expressed in DA midbrain neurons, the active phase of locomotor day rhythms was extended following CNO administration. This result resembled the same hyperdopaminergic phenotype that has been observed following amphetamine administration or modeled in the DAT-knockout (KO) mouse (Blum et al., 2014). With the use of a dual viral strategy to allow better spatial resolution of specific DA projections, analogous results were demonstrated in rats (Boender et al., 2014). Here, projection-specific expression of hM3Dq in VTA-NAc neurons was obtained by combined use of the retrogradely transported CAV2-Cre virus injected into the NAc and an AAV-encoding Cre-dependent DREADD vector injected into the VTA of WT rats. This study also showed long-lasting increases in locomotor activity (3–7 h). These findings were subsequently substantiated by selective expression of viral hM3Dq in TH-Cre rats, and engagement of TH-positive (i.e., considered DAergic) neurons of the VTA, but not the SN pars compacta, demonstrated similar long-lasting hyperlocomotion (Boekhoudt et al., 2016). This is consistent with older pharmacological data and literature supporting a key role of VTA DA neurons in the regulation of hyperactivity, while SN DA neurons appear to be more involved in other aspects of locomotion such as motor coordination (Delfs et al., 1990; Barter et al., 2014). We also recently demonstrated clear bidirectional regulation of locomotor activity during habituation to a novel environment in TH-Cre mice injected in VTA with Cre-dependent rM3Ds and hM4Di (Runegaard et al., 2018). In this study, CNO pretreatment increased and reduced locomotor activity in rM3Ds- and hM4Di-expressing mice, respectively, while CNO had no effects in WT animals (Fig. 4).

Exploratory locomotor activity has also been shown to be reduced by SalB or CNO in WT rats expressing KORD or hM4Di, respectively, in the midbrain (Marchant et al., 2016b). Locomotor activity can be indirectly manipulated by expressing KORD in midbrain GABAergic interneurons, which upon SalB administration resulted in disinhibition of VTA DA neurons and increased locomotor activity (Vardy et al., 2015). Summarized, these studies substantiate already compelling evidence that DA signaling directly regulates locomotor activity and confirms well-established pharmacological evidence that the mesolimbic pathway, comprising VTA to NAc projecting DA neurons, enforces this effect (Delfs et al., 1990; dela Peña et al., 2015). Also, the studies support local circuits within the VTA that act as central regulatory components of locomotor activity.

2. The Relative Output-Activity from Direct Medium Spiny Neurons and Indirect Medium Spiny Neurons Determines Locomotor Activity. A simple direct relationship between DA and locomotor activity, as described above, is, however, challenged by the hypofunctional DA system seen in ADHD patients, although this has been suggested to be explained by alterations in post-synaptic DA receptor signaling (Volkow et al., 2009; Napolitano et al., 2010). Accordingly, many chemogenetic studies have focused on mimicking and/or manipulating DAergic receptor signaling directly in striatal MSN subpopulations, encompassing Gs-coupled D₁R expressing striatonigral neurons (dMSNs), belonging to the direct pathway, and Gi-coupled D₂R expressing striatopallidal neurons (iMSNs), belonging to the indirect pathway. In a transgenic mouse expressing Gs-coupled rM3Ds throughout all iMSNs, exploratory locomotor activity was strongly reduced upon CNO administration (Farrell et al., 2013; Chu et al., 2017; Bouabid and Zhou, 2018). This was recently linked to inhibition of globus pallidus neuronal firing as a consequence of rM3Ds-mediated increased excitation of the iMSNs (Bouabid and Zhou, 2018). This suggests that AC/cAMP signaling in MSNs plays a significant role in regulating locomotor activity, which is in line with a chemogenetic study demonstrating an important role of Gi-coupled signaling in dMSNs in regulating locomotion (Moehle et al., 2017). It was correspondingly shown that hM4Di stimulation in iMSN of anatomically defined striatal areas, including the dorsomedial striatum and NAc, increased locomotor activity (Carvalho Poyraz et al., 2016). Moreover, acute CNO administration led to increased and decreased locomotor activity in mice with dorsostriatal expression of hM3Dq in dMSNs of D₁R-Cre mice and in iMSNs of D₂R-Cre or A2a-Cre mice, respectively (Bellocchio et al., 2016; Alcacer et al., 2017). Interestingly, dissection of accumbal efferents revealed a stronger influence of the indirect pathway on locomotor activity. While significant bidirectional control of basal locomotor activity was

Fig. 4. Bidirectional effects on locomotor habituation by hM4Di and rM3Ds stimulation in VTA [modified and adapted from Runegaard et al. (2018)]. (A) AAV carrying hM4Di or rM3Ds was injected into the VTA of TH-Cre or WT mice. (B) Experimental timeline to assess habituation of novelty-induced locomotor activity. CNO was injected intraperitoneally in the home cage 30 min prior to placement into the center of an open field where locomotor activity was recorded and tracked for 90 min. (C) Time course of the 90-min habituation of novelty-induced locomotor activity in an open field 30 min after CNO injections in control (black), hM4Di (magenta), or rM3Ds (blue) expressing mice. Control mice injected with vehicle were included and compared with CNO-treated control mice to assess the effect of CNO alone, as shown by a dashed gray line. CNO demonstrated bidirectional control of novelty-induced exploratory activity during habituation in hM4Di and rM3Ds mice while leaving WT mice unaffected (**** $P < 0.0001$; *** $P < 0.001$; ** $P < 0.01$; * $P < 0.05$ relative to control). (D) Representative tracks and graphs of total distance traveled during the habituation. Analysis of the total distance traveled showed significant reduction and increase in novelty-induced exploration following CNO in hM4Di and rM3Ds mice, respectively, compared with WT controls (* $P < 0.05$; ** $P < 0.01$ relative to CNO control, $N = 15$ hM4Di CNO, 12 rM3Ds CNO, 13 control CNO). Data are shown as means + S.E.M.

demonstrated by NAc iMSNs with reduced and increased activity following hM3Dq or hM4Di stimulation, respectively, only slight increases in locomotion were observed with hM3Dq activation of dMSNs in NAc (Zhu et al., 2016). These observations collectively support a major influence of D_2R activation in iMSNs on the activity of dMSNs and thereby on locomotor activity. Notably, this is consistent with a new theory of lateral inhibition of iMSNs onto dMSNs (Dobbs et al., 2016) and also with evidence from rodent KO studies, indicating that MSN D_1R and D_2R interactions control locomotor activity in a synergistic, balanced manner (Welter et al., 2007). For a comprehensive review of DA receptor KO mouse behaviors, see Holmes et al. (2004).

3. Locomotor Activity—A Simple Behavior with Multiple Levels of Regulation. Midbrain DA neurons receive and integrate inputs from multiple brain regions, as well as from local interneurons. DREADD-induced

manipulations of different midbrain DA-projection neurons confirm the importance of these neurons for regulating DA activity, release, and behaviors including locomotion. As mentioned above, chemogenetic inhibition by KORD in VTA GABAergic interneurons increased locomotor activity as a consequence of VTA DA disinhibition (Vardy et al., 2015). GABAergic neurons residing within the lateral hypothalamus were recently found to alter locomotor activity via direct and indirect regulation of the VTA, i.e., hM3Dq stimulation of both VTA-projecting (i.e., GABA-expressing) neurons and non-VTA-projecting (i.e., galanin-expressing) neurons increased locomotor activity (Qualls-Creekmore et al., 2017). Similarly, CNO-induced release of neurotensin in the mouse VTA, with hM3Dq expression confined to lateral hypothalamic neurons expressing neurotensin, induced prolonged DA-dependent locomotor activity and DA release in the NAc via neurotensin receptor

1 expressed on VTA DA neurons (Patterson et al., 2015). Furthermore, morphine-induced hyperlocomotion was regulated by DREADD-mediated modulation of rostromedial tegmental nucleus GABA interneurons targeting NAc-projecting DA neurons (Wasserman et al., 2016). While there was no effect of either DREADDs on basal locomotor activity, hM4Di and hM3Dq prevented morphine-induced hyperlocomotion, possibly by changing the firing thresholds for GABA interneurons and thereby altering their inhibitory effect on NAc-projecting DA neurons (Wasserman et al., 2016). Note that chemogenetic interventions of locomotor response to other drugs such as cocaine and amphetamine are addressed in section II.C.1.

4. Dysfunctional Motor Control in Parkinson's Disease. In contrast to locomotor activity, which appears to be regulated predominantly by midbrain DA neurons projecting to ventral parts of the striatum, motor control and initiation of movement are more dependent on dorsal striatal DA neurotransmission. This is particularly evident in PD, where major symptoms include bradykinesia, akinesia, tremors, and movement rigidity (Vazey and Aston-Jones, 2013). These motor disabilities are associated with major loss of DAergic neurons within the SN, resulting in progressive loss of dorsostriatal DA input. A classic and commonly used rodent/primate model of PD is unilateral 6-hydroxydopamine (6-OHDA)-induced lesion of striatal terminal neurons that is believed to recapitulate key changes in basal ganglia circuitry and pharmacology observed in patients with PD (Ungerstedt and Arbuthnott, 1970). The resulting hemi-parkinsonian rodents exhibit severe behavioral imbalances that can be restored by L-3, 4-dihydroxyphenylalanine (L-DOPA), the DA precursor applied for therapeutic use in PD patients. In nonlesioned control mice, unilateral expression and stimulation of hM3Dq in dMSNs or iMSNs of the dorsal striatum induced contra- and ipsilateral rotations, respectively (Alcacer et al., 2017). In lesioned hemi-parkinsonian mice, unilateral stimulation of dMSN in the DA-denervated striatum mimicked the therapeutic effect of L-DOPA on contralateral forelimb use, as assessed in a cylinder test, while stimulation of iMSN prevented L-DOPA's effect in these mice (Alcacer et al., 2017). The findings are noteworthy, and perhaps relevant for future clinical intervention, as prolonged L-DOPA treatment is associated with dyskinesia, a debilitating and treatment-limiting condition affecting up to 80% of PD patients within 10 years (Manson et al., 2012). In the Alcacer et al. (2017) study, it was also shown that L-DOPA-induced dyskinesia or involuntary movements in hemi-parkinsonian mice were reduced by hM3Dq expressed in iMSNs on the denervated striatal site. However, stimulation of rM3Ds or hM3Dq in dMSNs of the denervated striatal site induced abnormal involuntary movements, with the most severe effect seen upon rM3Ds stimulation that

resulted in L-DOPA-like induced dyskinesia (Alcacer et al., 2017). This result fully complements recent findings using a C-fos driver mouse line and optogenetics to probe the origin of L-DOPA-induced dyskinesia (Girasole et al., 2018).

Fetal DA neuron transplantation trials in PD patients suggest that motor disturbances can be normalized in some patients where L-DOPA treatment is inefficient or associated with dyskinesia. The efficiency of the neuronal transplantation, however, is highly variable (Barker et al., 2013). Chemogenetics were recently employed in rodent models to investigate whether the therapeutic efficacy can be improved by activating the excitatory hM3Dq within engrafted neurons. Strikingly, such manipulations did lead to increased excitability and DA release within induced or neuroblastic DA neurons transplanted into the dorsal striatum of 6-OHDA-lesioned mice. Moreover, the manipulations rescued motor deficits in mice (Chen et al., 2016) and rats (Dell'Anno et al., 2014; Aldrin-Kirk et al., 2016). Aldrin-Kirk et al. (2016) even went on to simultaneously express and activate hM3Dq and rM3Ds in the grafted cells. With this, they induced dyskinesia similar to what develops as a side-effect in some patients following transplantation of DAergic neurons. They suggested a mechanism for graft-induced dyskinesia that involves 5-HT₆ receptors causing dysplastic DA release from transplants by activating Gs-signaling that in turn counteracts autoinhibitory control by D₂Rs (Aldrin-Kirk et al., 2016).

A recent chemogenetic study adds further support to the hypothesis that parkinsonian-related motor deficits are caused by maladaptive, relative regulation of the direct and indirect pathways of basal ganglia circuitry, rather than by reduced absolute DA levels (Parker et al., 2016). The severity of motor deficits observed in 6-OHDA-lesioned mice was correlated with a reversed relative strength of excitatory thalamic drive onto direct and indirect pathway MSNs with a bias toward the indirect pathway. Inhibition of this thalamostriatal input, which drives maladaptive asymmetric activation of basal ganglia by activating hM4Di in thalamostriatal terminals, restored motor function in DA-depleted mice (Parker et al., 2016). Likewise, in 6-OHDA-lesioned rats, stimulation of cholinergic interneurons in the striatum via hM3Dq/rM3Ds potentiated the therapeutic effects of low-dose L-DOPA (i.e., 1 mg/kg) and facilitated L-DOPA-induced abnormal involuntary movement after moderate and high doses (i.e., 3 and 9 mg/kg) of L-DOPA—effects that appeared to be driven primarily through the iMSN pathway since the same effect was induced by the D₂ agonist quinpirole (Aldrin-Kirk et al., 2018). Further support for maladaptive regulation between the direct and indirect pathways was achieved in a study in which bradykinesia was modeled in a genetic mouse model with selective deletion of the D₂R in iMSNs (Lemos et al., 2016). The degree of

bradykinesia correlated with the enhanced GABAergic transmission downstream of iMSNs, and GABAergic tone and locomotor control were restored through stimulation of Gi-signaling in iMSNs by activating hM4Di expressed in iMSNs of the dorsal striatum and NAc to mimic D₂R function (Lemos et al., 2016). Lastly, a chemogenetic approach has been applied to investigate the role of striatal striosome and matrix compartments in procedural learning and use of motor skill. The study supported that there is no synaptic connectivity between these two compartments. Nevertheless, by selective transduction of matrix neurons only with hM4Di, evidence was obtained for a role of dorsolateral striatal matrix neurons in forepaw use (Lopez-Huerta et al., 2016).

B. Motivation and Reward-Guided Behaviors

DAergic-mediated regulation of motivation and reward-guided behaviors has garnered much attention not least due to its importance in pathologic states such as addiction and obesity (Baik, 2013). The below mentioned DREADD studies have assessed DA behavioral regulation in rodent test models in both homeostatic and maladaptive reward-related states that take advantage of an organism's desire to survive (Salamone et al., 2012). As such, operant conditioning and learning tasks involve reward reinforcement and punishment processes that lead to behavioral acquisition, which regulates the probability, proximity and availability of relevant stimuli in rodent tasks such as progressive ratio scheduling (Richardson and Roberts, 1996), operant learning (Todd et al., 2014), and cue/sign-tracking (Salamone et al., 2012).

1. *Substantiating Dopamine as a Key Regulator of Motivation.* Motivation deficits represent a key feature of psychiatric disorders with DA being a critical neural substrate (Baik, 2013). Through the use of progressive ratio scheduling to assess motivational behaviors, Boender et al. (2014) demonstrated that hM3Dq-mediated activation of the mesolimbic DAergic pathway, as targeted through use of a Cre-recombinase expressing CAV2 vector, increased progressive ratio performance, and hence effort exertion. The same research group refined their methodological approach to show that this enhanced motivation was specifically due to increased action response initiation and not due to other forms of motivational behaviors (Boekhoudt et al., 2018).

2. *The Role of Direct Medium Spiny Neurons and Indirect Medium Spiny Neurons in Regulating Reward-Motivated Behaviors.* Given its importance in mediating DAergic neurotransmission, the dorsal striatum is known to play a critical role in reward-guided decision making, but the precise circuits involved have not been fully elucidated. In one study by Ferguson and Neumaier (2012), it was observed that transient hM4Di activation of dMSNs within the rat dorsal striatum

impaired acquisition of an operant learning task assessing reward reinforcement learning, as supported by significantly lower lever pressing rates for a reward. In another study, the same group further investigated the role of the dorsomedial striatum in action-outcome learning using a high-versus-low reward discrimination task (Ferguson et al., 2013). Rats learned to associate lever presses with one daily alternating lever for a high-reward outcome and the other for a low-reward outcome. Upon chemogenetic modulation of dMSN signaling via activation of hM4Di or rM3Ds during training sessions, no effect was seen on task acquisition or subsequent performance, i.e., 24 h after the last training session. However, hM4Di activation during training sessions significantly impaired stable retention of task information, as these rats demonstrated a reduced preference for the high-reward lever and needed more trials to reach criterion, that being the choice of the high-reward lever 75% of all trials, 1 week after the last training session. Activation of rM3Ds in dMSNs produced the opposite effects and improved stable retention of this decision-making strategy 1-week posttraining. These findings provided strong evidence that activation of dMSNs in the dorsomedial striatum is essential for retaining test-specific information regarding reward-guided behaviors. Moreover, the fact that these chemogenetic manipulations of dMSNs affected behaviors 1 week post-CNO administration suggested that significant neuronal plasticity changes took place within these neurons during acquisition of this task (Ferguson et al., 2013).

DREADD approaches, together with an operant lever-pressing task with progressive ratio schedules, were used to assess the influence of iMSNs in NAc and dorsomedial striatum on motivational processes. Activation of hM4Di expressed in iMSNs of either the dorsomedial striatum or NAc reversed motivational deficits in mice genetically manipulated to overexpress D₂R_s and also in WT controls. These effects were due to enhanced response initiation without goal-guided efficiency, motivational vigor, or sensitivities relating to up- or down-valuation of reward being affected (Carvalho Poyraz et al., 2016). Hence, Gi-signaling in iMSNs appeared to increase motivation to work for reward, probably as a result of reduced GABAergic tone in target areas or directly onto neighboring dMSNs causing disinhibition (Dobbs et al., 2016). This is further supported by a study by Gallo et al substantiating a role of the ventral pallidum. It was shown in this study that activation of hM4Di expressed in NAc iMSNs projecting to the ventral pallidum increased motivation to work for a salient reward in a lever-pressing task with progressive ratio schedule (Gallo et al., 2018). Of additional interest, selective hM3Dq-stimulation of NAc interneurons expressing nitric oxide synthase was demonstrated to increase sucrose seeking after extinction and cue-induced reinstatement (Bobadilla et al., 2017).

Together, these studies suggest that DA-dependent, reward-motivated behaviors are regulated by multiple pathways that depend on the output of both dorsal striatal and NAc MSNs. Importantly, while the chemogenetic manipulations of MSN activity described above tend to be considered a direct reflection of DA transmission with the Gi- and Gs-coupled DREADDs directly modulating DA receptor signaling, it should be noted that these neurons express multiple different GPCRs that signal through similar cascades. Accordingly, the behaviors studied are not regulated solely by DA transmission but also by other transmitter systems acting through different sets of GPCRs.

3. Feeding. Food activates neural circuits associated with reward and DAergic signaling, particularly in the NAc, midbrain, and amygdala. It is widely believed that the DA mesolimbic pathway facilitates associative learning between food-related rewards and feeding-related environments, with food or food-related cues initiating phasic mesolimbic DAergic firing and behaviors associated with acquiring food (Baik, 2013). Boekhoudt et al. (2017a) investigated whether chemogenetic activation of midbrain DAergic neurons promoted or reduced feeding to establish a causal relationship between DAergic activity and food intake. They measured home-cage feeding patterns while concurrently stimulating hM3Dq in the SN pars compacta or selective VTA subpopulation projections toward the NAc, amygdala, or PFC. They found that general stimulation of hM3Dq in VTA reduced meal size, while at the same time increasing locomotor activity. Specific mesolimbic pathway DA activation disrupted feeding patterns by reducing mean meal size and duration but increased the frequency of meal initiations (Boekhoudt et al., 2017a). Although, the mesolimbic DA pathway seemed to both promote and reduce food intake in that study by facilitating feeding initiation and cessation behaviors (Boekhoudt et al., 2017a), chemogenetic stimulation in the same mesolimbic pathway in another study prevented anorexic weight loss in an activity-based rat model of anorexia (Foldi et al., 2017). When rats were given time-limited daily access to food, hM3Dq stimulation of VTA DAergic neurons projecting to the NAc increased food intake and food anticipatory activity and blocked anhedonia-associated weight loss and subsequent death (Foldi et al., 2017).

Specific chemogenetic activation of dMSNs or iMSNs within the NAc was done to assess food intake under both low and high-energy expenditure (i.e., a running wheel was placed in the home cage) conditions. The data showed that hM3Dq and hM4Di activation of NAc dMSNs enhanced and diminished food intake, respectively, in high-expenditure conditions when the running wheel was placed in the animals' home cages for 14 days. Interestingly, the same study found that NAc dMSN hM3Dq-stimulation reduced overall food intake in low expenditure conditions (i.e., open field test) over the

course of 24 h (Zhu et al., 2016). These studies support and complement the hypothesis that DAergic pathways, particularly the mesolimbic, regulate homeostatic energy expenditure and feeding patterns. In addition, altered DAergic signaling patterns, induced by chemogenetic manipulations, were consistently found to disrupt feeding patterns, indicating that tightly controlled and optimal DA neurotransmission exists to regulate feeding.

As outlined below, there is strong evidence suggesting that serotonin directly acts on VTA DA neurons to reduce food intake, linking the two transmitters together in regulation of feeding behaviors. As well as finding that stimulation of hM3Dq expressed in VTA DA neurons inhibited binge-like eating in mice on a high-fat diet, Xu et al. (2017) demonstrated that the 5-HT_{2C} receptor agonist lorcaserin, which has been approved for obesity treatment (Martin et al., 2011), and the 5-HT-releasing agent fenfluramine inhibited these eating patterns via 5-HT_{2C} receptor activation. This effect was directly linked to an increase in VTA DAergic signaling and further supported by the finding that VTA DA hM4Di activation prevented fenfluramine from suppressing binge-like eating patterns (Xu et al., 2017). The same group subsequently demonstrated that hM3Dq-mediated activation of VTA 5-HT_{2C} receptor-expressing neurons reduced ad libitum food intake, as well as lowered progressive ratio responding for sugar pellets and incentive value for food (Valencia-Torres et al., 2017). The studies substantiate that 5-HT_{2C} agonism inhibits food-related motivational processes at least in part via direct effects on midbrain DAergic transmission.

Application of chemogenetics has also shown that neurons within the nucleus tractus solitarius in the brain stem can regulate food intake by affecting VTA DA neuronal activity. Activation of hM3Dq in these neurons suppressed high-fat chow diet intake by diminishing signaling of VTA DAergic neuron projecting to the NAc shell (Wang et al., 2015). Notably, it was found that this likely was the result of increased release of glucagon-like-peptide 1 acting through GLP-1 receptors on DA VTA neurons, leading to reduced excitatory synaptic strength (Wang et al., 2015). Likewise, hM3Dq-mediated activation of the lateral hypothalamic GABAergic neurons caused an overall increase in chow diet consumption as a consequence of direct VTA innervation (Qualls-Creekmore et al., 2017). These studies highlight how different brain regions and neurochemical systems can modulate DA-dependent signaling and behaviors and how chemogenetics can aid elucidation of previously unknown pathways.

4. Mating. During sexual behaviors, VTA μ -opioid receptor-expressing neurons are activated and DA is released into NAc and striatum (Damsma et al., 1992; Bassareo and Di Chiara, 1999; Balfour et al., 2004). In rodents, sexual behavior is normally assessed by

pairing and allowing a male to mate with a female subject. The latency and number of investigations, intromissions, and ejaculations allow for measurement of sexual motivation, performance, memory, and reward. In rats with hM4Di expression confined to TH-positive neurons in the VTA, CNO did not affect, however, such sexual behavior (Beloate et al., 2016). Moreover, although CNO prevented VTA DA neuron activation and C-fos induction in these neurons in response to sexual behaviors, CNO had no effect on mating-induced C-fos expression in VTA projection areas, including NAc, PFC, and basolateral amygdala. CNO was nonetheless found to impair sexual experience-induced cross-sensitization of amphetamine conditioned place preference (CPP). Thus, activation of VTA DA neurons appears not to be critical for initiation or expression of sexual behavior. Rather such activation, and thereby mesolimbic DA, increases the sensitivity to addictive drugs upon loss of the natural reward via mesolimbic plasticity established during the natural reward experience. Together, the study substantiates how DAergic VTA neurons can regulate the cross-sensitization of natural (i.e., sexual experience) and drug-related rewards (Beloate et al., 2016).

In another study, the importance during sexual behavior of dorsostriatal dMSN and iMSN activity was assessed by using genetically encoded, fluorescence resonance energy transfer-based PKA and ERK sensors together with DREADDs. The sensors were expressed in dMSNs and iMSNs, and it was found that PKA and ERK activities in these neurons rapidly switched when the male mice shifted between active and inactive phases of mating reactions. To investigate the mechanism underlying this rapid change, hM4Di or rM3Ds DREADDs were expressed in striatal dMSNs and iMSNs together with the PKA biosensor. In hM4Di-expressing dMSNs or iMSNs, CNO reduced PKA activity and concomitantly caused suppression or augmentation of positive mating reactions, respectively. In rM3Ds-expressing dMSNs or iMSNs, CNO increased PKA activity concomitantly with enhancement or suppression of positive mating reactions, respectively. Hence, the dMSNs and iMSNs of the dorsal striatum appear to differentially control mating reactions in male mice by directly regulating PKA activity (Goto et al., 2015).

5. Sleep-Wake Patterns and Arousal. Activation of VTA DAergic neurons by hM3Dq was shown to increase wakefulness periods while reducing sleep-associated behaviors, such as nest building, in both stages of the light/dark cycle in a D₂R-dependent manner (Oishi et al., 2017a; Sun et al., 2017). Further investigations revealed that NAc D₁R- and NAc core adenosine A_{2a} receptor-expressing neurons also regulated these sleep-wake patterns (Oishi et al., 2017b; Luo et al., 2018). Activation of D₁R expressing neurons in the NAc by hM3Dq-stimulation prolonged wakefulness and concurrently

reduced food intake, while stimulation of the inhibitory hM4Di suppressed arousal and induced sleep behaviors (Luo et al., 2018). Another approach, albeit with the same chemogenetic tools, demonstrated that adenosine A_{2a} receptor-expressing indirect pathway neurons in the NAc core can regulate slow-wave sleep, as hM3Dq or hM4Di stimulation of these neurons either induced or prevented slow-wave sleep, respectively (Oishi et al., 2017b). The studies highlight the importance of mid-brain DA neuronal circuits for regulating arousal, both discovering new DA-dependent neural circuits critical for the induction and maintenance of wakefulness, projecting to the lateral hypothalamus (Luo et al., 2018) and ventral pallidum (Oishi et al., 2017b), respectively.

It was furthermore observed that acute hM3Dq-activation in dorsostriatal dMSNs increased wakefulness while reducing the time spent in the slow-wave and rapid eye movement sleep stages. Acute hM3Dq-activation of iMSNs, however, increased slow-wave phase sleep while chronic activation resulted in enhanced wakefulness (Bellocchio et al., 2016). Expectedly, one's arousal state affects motivation with the above studies revealing a direct effect of DA activity on wakefulness and sleep-wake patterns. More direct evidence for DA as a substrate integrating reward-based motivation and arousal was obtained from data indicating that DA is critical for maintaining wakefulness even in the presence of relevant salient stimuli (Eban-Rothschild et al., 2016; Cho et al., 2017).

By using fiber photometry to measure calcium activity and polysomnographic recordings to assess wakefulness, it was observed that hM4Di-mediated inhibition of DAergic neurons in the VTA and dorsal raphe nuclei decreased wakefulness and increased sleep-related behaviors and length of sleep. This decrease in arousal occurred even when stimuli, such as accessible palatable food, a female mouse, or predator odors, were present in the same or in a neighboring cage (Eban-Rothschild et al., 2016; Cho et al., 2017). The data emphasize a plausible role governed by the DAergic system in regulating the interaction between sleep-wake patterns and reward-based motivation.

6. Immunity. When VTA neurons were acutely activated via stimulation of hM3Dq receptors expressed in these neurons 24 h before an *Escherichia coli* immune challenge, enhanced innate and adaptive immune responses were exhibited 3- and 7-days postchallenge. In addition to increased host defense capabilities, the same VTA DA neuronal activation resulted in enhanced CPP toward the conditioned chamber. These findings indicate that VTA DAergic neurons balance their control of reward-guided behaviors with immunity or, at least, that there might be a causal relationship between enhanced VTA activity and the host's immune response to a bacterial infection (Ben-Shaanan et al., 2016).

C. Addiction

Drugs of abuse, including the psychostimulants cocaine and amphetamine that act directly on the DA system, induce substantial synaptic modifications that hijack the natural reward system and cause the transition to substance addiction (Baik, 2013). Increased DA levels within the NAc have been linked to hyperlocomotion as well as to an initial perception of reward and subsequent reward prediction error guiding goal-directed behaviors (Schultz, 2007). Although the elevations in accumbal DA levels are related to both psychostimulant-induced hyperactivity (acute as well as sensitized) and general reward perception (which in rodents can be assessed indirectly by CPP or self-administration paradigms), these behaviors are likely segregated because not all addictive drugs induce hyperactivity. It has been hypothesized, however, that psychostimulant-induced locomotor hyperactivity in rodents reflects the perception of reward and could be reflective of addiction vulnerability (Deminere et al., 1989; Piazza et al., 1989). Consistent with this hypothesis, a greater acute locomotor response to psychostimulants was found to be predictive of psychostimulant self-administration (Piazza et al., 1989) and high drug intake (Piazza et al., 2000). The rewarding effect/addictive properties of drugs of abuse and their influence on subsequent drug-related behaviors, including drug-seeking and reinstatement, are commonly assessed by CPP and self-administration paradigms. In addition to assessing relapse following withdrawal (induced by either a cue or a prime-injection of the rewarding stimuli), the self-administration paradigm may assess compulsive drug intake (continued self-administration despite negative consequences, such as foot shock), which is considered a major hallmark of addiction. Of further interest, we recently published a single exposure place preference protocol that allows assessment of the initial perception of reward in rodents and thereby represents an attractive model for further investigating initial drug effects and dissect reward and locomotor-enhancing responses various drugs (Runegaard et al., 2017a). Chemogenetics has been used widely in various experimental set-ups to further dissect and shed light on behaviors and underlying changes to neural networks related to addiction including acute locomotor response, behavioral sensitization, CPP, and self-administration.

1. Locomotor Sensitization. Locomotor sensitization is a lasting, increased locomotor response observed following repeated, noncontingent administration of psychostimulants. This is believed to reflect the synaptic adaptive changes that occur with repeated drug intake and to be associated with the transition from initial drug use to addiction (Hooks et al., 1991a,b; Piazza et al., 1991; Belin and Deroche-Gamonet, 2012). Not surprisingly, DREADDs have been applied to the DA system to examine the influence of dMSNs and

iMSNs during repeated drug administrations and after drug-induced behavioral sensitization. The results confirmed previous pharmacological and transgenic pieces of evidence on the opposing roles of the indirect and direct MSN pathways (Lobo and Nestler, 2011). Indeed, activation of hM4Di in iMSNs or dMSNs enhanced or attenuated, respectively, behavioral sensitization to amphetamine (Ferguson et al., 2011). Similarly, behavioral sensitization was prevented when Gi-induced signaling was opposed by activating rM3Ds in iMSNs (Farrell et al., 2013).

The findings support that iMSN activity is a driving force underlying behavioral sensitization (Farrell et al., 2013), that is, the experiments indicate that locomotor sensitization, similarly to general locomotor activity (see section II.A.2) depends on a relative output signal from striatal MSNs following repeated drug administration. Interestingly, acute and sensitized responses to psychostimulants seem to be affected differently by the same DREADD manipulations. Activation of hM4Di in VTA neurons strongly inhibited the acute response to cocaine, but this neither prevented the development nor the expression of a sensitized response (Runegaard et al., 2018). On the contrary, hM4Di activation in iMSNs during amphetamine sensitization had no effect on the acute response or during initiation, but prevented expression of behavioral sensitization during a challenge dose 1 week later (Ferguson et al., 2011). This suggests that the responses are driven by separate and to some extent independent mechanisms, possibly segregated by pre- and postsynaptic effects. Moreover, the potential influence and role of coreleased neurotransmitters from heterogeneous populations of midbrain DA neurons must be taken into consideration to understand fully the mechanisms underlying the segregation of these behaviors (Poulin et al., 2018).

2. Reward Perception, Drug-Cue Associations, and Relapse Behaviors. The perception of reward is linked to increases in DA levels, and such increases are considered the common mechanism responsible for the addictive properties of drugs of abuse (Wise and Bozarth, 1985; Di Chiara and Imperato, 1988; Koob and Le Moal, 1997). How the initial increase in DA conveys into adaptive changes that only in a subpopulation of subjects induce the transition from recreational use to addiction is still poorly understood. Based on the fact that females tend to transition faster to addiction and experience greater difficulties remaining abstinent (Becker and Koob, 2016), Calipari et al. (2017) investigated the influence of the estradiol-cycle on VTA signaling and cocaine reward using the CPP paradigm. Their results revealed an estradiol-mediated increase in the activity of VTA DAergic neurons that promoted phosphorylation of DAT, which in turn was proposed to increase the ability of cocaine to inhibit its function. By use of chemogenetics, they moreover obtained data indicating a possible mechanistic link between the

increased activity of VTA DA neurons, via hM3Dq activation, and the enhanced affinity of cocaine for DAT, subsequent reward processing, and increased CPP (Calipari et al., 2017). In an earlier study, the same research group linked chronic cocaine exposure to dysregulated dMSNs signaling, impaired extinction, and facilitated cocaine-primed reinstatement. Specifically, it was found that hM4Di-mediated inhibition of dMSNs attenuated the strength of drug association and prevented relapse (Calipari et al., 2016).

The bed nucleus of the stria terminalis (BNST) has been implicated in cue-induced cocaine-seeking behaviors, and hM4Di-mediated inhibition of this region suggested that it is involved in ethanol-seeking behaviors (Pina et al., 2015). By using a dual viral chemogenetic approach, it was demonstrated that BNST neurons projecting to the VTA are involved in cue-induced ethanol-seeking behavior, as hM4Di activation in these neurons prevented alcohol CPP (Pina and Cunningham, 2017). Although ethanol has a different pharmacological profile compared with psychostimulants (Vengeliene et al., 2008; Kumar et al., 2009), the influence of DA in mediating the transition to addiction appears similar (Soderpalm and Ericson, 2011; Sulzer, 2011). Interestingly, Cheng et al. (2017) demonstrated an alcohol-induced enhancement of glutamatergic and GABAergic transmission onto dMSNs and iMSNs, respectively. That is, *in vivo* chemogenetic manipulation of dMSN or iMSN via hM3Dq or hM4Di, mimicking glutamatergic or GABAergic enhancement, respectively, affected alcohol consumption with hM3Dq activation in dMSNs or hM4Di activation in iMSNs promoting alcohol consumption, and oppositely reducing consumption through hM3Dq activation in iMSNs or hM4Di in dMSNs (Cheng et al., 2017). Anatomically distinct functions of the NAc core and shell in alcohol consumption were revealed in a limited access mouse model where alcohol consumption was reduced or increased following hM3Dq or hM4Di stimulation, respectively, of the NAc core but not accumbal shell (Purohit et al., 2018). It has also been observed that activation of KORD in neurons projecting to the NAc from the ventral subiculum decreased context-induced relapse of alcohol self-administration (Marchant et al., 2016a). While the mechanism of action of inhalant drugs of abuse has lagged behind, one recent chemogenetic study revealed an important role of infralimbic cortical projections to the NAc (Wayman and Woodward, 2018). The study described a toluene-induced (i.e., glue sniffing) reduction of excitability in rat infralimbic cortical neurons projecting to the NAc shell and inhibition of toluene-induced place preference after stimulating hM4Dq in this pathway (Wayman and Woodward, 2018).

Chemogenetics has further been employed to obtain evidence that astrocytes modulate the sensitivity and behavioral responses to reinforcing stimuli. Activating

hM3Dq specifically in the astrocytes of the NAc core increased intracranial self-stimulation, as well as reduced motivation to self-administer ethanol after 3 weeks abstinence (Bull et al., 2014). Similar chemogenetic activation of astrocytes expressing hM3Dq was shown to inhibit cue-induced reinstatement of cocaine seeking (Scofield et al., 2015). By the use of glutamate biosensors, Scofield et al. (2015) found that hM3Dq activation in the NAc astrocytes elevated glutamate levels as well as they obtained evidence that this glutamatergic gliotransmission inhibited cue-induced reinstatement of cocaine seeking as a result of activating neurotransmitter release-regulating metabotropic glutamate 2/3 autoreceptors.

The relative output from dMSNs and iMSNs, residing within the NAc, is likely directly to influence relapse since both rM3Ds-mediated stimulation of dMSN and hM4Di-mediated inhibition of iMSNs led to an increase in cue-induced reinstatement of cocaine self-administration (Heinsbroek et al., 2017). This effect was reversed with simultaneous hM4Di-mediated inhibition of ventral pallidum neurons (Heinsbroek et al., 2017). Consistent with these findings, ventral pallidum neurons have been associated with reward-cue pairing and reinstatement, and chemogenetic inhibition of ventral pallidum neurons impaired the ability of rats to acquire sign-tracking behaviors in response to a learned natural reward-associated cue (Chang et al., 2015). Interestingly, different areas of the ventral pallidum seem to regulate different forms of reinstatement. Local application of CNO in VTA of mice expressing hM4Di in either the rostral or caudal ventral pallidum revealed that only cue-induced reinstatement was suppressed when inhibiting rostral ventral pallidum-VTA projecting neurons, whereas caudal ventral pallidum-VTA inhibition had no effect on reinstatement. Conversely, systemic CNO was found to block primed reinstatement in caudal ventral pallidum-hM4Di rats, indicating that although caudal ventral pallidum is necessary for cocaine-primed reinstatement, its VTA targets are not essential for this action (Mahler et al., 2014). This suggests a complex contribution of specific subnuclei projections and circuits involving the NAc and ventral pallidum in drug-seeking behaviors. In addition, application of chemogenetics has also pointed to a role of midline thalamic nuclei neurons in drug-seeking behaviors. Interestingly, reducing activity of midline thalamic nuclei neurons by hM4Di stimulation inhibited both cue-induced and cocaine-primed reinstatement, whereas chemogenetic inhibition specifically of efferent projections from the anterior portion of midline thalamic nuclei to the NAc only blocked cocaine-primed reinstatement and enhanced cue-induced reinstatement (Wunsch et al., 2017). Similar to lesions of the NAc, suppressing the activity of accumbal neurons with hM4Di reduced ethanol consumption in a limited access model. In contrast,

increasing NAc activity with hM3Dq had no effect on ethanol intake (Cassataro et al., 2014). In an alcohol self-administration paradigm, the ventral pallidum was again demonstrated to influence reinstatement with hM4Di or rM3Ds stimulation, resulting in reduced or increased context-induced, as well as prime-induced, reinstatement of extinguished alcohol seeking, respectively (Prasad and McNally, 2016).

Lastly, the medial PFC is an area implicated in compulsive behaviors and has also been associated with relapse behaviors (Kalivas, 2008). Functional dissection of the ventro- and dorsomedial PFC areas has shown anatomically distinct influences on relapse behaviors. For example, the dorsomedial PFC region promoted relapse, while the ventromedial PFC region regulated extinction-like reductions in cocaine seeking (Peters et al., 2009; Moorman et al., 2015). In cocaine self-administering rats, hM4Di activation of medial PFC afferents to the NAc had no effect on motivation for taking cocaine during progressive ratio scheduling but increased subsequent drug-seeking behavior during extinction. Furthermore, primed reinstatement was significantly higher in rats that had been treated with CNO during progressive ratio scheduling. Strikingly, CNO treatment prior to a priming injection attenuated this response (Kerstetter et al., 2016). The authors suggested that this indicated a strengthening of the associative processes that linked cocaine with surrounding stimuli, which may seem counterintuitive when dampening medial PFC afferent activity to the NAc. However, the authors further speculated that late-stage beta-arrestin-mediated recruitment of mitogen-activated protein kinase/extracellular signal-regulated kinase (ERK) signaling modulated long-term alterations in neuronal plasticity and memory formation (Kerstetter et al., 2016). The specific role of the ventromedial PFC in extinction behaviors was confirmed through a chemogenetic approach aiming at activating ventromedial PFC neurons in a self-administration reinstatement model (Augur et al., 2016). By activating hM3Dq in ventromedial PFC neurons projecting to the NAc shell, Augur et al. (2016) demonstrated that these neurons played a role following cue-extinction learning, as they observed that cue-, but not prime-induced, reinstatement of cocaine seeking was reduced postextinction. Furthermore, they observed no effect following abstinence where the rats had not been through the “de-learning” phase of the cue-drug pairing (Augur et al., 2016). Altogether, neuronal adaptations seem to occur during extinction, which involves an active process in terms of context extinction memory, making drug-seeking behaviors dependent on ventromedial PFC activity.

3. Compulsive Drug Taking. The resistance to punishment after the acquisition of drug self-stimulation was recently found to be associated with enhanced neural activity in the orbitofrontal cortex. Chemogenetic

inhibition of this region reduced this form of compulsivity and prevented persevered self-stimulation despite punishment (Pascoli et al., 2015). Previously, cocaine-induced perseverative behaviors in mice were associated with altered synaptic plasticity in accumbal iMSNs. A potentiation of glutamatergic input to iMSNs was, contrary to potentiation of dMSNs, observed only in mice that showed resilience toward compulsive cocaine seeking. Chemogenetic inhibition of NAc iMSNs enhanced this motivation to obtain cocaine without affecting drug intake (Bock et al., 2013). The data provide a possible input-output link of a natural protective mechanism against compulsive drug intake, with cocaine inducing an enhanced activity of orbitofrontal cortex circuits synapsing onto iMSNs within the NAc.

Summarized, the studies described above on the role of the DA system in addiction reveal how certain neuronal populations and regions already associated with the DA reward circuitry system can influence addictive behaviors. Although several findings are not surprising as they support previous hypotheses, they provide a novel and unique spatial-selective map. Since regions and specific neuronal populations can be linked together, this combination reveals an immensely complex reward system, with pathways projecting from multiple brain regions that either directly or indirectly regulate prime regions of DA transmission including the VTA and NAc.

D. Mood, Nociception, and Cognition

Anxiety and depression are among the most common neuropsychiatric disorders, but the underlying neurobiology is poorly understood. Although 5-HT dysfunction is classically associated with mood disorders, the possible contribution of DAergic neurons in mood regulation has been examined using DREADDs. Zhong et al. (2014) induced an anxiogenic-like and depressive-like phenotype by selectively deleting expression of Cdk5, a kinase involved in the phosphorylation of TH, in the VTA of mice. This deletion caused reduced TH phosphorylation and VTA cAMP levels and NAc DA release. Selective rM3Ds activation of VTA DA neurons in this genetically modified mouse strain restored TH phosphorylation and DA levels, as well as rescued the behavioral deficits induced by deletion of the kinase (Zhong et al., 2014). DAergic neurons were also implicated in depressive-like behaviors under systemic inflammation conditions. When discomfort associated with a systemically induced inflammation state was investigated in the CPP paradigm, mice demonstrated an aversion to the inflammation-paired chamber and preferred the neutral chamber (Fritz et al., 2016). Chemogenetic activation of hM3Dq-expressing mice in midbrain DAergic neurons abolished this systemic inflammation place aversion, suggesting that this aversion is mediated via or at least integrated into midbrain DAergic neurotransmission.

Other studies have focused on the regulation of DAergic neurons under pain conditions. Chemogenetic suppression of VTA DAergic activity offset the phenomenon of exercise-induced hypoalgesia in partially sciatic nerve-ligated mice (Wakaizumi et al., 2016). Furthermore, specific hM3Dq activation of DAergic neurons in the ventral periaqueductal gray/dorsal raphe neurons was found to induce an anti-nociceptive effect in the hotplate, but not the tail flick assay, proposing that DA released from the periaqueductal gray/dorsal raphe is associated with supraspinal, but not spinal, nociception (Li et al., 2016).

DA was also shown to play a crucial role in fear learning, and therefore anxiety disorders and posttraumatic stress disorder (Lee et al., 2016; Bouchet et al., 2018; Mayer et al., 2018). Yet how DAergic neurons modulate these processes is unclear. Chemogenetic tools have facilitated investigations into this subject area, with hM4Di-mediated inhibition of a specific part of the mesolimbic system, projecting from the posterior medial VTA to the NAc shell, blocking relief learning following an aversive electrical stimulus cue (Mayer et al., 2018). There is ever growing data implicating the nigrostriatal pathway in this form of emotion since hM3Dq-mediated activation of SN DAergic neurons improved fear extinction memories and prevented future fear renewals (Bouchet et al., 2018). These early data propose an important role of DAergic pathways in the processing of signals predicting fear and relief, and therefore may represent a potential target for improving extinction-based therapies in anxiety and trauma-related disorders (Lee et al., 2016).

DA is known to play key roles in PFC-mediated cognition with system deficits apparent in ADHD and schizophrenia (Arnsten and Pliszka, 2011). To elucidate the cellular mechanisms governing D₂R-mediated enhancement of PFC pyramidal neuron excitability, each CNO-sensitive DREADD was probed in an in vitro electrophysiological set up. Strikingly, the results showed that only Gs-DREADD stimulation of the medial PFC mimicked this phenomenon implicated in schizophrenia (Robinson and Sohal, 2017). It is not only the higher cortical areas where DREADDs have been applied to study DA-regulated cognition. The importance of D₁Rs located in the lateral nucleus of the cerebellum was demonstrated using hM4Di-mediated cerebellar stimulation in D₁R-Cre mice to induce deficits on working memory and prepulse inhibition of the acoustic startle response (Locke et al., 2018).

The five-choice serial reaction time task assesses attentional, impulsive, and motivational control in rodents, and it is well-known that the DAergic system is centrally implicated in regulating all these behaviors, as also previously reviewed (Robbins, 2002; Jørgensen et al., 2017). In short, food-restricted subjects must wait and attend to a brief light stimulus in one of five apertures to receive a reward. The number of premature

responses, before the light appears, represents a measure of impulsivity, while incorrect or omitted responses are considered a measure of attention impairments. Interestingly, activation of hM3Dq in the rat VTA and SN pars compacta was shown to cause an increase in omitted responses, while only SN pars compacta activation decreased the number of correctly responded trials and mean correct latency (Boekhoudt et al., 2017b). As no effect on premature responses was observed, the authors suggested that increased midbrain DA neuronal activity caused attentional, but not impulsive, deficits.

III. Discussion and Future Directions

Summarized, it seems very clear that use of DREADDs in the area of DA research has advanced substantially our understanding of the DA system and led to many new discoveries. Indeed, the DAergic system is complex, consisting of a heterogeneous population of DA neurons and output cells that are intermingled and difficult to target in behavioral studies. By providing the spatial specificity that standard pharmacologic interventions lack, the chemogenetic toolbox has shown its potential to dissect the DA system and enlighten its influence on behavior. The DREADDs technique allows for relatively noninvasive regulation of neuronal signaling and depends largely on the intrinsic ability of neurons to regulate their own activity, as DREADDs work through endogenous intracellular signaling mechanisms. Once a chemogenetic receptor is expressed in neuronal populations/projections of interest, systemic administration of the inert ligand, CNO/clozapine or SalB, permits transient and repeated perturbations of neural activity. This can even be accomplished bilaterally in large or elongated brain structures, which can be extremely difficult with micro-injections of pharmacologic compounds or other molecular-genetic approaches such as optogenetics (Smith et al., 2016).

Despite its far-reaching usefulness and potential, the DREADDs technology is still in its infancy, and we are only beginning to realize the true nature of this toolbox's contents. However, as any other technique, chemogenetic interventions are not without limitations and challenges, and a lot of effort should be put into understanding the actual effects of injecting, expressing, and activating the various DREADDs, as well as possible adaptive changes occurring due to repeated activation.

A. Designer Receptors Exclusively Activated By Designer Drugs Effects in Theory and in Reality

1. *Designer Receptors Exclusively Activated By Designer Drugs Expression—The Meaning of Inertness.* The chemogenetic toolbox contains mutated GPCRs that are inert until application of their agonist (see

Fig. 1). Supposing that the DREADD agonists CNO and SalB are also inert and do not target endogenous receptors, their use provides the neuroscience field with, in theory, an ideal system for pharmacological remote control of any neuronal population of interest. However, there are no guarantees that this inertness applies in reality, and, as mentioned in the introduction (section I.2), recent evidence has emerged, indicating that at least under some circumstances in certain systems, extra caution should be exerted when interpreting chemogenetic experimental data (Maclaren et al., 2016; Gomez et al., 2017; Padovan-Hernandez and Knackstedt, 2018). CNO was previously reported to back-metabolize to clozapine in humans, nonhuman primates, and guinea pigs, but has generally been considered inert in rats and mice (Jann et al., 1994; Chang et al., 1998; Guettier et al., 2009; Loffler et al., 2012). More recently, however, MacLaren et al. (2016) described multiple dose-dependent effects in vivo in rats as a consequence of the back-metabolism of CNO to its bioactive parent compounds clozapine and *N*-desmethylozapine. Still, at CNO doses most commonly applied (1 to 2 mg/kg), there were no measurable effects on basal locomotion, nor amphetamine-induced DA release or hyperactivity. The acoustic startle reflex was the only tested measure that was affected by 1 or 2 mg/kg CNO (Maclaren et al., 2016). In the last decade, DREADDs have been applied in hundreds of laboratories to study various different neuronal circuits reporting no effects of CNO in non-DREADD-expressing mice. Accordingly, the concerns regarding the use of DREADDs to functional studies in vivo seem to be replaced with prudent caution and refinement (Mahler and Aston-Jones, 2018). To what extent SalB is inert remains to be determined. As a general recommendation, researchers should include and carefully plan control studies to exclude potential undesirable effects caused by DREADD ligands themselves.

So far, there have been no reports of constitutive activity of DREADDs expressed within neurons. Nonetheless, logically such activity should correlate with not just the number of DREADDs but also the endogenous signal transduction machinery, which varies between different populations of cells and neurons. Some constitutive activity of the rM3Ds DREADD in transfected cells has been reported and possibly explains the modest basal phenotype described in pancreatic beta cells (Guettier et al., 2009). Hence, chemogenetic receptors may not have profound basal activity in neuronal systems tested, but it cannot be ruled out that they can present with constitutive activity in certain cells of the CNS or when expressed by certain approaches. Low levels of constitutive activity might not cause a detectable phenotype right away but instead induce subtle alterations in homeostasis and long-term behaviors.

Another important issue is potential concerns relating to the method used to introduce DREADDs into

neurons and how this might differentially affect the function of the cells. It has, for example, been shown in optogenetic studies that the route of administration of the exact same opsin using identical AAV vector backbone, but encapsulated with various capsids (i.e., serotypes 1, 5, 8, 9), affected the efficacy of neuronal activity differently (Jackman et al., 2014). If this also applies to DREADDs, then even the choice of AAV serotype can determine how neurons respond to DREADD ligands. While such administration-related phenomena might not be a concern for interpreting one single DREADD study, it could impede comparisons between studies across the same scientific objectives in which the expression of DREADDs in identical cells were obtained by different viral serotypes and/or transgenic animals.

2. Interpretation of Activating Designer Receptors Exclusively Activated By Designer Drugs with Clozapine N-Oxide. Interpretation of chemogenetic manipulations affecting behavior is often condensed to activation or silencing of a specific neuron population/projection by hM3Dq/rM3Ds and/or hM4Di/KORD, respectively, but this is possibly an extreme simplification of what CNO actually does to neurons expressing DREADDs. Firstly, the effect of activating a certain DREADD may differ between neurons and even subpopulations of neurons, depending on the intracellular signal machinery, as mentioned above. Hence, the effect of activating hM4Di in different populations of neurons or even within the same neurons at different times can be quite different. For example, the activation of neurons by hM4Di inhibits presynaptic firing by hyperpolarizing neurons via GIRK (Roth, 2016), but some neurons may possess low expression of GIRK, and, consequently, the effect of CNO in these neurons may be negligible or dominated by other downstream mediators. In addition, GIRK levels may change because of previous drug exposure. Both cocaine and methamphetamine have been reported to inhibit midbrain GIRK-mediated currents in DAergic brain neurons for up to 2 weeks post-psychostimulant administration (Kobayashi et al., 2007; Padgett et al., 2012; Sharpe et al., 2014). This psychostimulant-related effect on GIRK is important to consider when utilizing chemogenetic tools, especially hM4Di, in DA-based research, given how psychostimulants alter DAergic neural circuitry and contribute to locomotion and reward-related processes. Recently, we demonstrated that following withdrawal of repeated cocaine, hM4Di-stimulation of VTA DA neurons no longer influenced cocaine-induced locomotion (Runegaard et al., 2018). Therefore, it will be important to investigate further whether and to what degree various drugs, including cocaine and amphetamine, affect the efficacy of chemogenetic modulation on neuronal activity and behavior, including how DREADD expression affects ligand-receptor binding interactions.

3. *Considerations for Repeated Activation of Designer Receptors Exclusively Activated By Designer Drugs.* CNO-induced activation of DREADDs depends on endogenous GPCR signaling cascades, and although these designer receptors are always available and should not desensitize, downstream effectors may limit the neuronal effect upon repeated activation (e.g., GIRK desensitization) (Roth, 2016). Due to the high expression levels of chemogenetic receptors, and the subsequent amount of receptor reserve, substantial desensitization by CNO or SalB causing a blunted or absent effect is obviously less likely. However, this is an issue that has yet not been satisfactorily addressed (Roman et al., 2016; Roth, 2016). So far, it has been shown that mice expressing hM4Di in the dorsomedial striatum or NAc developed a tolerance toward CNO following 2 weeks of chronic administration. However, that tolerance disappeared 48 h after chronic administration stopped (Carvalho Poyraz et al., 2016). Bellocchio et al. (2016) also found that chronic activation of hM3Dq with a high CNO dose of 10 mg/kg altered behavioral responses and induced cell death in vivo, possibly due to the unnatural, intense signal cascades activated by CNO, which caused neuronal damage. They discovered that chronic hM3Dq activation in iMSN caused enhanced locomotor activity and neuronal death via the c-Jun N-terminal kinase pathway (Bellocchio et al., 2016). Accordingly, it is important when designing long-lasting DREADD experiments to exclude possible adaptations caused by repeated administration of DREADD ligands.

4. *Can Selective Dopamine Expression Be Achieved?* As previously mentioned, one major advantage of DREADDs, but not limited to this specific technology, is the spatial selectivity by which the manipulation can be performed. Although the theoretical foundation of selective DREADD expression by Cre-dependency and/or promoter-regulated expression appears solid, practical issues such as leaky expression of Cre and concerns regarding promoter specificity may hamper the actual specificity and confound interpretations. In addition, the transgenic Cre-driver lines applied to specify DREADDs expression have not all been validated, which could also confound interpretation of DREADDs data. Thus, it seems important that researchers validate their transgenic Cre-driver line before proceeding with chemogenetic manipulations—in particular when WT littermates (non-DREADD expressing) are included as a control for CNO's own effects. Importantly, we recently characterized the commonly used TH-Cre driver mouse line (Savitt et al., 2005) and found neither alterations in DA system-related function nor in behavior (Runegaard et al., 2017b).

Recent studies have revealed a higher degree of structural, as well as functional heterogeneity, in the DA system than previously assumed. This calls for increased anatomic specificity when manipulating

specific DA neurons and projections to provide reliable and accurate interpretations. Indeed, concerns have been raised regarding the use of a single gene marker to define the DAergic phenotype (Björklund and Dunnett, 2007; Apuschkin et al., 2015; Lammel et al., 2015; Stuber et al., 2015), and the DAT- and TH-Cre driver lines used to specify expression to DAergic neurons may target different but overlapping populations of DAergic neurons. Furthermore, recent evidence suggests that effector gene expression in transgenic TH-Cre mice is not limited to DAergic neurons but also seen ectopically such as in TH-transcript positive GABAergic VTA neurons projecting to the lateral habenula (Lammel et al., 2015; Stuber et al., 2015). This knowledge emphasizes the need for an updated characterization of the DA phenotype, or rather a dissection of the different phenotypes of this heterogeneous population of neurons. Specifically, it might be necessary to redefine the hallmarks of a DAergic neuron, considering whether it is defined by its synthesis, release, and/or reuptake mechanisms (Berke, 2018). It is, moreover, essential to achieve a better understanding of the DAergic subpopulations coreleasing other neurotransmitters such as glutamate (Morales and Root, 2014) and GABA (Barker et al., 2016) to guide us in our understanding of DAergic neuronal and projection patterns.

5. *Can Designer Receptors Exclusively Activated By Designer Drugs Help Dissect the Dopaminergic Pathways?* As emphasized above, a fair interpretation of the chemogenetic manipulations relies on the thorough validation of targeted neurons, projections, and an understanding of their characteristics. Independent of the definition of DAergic neurons, refinements of the chemogenetic technique are also important. These include improvements in projection-specific expression by the use of retrogradely transported viral vectors (e.g., CAV-2-Cre) (Boender et al., 2014), recombinant AAVs carrying tissue- or cell-specific promoters (Gompf et al., 2015), or by double-recombination approaches through the introduction of an additional recombinase, such as Flp (Stamatakis et al., 2013). This latter approach could be even more sophisticated if introduced under an activity-dependent promoter, e.g., mediated by C-fos activation (Garner et al., 2012; Kawashima et al., 2014; Roy et al., 2016). Additionally, the temporal and spatial resolution can be improved further via photo-caging DREADDs or creating DREADD antagonists, which could become highly useful and emphasize the continuous development and refinement of the DREADD technology (Roth, 2016). Of further interest, refinement of viral genetic tracing techniques has recently allowed systematic mapping of the input-output circuitry of neuronal nuclei (Beier et al., 2015; Schwarz et al., 2015; Cardozo Pinto and Lammel, 2018). Beier et al. (2015) demonstrated that VTA DA neurons projecting to different forebrain regions exhibited specific biases in

their input selection, and VTA DA neurons projecting to the lateral and medial NAc innervated largely non-overlapping striatal targets, with the latter also sending extrastriatal axon collaterals.

B. Have Chemogenetic Studies Advanced Our Understanding of Dopamine-Related Behaviors?

While the chemogenetic toolbox has facilitated studies previously not possible, it is evident that some of the results obtained by application of chemogenetics do not provide true novel insights but rather confirm what has previously been shown by lesion studies and by use of transgenic or pharmacologic approaches. However, as described in this review, the improved specificity of the chemogenetic manipulations has in many cases enabled much more refined conclusions in terms of spatial and specific influences of distinct neuronal populations on behavioral outputs. We believe accordingly that future chemogenetic studies have the potential to significantly further advance our understanding of the complex role of DA in regulating the multiple behaviors discussed here. Indeed, the wide use of chemogenetics in laboratories all over the world directly reflects the vast potential of the technology to progress our understanding of neuronal signaling and behavior. Nonetheless, when interpreting chemogenetic data with a wish to conclude about DA-related behaviors, it is highly important to consider the immense heterogeneity of the DA system. Not least, the still poorly understood importance of coreleased neurotransmitters, such as GABA and glutamate, must be taken into account (Poulin et al., 2018). Moreover, and as also mentioned earlier, it has to be considered that when chemogenetically manipulating, e.g., MSNs, the outcome might not necessarily solely reflect DA transmission, because neurons receive metabotropic input from multiple GPCRs and not only G protein-coupled DRs. In context of these challenges, it will be highly important to combine the specificity by which DREADDs can be expressed and activated with advanced techniques assessing neurotransmitter release, activity, and dynamics, such as the recently published DA biosensors (Patriarchi et al., 2018; Sun et al., 2018) or two-photon excitation microscopy technologies (Svoboda and Yasuda, 2006). Such approaches should indeed enable better comprehension of the multifaceted action governed by DA and its circuitry in the near future.

C. Future Directions

Besides the use of DREADDs in biologic and disease research, this technique possesses tremendous potential in translational research and clinical therapies (Roth, 2016). As well as developing multiple designer receptors and agonists, the fast-paced DREADDs field has dissected the mesocorticolimbic network with positron emission tomography (Michaelides et al., 2013) and functional magnetic resonance imaging (Roelofs

et al., 2017) and has already been used in nonhuman primates (Eldridge et al., 2016; Grayson et al., 2016). While the idea of gene therapy and thereby expressing DREADDs in humans may not be a major issue, it is the ligand CNO that arouses most worry because of the back-metabolization of CNO to clozapine in several species, potentially limiting its use in patients. However, the newer KORD variant is activated by a different ligand, SalB, which has not yet been associated with such issues (Vardy et al., 2015). In addition, structure-activity relationship studies were applied to develop alternative ligands for activating DREADDs, leading to the identification of a new potent hM3Dq agonist that is neither back-metabolized nor activating endogenous receptors (Chen et al., 2015). Development of more sophisticated DREADD variants will thereby probably allow for the use of DREADDs in humans in the future (Urban and Roth, 2015; Smith et al., 2016).

IV. Conclusion

A plethora of novel techniques has become available within the field of neuroscience in recent years. Together these have revolutionized neuroscience and dramatically improved our understanding of complex processes taking place in the brain and how such processes might change during disease states. One technique of particular interest is chemogenetics that, by allowing pharmacological remote control of neuronal activity, have opened up for entirely new possibilities for dissecting neuronal circuits and deciphering the role of diverse neuronal populations. Chemogenetics has already been extensively used for studying the DA system, and, as reviewed here, multiple different components of the DA system, including DA-releasing neurons, striatal postsynaptic MSNs conveying DAergic signals, and other neural regions/substances synapsing onto the DA neurons, have been targeted by DREADDs to investigate regulation of discrete and multiple behavioral functions. One important lesson is that we should learn and appreciate that behavior is not simply driven by just one single neuronal population or projection pattern, but rather determined by a tightly regulated integration of multiple neuronal inputs. Indeed, behavior is complex and should not be simplified to a binary constituent taking place in response to “activation” or “inhibition” of specific neurons.

The major advancement of chemogenetics, compared with traditional pharmacological interventions, has been the increased anatomic specificity by which the manipulations can be undertaken and thereby dissect not just a biologic heterogeneous population of DA neurons, but also an anatomically important complex projection system. There are no doubts that the potential of chemogenetics is huge, but as with any other technique, there are concerns, some of which need to be further addressed. These include how to target a

heterogeneous DAergic population, the dependency of DREADDs-related effects on endogenous GPCR signaling machinery (which may differ between different neuronal subpopulations), expression-induced adaptations, as well as possible alterations in DREADDs expression and/or adaptive cellular responses with repeated or chronic CNO administration. It should nonetheless be appreciated that the chemogenetic technique is still at a relatively early stage, and there is no reason to believe that many of these obstacles cannot be addressed and the technique further developed. With the progress of the CRISPR technology (Wang et al., 2016), one might even consider that DREADDs-like manipulations of any endogenous receptor/transporter of interests soon might become part of the commonly used methodological repertoire. In relation to direct translational application of the DREADD technology, searches for new ligands (Chen et al., 2015; Roth, 2016) and development of new designer receptor variants activated by other ligands (e.g., KORD) (Vardy et al., 2015), may overcome the problem of extensive metabolism of the activating ligand CNO in primates, and thus, chemogenetic therapy to treat neurologic and psychiatric disorders may not be an unrealistic or distant goal.

Authorship Contributions

Wrote or contributed to the writing of the manuscript: Runegaard, Fitzpatrick, Woldbye, Andreassen, Sørensen, and Gether.

References

- Alcacer C, Andreoli L, Sebastianutto I, Jakobsson J, Fieblinger T, and Cenci MA (2017) Chemogenetic stimulation of striatal projection neurons modulates responses to Parkinson's disease therapy. *J Clin Invest* **127**:720–734.
- Aldrin-Kirk P, Heuer A, Rylander Ottosson D, Davidsson M, Mattsson B, and Björklund T (2018) Chemogenetic modulation of cholinergic interneurons reveals their regulating role on the direct and indirect output pathways from the striatum. *Neurobiol Dis* **109** (Pt A):148–162.
- Aldrin-Kirk P, Heuer A, Wang G, Mattsson B, Lundblad M, Parmar M, and Björklund T (2016) DREADD modulation of transplanted DA neurons reveals a novel Parkinsonian dyskinesia mechanism mediated by the serotonin 5-HT₆ receptor. *Neuron* **90**:955–968.
- Alexander GM, Rogan SC, Abbas AI, Armbruster BN, Pei Y, Allen JA, Nonneman RJ, Hartmann J, Moy SS, Nicoletis MA, et al. (2009) Remote control of neuronal activity in transgenic mice expressing evolved G protein-coupled receptors. *Neuron* **63**:27–39.
- Apushkin M, Stilling S, Rahbek-Clemmensen T, Sørensen G, Fortin G, Herborg Hansen F, Eriksen J, Trudeau LE, Egerod K, Gether U, et al. (2015) A novel dopamine transporter transgenic mouse line for identification and purification of midbrain dopaminergic neurons reveals midbrain heterogeneity. *Eur J Neurosci* **42**:2438–2454.
- Arias-Carrión O and Pöppel E (2007) Dopamine, learning, and reward-seeking behavior. *Acta Neurobiol Exp (Warsz)* **67**:481–488.
- Armbruster BN, Li X, Pausch MH, Herlitze S, and Roth BL (2007) Evolving the lock to fit the key to create a family of G protein-coupled receptors potentially activated by an inert ligand. *Proc Natl Acad Sci USA* **104**:5163–5168.
- Arntsen AFT and Pliszka SR (2011) Catecholamine influences on prefrontal cortical function: relevance to treatment of attention deficit/hyperactivity disorder and related disorders. *Pharmacol Biochem Behav* **99**:211–216.
- Augur IF, Wyckoff AR, Aston-Jones G, Kalivas PW, and Peters J (2016) Chemogenetic activation of an extinction neural circuit reduces cue-induced reinstatement of cocaine seeking. *J Neurosci* **36**:10174–10180.
- Bäckman CM, Malik N, Zhang Y, Shan L, Grinberg A, Hoffer BJ, Westphal H, and Tomac AC (2006) Characterization of a mouse strain expressing Cre recombinase from the 3' untranslated region of the dopamine transporter locus. *Genesis* **44**:383–390.
- Baik J-H (2013) Dopamine signaling in reward-related behaviors. *Front Neural Circuits* **7**:152.
- Balfour ME, Yu L, and Coolen LM (2004) Sexual behavior and sex-associated environmental cues activate the mesolimbic system in male rats. *Neuropsychopharmacology* **29**:718–730.
- Barker DJ, Root DH, Zhang S, and Morales M (2016) Multiplexed neurochemical signaling by neurons of the ventral tegmental area. *J Chem Neuroanat* **73**:33–42.
- Barker RA, Barrett J, Mason SL, and Björklund A (2013) Fetal dopaminergic transplantation trials and the future of neural grafting in Parkinson's disease. *Lancet Neurol* **12**:84–91.
- Barter JW, Castro S, Sukharnikova T, Rossi MA, and Yin HH (2014) The role of the substantia nigra in posture control. *Eur J Neurosci* **39**:1465–1473.
- Bassareo V and Di Chiara G (1999) Differential responsiveness of dopamine transmission to food-stimuli in nucleus accumbens shell/core compartments. *Neuroscience* **89**:637–641.
- Beaulieu JM, Espinoza S, and Gainetdinov RR (2015) Dopamine receptors - IUPHAR review 13. *Br J Pharmacol* **172**:1–23.
- Beaulieu J-M and Gainetdinov RR (2011) The physiology, signaling, and pharmacology of dopamine receptors. *Pharmacol Rev* **63**:182–217.
- Beaulieu JM, Sotnikova TD, Marion S, Lefkowitz RJ, Gainetdinov RR, and Caron MG (2005) An Akt/beta-arrestin 2/PP2A signaling complex mediates dopaminergic neurotransmission and behavior. *Cell* **122**:261–273.
- Beaulieu JM, Sotnikova TD, Yao WD, Kockeritz L, Woodgett JR, Gainetdinov RR, and Caron MG (2004) Lithium antagonizes dopamine-dependent behaviors mediated by an AKT/glycogen synthase kinase 3 signaling cascade. *Proc Natl Acad Sci USA* **101**:5099–5104.
- Beaulieu JM, Tirota E, Sotnikova TD, Masri B, Salahpour A, Gainetdinov RR, Borrelli E, and Caron MG (2007) Regulation of Akt signaling by D2 and D3 dopamine receptors in vivo. *J Neurosci* **27**:881–885.
- Becker JB and Koob GF (2016) Sex differences in animal models: focus on addiction. *Pharmacol Rev* **68**:242–263.
- Beier KT, Steinberg EE, DeLoach KE, Xie S, Miyamichi K, Schwarz L, Gao XJ, Kremer EJ, Malenka RC, and Luo L (2015) Circuit architecture of VTA dopamine neurons revealed by systematic input-output mapping. *Cell* **162**:622–634.
- Belin D and Deroche-Gammonet V (2012) Responses to novelty and vulnerability to cocaine addiction: contribution of a multi-symptomatic animal model. *Cold Spring Harb Perspect Med* **2**:a011940.
- Bellochio L, Ruiz-Calvo A, Chiarlone A, Cabanas M, Resel E, Cazalets J-R, Blázquez C, Cho YH, Galve-Roperh I, and Guzmán M (2016) Sustained Gq-protein signaling disrupts striatal circuits via JNK. *J Neurosci* **36**:10611–10624.
- Belout LN, Omrani A, Adan RA, Webb JC, and Coolen LM (2016) Ventral tegmental area dopamine cell activation during male rat sexual behavior regulates neuroplasticity and D-amphetamine cross-sensitization following sex abstinence. *J Neurosci* **36**:9949–9961.
- Ben-Shaanan TL, Azuly-Debby H, Dubovik T, Starosvetsky E, Korin B, Schiller M, Green NL, Admon Y, Hakim F, Shen-Orr SS, et al. (2016) Activation of the reward system boosts innate and adaptive immunity. *Nat Med* **22**:940–944.
- Berke JD (2018) What does dopamine mean? *Nat Neurosci* **21**:787–793.
- Berrios J, Stamatakis AM, Kantak PA, McElligott ZA, Judson MC, Aita M, Rougie M, Stubbs GD, and Philpot BD (2016) Loss of UBE3A from TH-expressing neurons suppresses GABA co-release and enhances VTA-NAc optical self-stimulation. *Nat Commun* **7**:10702.
- Björklund A and Dunnett SB (2007) Dopamine neuron systems in the brain: an update. *Trends Neurosci* **30**:194–202.
- Blum ID, Zhu L, Moquin L, Kokoeva MV, Gratton A, Giros B, and Storch KF (2014) A highly tunable dopaminergic oscillator generates ultradian rhythms of behavioral arousal. *eLife* **3**:e05105.
- Bobadilla AC, Garcia-Keller C, Heinsbroek JA, Scofield MD, Chareunsouk V, Monforton C, and Kalivas PW (2017) Accumbens mechanisms for cued sucrose seeking. *Neuropsychopharmacology* **42**:2377–2386.
- Bock R, Shin JH, Kaplan AR, Dobi A, Markey E, Kramer PF, Gremel CM, Christensen CH, Adrover MF, and Alvarez VA (2013) Strengthening the accumbal indirect pathway promotes resilience to compulsive cocaine use. *Nat Neurosci* **16**:632–638.
- Boekhoudt L, Omrani A, Luijckx MCM, Wolterink-Donselaar IG, Wijbrans EC, van der Plasse G, and Adan RAH (2016) Chemogenetic activation of dopamine neurons in the ventral tegmental area, but not substantia nigra, induces hyperactivity in rats. *Eur Neuropsychopharmacol* **26**:1784–1793.
- Boekhoudt L, Roelofs TJM, de Jong JW, de Leeuw AE, Luijckx MCM, Wolterink-Donselaar IG, van der Plasse G, and Adan RAH (2017a) Does activation of midbrain dopamine neurons promote or reduce feeding? *Int J Obes (Lond)* **41**:1131–1140.
- Boekhoudt L, Voets ES, Flores-Dourojeanni JP, Luijckx MC, Vanderschuren LJ, and Adan RAH (2017b) Chemogenetic activation of midbrain dopamine neurons affects attention, but not impulsivity, in the five-choice serial reaction time-task in rats. *Neuropsychopharmacology* **42**:1315–1325.
- Boekhoudt L, Wijbrans EC, Man JHK, Luijckx MCM, de Jong JW, van der Plasse G, Vanderschuren LJMJ, and Adan RAH (2018) Enhancing excitability of dopamine neurons promotes motivational behaviour through increased action initiation. *Eur Neuropsychopharmacol* **28**:171–184.
- Boender AJ, de Jong JW, Boekhoudt L, Luijckx MCM, van der Plasse G, and Adan RAH (2014) Combined use of the canine adenovirus-2 and DREADD-technology to activate specific neural pathways in vivo. *PLoS One* **9**:e95392.
- Bouabid S and Zhou FM (2018) Cyclic AMP-producing chemogenetic activation of indirect pathway striatal projection neurons and the downstream effects on the globus pallidus and subthalamic nucleus in freely moving mice. *J Neurochem* **145**:436–448.
- Bouchet CA, Miner MA, Loetz EC, Rosberg AJ, Hake HS, Farmer CE, Ostrovskyy M, Gray N, and Greenwood BN (2018) Activation of nigrostriatal dopamine neurons during fear extinction prevents the renewal of fear. *Neuropsychopharmacology* **43**:665–672.
- Bull C, Freitas KCC, Zou S, Poland RS, Syed WA, Urban DJ, Minter SC, Shelton KL, Hauser KF, Negus SS, et al. (2014) Rat nucleus accumbens core astrocytes modulate reward and the motivation to self-administer ethanol after abstinence. *Neuropsychopharmacology* **39**:2835–2845.
- Calipari ES, Bagot RC, Purushothaman I, Davidson TJ, Yorgason JT, Peña CJ, Walker DM, Pirpinias ST, Guise KG, Ramakrishnan C, et al. (2016) In vivo imaging identifies temporal signature of D1 and D2 medium spiny neurons in cocaine reward. *Proc Natl Acad Sci USA* **113**:2726–2731.

- Calipari ES, Juarez B, Morel C, Walker DM, Cahill ME, Ribeiro E, Roman-Ortiz C, Ramakrishnan C, Deisseroth K, Han M-H, et al. (2017) Dopaminergic dynamics underlying sex-specific cocaine reward. *Nat Commun* **8**:13877.
- Cardozo Pinto DF and Lammel S (2018) Viral vector strategies for investigating midbrain dopamine circuits underlying motivated behaviors. *Pharmacol Biochem Behav* **174**:23–32.
- Carlsson A and Waldeck B (1958) A fluorimetric method for the determination of dopamine (3-hydroxytyramine). *Acta Physiol Scand* **44**:293–298.
- Carter ME, Soden ME, Zweifel LS, and Palmiter RD (2013) Genetic identification of a neural circuit that suppresses appetite. *Nature* **503**:111–114.
- Carvalho Poyraz F, Holzner E, Bailey MR, Meszaros J, Kenney L, Kheirbek MA, Balsam PD, and Kellendonk C (2016) Decreasing striatopallidal pathway function enhances motivation by energizing the initiation of goal-directed action. *J Neurosci* **36**:5988–6001.
- Cassaturo D, Bergfeldt D, Malekian C, Van Snellenberg JX, Thanos PK, Fishell G, and Sjulson L (2014) Reverse pharmacogenetic modulation of the nucleus accumbens reduces ethanol consumption in a limited access paradigm. *Neuropsychopharmacology* **39**:283–290.
- Chan KY, Jang MJ, Yoo BB, Greenbaum A, Ravi N, Wu WL, Sánchez-Guardado L, Lois C, Mazmanian SK, Deverman BE, et al. (2017) Engineered AAVs for efficient noninvasive gene delivery to the central and peripheral nervous systems. *Nat Neurosci* **20**:1172–1179.
- Chang SE, Todd TP, Buccini DJ, and Smith KS (2015) Chemogenetic manipulation of ventral pallidal neurons impairs acquisition of sign-tracking in rats. *Eur J Neurosci* **42**:3105–3116.
- Chang WH, Lin SK, Lane HY, Wei FC, Hu WH, Lam YWF, and Jann MW (1998) Reversible metabolism of clozapine and clozapine N-oxide in schizophrenic patients. *Prog Neuropsychopharmacol Biol Psychiatry* **22**:723–739.
- Chen X, Choo H, Huang XP, Yang X, Stone O, Roth BL, and Jin J (2015) The first structure-activity relationship studies for designer receptors exclusively activated by designer drugs. *ACS Chem Neurosci* **6**:476–484.
- Chen Y, Xiong M, Dong Y, Haberman A, Cao J, Liu H, Zhou W, and Zhang S-C (2016) Chemical control of grafted human PSC-derived neurons in a mouse model of Parkinson's disease. *Cell Stem Cell* **18**:817–826.
- Cheng Y, Huang CCY, Ma T, Wei X, Wang X, Lu J, and Wang J (2017) Distinct synaptic strengthening of the striatal direct and indirect pathways drives alcohol consumption. *Biol Psychiatry* **81**:918–929.
- Cho JR, Treweek JB, Robinson JE, Xiao C, Bremner LR, Greenbaum A, and Gradinaru V (2017) Dorsal raphe dopamine neurons modulate arousal and promote wakefulness by salient stimuli. *Neuron* **94**:1205–1219.e8.
- Chu HY, McIver EL, Kovaleski RF, Atherton JF, and Bevan MD (2017) Loss of hyperdirect pathway cortico-subthalamic inputs following degeneration of mid-brain dopamine neurons. *Neuron* **95**:1306–1318.e5.
- Chuhma N, Zhang H, Masson J, Zhuang X, Sulzer D, Hen R, and Rayport S (2004) Dopamine neurons mediate a fast excitatory signal via their glutamatergic synapses. *J Neurosci* **24**:972–981.
- Conklin BR, Hsiao EC, Claysen S, Dumuis A, Srinivasan S, Forsayeth JR, Guettier JM, Chang WC, Pei Y, McCarthy KD, et al. (2008) Engineering GPCR signaling pathways with RASLS. *Nat Methods* **5**:673–678.
- Damsma G, Pfau JG, Wenkstern D, Phillips AG, and Fibiger HC (1992) Sexual behavior increases dopamine transmission in the nucleus accumbens and striatum of male rats: comparison with novelty and locomotion. *Behav Neurosci* **106**:181–191.
- dela Peña I, Gevorkiana R, and Shi W-X (2015) Psychostimulants affect dopamine transmission through both dopamine transporter-dependent and independent mechanisms. *Eur J Pharmacol* **764**:562–570.
- Delfs JM, Schreiber L, and Kelley AE (1990) Microinjection of cocaine into the nucleus accumbens elicits locomotor activation in the rat. *J Neurosci* **10**:303–310.
- Dell'Anno MT, Caiazzo M, Leo D, Dvoretzskova E, Medrihan L, Colasante G, Giannelli S, Theka I, Russo G, Mus L, et al. (2014) Remote control of induced dopaminergic neurons in parkinsonian rats. *J Clin Invest* **124**:3215–3229.
- Demiñiere JM, Piazza PV, Le Moal M, and Simon H (1989) Experimental approach to individual vulnerability to psychostimulant addiction. *Neurosci Biobehav Rev* **13**:141–147.
- Di Chiara G and Imperato A (1988) Drugs abused by humans preferentially increase synaptic dopamine concentrations in the mesolimbic system of freely moving rats. *Proc Natl Acad Sci USA* **85**:5274–5278.
- Dobbs LKK, Kaplan ARR, Lemos JCC, Matsui A, Rubinstein M, and Alvarez VAA (2016) Dopamine regulation of lateral inhibition between striatal neurons gates the stimulant actions of cocaine. *Neuron* **90**:1100–1113.
- Eban-Rothschild A, Rothschild G, Giardino WJ, Jones JR, and de Lecea L (2016) VTA dopaminergic neurons regulate ethologically relevant sleep-wake behaviors. *Nat Neurosci* **19**:1356–1366.
- Eldridge MAG, Lerchner W, Saunders RC, Kaneko H, Krausz KW, Gonzalez FJ, Ji B, Higuchi M, Minamimoto T, and Richmond BJ (2016) Chemogenetic disconnection of monkey orbitofrontal and rhinal cortex reversibly disrupts reward value. *Nat Neurosci* **19**:37–39.
- Farrell MS, Pei Y, Wan Y, Yadav PN, Daigle TL, Urban DJ, Lee H-M, Siacky N, Simmons A, Noneman RJ, et al. (2013) A GAS DREADD mouse for selective modulation of cAMP production in striatopallidal neurons. *Neuropsychopharmacology* **38**:854–862.
- Ferguson SM, Eskenazi D, Ishikawa M, Wanat MJ, Phillips PEM, Dong Y, Roth BL, and Neumaier JF (2011) Transient neuronal inhibition reveals opposing roles of indirect and direct pathways in sensitization. *Nat Neurosci* **14**:22–24.
- Ferguson SM and Neumaier JF (2012) Grateful DREADDs: engineered receptors reveal how neural circuits regulate behavior. *Neuropsychopharmacology* **37**:296–297.
- Ferguson SM, Phillips PEM, Roth BL, Wess J, and Neumaier JF (2013) Direct-pathway striatal neurons regulate the retention of decision-making strategies. *J Neurosci* **33**:11668–11676.
- Foldi CJ, Milton LK, and Oldfield BJ (2017) The role of mesolimbic reward neurocircuitry in prevention and rescue of the activity-based anorexia (ABA) phenotype in rats. *Neuropsychopharmacology* **42**:2292–2300.
- Fritz M, Klawonn AM, Nilsson A, Singh AK, Zajdel J, Wilhelms DB, Lazarus M, Löfberg A, Jaarola M, Kugelberg UO, et al. (2016) Prostaglandin-dependent modulation of dopaminergic neurotransmission elicits inflammation-induced aversion in mice. *J Clin Invest* **126**:695–705.
- Gallo EF, Meszaros J, Sherman JD, Chohan MO, Teboul E, Choi CS, Moore H, Javitch JA, and Kellendonk C (2018) Accumbens dopamine D2 receptors increase motivation by decreasing inhibitory transmission to the ventral pallidum. *Nat Commun* **9**:1086.
- Garner AR, Rowland DC, Hwang SY, Baumgaertel K, Roth BL, Kentros C, and Mayford M (2012) Generation of a synthetic memory trace. *Science* **335**:1513–1516.
- German CL, Baladi MG, McFadden LM, Hanson GR, and Fleckenstein AE (2015) Regulation of the dopamine and vesicular monoamine transporters: pharmacological targets and implications for disease. *Pharmacol Rev* **67**:1005–1024.
- Girasole AE, Lum MY, Nathaniel D, Bair-Marshall CJ, Guenther CJ, Luo L, Kreitzer AC, and Nelson AB (2018) A subpopulation of striatal neurons mediates levodopa-induced dyskinesia. *Neuron* **97**:787–795.e6.
- Gomez JL, Bonaventura J, Lesniak W, Mathews WB, Sysa-Shah P, Rodriguez LA, Ellis RJ, Richie CT, Harvey BK, Dannals RF, et al. (2017) Chemogenetics revealed: DREADD occupancy and activation via converted clozapine. *Science* **357**:503–507.
- Gompf HS, Budygin EA, Fuller PM, and Bass CE (2015) Targeted genetic manipulations of neuronal subtypes using promoter-specific combinatorial AAVs in wild-type animals. *Front Behav Neurosci* **9**:152.
- Goto A, Nakahara I, Yamaguchi T, Kamioka Y, Sumiyama K, Matsuda M, Nakanishi S, and Funabiki K (2015) Circuit-dependent striatal PKA and ERK signaling underlies rapid behavioral shift in mating reaction of male mice. *Proc Natl Acad Sci USA* **112**:6718–6723.
- Grayson DS, Bliss-Moreau E, Machado CJ, Bennett J, Shen K, Grant KA, Fair DA, and Amaral DG (2016) The rhesus monkey connectome predicts disrupted functional networks resulting from pharmacogenetic inactivation of the amygdala. *Neuron* **91**:453–466.
- Guettier J-M, Gautam D, Scarselli M, Ruiz de Azua I, Li JH, Rosemond E, Ma X, Gonzalez FJ, Armbruster BN, Lu H, et al. (2009) A chemical-genetic approach to study G protein regulation of beta cell function in vivo. *Proc Natl Acad Sci USA* **106**:19197–19202.
- Heinsbroek JA, Neuhof DN, Griffin WC III, Siegel GS, Bobadilla AC, Kupchik YM, and Kalivas PW (2017) Loss of plasticity in the D2-accumbens pallidal pathway promotes cocaine seeking. *J Neurosci* **37**:757–767.
- Hnasko TS, Chuhma N, Zhang H, Goh GY, Sulzer D, Palmiter RD, Rayport S, and Edwards RH (2010) Vesicular glutamate transport promotes dopamine storage and glutamate corelease in vivo. *Neuron* **65**:643–656.
- Hnasko TS, Perez FA, Scouras AD, Stoll EA, Gale SD, Luquet S, Phillips PEM, Kremer EJ, and Palmiter RD (2006) Cre recombinase-mediated restoration of nigrostriatal dopamine in dopamine-deficient mice reverses hypophagia and bradykinesia. *Proc Natl Acad Sci USA* **103**:8858–8863.
- Holmes A, Lachowicz JE, and Sibley DR (2004) Phenotypic analysis of dopamine receptor knockout mice; recent insights into the functional specificity of dopamine receptor subtypes. *Neuropharmacology* **47**:1117–1134.
- Hooks MS, Jones GH, Smith AD, Neill DB, and Justice JB Jr. (1991a) Individual differences in locomotor activity and sensitization. *Pharmacol Biochem Behav* **38**:467–470.
- Hooks MS, Jones GH, Smith AD, Neill DB, and Justice JB Jr. (1991b) Response to novelty predicts the locomotor and nucleus accumbens dopamine response to cocaine. *Synapse* **9**:121–128.
- Hornykiewicz O (1966) Dopamine (3-hydroxytyramine) and brain function. *Pharmacol Rev* **18**:925–964.
- Jackman SL, Beneduce BM, Drew IR, and Regehr WG (2014) Achieving high-frequency optical control of synaptic transmission. *J Neurosci* **34**:7704–7714.
- Jacobson KA, Gao ZG, Chen A, Barak D, Kim SA, Lee K, Link A, Rompaey PV, van Calenbergh S, and Liang BT (2001) Neocortex concept based on molecular complementarity in GPCRs: a mutant adenosine A(3) receptor with selectively enhanced affinity for amine-modified nucleosides. *J Med Chem* **44**:4125–4136.
- Jann MW, Lam YW, and Chang WH (1994) Rapid formation of clozapine in guinea-pigs and man following clozapine-N-oxide administration. *Arch Int Pharmacodyn Ther* **328**:243–250.
- Jones SR, Gainetdinov RR, Jaber M, Giros B, Wightman RM, and Caron MG (1998) Profound neuronal plasticity in response to inactivation of the dopamine transporter. *Proc Natl Acad Sci USA* **95**:4029–4034.
- Jørgensen SH, Fitzpatrick CM, Gether U, Woldbye DPD, and Sørensen AT (2017) Chemogenetic modulation of G protein-coupled receptor signalling in visual attention research. *Basic Clin Pharmacol Toxicol* **121**:373–381.
- Kalivas PW (2008) Addiction as a pathology in prefrontal cortical regulation of corticostriatal habit circuitry. *Neurotox Res* **14**:185–189.
- Kawashima T, Okuno H, and Bito H (2014) A new era for functional labeling of neurons: activity-dependent promoters have come of age. *Front Neural Circuits* **8**:37.
- Kerstetter KA, Wunsch AM, Nakata KG, Donckels E, Neumaier JF, and Ferguson SM (2016) Corticostriatal afferents modulate responsiveness to psychostimulant drugs and drug-associated stimuli. *Neuropsychopharmacology* **41**:1128–1137.
- Kim JI, Ganesan S, Luo SX, Wu YW, Park E, Huang EJ, Chen L, and Ding JB (2015) Aldehyde dehydrogenase 1a1 mediates a GABA synthesis pathway in midbrain dopaminergic neurons. *Science* **350**:102–106.
- Kobayashi T, Nishizawa D, Iwamura T, and Ikeda K (2007) Inhibition by cocaine of G protein-activated inwardly rectifying K⁺ channels expressed in *Xenopus* oocytes. *Toxicol In Vitro* **21**:656–664.
- Koob GF and Le Moal M (1997) Drug abuse: hedonic homeostatic dysregulation. *Science* **278**:52–58.

- Kristensen AS, Andersen J, Jørgensen TN, Sørensen L, Eriksen J, Loland CJ, Strømgaard K, and Gether U (2011) SLC6 neurotransmitter transporters: structure, function, and regulation. *Pharmacol Rev* **63**:585–640.
- Kumar S, Porcu P, Werner DF, Matthews DB, Diaz-Granados JL, Helfand RS, and Morrow AL (2009) The role of GABA(A) receptors in the acute and chronic effects of ethanol: a decade of progress. *Psychopharmacology (Berl)* **205**:529–564.
- Kupchik YM, Brown RM, Heinsbroek JA, Lobo MK, Schwartz DJ, and Kalivas PW (2015) Coding the direct/indirect pathways by D1 and D2 receptors is not valid for accumbens projections. *Nat Neurosci* **18**:1230–1232.
- Kuzhikandathil EV, Yu W, and Oxford GS (1998) Human dopamine D3 and D2L receptors couple to inward rectifier potassium channels in mammalian cell lines. *Mol Cell Neurosci* **12**:390–402.
- Lammel S, Hetzel A, Häckel O, Jones I, Liss B, and Roeper J (2008) Unique properties of mesoprefrontal neurons within a dual mesocorticolimbic dopamine system. *Neuron* **57**:760–773.
- Lammel S, Lim BK, and Malenka RC (2014) Reward and aversion in a heterogeneous midbrain dopamine system. *Neuropharmacology* **76** (Pt B):351–359.
- Lammel S, Steinberg EE, Földy C, Wall NR, Beier K, Luo L, and Malenka RC (2015) Diversity of transgenic mouse models for selective targeting of midbrain dopamine neurons. *Neuron* **85**:429–438.
- Lavine N, Ethier N, Oak JN, Pei L, Liu F, Trieu P, Rebois RV, Bouvier M, Hebert TE, and Van Tol HH (2002) G protein-coupled receptors form stable complexes with inwardly rectifying potassium channels and adenylyl cyclase. *J Biol Chem* **277**:46010–46019.
- Lee JC, Wang LP, and Tsien JZ (2016) Dopamine rebound-excitation theory: putting brakes on PTSD. *Front Psychiatry* **7**:163.
- Lemos JC, Friend DM, Kaplan AR, Shin JH, Rubinstein M, Kravitz AV, and Alvarez VA (2016) Enhanced GABA transmission drives bradykinesia following loss of dopamine D2 receptor signaling. *Neuron* **90**:824–838.
- Li C, Sugam JA, Lowery-Gionta EG, McElligott ZA, McCall NM, Lopez AJ, McKlveen JM, Pleil KE, and Kash TL (2016) Mu opioid receptor modulation of dopamine neurons in the periaqueductal gray/dorsal raphe: a role in regulation of pain. *Neuropsychopharmacology* **41**:2122–2132.
- Lindeberg J, Usoskin D, Bengtsson H, Gustafsson A, Kylberg A, Söderström S, and Ebendal T (2004) Transgenic expression of Cre recombinase from the tyrosine hydroxylase locus. *Genesis* **40**:67–73.
- Lindgren N, Usiello A, Goins M, Haycock J, Erbs E, Greengard P, Hokfelt T, Borrelli E, and Fisone G (2003) Distinct roles of dopamine D2L and D2S receptor isoforms in the regulation of protein phosphorylation at presynaptic and postsynaptic sites. *Proc Natl Acad Sci USA* **100**:4305–4309.
- Lobo MK and Nestler EJ (2011) The striatal balancing act in drug addiction: distinct roles of direct and indirect pathway medium spiny neurons. *Front Neuroanat* **5**:41.
- Locke TM, Soden ME, Miller SM, Hunker A, Knakal C, Licholai JA, Dhillion KS, Keene CD, Zweifel LS, and Carlson ES (2018) Dopamine D1 receptor-positive neurons in the lateral nucleus of the cerebellum contribute to cognitive behavior. *Biol Psychiatry* **84**:401–412.
- Löffler S, Körber J, Nubbemeyer U, and Fehsel K (2012) Comment on “Impaired respiratory and body temperature control upon acute serotonergic neuron inhibition”. *Science* **337**:646–, author reply 646.
- Lopez-Huerta VG, Nakano Y, Bausenwein J, Jaidar O, Lazarus M, Cherasse Y, Garcia-Munoz M, and Arbutnot G (2016) The neostriatum: two entities, one structure? *Brain Struct Funct* **221**:1737–1749.
- Luo Y-J, Li Y-D, Wang L, Yang S-R, Yuan X-S, Wang J, Cherasse Y, Lazarus M, Chen J-F, Qu W-M, et al. (2018) Nucleus accumbens controls wakefulness by a subpopulation of neurons expressing dopamine D₁ receptors. *Nat Commun* **9**:1576.
- Maclaren DA, Browne RW, Shaw JK, Krishnan Radhakrishnan S, Khare P, España RA, and Clark SD (2016) Clozapine N-oxide administration produces behavioral effects in Long-Evans rats: implications for designing DREADD experiments. *eNeuro* **3**.
- Mahler SV and Aston-Jones G (2018) CNO evil? considerations for the use of DREADDs in behavioral neuroscience. *Neuropsychopharmacology* **43**:934–936.
- Mahler SV, Vazey EM, Beckley JT, Keistler CR, McGlinchey EM, Kauffling J, Wilson SP, Deisseroth K, Woodward JJ, and Aston-Jones G (2014) Designer receptors show role for ventral pallidum input to ventral tegmental area in cocaine seeking. *Nat Neurosci* **17**:577–585.
- Manson A, Stirpe P, and Schrag A (2012) Levodopa-induced-dyskinesias clinical features, incidence, risk factors, management and impact on quality of life. *J Parkinsons Dis* **2**:189–198.
- Marchant NJ, Campbell EJ, Whitaker LR, Harvey BK, Kaganovsky K, Adhikary S, Hope BT, Heins RC, Priszczano TE, Vardy E, et al. (2016a) Role of ventral subiculum in context-induced relapse to alcohol seeking after punishment-imposed abstinence. *J Neurosci* **36**:3281–3294.
- Marchant NJ, Whitaker LR, Bossert JM, Harvey BK, Hope BT, Kaganovsky K, Adhikary S, Priszczano TE, Vardy E, Roth BL, et al. (2016b) Behavioral and physiological effects of a novel kappa-opioid receptor-based DREADD in rats. *Neuropsychopharmacology* **41**:402–409.
- Martin CK, Redman LM, Zhang J, Sanchez M, Anderson CM, Smith SR, and Ravussin E (2011) Lorcaserin, a 5-HT(2C) receptor agonist, reduces body weight by decreasing energy intake without influencing energy expenditure. *J Clin Endocrinol Metab* **96**:837–845.
- Mayer D, Kahl E, Uzuneser TC, and Fendt M (2018) Role of the mesolimbic dopamine system in relief learning. *Neuropsychopharmacology* **43**:1651–1659.
- Michaelides M, Anderson SAR, Ananth M, Smirnov D, Thanos PK, Neumaier JF, Wang GJ, Volkow ND, and Hurd YL (2013) Whole-brain circuit dissection in free-moving animals reveals cell-specific mesocorticolimbic networks. *J Clin Invest* **123**:5342–5350.
- Moehle MS, Pancani T, Byun N, Yohn SE, Wilson GH III, Dickerson JW, Remke DH, Xiang Z, Niswender CM, Wess J, et al. (2017) Cholinergic projections to the substantia nigra pars reticulata inhibit dopamine modulation of basal ganglia through the M4 muscarinic receptor. *Neuron* **96**:1358–1372.e4.
- Moorman DE, James MH, McGlinchey EM, and Aston-Jones G (2015) Differential roles of medial prefrontal subregions in the regulation of drug seeking. *Brain Res* **1628** (Pt A):130–146.
- Morales M and Margolis EB (2017) Ventral tegmental area: cellular heterogeneity, connectivity and behaviour. *Nat Rev Neurosci* **18**:73–85.
- Morales M and Root DH (2014) Glutamate neurons within the midbrain dopamine regions. *Neuroscience* **282**:60–68.
- Nair-Roberts RG, Chatelain-Badie SD, Benson E, White-Cooper H, Bolam JP, and Ungless MA (2008) Stereological estimates of dopaminergic, GABAergic and glutamatergic neurons in the ventral tegmental area, substantia nigra and retrorubral field in the rat. *Neuroscience* **152**:1024–1031.
- Nakajima K and Wess J (2012) Design and functional characterization of a novel, arrestin-biased designer G protein-coupled receptor. *Mol Pharmacol* **82**:575–582.
- Napolitano F, Bonito-Oliva A, Federici M, Carta M, Errico F, Magara S, Martella G, Nisticò R, Centonze D, Pisani A, et al. (2010) Role of aberrant striatal dopamine D1 receptor/cAMP/protein kinase A/DARPP32 signaling in the paradoxical calming effect of amphetamine. *J Neurosci* **30**:11043–11056.
- Neve KA, Seamans JK, and Trantham-Davidson H (2004) Dopamine receptor signaling. *J Recept Signal Transduct Res* **24**:165–205.
- Oishi Y, Suzuki Y, Takahashi K, Yonezawa T, Kanda T, Takata Y, Cherasse Y, and Lazarus M (2017a) Activation of ventral tegmental area dopamine neurons produces wakefulness through dopamine D₂-like receptors in mice. *Brain Struct Funct* **222**:2907–2915.
- Oishi Y, Xu Q, Wang L, Zhang BJ, Takahashi K, Takata Y, Luo YJ, Cherasse Y, Schiffmann SN, de Kerchove d'Exaerde A, et al. (2017b) Slow-wave sleep is controlled by a subset of nucleus accumbens core neurons in mice. *Nat Commun* **8**:734.
- Padgett CL, Lalive AL, Tan KR, Terunuma M, Munoz MB, Pangalos MN, Martínez-Hernández J, Watanabe M, Moss SJ, Luján R, et al. (2012) Methamphetamine-evoked depression of GABA(B) receptor signaling in GABA neurons of the VTA. *Neuron* **73**:978–989.
- Padovan-Hernandez Y and Knackstedt LA (2018) Dose-dependent reduction in cocaine-induced locomotion by clozapine-N-oxide in rats with a history of cocaine self-administration. *Neurosci Lett* **674**:132–135.
- Parker PRL, Lalive AL, and Kreitzer AC (2016) Pathway-specific remodeling of thalamostriatal synapses in Parkinsonian mice. *Neuron* **89**:734–740.
- Pascoli V, Terrier J, Hiver A, and Lüscher C (2015) Sufficiency of mesolimbic dopamine neuron stimulation for the progression to addiction. *Neuron* **88**:1054–1066.
- Patriarchi T, Cho JR, Merten K, Howe MW, Marley A, Xiong WH, Folk RW, Broutchard G, Liang R, Jang MJ, et al. (2018) Ultrafast neuronal imaging of dopamine dynamics with designed genetically encoded sensors. *Science* **360**:eaat4422.
- Patterson CM, Wong JM, Leininger GM, Allison MB, Mabrouk OS, Kasper CL, Gonzalez JE, Mackenzie A, Jones JC, Kennedy RT, et al. (2015) Ventral tegmental area neurotransmitter signaling links the lateral hypothalamus to locomotor activity and striatal dopamine efflux in male mice. *Endocrinology* **156**:1692–1700.
- Peters J, Kalivas PW, and Quirk GJ (2009) Extinction circuits for fear and addiction overlap in prefrontal cortex. *Learn Mem* **16**:279–288.
- Piazza PV, Deminière JM, Le Moal M, and Simon H (1989) Factors that predict individual vulnerability to amphetamine self-administration. *Science* **245**:1511–1513.
- Piazza PV, Deroche-Gamont V, Rouge-Pont F, and Le Moal M (2000) Vertical shifts in self-administration dose-response functions predict a drug-vulnerable phenotype predisposed to addiction. *J Neurosci* **20**:4226–4232.
- Piazza PV, Rouge-Pont F, Deminière JM, Kharoubi M, Le Moal M, and Simon H (1991) Dopaminergic activity is reduced in the prefrontal cortex and increased in the nucleus accumbens of rats predisposed to develop amphetamine self-administration. *Brain Res* **567**:169–174.
- Pijnenburg AJ, Honig WM, Van der Heyden JA, and Van Rossum JM (1976) Effects of chemical stimulation of the mesolimbic dopamine system upon locomotor activity. *Eur J Pharmacol* **35**:45–58.
- Pina MM and Cunningham CL (2017) Ethanol-seeking behavior is expressed directly through an extended amygdala to midbrain neural circuit. *Neurobiol Learn Mem* **137**:83–91.
- Pina MM, Young EA, Ryabinin AE, and Cunningham CL (2015) The bed nucleus of the stria terminalis regulates ethanol-seeking behavior in mice. *Neuropharmacology* **99**:627–638.
- Poulin JF, Caronia G, Hofer C, Cui Q, Helm B, Ramakrishnan C, Chan CS, Dombeck DA, Deisseroth K, and Awatramani R (2018) Mapping projections of molecularly defined dopamine neuron subtypes using intersectional genetic approaches. *Nat Neurosci* **21**:1260–1271.
- Prasad AA and McNally GP (2016) Ventral pallidum output pathways in context-induced reinstatement of alcohol seeking. *J Neurosci* **36**:11716–11726.
- Purohit K, Parekh PK, Kern J, Logan RW, Liu Z, Huang Y, McClung CA, Crabbe JC, and Ozburn AR (2018) Pharmacogenetic manipulation of the nucleus accumbens alters binge-like alcohol drinking in mice. *Alcohol Clin Exp Res* **42**:879–888.
- Qualls-Creekmore E, Yu S, Francois M, Hoang J, Huesing C, Bruce-Keller A, Burk D, Berthoud HR, Morrison CD, and Münzberg H (2017) Galanin-expressing GABA neurons in the lateral hypothalamus modulate food reward and noncompulsive locomotion. *J Neurosci* **37**:6053–6065.
- Richardson NR and Roberts DC (1996) Progressive ratio schedules in drug self-administration studies in rats: a method to evaluate reinforcing efficacy. *J Neurosci Methods* **66**:1–11.
- Richfield EK, Young AB, and Penney JB (1987) Comparative distribution of dopamine D-1 and D-2 receptors in the basal ganglia of turtles, pigeons, rats, cats, and monkeys. *J Comp Neurol* **262**:446–463.
- Robbins TW (2002) The 5-choice serial reaction time task: behavioural pharmacology and functional neurochemistry. *Psychopharmacology (Berl)* **163**:362–380.
- Robinson SE and Sohal VS (2017) Dopamine D2 receptors modulate pyramidal neurons in mouse medial prefrontal cortex through a stimulatory G-protein pathway. *J Neurosci* **37**:10063–10073.

- Roelofs TJM, Verharen JPH, van Tilborg GAF, Boekhoudt L, van der Toorn A, de Jong JW, Luijendijk MCM, Otte WM, Adan RAH, and Dijkhuizen RM (2017) A novel approach to map induced activation of neuronal networks using chemogenetics and functional neuroimaging in rats: a proof-of-concept study on the mesocorticolimbic system. *Neuroimage* **156**:109–118.
- Roman CW, Derkach VA, and Palmiter RD (2016) Genetically and functionally defined NTS to PBN brain circuits mediating anorexia. *Nat Commun* **7**:11905.
- Roth BL (2016) DREADDs for neuroscientists. *Neuron* **89**:683–694.
- Roy DS, Arons A, Mitchell TI, Pignatelli M, Ryan TJ, and Tonegawa S (2016) Memory retrieval by activating engram cells in mouse models of early Alzheimer's disease. *Nature* **531**:508–512.
- Runegaard AH, Jensen KL, Dencker D, Wörtwein G, and Gether U (2017a) Subjective perception of cocaine reward in mice assessed by a single exposure place preference (sePP) paradigm. *J Neurosci Methods* **289**:85–92.
- Runegaard AH, Jensen KL, Fitzpatrick CM, Dencker D, Weikop P, Gether U, and Rickhag M (2017b) Preserved dopaminergic homeostasis and dopamine-related behaviour in hemizygous TH-Cre mice. *Eur J Neurosci* **45**:121–128.
- Runegaard AH, Sørensen AT, Fitzpatrick CM, Jørgensen SH, Petersen AV, Hansen NW, Weikop P, Andreasen JT, Mikkelsen JD, Perrier JF, et al. (2018) Locomotor- and reward-enhancing effects of cocaine are differentially regulated by chemogenetic stimulation of Gi-signaling in dopaminergic neurons. *eNeuro* **5**.
- Salamone JD, Correa M, Nunes EJ, Randall PA, and Pardo M (2012) The behavioral pharmacology of effort-related choice behavior: dopamine, adenosine and beyond. *J Exp Anal Behav* **97**:125–146.
- Sandhu EC, Fernando ABP, Irvine EE, Tossell K, Kokkinou M, Glegola J, Smith MA, Howes OD, Withers DJ, and Ungless MA (2018) Phasic stimulation of midbrain dopamine neuron activity reduces salt consumption. *eNeuro* **5**:ENEURO.0064-18.2018.
- Savitt JM, Jang SS, Mu W, Dawson VL, and Dawson TM (2005) Bcl-x is required for proper development of the mouse substantia nigra. *J Neurosci* **25**:6721–6728.
- Schultz W (2007) Multiple dopamine functions at different time courses. *Annu Rev Neurosci* **30**:259–288.
- Schwarz LA, Miyamichi K, Gao XJ, Beier KT, Weissbourd B, DeLoach KE, Ren J, Ibanes S, Malenka RC, Kremer EJ, et al. (2015) Viral-genetic tracing of the input-output organization of a central noradrenergic circuit. *Nature* **524**:88–92.
- Scofield MD, Boger HA, Smith RJ, Li H, Haydon PG, and Kalivas PW (2015) Gq-DREADD selectively initiates glial glutamate release and inhibits cue-induced cocaine seeking. *Biol Psychiatry* **78**:441–451.
- Sharpe AL, Varela E, Bettinger L, and Beckstead MJ (2014) Methamphetamine self-administration in mice decreases GIRK channel-mediated currents in midbrain dopamine neurons. *Int J Neuropsychopharmacol* **18**:1–10.
- Smith KS, Bucci DJ, Luikart BW, and Mahler SV (2016) DREADDs: use and application in behavioral neuroscience. *Behav Neurosci* **130**:137–155.
- Söderpalm B and Ericson M (2013) Neurocircuitry involved in the development of alcohol addiction: the dopamine system and its access points. *Curr Top Behav Neurosci* **13**:127–161.
- Stamatakis AM, Jennings JH, Ung RL, Blair GA, Weinberg RJ, Neve RL, Boyce F, Mattis J, Ramakrishnan C, Deisseroth K, et al. (2013) A unique population of ventral tegmental area neurons inhibits the lateral habenula to promote reward. *Neuron* **80**:1039–1053.
- Stauffer WR, Lak A, Yang A, Borel M, Paulsen O, Boyden ES, and Schultz W (2016) Dopamine neuron-specific optogenetic stimulation in rhesus macaques. *Cell* **166**:1564–1571.e6.
- Strader CD, Gaffney T, Sugg EE, Candelore MR, Keys R, Patchett AA, and Dixon RA (1991) Allele-specific activation of genetically engineered receptors. *J Biol Chem* **266**:5–8.
- Stuber GD, Stamatakis AM, and Katak PA (2015) Considerations when using cre-driver rodent lines for studying ventral tegmental area circuitry. *Neuron* **85**:439–445.
- Stuber GD, Hnasko TS, Britt JP, Edwards RH, and Bonci A (2010) Dopaminergic terminals in the nucleus accumbens but not the dorsal striatum corelease glutamate. *J Neurosci* **30**:8229–8233.
- Sulzer D (2011) How addictive drugs disrupt presynaptic dopamine neurotransmission. *Neuron* **69**:628–649.
- Sun F, Zeng J, Jing M, Zhou J, Feng J, Owen SF, Luo Y, Li F, Wang H, Yamaguchi T, et al. (2018) A genetically encoded fluorescent sensor enables rapid and specific detection of dopamine in flies, fish, and mice. *Cell* **174**:481–496.e19.
- Sun HX, Wang DR, Ye CB, Hu ZZ, Wang CY, Huang ZL, and Yang SR (2017) Activation of the ventral tegmental area increased wakefulness in mice. *Sleep Biol Rhythms* **15**:107–115.
- Svoboda K and Yasuda R (2006) Principles of two-photon excitation microscopy and its applications to neuroscience. *Neuron* **50**:823–839.
- Tervo DGR, Hwang B-Y, Viswanathan S, Gaj T, Lavzin M, Ritola KD, Lindo S, Michael S, Kuleshova E, Ojala D, et al. (2016) A designer AAV variant permits efficient retrograde access to projection neurons. *Neuron* **92**:372–382.
- Todd TP, Vurbic D, and Bouton ME (2014) Behavioral and neurobiological mechanisms of extinction in Pavlovian and instrumental learning. *Neurobiol Learn Mem* **108**:52–64.
- Torres GE and Amara SG (2007) Glutamate and monoamine transporters: new visions of form and function. *Curr Opin Neurobiol* **17**:304–312.
- Tritsch NX, Oh WJ, Gu C, and Sabatini BL (2014) Midbrain dopamine neurons sustain inhibitory transmission using plasma membrane uptake of GABA, not synthesis. *eLife* **3**:e01936.
- Tritsch NX and Sabatini BL (2012) Dopaminergic modulation of synaptic transmission in cortex and striatum. *Neuron* **76**:33–50.
- Ungerstedt U and Arbuthnott GW (1970) Quantitative recording of rotational behavior in rats after 6-hydroxy-dopamine lesions of the nigrostriatal dopamine system. *Brain Res* **24**:485–493.
- Urban DJ and Roth BL (2015) DREADDs (designer receptors exclusively activated by designer drugs): chemogenetic tools with therapeutic utility. *Annu Rev Pharmacol Toxicol* **55**:399–417.
- Valencia-Torres L, Olarte-Sánchez CM, Lyons DJ, Georgescu T, Greenwald-Yarnell M, Myers MG Jr., Bradshaw CM, and Heisler LK (2017) Activation of ventral tegmental area 5-HT2C receptors reduces incentive motivation. *Neuropsychopharmacology* **42**:1511–1521.
- Vardy E, Robinson JE, Li C, Olsen RHJ, DiBerto JF, Giguere PM, Sassano FM, Huang XP, Zhu H, Urban DJ, et al. (2015) A new DREADD facilitates the multiplexed chemogenetic interrogation of behavior. *Neuron* **86**:936–946.
- Vazey EM and Aston-Jones G (2013) New tricks for old dogmas: optogenetic and designer receptor insights for Parkinson's disease. *Brain Res* **1511**:153–163.
- Vengeliene V, Bilbao A, Molander A, and Spanagel R (2008) Neuropharmacology of alcohol addiction. *Br J Pharmacol* **154**:299–315.
- Volkow ND, Fowler JS, Wang GJ, and Swanson JM (2004) Dopamine in drug abuse and addiction: results from imaging studies and treatment implications. *Mol Psychiatry* **9**:557–569.
- Volkow ND, Fowler JS, Wang G, Swanson JM, and Telang F (2007) Dopamine in drug abuse and addiction: results of imaging studies and treatment implications. *Arch Neurol* **64**:1575–1579.
- Wakaizumi K, Kondo T, Hamada Y, Narita M, Kawabe R, Narita H, Watanabe M, Kato S, Senba E, Kobayashi K, et al. (2016) Involvement of mesolimbic dopaminergic network in neuropathic pain relief by treadmill exercise: a study for specific neural control with Gi-DREADD in mice. *Mol Pain* **12**:1–11.
- Wang H, La Russa M, and Qi LS (2016) CRISPR/Cas9 in genome editing and beyond. *Annu Rev Biochem* **85**:227–264.
- Wang S, Tan Y, Zhang JE, and Luo M (2013) Pharmacogenetic activation of midbrain dopaminergic neurons induces hyperactivity. *Neurosci Bull* **29**:517–524.
- Wang X-F, Liu J-J, Xia J, Liu J, Mirabella V, and Pang ZP (2015) Endogenous glucagon-like peptide-1 suppresses high-fat food intake by reducing synaptic drive onto mesolimbic dopamine neurons. *Cell Reports* **12**:726–733.
- Wasserman DI, Tan JMJ, Kim JC, and Yeomans JS (2016) Muscarinic control of rostromedial tegmental nucleus GABA neurons and morphine-induced locomotion. *Eur J Neurosci* **44**:1761–1770.
- Wayman WN and Woodward JJ (2018) Chemogenetic excitation of accumbens-projecting infralimbic cortical neurons blocks toluene-induced conditioned place preference. *J Neurosci* **38**:1462–1471.
- Welter M, Vallone D, Samad TA, Meziane H, Usiello A, and Borrelli E (2007) Absence of dopamine D2 receptors unmasks an inhibitory control over the brain circuitries activated by cocaine. *Proc Natl Acad Sci USA* **104**:6840–6845.
- Whissell PD, Tohyama S, and Martin LJ (2016) The use of DREADDs to deconstruct behavior. *Front Genet* **7**:70.
- Wise RA and Bozarth MA (1985) Brain mechanisms of drug reward and euphoria. *Psychiatr Med* **3**:445–460.
- Wunsch AM, Yager LM, Donckels EA, Le CT, Neumaier JF, and Ferguson SM (2017) Chemogenetic inhibition reveals midline thalamic nuclei and thalamo-accumbens projections mediate cocaine-seeking in rats. *Eur J Neurosci* **46**:1850–1862.
- Xu P, He Y, Cao X, Valencia-Torres L, Yan X, Saito K, Wang C, Yang Y, Hinton A Jr., Zhu L, et al. (2017) Activation of serotonin 2C receptors in dopamine neurons inhibits binge-like eating in mice. *Biol Psychiatry* **81**:737–747.
- Yan Z, Song WJ, and Surmeier J (1997) D2 dopamine receptors reduce N-type Ca²⁺ currents in rat neostriatal cholinergic interneurons through a membrane-delimited, protein-kinase-C-insensitive pathway. *J Neurophysiol* **77**:1003–1015.
- Zhong P, Liu X, Zhang Z, Hu Y, Liu SJ, Lezama-Ruiz M, Joksimovic M, and Liu Q-S (2014) Cyclin-dependent kinase 5 in the ventral tegmental area regulates depression-related behaviors. *J Neurosci* **34**:6352–6366.
- Zhou FM, Liang Y, Salas R, Zhang L, De Biasi M, and Dani JA (2005) Corelease of dopamine and serotonin from striatal dopamine terminals. *Neuron* **46**:65–74.
- Zhu X, Ottenheimer D, and DiLeone RJ (2016) Activity of D1/2 receptor expressing neurons in the nucleus accumbens regulates running, locomotion, and food intake. *Front Behav Neurosci* **10**:66.