ASSOCIATE EDITOR: ERIC BARKER

E1 Enzymes as Therapeutic Targets in Cancer^S

Samir H. Barghout and Aaron D. Schimmer

Department of Medical Biophysics, Faculty of Medicine, University of Toronto, Toronto, Ontario, Canada (S.H.B., A.D.S.); Princess Margaret Cancer Centre, University Health Network, Toronto, Ontario, Canada (S.H.B., A.D.S.); and Department of Pharmacology and Toxicology, Faculty of Pharmacy, Tanta University, Tanta, Egypt (S.H.B.)

Abstract	3
Significance Statement	3
I. Ubiquitin-Like Protein Conjugation System	3
II. E1 Enzymes	4
A. Members of the E1 Enzyme Class	4
B. Biochemical and Structural Mechanisms of E1-Catalyzed Ubiquitin-Like Protein Activation	6
C. Druggability of E1 Enzymes	7
D. Rationale for Targeting E1 Enzymes in Cancer	8
E. Biologic Functions of E1 Enzymes and Their Implications in Cancer Therapy	10
1. Proliferative Signaling and Growth Suppressors	10
2. Cell Cycle Progression	11
3. DNA Damage Response	11
4. Apoptosis and Autophagy	11
5. Inflammation, Immune Responses, and Tumor Microenvironment	11
6. Angiogenesis and Metastasis	12
7. Cellular Stress Phenotypes	12
F. Targeting E1 Versus E3 Enzymes	12
III. Tools and Assays for Exploring E1 Biology and Drug Discovery	12
A. ATP:PPi and ATP:AMP Exchange Assays	12
B. E1~ Ubiquitin-Like Protein Thioester Formation and Transthiolation Assays	13
C. Substrate Ubiquitin-Like Protein Conjugation Assays	13
D. Detection of Ubiquitin-Like Protein–Drug Adduct Formation	13
IV. Canonical E1 Enzymes	14
A. Neural Precursor Cell–Expressed Developmentally Downregulated Protein 8–Activating Enzyme	14
1. Role of Neural Precursor Cell–Expressed Developmentally Downregulated Protein 8–Activating Enzyme in Cancer	14
2. Neural Precursor Cell–Expressed Developmentally Downregulated Protein 8–Activating Enzyme Inhibitors	14
a. Pevonedistat	15
b. TAS4464	22

Address correspondence to: Dr. Aaron Schimmer, Princess Margaret Cancer Centre, University Health Network, Princess Margaret Cancer Research Tower, Rm. 8-706, 101 College St, Toronto, ON, M5G 1L7, Canada. E-mail: aaron.schimmer@uhn.ca

This work was supported by the Leukemia & Lymphoma Society of Canada, the Canadian Institutes of Health Research (CIHR), the Princess Margaret Cancer Centre Foundation, and the Ontario Ministry of Health and Long-Term Care. A.D.S. holds the Ronald N. Buick Chair in Oncology Research. S.H.B. was supported by the Ontario Trillium Scholarship, Department of Medical Biophysics Fellowship, Graduate Studies Endowment Fund (GSEF) Scholarship, and Queen Elizabeth II Scholarship from the Faculty of Medicine, University of Toronto, Canada.

A.D.S. has received honorariums or consulting fees from Novartis, Jazz, Otsuka, and Takeda Pharmaceuticals and research support from Medivir AB and Takeda Pharmaceuticals. A.D.S. owns stock in Abbvie Pharmaceuticals and is named on a patent application for the use of DNT cells for the treatment of leukemia.

^SThis article has supplemental material available at [pharmrev.aspetjournals.org](https://doi.org/10.1124/pharmrev.120.000053).
<https://doi.org/10.1124/pharmrev.120.000053>.

c.	Neural Precursor Cell–Expressed Developmentally Downregulated Protein 8 Adenylate Analogs and Neural Precursor Cell–Expressed Developmentally Downregulated Protein 8–Mimicking Peptides	22
d.	Natural and Semisynthetic Neural Precursor Cell–Expressed Developmentally Downregulated Protein 8–Activating Enzyme Inhibitors	22
e.	Other Natural and Semisynthetic Neural Precursor Cell–Expressed Developmentally Downregulated Protein 8–Activating Enzyme Inhibitors	23
B.	Ubiquitin-Like Modifier–Activating Enzyme 1	23
1.	Role of Ubiquitin-Like Modifier–Activating Enzyme 1 in Cancer	24
2.	Ubiquitin-Like Modifier–Activating Enzyme 1 Inhibitors	25
a.	TAK-243	25
b.	Ubiquitin Adenylate Analogs	27
c.	Natural Compounds	27
d.	Nitropyrazones and Nitropiperazines	28
e.	Disulfides	28
C.	Small Ubiquitin-Like Modifier–Activating Enzyme	28
1.	Role of Small Ubiquitin-Like Modifier–Activating Enzyme in Cancer	29
2.	Role of SUMOylation in Differentiation Therapy of Acute Promyelocytic Leukemia	29
3.	Antitumor Immunomodulatory Roles of SUMOylation	30
4.	Small Ubiquitin-Like Modifier–Activating Enzyme Inhibitors	32
a.	ML-792 and TAK-981	32
b.	COH000	33
c.	Small Ubiquitin-Like Modifier Adenylate Analogs and Small Ubiquitin-Like Modifier–Mimicking Peptides	33
d.	Natural Small Ubiquitin-Like Modifier–Activating Enzyme Inhibitors	33
e.	Urea-Based Small Ubiquitin-Like Modifier–Activating Enzyme Inhibitors	34
D.	Ubiquitin-Like Modifier–Activating Enzyme 6	35
1.	Role of Ubiquitin-Like Modifier–Activating Enzyme 6 in Cancer	36
2.	Ubiquitin-Like Modifier–Activating Enzyme 6 Inhibitors	36
E.	Ubiquitin-Like Modifier–Activating Enzyme 7	36
1.	Role of Ubiquitin-Like Modifier–Activating Enzyme 7 in Cancer	37
2.	Ubiquitin-Like Modifier–Activating Enzyme 7 Inhibitors	38
V.	Noncanonical E1 Enzymes	38
A.	Autophagy-Related Protein 7	38
1.	Role of Autophagy-Related Protein 7 in Cancer	38
2.	Autophagy-Related Protein 7 Inhibitors	39
B.	Ubiquitin-Like Modifier–Activating Enzyme 4	39
1.	Role of Ubiquitin-Like Modifier–Activating Enzyme 4 in Cancer	40
2.	Ubiquitin-Like Modifier–Activating Enzyme 4 Inhibitors	40
C.	Ubiquitin-Like Modifier–Activating Enzyme 5	40

ABBREVIATIONS: AAD, active adenylation domain; ABP, activity-based probe; ABPA3, activity-based probe A3; AD, adenylation domain; ADS, adenosine 5'-sulfamate; AML, acute myeloid leukemia; AMSN, 5'-(sulfonaminodeoxy)adenosine; APL, acute promyelocytic leukemia; APN, adenosyl-phospho-NEDD8; APU, adenosyl-phospho-ubiquitinol; ATG, autophagy-related protein; ATO, arsenic trioxide; ATRA, all-trans retinoic acid; CCD, catalytic cysteine domain; CDT1, chromatin licensing and DNA replication factor 1; cGAS, cGMP-AMP synthase; CNV, copy number variation; CRL, cullin-RING ubiquitin ligase; CTD, C-terminal domain; DUB, deubiquitylase; ECTD, extreme C-terminal domain; EGFR, epidermal growth factor receptor; ER, endoplasmic reticulum; FAT10, human leukocyte antigen F-associated transcript 10; HDAC, histone deacetylase; HIF, hypoxia-inducible factor; hLRH-1, human liver receptor homolog-1; HTS, high-throughput screen; IAD, inactive adenylation domain; I κ B α , inhibitor of NF- κ B α ; IFN, interferon; IFNAR, IFN- α receptor; ISG, interferon-stimulated gene; MDS, myelodysplastic syndrome; MM, multiple myeloma; MOCS3, molybdenum cofactor synthesis protein 3; NAE, NEDD8-activating enzyme; NB, nuclear body; NEDD8, neural precursor cell-expressed developmentally downregulated protein 8; NEMO, NF- κ B essential modulator; NF- κ B, nuclear factor κ B; NTD, N-terminal domain; PCNA, proliferating cell nuclear antigen; PE, phosphatidylethanolamine; PK, pharmacokinetic; PML, promyelocytic leukemia; RanGAP1, Ran GTPase-activating protein 1; RARA, retinoic acid receptor α ; RHD, rhodanese homology domain; RING, really interesting new gene; ROS, reactive oxygen species; SAE, SUMO-activating enzyme; SAR, structure-activity relationship; SENP, SUMO-specific protease; SUMO, small ubiquitin-like modifier; $t_{1/2}$, terminal elimination half-life time; TRAIL, tumor necrosis factor-related apoptosis-inducing ligand; UBA, ubiquitin-like modifier-activating enzyme; UBL, ubiquitin-like protein; UFC1, UFM1-conjugating enzyme 1; UFD, ubiquitin-fold domain; UFM1, ubiquitin-fold modifier 1; UFSP, UFM1-specific protease; UIS, UFM1-interacting sequence; ULP, UBL-specific protease; UPR, unfolded protein response; UPS, ubiquitin-proteasome system; URM1, ubiquitin-related modifier 1; USP, ubiquitin-specific protease.

1. Role of Ubiquitin-Like Modifier–Activating Enzyme 5 in Cancer	41
2. Ubiquitin-Like Modifier–Activating Enzyme 5 Inhibitors.....	41
VI. Dual and Multi-E1 Inhibitors	42
A. Multi-E1 Adenosine Sulfamate Inhibitors.....	43
B. LP0040	44
VII. Adenosine Sulfamate E1 Inhibitors.....	44
A. Discovery and Structure-Activity Relationship	44
B. Substrate-Assisted Inhibition.....	44
C. Selectivity and Potency of Adenosine Sulfamate E1 Inhibitors.....	45
D. Resistance to Adenosine Sulfamate E1 Inhibitors	46
E. Pharmacodynamic Activity of Adenosine Sulfamate E1 Inhibitors	47
F. Cell Cycle Effects of Adenosine Sulfamate E1 Inhibitors	48
G. Pharmacokinetics of Adenosine Sulfamate E1 Inhibitors	48
H. Insights into Clinical Trials of E1 Inhibitors	49
VIII. Concluding Remarks.....	49
References.....	49

Abstract—Post-translational modifications of cellular substrates with ubiquitin and ubiquitin-like proteins (UBLs), including ubiquitin, SUMOs, and neural precursor cell-expressed developmentally downregulated protein 8, play a central role in regulating many aspects of cell biology. The UBL conjugation cascade is initiated by a family of ATP-dependent enzymes termed E1 activating enzymes and executed by the downstream E2-conjugating enzymes and E3 ligases. Despite their druggability and their key position at the apex of the cascade, pharmacologic modulation of E1s with potent and selective drugs has remained elusive until 2009. Among the eight E1 enzymes identified so far, those initiating ubiquitylation (UBA1), SUMOylation (SAE), and neddylation (NAE) are the most characterized and are implicated in various aspects of cancer biology. To date, over 40 inhibitors have been reported to target UBA1, SAE, and NAE, including the NAE inhibitor pevonedistat, evaluated in more than 30

clinical trials. In this Review, we discuss E1 enzymes, the rationale for their therapeutic targeting in cancer, and their different inhibitors, with emphasis on the pharmacologic properties of adenosine sulfamates and their unique mechanism of action, termed substrate-assisted inhibition. Moreover, we highlight other less-characterized E1s—UBA6, UBA7, UBA4, UBA5, and autophagy-related protein 7—and the opportunities for targeting these enzymes in cancer.

Significance Statement—The clinical successes of proteasome inhibitors in cancer therapy and the emerging resistance to these agents have prompted the exploration of other signaling nodes in the ubiquitin-proteasome system including E1 enzymes. Therefore, it is crucial to understand the biology of different E1 enzymes, their roles in cancer, and how to translate this knowledge into novel therapeutic strategies with potential implications in cancer treatment.

I. Ubiquitin-Like Protein Conjugation System

Ubiquitin is an 8.5-kDa, 76–amino acid polypeptide that serves as a post-translational modifier of cellular substrates (Bedford et al., 2011). Similar to other post-translational modifications, the process of reversible enzymatic attachment of ubiquitin to proteins is termed ubiquitylation (also known as ubiquitination). This post-translational modification is involved in modulating turnover, function, interaction, and/or localization of cellular proteins and therefore regulates a wide range of biologic processes (Rape, 2018). These include, among other functions, protein homeostasis, cell cycle regulation, DNA repair, transcriptional regulation, and endocytosis (Haglund et al., 2003; Ciechanover, 2005; Dhananjayan et al., 2005; Huang and D’Andrea, 2006; Gilberto and Peter, 2017). Given its indispensable biologic roles, ubiquitin is evolutionarily conserved among eukaryotes and, as the name suggests, is ubiquitously expressed in most tissues (Goldstein et al., 1975).

In eukaryotes, a group of proteins collectively known as ubiquitin-like proteins (UBLs) share sequence homology and have a similar three-dimensional structure to ubiquitin (Hochstrasser, 2009; van der Veen and Ploegh, 2012). There are more than a dozen UBLs, which are classified into eight families, including neural precursor cell-expressed developmentally down-regulated protein 8 (NEDD8), small ubiquitin-like modifier (SUMO), human leukocyte antigen F-associated transcript 10 (FAT10/ubiquitin D), interferon-stimulated gene (ISG) 15, autophagy-related protein (ATG) 8, ATG12, ubiquitin-fold modifier 1 (UFM1), and ubiquitin-related modifier 1 (URM1) protein families. These UBLs, known as type I UBLs, are similarly involved in post-translational modifications that modulate a wide variety of cellular processes (Cappadocia and Lima, 2018). In contrast, type II UBLs are not conjugated to cellular substrates but rather exist as a part of proteins with multiple domains, and these include ubiquitin-like 5 (UBL5)/HTLV-I U5RE-binding protein 1 (HUB1) and *fau* and its ubiquitin-like domain (FUBI) (Rossman

et al., 2003; Oka et al., 2014). Here, our focus will be on ubiquitin and type I UBLs, which we will simply refer to as UBLs.

Post-translational modifications with UBLs are brought about by a sequential enzymatic cascade that involves three classes of enzymes: E1 activating enzymes, E2 conjugating enzymes, and E3 ligases (Nalepa et al., 2006) (Fig. 1). The function of E1 enzymes is to activate UBLs in an ATP-dependent multistep enzymatic reaction whereby the UBL becomes attached to the catalytic cysteine residue of the E1 enzyme via a high-energy thioester bond [symbolized by a tilde (~)], forming a UBL-loaded E1 enzyme (E1~UBL). The E1-bound UBL is then transferred from the catalytic cysteine of the E1 enzyme to the catalytic cysteine of a cognate E2 enzyme via a transthioesterification reaction, forming a UBL-loaded E2 enzyme (E2~UBL) (Cappadocia and Lima, 2018). In concert with their cognate E3 ligases, the UBL is conjugated to cellular proteins and/or phospholipids by the E2 enzyme, forming a UBL-modified substrate usually via an isopeptide linkage with a lysine residue (Schulman and Harper, 2009). E3 ligases may function as either adaptors that bring E2s in close proximity to cellular substrates for UBL conjugation (represented by U-box E3s and RING finger E3s) or UBL acceptors that form thioester intermediates with UBLs for subsequent transfer to their substrates (represented by homologous to the E6-AP carboxyl terminus [HECT] domain E3s) (Nakayama and Nakayama, 2006). In the human proteome, there exist eight known E1s, >40 E2s, and >600 E3s, with the E3 ligases controlling substrate specificity (Clague et al., 2015). As UBL conjugation is a reversible process, the UBL signal can be removed by another set of enzymes, known collectively as deubiquitylases (DUBs) and UBL-specific proteases (ULPs), which catalyze proteolytic cleavage of these post-translational modifications from their substrates. There exist more than 100 DUBs and ULPs in the human proteome (Clague et al., 2015; Ronau et al., 2016). A subset of ubiquitylated proteins, specifically those tagged with K48- and K11-linked polyubiquitin chains, are identified by the proteasome, deubiquitylated, and degraded into peptides to maintain protein homeostasis in the cell (Swatek and Komander, 2016). The cellular machinery orchestrating ubiquitin-dependent proteasomal degradation is collectively known as the ubiquitin-proteasome system (UPS) (Nalepa et al., 2006). Apart from degradative ubiquitylation, conjugation with other forms of ubiquitin (e.g., K63-linked polyubiquitin chains and monoubiquitin) and UBLs serves nondegradative functions in many aspects of cellular signaling (Bedford et al., 2011; Swatek and Komander, 2016).

Historically, drug discovery within the ubiquitin and UBL systems started with two parallel programs targeting the proteasome and E3 ligases. Despite the nonselective nature of proteasome inhibition, this

Fig. 1. Ubiquitin and UBL conjugation system. The UBL conjugation system comprises three enzyme classes that act sequentially to catalyze UBL conjugation: UBL-activating enzymes (E1), UBL-conjugating enzymes (E2), and UBL E3 ligases. Protein substrates are conjugated with different forms of UBLs. Proteins conjugated with Lys48 (K48)- or K11-linked polyubiquitin chains are recognized by the proteasome, the major proteolytic machinery in the cell that degrades such proteins into smaller peptides. Other forms of UBL conjugation are involved in various cellular signaling pathways. UBL conjugation is reversed by other classes of enzymes, including DUBs, which deconjugate ubiquitin signals, and ULPs, which deconjugate other UBL signals.

program has led to the discovery of clinically useful drugs (the prototype of which is bortezomib), whereas the E3 ligase program is making slower progress (Mattern et al., 2014). Moreover, researchers have explored the therapeutic targeting of other signaling nodes in these systems including E1 enzymes that lie at the apex of the UBL conjugation cascade and thereby control degradative as well as nondegradative signaling processes in the cell (Schulman and Harper, 2009).

In this Review, we highlight different E1 enzymes, their biologic roles, and pathobiologic alterations, particularly in the context of cancer. In addition, we discuss in more detail the biochemical mechanisms of UBL activation and therapeutic strategies known so far to target different E1 enzymes with a focus on mechanism-based E1 inhibitors that have been advanced to clinical trials.

II. E1 Enzymes

A. Members of the E1 Enzyme Class

In the human proteome, eight E1 enzymes are known to activate UBLs (Schulman and Harper, 2009; Clague et al., 2015). These include ubiquitin-like modifier-activating enzyme (UBA) 1, NEDD8-activating enzyme (NAE), SUMO-activating enzyme (SAE), UBA6, UBA7, UBA4 (MOCS3), UBA5, and autophagy-related protein (ATG) 7. Based on their structural and biochemical properties, E1s have been subdivided into canonical (UBA1, NAE, SAE, UBA6, UBA7) and noncanonical

Fig. 2. E1 enzymes and their structural domains. There are eight E1 enzymes known so far, including canonical (UBA1, UBA6, UBA7, SAE, and NAE) and noncanonical (UBA4, UBA5, and ATG7) E1 enzymes. Of these, UBA1, UBA6, and UBA7 are monomeric; SAE and NAE are heterodimeric; and UBA4, UBA5, and ATG7 are homodimeric. The AAD, catalytic cysteine half-domains (FCCH and SCCD), and UFD are conserved among canonical E1s. Noncanonical E1s do not possess discernible CCD, and the catalytic cysteine is located on the AD. They also comprise noncanonical E1-specific domains and sequences such as the RHD of UBA4. CC, catalytic cysteine; FCCH, first catalytic cysteine half-domain; SCCD, second catalytic cysteine half-domain; UFCBS, UFC1-binding sequence.

(UBA4, UBA5, and ATG7) E1s (Schulman and Harper, 2009). Structurally, E1 enzymes adopt a monomeric (UBA1, UBA6, and UBA7), heterodimeric (NAE and SAE), or homodimeric (UBA4, UBA5, and ATG7) architecture (Fig. 2).

To carry out their multistep catalytic function, eukaryotic canonical E1 enzymes possess multiple domains, including the adenylation domain (AD), the catalytic cysteine domain (CCD), and the ubiquitin-fold domain (UFD), with several variations that exist in the noncanonical E1 enzymes (Huang et al., 2004b).

In canonical E1s, the AD is pseudosymmetric, with one active adenylation domain (AAD) involved in identifying and adenylating the C terminus of cognate UBLs and another inactive adenylation domain (IAD) that may provide structural stability (Schulman and Harper, 2009; Lv et al., 2017b). Similarly, the CCD is divided into the first catalytic cysteine half-domain and

the second catalytic cysteine half-domain and is involved in thioester bond formation with UBLs (Lv et al., 2017b). The UFD is involved in the interaction with cognate E2s for UBL transfer via transthiolation. Of these, the AD is the most conserved domain and is homologous to adenylation domains in the ancestral prokaryotic enzymes molybdopterin biosynthetic enzyme B (MoeB) and thiamine biosynthesis protein F (ThiF), involved in molybdopterin and thiamine biosynthesis, respectively, suggesting that adenylation of UBLs is the most conserved functionality of all E1s (Huang et al., 2004b). Other domains might have evolved to accommodate the more complex catalytic activity carried out by eukaryotic E1s as opposed to their prokaryotic homologs (Cappadocia and Lima, 2018).

On the other hand, noncanonical E1s are homodimeric with symmetric ADs as well as other noncanonical E1-specific domains. In addition, their catalytic

TABLE 1
E1 enzymes, their cognate E2 enzymes, UBLs, and UBL proteases (Groettrup et al., 2008; Schulman and Harper, 2009)

E1	Structure	% Identity ^a	UBL	E2s	UBL Proteases	Amino Acid Number	Molecular Mass (kDa)
UBA1 (UBE1)	Monomer	100%	Ubiquitin	Multiple	DUBs	1058	117.85
UBA6 (UBE1L2)	Monomer	40%	Ubiquitin	USE1	DUBs	1052	117.97
UBA7 (UBE1L)	Monomer	46%	FAT10 ISG15	(UBE2Z) UBCH8 UBCH6	Unknown USP18	1012	111.69
SAE (SAE1-UBA2)	Heterodimer	SAE1: 30% UBA2: 17%	SUMO 1/2/3	UBC9	SENPs	SAE1: 346 UBA2: 640	SAE1: 38.45 UBA2: 71.22
NAE (NAE1-UBA3)	Heterodimer	NAE1: 17% UBA3: 21%	NEDD8	UBC12 UBE2Fs	CSN5 DEN1	NAE1: 534 UBA3: 463	NAE1: 60.25 UBA3: 51.85
UBA4 (MOCS3)	Homodimer	17%	URM1	—	—	460	49.67
UBA5 (UBE1DC1)	Homodimer	13%	UFM1	UFC1	UFSP1 UFSP2	404	44.86
ATG7	Homodimer	6%	ATG12 ATG8 family	ATG10 ATG3	—	703	77.96

UBE1, ubiquitin-activating enzyme E1.

^aPercent amino acid identity relative to UBA1.

Fig. 3. Cascade of canonical E1-catalyzed UBL activation and substrate-assisted inhibition by adenosine sulfamates. (A) Canonical E1-catalyzed UBL activation is a multistep catalytic process that involves UBL adenylation in an ATP-dependent manner forming an E1-AMP~UBL intermediate and releasing PP_i. The UBL at the adenylation domain is then attacked by the sulfhydryl (-SH) group to form an E1-S~UBL thioester intermediate associated with the release of AMP. The adenylation step is then repeated with another ATP molecule to form E1 doubly loaded with two UBL molecules at two distinct sites. The UBL at the catalytic cysteine is then attacked by the -SH group of the cognate E2 enzyme to transfer the UBL in a transthiolation reaction. ADSs inhibit E1 enzymes by attacking the E1-S~UBL intermediate (highlighted in a green box) and forming an ADS-UBL covalent adduct, which binds to the nucleotide-binding site of E1, preventing its utilization in subsequent reactions. R and R' correspond to the side chains in different ADSs. (B) The four catalytic steps of UBL activation and transfer to cognate E2s (first UBL adenylation, E1~UBL thioester formation, second UBL adenylation, and E1-E2 transthiolation) are illustrated.

cysteines are situated close to the adenylation pocket without having dedicated CCDs as opposed to canonical E1s (Cappadocia and Lima, 2018). UBA4 possesses a distinctive rhodanese homology domain (RHD) at its C terminus, which is involved in the sulfur transfer to URM1, the cognate UBL of UBA4 (Termathe and Leidel, 2018). Each of these E1s activates specific UBL(s) and transfers them to one or more of its cognate E2s, establishing distinct UBL→E1→E2→E3→substrate cascades to influence a broad range of cellular functions. Of these, the UBA1-initiated cascade is the most branched, with tens of E2s, hundreds of E3, and thousands of substrates (Clague et al., 2015) (Table 1).

B. Biochemical and Structural Mechanisms of E1-Catalyzed Ubiquitin-Like Protein Activation

UBL activation is a multistep ATP-dependent enzymatic process whereby the UBL moiety is attached to the catalytic cysteine of E1 enzyme via a thioester bond

for subsequent transfer to cognate E2s (Schulman and Harper, 2009). With their multiple functional domains, particularly the AD, E1s are catalytically competent to perform such activation as opposed to the simpler E2 enzymes (Stewart et al., 2016). The catalytic steps of UBL activation have been well characterized with canonical E1s, particularly the archetypal UBA1 enzyme (Bohnsack and Haas, 2003). There exist, however, some variations in the UBL activation mechanisms by noncanonical E1s (Schulman and Harper, 2009; Cappadocia and Lima, 2018). In general, the catalytic cascade of UBL activation and transfer by canonical E1s includes four major steps: 1) first adenylation, 2) thioester formation, 3) second adenylation, and 4) UBL transfer to E2 by transthiolation.

UBL activation starts with the C-terminal adenylation of a free UBL molecule in the presence of ATP and Mg²⁺, followed by binding of the adenylated UBL to the AAD of E1, forming an E1-bound UBL~adenylate (UBL~AMP)

Fig. 4. Inhibitors of E1 enzymes and their sites of action. A diagram of the E1 enzyme showing different E1 inhibitors and the structural domains or active sites they are reported to target.

intermediate and releasing PP_i . This is followed by a nucleophilic interdomain reaction whereby the catalytic cysteine of the CCD attacks the adenylated UBL, forming a UBL-bound E1 via a covalent thioester bond (E1~UBL), coupled with the release of free AMP. Subsequently, E1 catalyzes another round of adenylation of a second UBL molecule, releasing PP_i and forming a double-loaded E1 with two UBL molecules at two sites: one bound to the CCD via a covalent thioester bond and another bound to the AAD via a noncovalent bond (Haas and Rose, 1982). From energetic and conformational perspectives, this form is likely more competent for subsequent transfer of UBL (Schulman and Harper, 2009; Cappadocia and Lima, 2018). The double-loaded E1 then interacts with the cognate E2 enzyme through the UFD to transfer the thiol-bound UBL molecule to the corresponding thiol in the CCD of E2 via a transthioleation reaction, leaving an E1-bound UBL~AMP intermediate that is used in other rounds of UBL activation (Schulman and Harper, 2009) (Fig. 3).

The catalytic activity of E1s is enabled by several conformational changes that facilitate different steps of UBL activation. Of particular importance is the thioester bond formation facilitated by the rotation of the CCD to come in close proximity to the UBL~AMP and attack the UBL moiety (Olsen et al., 2010). The rotation is associated with other conformational changes that lead to remodeling of the AAD of E1 and switching from an open to a closed conformation (Streich and Lima, 2014). These conformational changes are transient, as the E1 restores the open conformation after forming the E1~UBL thioester. These structural insights were partly derived by using E1 inhibitors that serve as UBL~AMP mimetics, such as SUMO-AMSN and SUMO-AVSN (Olsen et al., 2010). Although all steps of UBL activation are bidirectional, the progression of these reactions is maintained in one direction by several biochemical and structural factors, including the

abundance of ATP, active site remodeling to promote thioester bond formation and PP_i release, and double loading with a second UBL molecule to regain the open conformation of the AAD and drive subsequent UBL transfer to cognate E2s (Cappadocia and Lima, 2018).

C. Druggability of E1 Enzymes

E1 enzymes catalyze the multistep UBL activation exploiting their multidomain structure, with at least two active sites that are readily amenable to therapeutic targeting (Fig. 4). The first active site is the nucleotide-binding pocket in the AAD to which the UBL~AMP binds. Historically, small-molecule ATP mimetics targeting mutant tyrosine kinases have been among the first classes of molecularly targeted agents to be developed for cancer therapy (Capdeville et al., 2002; Cohen, 2002). Therefore, E1 enzymes with such ATP-binding pockets can serve as typical druggable targets for modulating different UBL conjugation pathways. Moreover, the unique catalytic mechanism involving the interaction of UBL~AMP rather than ATP to the nucleotide-binding pocket sets a clear distinction between E1s and other ATP-dependent enzymes and offers an opportunity to develop selective inhibitors with little impact on the kinome or other ATP-dependent targets (Hubbard and Till, 2000). The challenge lies, however, in defining selectivity among the eight E1 enzymes that use conserved or closely related mechanisms of UBL activation. In this respect, the involvement of UBLs in the interaction with the nucleotide-binding pocket can be perceived as another source of selectivity even among related E1s owing to the structural variations of different UBLs. Therefore, structural information on these UBLs can be exploited to develop inhibitors that selectively target individual E1s.

For example, semisynthetic UBL~AMP analogs have been used as selective chemical probes to inhibit and interrogate the structural biology of E1s (Wilkinson et al., 1990; Olsen et al., 2010). With more structural and biochemical information revealed on different E1s, small-molecule E1 inhibitors with more favorable drug-like properties have been developed. Standing out among these are the adenosine sulfamate E1 inhibitors that form a UBL~AMP-like intermediate in situ after permeation into the cells overcoming the drug delivery issue experienced with semisynthetic analogs (Ciavarri and Langston, 2017).

The second active site is the catalytic cysteine in the CCD to which the UBL binds via a thioester bond. This catalytic residue with its redox-sensitive thiol group offers another opportunity for developing thiol-reactive electrophilic inhibitors that covalently modify the active site (Visscher et al., 2016). Such cysteine-directed covalent agents are expected to exert irreversible and potent inhibition, which is desirable in several contexts, including cancer therapy. Although many of these

covalent inhibitors may have off-target effects because of their promiscuous reactivity, a number of Food and Drug Administration–approved agents, including the kinase inhibitors rociletinib and osimertinib, target cysteine residues with a high level of selectivity (Visscher et al., 2016). For E1 enzymes, several drugs that target the CCD have been reported, most prominently the nitropyrazone-based UBA1 inhibitors PYR-41 and PYZD-4409 (Yang et al., 2007; Xu et al., 2010). Interestingly, COH000 is a recently discovered SAE inhibitor that targets a cysteine residue in the AAD without affecting the catalytic cysteine (Lv et al., 2018b; Li et al., 2019).

The third, and yet more challenging, site to target in E1 enzymes is the UFD through which they interact with their cognate E2s for subsequent UBL transfer. The difficulty of targeting E1-E2 interaction, like other protein-protein interactions, lies in the large surface area involved and the lack of pockets to which small-molecule inhibitors can bind (Jin et al., 2014). Added to these difficulties, many inhibitors targeting protein-protein interactions are peptide-based with less favorable drug-like properties (Vlieghe et al., 2010). An example of these inhibitors is UBC12N26, a 26-amino acid peptide that inhibits the interaction between NAE and its cognate E2 enzyme, UBC12. UBC12N26 corresponds to the N terminus of UBC12 and has been used to gain structural insights into NAE-UBC12 interaction (Huang et al., 2004a).

With all these druggable sites on E1s, there is ample opportunity to target these enzymes with structurally, mechanistically, and pharmacologically diverse drugs, including irreversible inhibitors that are well-suited for use in cancer therapy.

D. Rationale for Targeting E1 Enzymes in Cancer

As a genetic disease, cancer is initiated and maintained by tumor-specific oncogenic alterations of cellular proteins, a state known as oncogene addiction (Weinstein and Joe, 2006). However, the cellular stresses created as a result of malignant transformation and growth, such as proteotoxic, replicative, oxidative and metabolic stresses, need to be supported by other nonmutated, broadly acting cellular machineries that are essential for both normal and cancer cells (Luo et al., 2009). With these stresses, cancer cells become much more dependent on these machineries compared with normal cells, a state known as nononcogene addiction, which results in higher vulnerability to therapeutic agents that target these machineries. Although molecularly targeted agents directed mostly against tumor-specific oncogenes are relatively safer compared with cytotoxic agents, the emergence of resistance is a common pitfall that compromises the efficacy of such agents (Dobbelstein and Moll, 2014). Therefore, it has become increasingly important to explore targeting nononcogene addictions of cancer with drugs that are potentially

more effective than molecularly targeted agents and less toxic compared with cytotoxic therapy (Luo et al., 2009; Dobbelstein and Moll, 2014).

Proteasome inhibitors constitute a classic example of these drugs (Dobbelstein and Moll, 2014). They target the proteasome, which is an essential cellular machinery with broad cellular functions in both normal and cancer cells, including a key role in protecting against proteotoxic stress, particularly in myelomas that are engaged in immunoglobulin production (Hoeller and Dikic, 2009). Despite this essentiality and the expected toxicity of proteasome inhibition, bortezomib—the prototype of this class—is effective in malignancies that are highly dependent on the UPS, such as multiple myeloma (MM) and mantle cell lymphoma, with a therapeutic window that allows for a clinically acceptable safety profile in selected patient cohorts (Manasanch and Orlowski, 2017).

Similarly, several E1 enzymes are essential for supporting cellular stresses of cancer cells, including proteotoxic and DNA damage stress (Schulman and Harper, 2009). Of all E1s, UBA1 is the most essential enzyme whose loss-of-function is anticipated to be most deleterious to the survival and growth of cancer cells (Clague et al., 2015; Groen and Gillingwater, 2015). As assessed by the analysis of publicly available datasets of cancer cell line dependencies, only *UBA1* and *UBA2* (encoding the active subunit of SAE) genes are considered to be essential upon knockdown in large pancancer RNA interference screens (Tsherniak et al., 2017). However, the analysis of CRISPR/Cas9 knockout screens extends essentiality to *UBA3* (encoding the active subunit of NAE), *UBA4* and *UBA5*, as well as *SAE1* and *NAE1*, encoding the nonactive subunits of SAE and NAE, respectively (Fig. 5; Supplemental Figs. 1 and 2).

As highlighted above, E1 enzymes lie at the apex of the UBL conjugation cascade, and their targeting is expected to disrupt a variable range of biologic processes regulated by different UBLs (Schulman and Harper, 2009). For instance, UBA1 activates ubiquitylation required for both degradative and nondegradative cellular functions (Ulrich and Walden, 2010). Compared with proteasome inhibitors, UBA1 inhibitors are anticipated to induce broader, and thus more efficacious, biologic effects, as they disrupt proteasomal degradation as well as ubiquitin-regulated signaling pathways, such as DNA repair and nuclear factor-kappa B (NF- κ B) signaling (Bedford et al., 2011). Such higher efficacy may be needed in malignancies in which proteasome inhibitors suffer from intrinsic or acquired resistance (Manasanch and Orlowski, 2017). Targeting SAE is similarly expected to induce broad, yet less profound, effects as UBA1 inhibitors because of the indispensable role of SUMOylation in many signaling pathways (Gareau and Lima, 2010; Seeler and Dejean, 2017). Since neddylation is required to regulate a subset of ubiquitin E3

Fig. 5. Cancer dependence of E1 genes. Essentiality of E1 enzymes was assessed by pancancer genome-wide loss-of-function screens conducted in 710 (RNAi) and 582 (CRISPR) cancer cell lines. Donut charts reveal the percentage of cell lines dependent on the indicated genes encoding E1 or E1 subunits. A cell line is regarded as “dependent” when it has a probability of dependence greater than 0.5. These data were obtained by the analysis of publicly available datasets on the depmap portal at <https://depmap.org/portal>. RNAi, RNA interference.

ligases, targeting NAE induces a narrower spectrum of ubiquitylation-dependent biologic effects and is perhaps associated with higher tolerability compared with UBA1 and SAE inhibitors. This is consistent with the development of an NAE inhibitor, pevonedistat, as the first E1-targeted drug to enter clinical trials (Soucy et al., 2009a).

Given the important biologic functions of UBL conjugation pathways initiated by E1 enzymes and the perceived therapeutic window resulting from the higher dependence of cancer cells on these pathways, E1 enzymes can thus serve as attractive therapeutic targets in cancer. These specific biologic roles and their relevance to cancer biology

Fig. 6. Biologic functions of E1 enzymes and their implications in cancer therapy. E1 enzymes comprise eight members: UBA1, NAE, SAE, UBA6, UBA7, UBA4, UBA5, and ATG7. They initiate more than a dozen post-translational modifications, including ubiquitylation, neddylation, SUMOylation, FATylation, ISGylation, URMylation, UFMylation, and ATGylation. These modifications influence the biologic hallmark capabilities and cellular stress phenotypes (e.g., oxidative and proteotoxic stress) of cancer by modulating the activity, expression, stability, and/or localization of a wide range of signaling molecules. Therefore, targeting E1 enzymes may elicit useful antitumor effects in a context-dependent manner.

are detailed below and in sections IV. *Canonical E1 Enzymes* and V. *Noncanonical E1 Enzymes*.

E. Biologic Functions of E1 Enzymes and Their Implications in Cancer Therapy

Cancer is supported by a number of biologic hallmark capabilities and cellular stress phenotypes that enable tumor development and progression (Luo et al., 2009; Hanahan and Weinberg, 2011). As E1 enzymes initiate more than a dozen post-translational modifications that influence a wide array of cellular substrates, and with the functional diversity and breadth of such modifications, it is not surprising that E1s play a paramount role in almost every aspect of cancer cell biology (Welchman et al., 2005) (Fig. 6). In this section, we briefly summarize the biologic functions of E1-initiated post-translational modifications and their impact on key signaling pathways and cellular processes of relevance to cancer.

1. *Proliferative Signaling and Growth Suppressors.* Over the course of tumor development and progression, tumor cells accumulate genetic aberrations and other alterations that promote their proliferation and allow them to evade growth suppressors (Hanahan and Weinberg, 2011). The E1-initiated post-translational modifications are implicated in such rewiring of cancer cell signaling via proteolytic and nonproteolytic

mechanisms (Pérez-Benavente et al., 2020). In this respect, the activity of several growth factor receptors (e.g., EGFR, Met, platelet-derived growth factor receptor, and prolactin receptor) and their downstream kinases is normally regulated by ubiquitylation and subsequent proteasomal degradation. Several oncogenic alterations involve the stabilization of these proteins by downregulating their degradation, resulting in constitutive mitogenic signaling (Huangfu and Fuchs, 2010). In these contexts, UBA1 and UBA6, which initiate ubiquitylation, and NAE, which initiates the neddylation of a subset of ubiquitin E3 ligases enhancing their function, are particularly important. Therapeutic inhibition of these E1 enzymes in cancers driven mainly by such alterations may cause further activation of mitogenic signaling, resulting in unfavorable outcomes. Conversely, the antitumor activity of several tumor-suppressive proteins is reduced by alterations that enhance their ubiquitin-dependent proteasomal degradation (Huangfu and Fuchs, 2010). In such contexts, therapeutic inhibition of the ubiquitin-activating E1s or NAE is anticipated to stabilize these tumor suppressors and slow tumor progression. For instance, the well known tumor suppressors p53 and retinoblastoma-associated protein (Rb) are susceptible to proteasomal degradation after interaction with the ubiquitin E3 ligase double minute 2 protein (MDM2),

which facilitates their ubiquitylation (Chène, 2003; Uchida *et al.*, 2005). SUMOylation has also been reported to regulate the activity of tumor suppressors [e.g., p53 and phosphatase and tensin homolog (PTEN)] via nonproteolytic mechanisms, including the alteration of cellular localization (Stindt *et al.*, 2011; Bassi *et al.*, 2013).

2. Cell Cycle Progression. Ubiquitylation, neddylation, and SUMOylation play highly coordinated roles in orchestrating DNA replication, chromosome segregation, and mitosis in a spatiotemporal manner and mainly through nonproteolytic mechanisms (Eifler and Vertegaal, 2015; Gilberto and Peter, 2017). These modifications span a broad spectrum of cell cycle regulators, including, among others, cyclin-dependent kinases, cyclins, topoisomerases, and E3 ligases (Nakayama and Nakayama, 2006; Teixeira and Reed, 2013; Cuijpers and Vertegaal, 2018). As cancer cells proliferate at much higher rates compared with normal cells, they are highly dependent on these UBL conjugation machineries including E1 enzymes to support their proliferation. Therefore, inhibition of UBA1, NAE, and SAE is anticipated to cause cell cycle defects, replicative stress, and mitotic stress, and if sustained and excessive, these will ultimately lead to cell death (Hoeller *et al.*, 2006; Luo *et al.*, 2009). Specific cell cycle defects ensuing from the inhibition of different E1s are detailed in Sections IV. *Canonical E1 Enzymes*, V. *Noncanonical E1 Enzymes*, VI. *Dual and Multi-E1 Inhibitors*, and VII. *Adenosine Sulfamate E1 Inhibitors*.

3. DNA Damage Response. After exposure to genotoxic stress, ubiquitylation and SUMOylation are particularly important for coordinating the DNA damage response that comprises lesion sensing, DNA repair, and/or damage tolerance (Bergink and Jentsch, 2009). They are actively implicated in the different DNA repair pathways, including base-excision repair, nucleotide-excision repair, double-strand break repair, Fanconi anemia pathway, and translesion synthesis (Ulrich and Walden, 2010; Schwertman *et al.*, 2016). These modifications alter protein stability and/or localization and guide repair proteins to the site of the lesion (Ulrich, 2014). Examples of DNA damage response proteins that are amenable to such modifications are proliferating cell nuclear antigen (PCNA), Fanconi anemia group D2 protein (FANCD2), MDC1, xeroderma pigmentosum group C-complementing protein (XPC), thymine-DNA glycosylase, and H2A/B (Huang and D'Andrea, 2006; Jackson and Durocher, 2013; Dantuma and van Attikum, 2016). Inhibition of UBA1 and SAE is associated with the disruption of the DNA damage response and induction of DNA damage stress (Luo *et al.*, 2009; Hyer *et al.*, 2018).

4. Apoptosis and Autophagy. The signaling pathways that govern different forms of cell death are also amenable to regulation by E1-initiated post-translational modifications, particularly ubiquitylation. In this respect, the levels of pro- and antiapoptotic proteins are

controlled by ubiquitin-dependent proteasomal degradation. Caspases are also regulated by nondegradative ubiquitylation (Broemer and Meier, 2009; Vucic *et al.*, 2011). Tumor necrosis factor-related apoptosis-inducing ligand (TRAIL) receptors DR4 and DR5, which are involved in the extrinsic apoptosis pathway, may be endocytosed and/or downregulated in response to their ubiquitylation (Huangfu and Fuchs, 2010). Therefore, the stabilization of proapoptotic proteins [e.g., Bcl-2-associated X protein (Bax), Bcl-2 homologous antagonist killer (Bak), Bcl-2-interacting mediator of cell death (Bim), BH3-interacting domain death agonist (Bid), p53 up-regulated modulator of apoptosis (Puma), Bcl-2-modifying factor (Bmf), Bcl-2-associated death promoter (Bad), Noxa, caspases, and DR4/5] upon UBA1 or NAE inhibition may be associated with useful anticancer effects in certain contexts (Chen *et al.*, 2016a; Hyer *et al.*, 2018). Post-translational modifications initiated by UBA1 and ATG7 are known to play a central role in autophagy, including the oncogenic and tumor-suppressive forms (Xiong, 2015; Gómez-Díaz and Ikeda, 2019); therefore, therapeutic targeting of these E1s may elicit autophagy-related anticancer effects in a context-dependent manner.

5. Inflammation, Immune Responses, and Tumor Microenvironment. Ubiquitylation plays an integral role in regulating inflammatory and immune signaling pathways, including those implicated in cancer pathogenesis (Bhoj and Chen, 2009; Hu and Sun, 2016; Fujita *et al.*, 2019). Of these pathways, the NF- κ B signaling pathway is particularly important. NF- κ B is a transcription factor that plays multifunctional roles in regulating survival, inflammation, and immune responses (Skaug *et al.*, 2009). The transcriptional activity of NF- κ B is kept in check by the inhibitor of NF- κ B ($\text{I}\kappa\text{B}\alpha$), which sequesters NF- κ B in an inactive state in the cytosol (Hoeller *et al.*, 2006). This inhibitory activity is released after ubiquitin-dependent proteasomal degradation of $\text{I}\kappa\text{B}\alpha$, resulting in the nuclear translocation and transcriptional activation of NF- κ B (DiDonato *et al.*, 2012). Therefore, UBA1 inhibition is anticipated to stabilize $\text{I}\kappa\text{B}\alpha$ and inhibit NF- κ B, leading to useful anticancer effects, particularly in malignancies with aberrant NF- κ B activity such as acute myeloid leukemia (AML) (Birkenkamp *et al.*, 2004; Bosman *et al.*, 2014). Of note, the NF- κ B signaling pathway is also regulated by nondegradative ubiquitylation and SUMOylation (Hoeller *et al.*, 2006; Skaug *et al.*, 2009). There is an emerging role of SUMOylation in modulating the innate and adaptive immune responses with potential implications in cancer therapy (see section IV. C. *Small Ubiquitin-Like Modifier-Activating Enzyme*). In addition to immune cells, other components of the tumor microenvironment, including cancer-associated fibroblasts and endothelial cells, are susceptible to regulation by UBL modifications including neddylation (Zhou *et al.*, 2019a).

6. Angiogenesis and Metastasis. E1-initiated post-translational modifications have been reported to regulate angiogenesis via direct and indirect mechanisms (Rahimi, 2012; Rabellino et al., 2020). In this respect, vascular endothelial growth factor receptor signaling is known to be regulated by ubiquitylation and SUMOylation (Simons et al., 2016; Zhou et al., 2018). In the context of cancer, ubiquitylation may indirectly promote tumor angiogenesis by enhancing p53 degradation and thus stabilizing hypoxia-inducible factor (HIF)-1 (Ravi et al., 2000). Similarly, ubiquitylation has been reported to regulate metastasis by multiple molecular mechanisms, such as the induction of Von Hippel-Lindau (VHL) degradation and thus HIF-1 stabilization (Jung et al., 2006; Gallo et al., 2017; Chen et al., 2019). Genetic and pharmacologic inhibition of ubiquitylation and neddylation have been reported to suppress tumor angiogenesis and metastasis (Tan *et al.*, 2014; Yao *et al.*, 2014; Jin *et al.*, 2018).

7. Cellular Stress Phenotypes. Cancer cells are characterized by several stress phenotypes, including proteotoxic, oxidative, metabolic, replicative, mitotic, and DNA damage stress (Luo et al., 2009). Such stresses are supported by normal cellular machineries, including the UBL conjugation systems, but to a greater extent in tumor versus normal cells, creating a myriad of therapeutic opportunities through stress overload and stress sensitization (Luo et al., 2009; Dobbstein and Moll, 2014). Therapeutic inhibition of UBA1 and NAE is known to induce unfolded protein response (UPR) and proteotoxic stress, particularly in hematologic malignancies (Soucy et al., 2009a; Hyer et al., 2018). Ubiquitylation and SUMOylation have been reported to respond to and modulate cellular production of reactive oxygen species (ROS) (Bossis and Melchior, 2006; Swords et al., 2010; Shang and Taylor, 2011; Graves et al., 2020; Ma et al., 2020). Thus, inhibition of UBA1 and NAE has been associated with the induction of oxidative stress (Nawrocki et al., 2013; Zhuang et al., 2019). Similar regulatory roles have been identified with cellular metabolism (Agbor and Taylor, 2008; Flick and Kaiser, 2012; Lavie et al., 2018; Thapa et al., 2020), replication, mitosis and DNA damage, as highlighted above. Dependent on the context, therapeutic targeting of E1s is anticipated to exacerbate one or more of these stresses, with potentially detrimental effects on cancer cell survival.

F. Targeting E1 Versus E3 Enzymes

E1 enzymes initiate the UBL conjugation cascade and thus lie upstream of the E3 enzymes that are responsible for substrate specificity (Schulman and Harper, 2009). Although targeting certain E3 enzymes is anticipated to elicit selective antitumor effects as a result of modulating one or a few substrates (Bedford et al., 2011; Huang and Dixit, 2016), targeting E1 enzymes offers several advantages over E3 enzymes. From a drug

discovery perspective, E1 enzymes possess well defined pockets that are readily druggable—as opposed to most E3 enzymes, which have multiple subunits and lack active pockets—and many E3s are not functionally annotated (Huang and Dixit, 2016). In addition, E3-oriented drug discovery endeavors have been mainly focused on targeting protein-protein interactions, which is a more challenging approach compared with targeting catalytic/active sites; therefore, the field witnessed slower progress until recently, as novel targeted protein degradation strategies that exploit ligands of E3 ligases to form heterobifunctional degraders have been developed (Cohen and Tcherpakov, 2010; Schapira et al., 2019; Barghout, 2020).

Although several ubiquitin E3 ligases are deregulated and thus implicated in oncogenic transformation and tumor progression, their selective therapeutic targeting may not be highly effective because of the functional redundancy among ubiquitin E3 ligases and the activation of compensatory signaling pathways (Nakayama and Nakayama, 2006; Senft et al., 2018; Wu et al., 2020). In contrast, targeting UBA1 that initiates the ubiquitylation cascade and regulates many ubiquitin-dependent signaling pathways may be more effective and less likely to suffer from the development of adaptive resistance, and this approach is useful in several contexts, particularly in malignancies highly dependent on ubiquitylation and where a sufficient therapeutic window can be attained (Dobbstein and Moll, 2014). This has been the case with proteasome inhibitors that, despite the lack of selectivity and perceived toxicity, proved to be more therapeutically useful compared with investigational E3 ligase inhibitors in several malignancies (Nakayama and Nakayama, 2006).

III. Tools and Assays for Exploring E1 Biology and Drug Discovery

E1 enzymes possess multiple domains that enable them to perform their multistep catalytic activity as UBL-activating enzymes (Schulman and Harper, 2009; Cappadocia and Lima, 2018). This catalytic activity comprises adenylation, thioester formation, and trans-thiolation and, in concert with downstream E2 and E3 enzymes, leads to the conjugation of specific UBL modifiers to selected substrates. Numerous assays and tools have been developed to interrogate the biology of E1 enzymes at different catalytic steps and to assist in the discovery and characterization of drugs and chemical probes that modulate the activity of these enzymes. In this section, we briefly highlight a number of these tools and their utility in E1 biology and the discovery of E1 inhibitors.

A. ATP:PPi and ATP:AMP Exchange Assays

E1 enzymes resemble aminoacyl-transfer RNA (tRNA) synthetases and other adenylate-forming enzymes in

their catalytic mechanisms (Dusha and Denes, 1977; Wilson and Aldrich, 2010). In this respect, they activate UBLs by catalyzing a three-step biochemical mechanism that involves the utilization of ATP and release of PP_i in the first and second adenylation steps and release of AMP in the thioester formation step (Fig. 3B).

In ATP:PP_i exchange assays, the catalytic reaction is assembled under cell-free conditions using a recombinant E1 enzyme, recombinant UBL, ATP, and radiolabeled PP_i. The incorporation of radiolabeled PP_i into ATP and formation of radiolabeled ATP, representing the reverse adenylation reaction, are quantitatively monitored using an appropriate detection method (Haas and Rose, 1982; Bruzzese et al., 2009). This assay can be used to assess the UBL specificity of E1 enzymes and to quantify their adenylate-forming activity by calculating several kinetic parameters, such as rate and equilibrium constants for the ternary E1-AMP~UBL complex formation and the turnover number for the enzyme—the number of E1-catalyzed reactions per unit time (Alontaga et al., 2012). On the other hand, ATP:AMP exchange assays are used to obtain information on both the adenylation and thioester formation steps of E1-catalyzed UBL activation (Alontaga et al., 2012). In these assays, the catalytic reaction is also assembled under cell-free conditions using recombinant E1 enzyme, recombinant UBL, ATP, PP_i, and radiolabeled AMP (Ciechanover et al., 1981; Wang et al., 2009). The generation of radiolabeled ATP from radiolabeled AMP is then monitored and quantified after the separation of both species by thin-layer chromatography. ATP:PP_i and ATP:AMP exchange assays have been widely used to characterize the catalytic activity of different E1 enzymes (Ciechanover et al., 1981, 1982; Haas and Rose, 1982; Bohnsack and Haas, 2003; Bruzzese et al., 2009; Wang et al., 2009). They have also been used in the context of biochemical characterization of novel E1 inhibitors to gain mechanistic insights into their mode of action and mechanisms of resistance (Brownell et al., 2010; Chen et al., 2011; Milhollen et al., 2012; Ungermannova et al., 2012a; Xu et al., 2014; Hyer et al., 2018).

B. E1~ Ubiquitin-Like Protein Thioester Formation and Transthioylation Assays

E1~UBL thioester formation assays determine the formation of E1~UBL thioesters under cell-free or cell-based conditions using detection methods such as immunoblotting with antibodies against either the E1 or UBL. In cell-free assays, the thioester formation reaction is assembled using recombinant E1, recombinant UBL, and ATP (Wang et al., 2007; Olsen et al., 2010). Since the thioester bond is sensitive to reducing agents such as dithiothreitol, the detection of E1~UBL thioesters requires nonreducing conditions (Alontaga et al., 2012). On the other hand, transthioylation assays determine the transfer of UBL from a certain E1 to

its cognate E2 enzyme to form an E2~UBL thioester. In addition to the reaction components used in the E1~UBL thioester formation assays, a recombinant E2 is also included to assess transthioylation. In addition, these assays are conducted in two steps and can provide quantitative information on both UBL activation and transfer (Alontaga et al., 2012). Recently, high-throughput luminescence-based assays have been developed to evaluate thioester formation with UBA1 and its E2, radiation sensitivity protein 6 (RAD6), in a quantitative manner (Fenteany et al., 2020). E1~UBL thioester formation and transthioylation assays have been used to identify and characterize the activity of different E1 inhibitors in various platforms including large-scale drug discovery screens (Yang et al., 2007; Brownell et al., 2010; Hyer et al., 2018; Lv et al., 2018b).

C. Substrate Ubiquitin-Like Protein Conjugation Assays

Substrate UBL conjugation assays determine the conjugation of specific UBLs to their target substrates and are conducted under cell-free or cell-based conditions. For example, candidate proteins such as histones H2A/H2B and PCNA are known to be common targets for monoubiquitylation, cullin-1 for neddylation, and Ran GTPase-activating protein 1 (RanGap1) for SUMOylation (Gareau and Lima, 2010; Enchev et al., 2015; Uckelmann and Sixma, 2017; Fenteany et al., 2020). In addition, the global changes in UBL conjugation to cellular proteins can be assessed by immunoblotting using antibodies against the UBL modifier or by large-scale proteomic approaches (Visconte et al., 2016). In cell-free assays, the whole UBL conjugation system including a recombinant E3 ligase needs to be assembled. However, this is not an absolute requirement, as for instance, RanGAP1 can be SUMOylated in the absence of an E3 ligase (Alontaga et al., 2012). Substrate UBL conjugation assays are not specific for E1 enzymes, as they evaluate the combined activity of E1, E2, and E3 enzymes. In addition to their use in drug discovery screens for E1 inhibitors, these assays are routinely used in evaluating the downstream biologic effects of potential E1 inhibitors on UBL conjugation to cellular substrates (Hyer et al., 2018; Barghout et al., 2019; Li et al., 2019).

D. Detection of Ubiquitin-Like Protein–Drug Adduct Formation

Adenosine sulfamates and related E1 inhibitors act by forming a covalent adduct with their cognate UBLs, and these adducts serve as the inhibitory species (Ciavarri and Langston, 2017). To detect the formation of these adducts, specific anti-UBL-drug antibodies have been generated and used in immunoblot-based assays to evaluate the pharmacodynamic profile of these drugs (Soucy et al., 2009a; Hyer et al., 2018). Similarly, a radiolabeled form of the adenosine aminosulfonamide TAS4464 has been used

to detect the formation of these adducts by immunoblotting and autoradiography (Yoshimura et al., 2019). However, it remains important to adapt these assays to provide quantitative information on the rate of adduct formation, which in turn depends on E1 activity.

In the following sections, we discuss in more detail the different E1s, their roles in cancer, the inhibitors developed so far to target these enzymes, and how these assays have been used to discover and characterize these inhibitors. In our discussion, we will emphasize mechanism-based E1 inhibitors, particularly those advanced to clinical trials, and the common principles shared among these agents. We will start with the NAE for which the prototypical clinical E1 inhibitor has been developed.

IV. Canonical E1 Enzymes

A. Neural Precursor Cell–Expressed Developmentally Downregulated Protein 8–Activating Enzyme

Neddylation is the process of conjugating NEDD8 to cellular substrates (Enchev et al., 2015). NEDD8 is 59% identical to ubiquitin—the highest level of similarity observed among all UBLs, albeit with structural differences that are sufficient to mediate distinct functions (Watson et al., 2011b; Enchev et al., 2015). Neddylation cascade is initiated by NAE that catalyzes NEDD8 activation. NAE is a heterodimeric enzyme composed of a regulatory subunit, NAE1 (amyloid β precursor protein-binding protein 1), and a catalytic subunit, NAE2 (UBA3), with similar domain structure as other canonical E1s (Fig. 2) (Schulman and Harper, 2009). Although the AAD is located on NAE2, the CCD half-domains are located on both subunits (Schulman and Harper, 2009; Cappadocia and Lima, 2018). Interestingly, the ubiquitin-specific UBA1 enzyme can activate NEDD8 under stress conditions, such as heat shock and oxidative stress (Leidecker et al., 2012). Two neddylation E2 enzymes have been reported in metazoans, UBC12 and UBE2F, which function in concert with E3 ligases including ubiquitin ligases (Enchev et al., 2015). Neddylation targets many substrates, of which cullins are the best characterized and the most established class of neddylated proteins (Enchev et al., 2015). Cullins serve as scaffold proteins upon which the largest class of RING ubiquitin E3 ligases, known as cullin-RING ubiquitin ligases (CRLs), are assembled (Watson et al., 2011b). Cullin neddylation plays a pivotal role in the ubiquitylation of a subset of cellular proteins by inducing conformational changes that increase the activity of CRLs and enhance ubiquitin conjugation (Lydeard et al., 2013; Baek et al., 2020). Other noncullin substrates include signaling molecules such as p53, p73, E2F1, epidermal growth factor receptor (EGFR), tumor growth factor β receptor, NF- κ B essential modulator (NEMO), Von Hippel–Lindau (VHL) tumor suppressor, and HIFs, which play diverse

roles in normal and cancer cell biology (Watson et al., 2011b; Enchev et al., 2015). Therefore, neddylation regulates numerous pathways, including proteasomal degradation, cell cycle progression, receptor tyrosine kinase signaling, apoptosis, DNA damage response, inflammatory/immune responses, oxidative stress, hypoxia, and nucleolar stress signaling (Soucy et al., 2009b; Mathewson et al., 2013, 2016; Brown et al., 2015; Enchev et al., 2015; Balachandran et al., 2016).

1. *Role of Neural Precursor Cell–Expressed Developmentally Downregulated Protein 8–Activating Enzyme in Cancer.* As neddylation plays a salient role in numerous signaling pathways, dysregulation of NAE and/or downstream components of neddylation cascade is anticipated to contribute to the development and progression of several malignancies (Soucy et al., 2010; Watson et al., 2011b). In most cases, expression data from published studies or publicly available datasets indicate a negative correlation between the expression of either NAE1 or NAE2 or both and clinical outcomes. For instance, NAE1/2 are upregulated in hepatocellular carcinoma and are associated with an aggressive phenotype and poor overall and relapse-free survival in patients (Barbier-Torres et al., 2015; Uhlen et al., 2017; Yu et al., 2018a). Similar data has been reported in lung cancer (Li et al., 2014b), glioblastoma (Hua et al., 2015), squamous cell carcinoma (Li et al., 2014b), intrahepatic cholangiocarcinoma (Gao et al., 2014), MM (Huang et al., 2015), esophageal squamous cell carcinoma (Chen et al., 2016a), uveal melanoma (Jin et al., 2018), pancreatic cancer (Misra et al., 2017; Li et al., 2018), clear cell renal cell carcinoma (Tong et al., 2017), and chronic myeloid leukemia (Liu et al., 2018). In contrast, patients with thyroid cancer show a favorable prognosis with high NAE1 expression (Uhlen et al., 2017). It appears, however, that no significant mutations are known to implicate NAE1/2 in cancer pathogenesis (Tate et al., 2019).

2. *Neural Precursor Cell–Expressed Developmentally Downregulated Protein 8–Activating Enzyme Inhibitors.* NAE is one of the first human E1 enzymes to have its crystal structure elucidated, and this provided valuable structural insights and a common framework for mechanisms of activation and transfer of NEDD8 as well as other UBLs (Walden et al., 2003a,b; Huang et al., 2004a, 2007). This structural information contributed, in part, to guiding drug discovery endeavors for E1 inhibitors, and it is not surprising that the first reported mechanism-based E1 inhibitor is targeted against NAE (Soucy et al., 2009a; Brownell et al., 2010). The hyperactive neddylation pathway or high reliance on one or more of its components in cancer is a therapeutic avenue that has garnered attention in recent years, particularly after the development of selective NAE inhibitors (Ying et al., 2018). Compounds targeting NAE comprise the adenosine sulfamate pevonedistat, the related adenosine aminosulfonamide

Fig. 7. Chemical structures of NAE inhibitors. NAE inhibitors belong to diverse classes, including adenosine sulfamates and related analogs (e.g., pevonedistat and TAS4464), natural compounds (e.g., 6,6''-Biapigenin), semisynthetic compounds (piperacillin), NEDD8 adenylyte analogs (e.g., adenosyl-phospho-NEDD8), NEDD8-mimicking peptides (e.g., pND20), and other inhibitors.

TAS4464 — both are selective clinical candidates — and several experimental inhibitors including NEDD8 adenylyte analogs and NEDD8-mimicking peptides (APN, pND20, and pND22), natural and semisynthetic NAE inhibitors (6,6''-biapigenin, flavokawain A, and piperacillin), as well as other inhibitors with diverse chemical structures [M22, LZ3, ZM223, and rhodium (III)-based complexes]. So far, pevonedistat is the most extensively studied E1 inhibitor, with a large number of preclinical and clinical studies in numerous malignancies. Apart from pevonedistat and TAS4464, most known NAE inhibitors have been discovered by *in silico* approaches and possess less favorable potency and selectivity profiles with activity in the micromolar range. Nonetheless, they can potentially serve as useful chemotypes to boost drug discovery endeavors and develop novel NAE inhibitors (Fig. 7; Table 2).

a. Pevonedistat. Pevonedistat (MLN4924/TAK-924) is the prototypical adenosine sulfamate E1 inhibitor (Soucy et al., 2009a). A high-throughput screen (HTS) identified N6-benzyl adenosine as an NAE inhibitor,

and further optimization by iterative medicinal chemistry led to the development of pevonedistat, a first-in-class mechanism-based NAE inhibitor (Soucy et al., 2009a; Ciavarrri and Langston, 2017). Mechanism-based inhibition is a form of enzyme inhibition whereby an analog of the enzyme substrate is processed through its normal catalytic mechanism, producing a species that inhibits the enzyme (Copp, 2003). Pevonedistat is structurally related to AMP and thereby acts as an AMP mimetic (Brownell et al., 2010). It exerts its NAE inhibitory activity by a form of mechanism-based inhibition termed “substrate-assisted inhibition,” whereby it forms a covalent adduct with NEDD8 in a reaction catalyzed by NAE itself. This covalent adduct tightly binds to the nucleotide-binding site and inhibits NAE in an ATP-competitive manner (Brownell et al., 2010). Substrate-assisted inhibition serves as a common mechanism of action among E1 adenosine sulfamate inhibitors (Ciavarrri and Langston, 2017).

Although compounds of this structural class may have inhibitory effects on adenylyte-forming enzymes

TABLE 2
NAE inhibitors, their chemical structures, and pharmacologic properties

Inhibitor	Type	Malignancies	Class	EC ₅₀ /IC ₅₀ /Dose	Reference
Pevonedistat (MLN4924; TAK-924)	Clinical: 33 clinical trials up to phase 3	Many solid and hematologic malignancies	Adenosine sulfamate	Cell-free thioester formation: 0.0047 μ M Cell-based neddylation: <0.1 μ M Cytotoxicity: 0.05–1.03 μ M In vivo dose: 30–90 mg/kg once daily or twice daily	Soucy et al., 2009a
TAS4464 (HY-128586)	Clinical: phase 1/2 clinical trial (MM and non-Hodgkin lymphoma)	Many solid and hematologic malignancies	Adenosine aminosulfonamide	Cell-free thioester formation: 0.955 nM Cytotoxicity $\leq$ 10 nM In vivo dose: 6.3–100 mg/kg once weekly	Yoshimura et al., 2019
6,6''-Biapigenin	Experimental	Cell-free and cellular activity in Caco-2 cells	Semisynthetic flavonoid	Cell-free UBC12~NEDD8 thioester formation: 20 μ M Cell-based UBC12~NEDD8 thioester formation: 5 μ M	Leung et al., 2011
Flavokawain A	Experimental	Prostate, bladder, and urothelial carcinoma	Chalcone flavonoid	Cell-free UBC12~NEDD8 thioester formation: 5 μ M	Zi and Simoneau, 2005; Liu et al., 2013; Li et al., 2015
M22	Experimental	Several cancer cell lines and a mouse model of gastric carcinoma	Piperidin-4-amine	Cytotoxicity: 8.98–26.6 μ M Dose in mice: 60 mg/kg, i.p., once daily	Lu et al., 2016
Piperacillin	Experimental	Cell-free and cellular activity in Caco-2 cells	Semisynthetic β -lactam antibiotic	Cell-free UBC12~NEDD8 thioester formation: 1 μ M	Zhong et al., 2014
Rhodium (III)-based complex	Experimental	Cell-free and cellular activity in Caco-2 cells	Cyclometallated rhodium(III) complex	Cell-free UBC12~NEDD8 thioester formation: 1.5 μ M Cytotoxicity in Caco-2 cells: 0.3 μ M	Zhong et al., 2012b
LZ3	Experimental	Cell-free and cellular activity in Caco-2, MCF7, and Bel-7402 cells	Sulfonamide derivative	Cell-free UBC12~NEDD8 thioester formation: 1.06 μ M Cytotoxicity: 12.3–29.5 μ M	Zhang et al., 2014
ZM223	Experimental	Cell-free and cellular activity in colon cancer and osteosarcoma cells	Benzothiazole derivative	Cytotoxicity: 0.1–1.22 μ M	Ma et al., 2017
Dipeptide-conjugated deoxy-vasicinone	Experimental	Cell-free and cellular activity in Caco-2 cells	Alkaloid derivative	Cell-free UBC12~NEDD8 thioester formation: 0.8 μ M Cell-based UBC12~NEDD8 thioester formation: 6 μ M	Zhong et al., 2012a
Adenosyl-phospho-NEDD8 (APN)	Experimental	Cell-free activity	NEDD8 adenylate	Cell-free NAE-UBC12 transthiolation: 0.05 μ M	Brownell et al., 2010
pND20 (VI LTFGG)	Experimental	Cell-free activity	Heptameric peptides	Cell-free NAE~NEDD8 thioester formation: 133 μ M	Zhao et al., 2013b
pND22 (VRLMFGG)	Experimental	Cell-free activity	Heptameric peptides	Cell-free NAE~NEDD8 thioester formation: 215 μ M	Zhao et al., 2013b
pN1 (VWLSYGG)	Experimental	Cell-free activity	Heptameric peptides	NA	Zhao et al., 2013a
pN7 (VMLFYGG)	Experimental	Cell-free activity	Heptameric peptides	NA	Zhao et al., 2013a
pN26 (VLALRGG)	Experimental	Cell-free activity	Heptameric peptides	NA	Zhao et al., 2013a

NA, not available.

TABLE 3
Preclinical and clinical studies of pevonedistat

Malignancy	Key Effects	Reference
Acute lymphoblastic leukemia	Preclinical efficacy in T-ALL models via induction of nucleolar stress signaling Preclinical efficacy in ALL models via ER stress and synergistic effects with dexamethasone, doxorubicin, and cytarabine Inhibition of the MEK/ERK pathway sensitizes ALL to pevonedistat in vitro and in vivo	Han et al., 2016 Ledere et al., 2016 Zheng et al., 2017a Swords et al., 2010
Acute myeloid leukemia	Preclinical efficacy via NF- κ B pathway inhibition, DNA damage, and ROS generation Sensitization of AML cell lines to retinoic acid differentiation therapy Preclinical efficacy in <i>FLT3-JTD</i> AML models via NF- κ B inhibition and downregulation of <i>miR-155</i> Antitumor effects in AML cells via Noxa upregulation as well as synergistic effects with Bcl-2 inhibitors Synergistic antileukemic effects with cytarabine in AML cells and mouse models via disruption of nucleotide metabolism Phase 1 clinical trial showing tolerability and modest clinical activity in patients with AML and MDS Synergistic effects with the TAP antagonist T-3256336 in cell lines and in vivo Synergistic combination with azacytidine in preclinical models of AML via antagonizing RRM2 Synergistic combination with belinostat in preclinical models of AML/MDS by reciprocal inhibition of specific DNA repair pathways Synergistic combination with the LSD1 inhibitor T-3775440 in preclinical models of AML via trans-differentiation and DNA re-replication Expanded safety analysis of pevonedistat in patients with AML/MDS with serious toxicity (e.g., hepatotoxicity and sepsis syndromes with multiorgan failure) only observed at doses beyond 100 mg/m ² Phase 1b clinical trial of pevonedistat in combination with azacytidine showing tolerability and potential benefit	Tan et al., 2011 Khalife et al., 2015 Knorr et al., 2015 Nawrocki et al., 2015 Swords et al., 2015 Sumi et al., 2016 Visconte et al., 2016 Zhou et al., 2016 Ishikawa et al., 2017 Swords et al., 2017 Swords et al., 2018 Yang et al., 2012 Lin et al., 2015 Gao et al., 2014 Godbersen et al., 2014 Paiva et al., 2015 Paiva et al., 2017 Liu et al., 2018 Lin et al., 2010 Picco et al., 2016
Breast cancer	Radiosensitization of breast cancer cells via induction of p21 accumulation Preclinical efficacy in models of CC as well as potentiation of cisplatin cytotoxicity	
Cervical carcinoma	Preclinical efficacy in ICC models and additive effects with cisplatin	
Cholangiocarcinoma	Antitumor effects in primary CLL B cells cocultured with stromal cells via NF- κ B pathway inhibition	
Chronic lymphocytic leukemia	Antitumor effects in primary CLL B cells cocultured with stromal cells via CDT1 accumulation and sensitization of cells to alkylating agents	
Chronic myeloid leukemia	Sensitization of primary CLL cells cocultured with stromal cells to death receptor agonists by enhancing death receptor signaling	
Colorectal cancer	Preclinical efficacy in CML models associated with p27 accumulation and overcoming mutation- and LSC-driven imatinib resistance Antitumor effects via DNA re-replication in CRC cell lines Preclinical efficacy in models of poorly differentiated, clinically aggressive CRC with a transcriptional signature associated with pevonedistat sensitivity	
Esophageal cancer	Sensitization of CRC cells to oxaliplatin via induction of DNA damage and CHK2 phosphorylation	Zheng et al., 2017b
Ewing sarcoma	Preclinical efficacy in ESCC models via ATF4-CHOP-DR5 axis-mediated extrinsic apoptosis	Chen et al., 2016a
Gastric cancer	Preclinical efficacy in ES models via WEE1 accumulation and deregulation of S-phase proteins Antitumor effects in gastric cancer cells associated with protective p27 upregulation	Mackintosh et al., 2013 Zhang et al., 2015b
Glioblastoma	Antitumor effects in gastric cancer cells associated with DNA damage induction, senescence, and autophagy Preclinical efficacy in models of GBM associated with induction of apoptosis and senescence Pevonedistat promotes migration of GBM cells via induction of caveolin-1 phosphorylation Upregulation of PD-L1 expression and synergism with anti-PD-L1 therapy in GBM models Antitumor effects in combination with TRAIL via promoting c-FLIP degradation in HNSCC cell lines	Lan et al., 2016 Hua et al., 2015 Park et al., 2018 Zhou et al., 2019b Zhao et al., 2011
Head and neck cancer	Preclinical efficacy in HNSCC models and radiosensitization of HNSCC cells	Vanderdys et al., 2018
Liver cancer	Preclinical efficacy associated with induction of protective autophagy and apoptosis in HCC models Preclinical efficacy in <i>Pbb1</i> -KO mouse models of HCC by destabilizing LKB1 and Akt	Luo et al., 2012 Barbier-Torres et al., 2015
Lung cancer	Synergistic effects with autophagy inhibitors in cell lines and mouse models of HCC Antitumor effects in <i>Kras</i> ^{G12D} -driven lung cancer models via inhibition of NF- κ B and mTOR pathways Preclinical efficacy and synergistic antitumor effects with platinum in lung adenocarcinoma models	Chen et al., 2015 Li et al., 2014a Li et al., 2014b
Lymphoma	Synergistic effects with the PARP inhibitor olaparib in NSCLC cell lines via inhibiting BRCA1 recruitment to DNA damage sites Antitumor effects in paclitaxel-resistant human lung adenocarcinoma cells with no synergism with paclitaxel Preclinical efficacy in DLBCL via NF- κ B pathway inhibition and DNA re-replication Antitumor effects in MCL cells via stabilization of Noxa Antitumor effects in lymphoma cell lines associated with apoptosis and senescence Preclinical efficacy in MCL models and additive/synergistic antitumor effects with conventional therapies Phase 1 clinical trial showing tolerability and pharmacodynamic effects in relapsed/refractory lymphoma Sensitization of DLBCL cells to the death receptor agonists TRAIL and FasL by enhancing death receptor signaling	Guo et al., 2017 Xu et al., 2018b Milhollen et al., 2010 Dengler et al., 2014 Wang et al., 2015 Czuczman et al., 2016 Shah et al., 2016 Paiva et al., 2017

(continued)

TABLE 3—Continued

Malignancy	Key Effects	Reference
Melanoma	A genome-wide siRNA screen identified the p21-dependent intra-S-phase checkpoint as a key mediator of pevonedistat cytotoxicity in melanoma cells independent of CDT1 stabilization	Blank et al., 2013
	Predelinear efficacy in melanoma models via DNA re-replication and senescence and synergistic effects with the BRAF kinase inhibitor vemurafenib	Benamar et al., 2016
Mesothelioma	Phase 1 clinical trial showing tolerability and durable stable disease in a subset of patients	Bhatia et al., 2016
	Predelinear efficacy and antimetastatic effects in UM models via antiangiogenic and anti-OSC effects	Jin et al., 2018
Myeloma	Predelinear efficacy in models of melanoma and gene expression differences between sensitive and resistant cell lines	Wong et al., 2017
	Predelinear efficacy in NF2-mutant mesothelioma models and synergistic effects with mTOR inhibitors	Cooper et al., 2017
Nasopharyngeal carcinoma	Predelinear efficacy and additive effects with conventional drugs in MM models	McMillin et al., 2012
	Antitumor effects in MM cells via upregulation of REDD1 and subsequent inhibition of AKT and mTOR signaling	Gu et al., 2014
Osteosarcoma	Antitumor effects in CKS1B-overexpressing MM cell lines via p21 stabilization	Huang et al., 2015
	Phase 1 clinical trial showing tolerability and pharmacodynamic effects in relapsed/refractory MM	Shah et al., 2016
Ovarian cancer	Synergistic cytotoxicity with pomalidomide using sequential treatment schedule in MM cells	Liu et al., 2019
	Predelinear efficacy in NPC models associated with stabilization of c-Jun and synergistic effects with radio- and chemotherapy	Xie et al., 2017
Pancreatic cancer	Predelinear efficacy in OS models associated with induction of senescence and apoptosis	Zhang et al., 2016
	Genetic and chemical inhibitors of Mcl-1 sensitize OS cells to pevonedistat	Zhang et al., 2017
Pediatric cancers	Synergistic effects with platinum in ovarian cancer cells	Jazaeri et al., 2013
	Synergistic effects with cisplatin in ovarian cancer cells and mouse xenograft models	Nawrocki et al., 2013
Prostate cancer	Antitumor effects alone and in combination with conventional therapies in EOC cells	Pan et al., 2013
	Synergistic effects with the IAP antagonist T-3256336 in an ovarian cancer cell line	Sumi et al., 2016
Renal cell carcinoma	Antagonistic effects with paclitaxel in ovarian cancer cells	Hong et al., 2019
	Radiosensitization of preclinical models of pancreatic cancer	Wei et al., 2012
Solid cancers	Antiangiogenic and antimetastatic effects in pancreatic cancer models	Yao et al., 2014
	Sensitization of preclinical models of PDAC to gemcitabine via accumulation of Noxa and ERBIN	Misra et al., 2017
Urothelial carcinoma	Potentiating the antitumor effects of pevonedistat by the Chk1 inhibitor SCH 90076 in preclinical models of pancreatic cancer	Li et al., 2018
	Predelinear efficacy in a subset of pediatric tumors tested in the Pediatric Preclinical Testing Program (PPTP)	Smith et al., 2012
Urothelial carcinoma	Antitumor effects in prostate cancer cells with apoptosis observed only at high concentrations	Rulina et al., 2016
	Radiosensitization of hormone-resistant prostate cancer cells via accumulation of WEE1, p21, and p27	Wang et al., 2016
Urothelial carcinoma	Pevonedistat promotes migration of prostate cancer cells via induction of caveolin-1 phosphorylation	Park et al., 2018
	Antitumor effects in RCC cell lines via induction of DNA damage and suppression of EMT	Tong et al., 2017
Urothelial carcinoma	Antitumor effects in CRC cell lines via upregulation of Noxa	Wang et al., 2017
	Predelinear efficacy in RCC models associated with the induction of cell cycle arrest, senescence, and apoptosis	Xu et al., 2018a
Urothelial carcinoma	Chemical/biochemical characterization and preclinical evaluation in models of CRC and lung cancer	Soucy et al., 2009a
	Antitumor effects via induction of senescence in CRC, lung, and GBM cell lines	Jia et al., 2011
Urothelial carcinoma	Synergistic effects with DNA interstrand cross-linking agents in CRC and breast cancer cell lines	Mihollen et al., 2011
	Inhibition of autophagy sensitizes solid cancer cell lines to pevonedistat	Kee et al., 2012
Urothelial carcinoma	Synergistic effects with mitomycin C, cisplatin, cytarabine, UV radiation, SN-38, and gemcitabine	Zhao et al., 2012
	Selective antitumor effects in solid CRC and melanoma cell lines as well as zebrafish embryos in a p53-based cyclotherapy to selectively target p53-mutant cells	Garcia et al., 2014
Urothelial carcinoma	Phase 1 clinical trial showing tolerability and pharmacodynamic effects of pevonedistat	Malhab et al., 2016
	Phase 1b clinical trial showing tolerability in combination with either docetaxel or carboplatin + paclitaxel with sustained clinical responses in the latter combination	Sarantopoulos et al., 2016
Urothelial carcinoma	Synergistic effects with cisplatin in urothelial cancer cell lines and mouse xenograft models	Lockhart et al., 2019
	Predelinear efficacy in UC models associated with ER stress induction, cell cycle arrest, and apoptosis	Ho et al., 2015
Urothelial carcinoma		Kuo et al., 2015

Akt, AKR thymoma protein; ALL, acute lymphoblastic leukemia; ATP4, activating transcription factor 4; Bcl-2, B-cell lymphoma 2; BRCA1, breast cancer type 1 susceptibility protein; CC, cervical carcinoma; CORCC, clear cell renal cell carcinoma; c-FLIP, cellular FADD-like IL-18-converting enzyme-inhibitory protein; CHK, checkpoint kinase; CHOP, C/EBP-homologous protein; CKS1B, cyclin-dependent kinases regulatory subunit 1; CLL, chronic lymphocytic leukemia; CML, chronic myeloid leukemia; CRC, colorectal cancer; CSC, cancer stem cell; DLBCL, diffuse large B-cell lymphoma; DR5, death receptor 5; EMT, epithelial-to-mesenchymal transition; EOC, epithelial ovarian cancer; ERBIN, ES, Ewing sarcoma; ESCC, esophageal squamous cell carcinoma; FasL, tumor necrosis factor receptor superfamily member 6 ligand; FLT3-ITD, Fms-like tyrosine kinase 3-internal tandem duplication; GBM, glioblastoma; HCC, hepatocellular carcinoma; HNSCC, head and neck squamous cell carcinoma; IAP, inhibitor of apoptosis protein; ICC, intrahepatic cholangiocarcinoma; KO, knockout; KRAS, Kirsten rat sarcoma viral oncogene homolog; LSC, leukemic stem cell; LSD1, lysine-specific demethylase 1A; MCL, mantle cell lymphoma; MEK, MAPK/ERK kinase; miR-155, microRNA-155; NF2, neurofibromatosis type 2; NPC, nasopharyngeal carcinoma; NSCLC, non-small-cell lung cancer; OS, osteosarcoma; PARP, poly (ADP-ribose) polymerase; PDAC, pancreatic ductal adenocarcinoma; PD-L1, programmed death-ligand 1; Phb1, prohibitin 1; RCC, renal cell carcinoma; REDD1, regulated in development and DNA damage responses 1; RRM2, ribonucleoside-diphosphate reductase; siRNA, small interfering RNA; T-ALL, T-cell acute lymphoblastic leukemia; UC, urothelial carcinoma; UM, uveal melanoma.

such as tRNA synthetases (Brown et al., 1999; Forrest et al., 2000; Lux et al., 2019), or potentially other ATP-dependent enzymes such as kinases, the action of pevonedistat is highly selective for NAE even compared with other E1 enzymes (Soucy et al., 2009a). In this context, it selectively inhibits neddylation with little or no effect on the conjugation of related UBLs with IC₅₀ values in the nanomolar range (Table 2).

Given the role of neddylation in regulating the activity of CRLs, pevonedistat induces a potent inhibition of these enzymes with a subsequent reduction in the turnover of a subset of cellular proteins (Soucy et al., 2009a,b). Prominent among these proteins is the DNA replication licensing factor CDT1, which is essential for prereplication complex assembly and plays a key role in DNA replication and mitosis (Milhollen et al., 2011). Stabilization of CDT1 in the S phase of the cell cycle is primarily responsible for the re-replication phenotype and mitotic defects observed after pevonedistat treatment (Soucy et al., 2009a; Lin et al., 2010; Milhollen et al., 2011). DNA re-replication is characterized by $\geq 4N$ DNA content as a result of unscheduled repeated rounds of DNA synthesis and leads to replication stress, DNA damage, and apoptosis (Petropoulos et al., 2019). Consistent with these cell cycle-specific effects, actively proliferating cells are more susceptible to pevonedistat cytotoxicity (Soucy et al., 2009a). TAK-243, the related UBA1 inhibitor, does not induce re-replication, possibly because of the stabilization of both CDT1 and its endogenous inhibitor geminin (Hyer et al., 2018).

With the narrower spectrum of proteins influenced by disrupting neddylation, it is conceivable that NAE inhibitors will exhibit a wider therapeutic window compared with proteasome and UBA1 inhibitors (Buac et al., 2013). Pevonedistat is well tolerated in vivo with broad antitumor activity in solid and hematologic malignancies (Table 3).

Apart from the induction of re-replication and DNA damage (Swords et al., 2010; Zhang et al., 2015b; Guo et al., 2017; Tong et al., 2017; Zheng et al., 2017b), a plethora of diverse mechanisms have been reported to mediate the antitumor effects of pevonedistat. These include the inhibition of the NF- κ B pathway (Milhollen et al., 2010; Swords et al., 2010; Godbersen et al., 2014; Li et al., 2014a; Khalife et al., 2015); stabilization and upregulation of proapoptotic proteins such as Noxa (Dengler et al., 2014; Knorr et al., 2015; Misra et al., 2017; Wang et al., 2017); generation of ROS (Swords et al., 2010), stabilization of cell cycle regulators such as p21, p27, and WEE1 (Yang et al., 2012; Blank et al., 2013; Mackintosh et al., 2013; Huang et al., 2015; Liu et al., 2018; Wang et al., 2016); inhibition of the mammalian/mechanistic target of rapamycin (mTOR) pathway (Gu et al., 2014; Li et al., 2014a); induction of endoplasmic reticulum (ER) stress (Kuo et al., 2015;

Leclerc et al., 2016); induction of immunomodulatory effects (Best et al., 2020); and activation of death receptor signaling (Chen et al., 2016a; Paiva et al., 2017). In addition to apoptosis, pevonedistat induces senescence in several malignancies (Jia et al., 2011; Wang et al., 2015; Benamar et al., 2016). Moreover, it exerts antiangiogenic and antimetastatic effects in preclinical models of pancreatic cancer (Yao et al., 2014) and uveal melanoma (Jin et al., 2018). While inhibiting migration in human urothelial (Kuo et al., 2015), clear cell renal (Tong et al., 2017) and gastric cancer cells (Lan et al., 2016), pevonedistat has been reported to display promigratory effects in glioblastoma and prostate cancer cells through induction of caveolin-1 phosphorylation (Park et al., 2018), suggesting control of migration by neddylation is context-dependent.

With the multiple effects exerted by pevonedistat, it is not surprising that additive or synergistic antitumor effects are observed with mechanistically diverse anti-neoplastic agents including conventional cytotoxic drugs (Kee et al., 2012; Jazaeri et al., 2013; Nawrocki et al., 2013; Garcia et al., 2014; Li et al., 2014b; Ho et al., 2015; Czuczman et al., 2016; Xu et al., 2018b), radiotherapy (Wei et al., 2012; Wang et al., 2016; Xie et al., 2017; Vanderdys et al., 2018), differentiation therapy (Tan et al., 2011), and targeted therapies such as Bcl-2 inhibitors (Knorr et al., 2015), epigenetic modulators (Visconte et al., 2016; Zhou et al., 2016), monoclonal antibodies (Czuczman et al., 2016), immunomodulatory drugs (Liu et al., 2019; Zhou et al., 2019b), poly (ADP-ribose) polymerase inhibitors (Guo et al., 2017), BRAF inhibitors (Benamar et al., 2016), and other investigational therapies (Chen et al., 2015; Sumi et al., 2016; Cooper et al., 2017; Ishikawa et al., 2017). In certain malignancies, the cytotoxicity of pevonedistat is ameliorated by the induction of cytoprotective autophagy (Luo et al., 2012), an effect that can be overcome by combination with autophagy inhibitors leading to synergistic antitumor effects (Zhao et al., 2012; Chen et al., 2015).

Since its discovery in 2009, pevonedistat has been and is currently evaluated in different phases of over 30 clinical trials alone and in combination with other drugs (Table 4). Approximately half of these are in AML and/or myelodysplastic syndromes (MDS).

Phase 1 clinical trials in AML/MDS has shown pevonedistat is generally well tolerated, with pyrexia as the most common adverse event and some dose-limiting toxicities, such as hepatotoxicity and sepsis syndromes with multiorgan failure, that are mostly observed at doses beyond 50 mg/m² (Swords et al., 2015, 2017). In AML, the rates of complete and partial responses ranged from 10% to 17%, with an overall response rate of 13% (Swords et al., 2015, 2017). When evaluated in combination with azacitidine, the overall response rate increased up to 83%, with a complete response/partial response rate of 80% in patients with

TABLE 4
Clinical trials of pevonedistat

Identifier	Title	Phase	Disease	Single/ Combination	Combined Drug(s)	Status	Reference
NCT01814826	Study of MLN4924 plus azacitidine in treatment-naïve patients with AML who are 60 y or older	1	AML	Combination	Azacitidine	C	Swords et al., 2018
NCT03772925	Pevonedistat and belinostat in treating patients with relapsed or refractory AML or MDS	1	AML/MDS	Combination	Belinostat	R	—
NCT01011530	Dose escalation study of MLN4924 in adults with melanoma	1	Melanoma	Single	—	C	Bhatia et al., 2016
NCT01862328	Dose escalation, multiarm study of MLN4924 plus docetaxel, gemcitabine, or combination of carboplatin and paclitaxel in patients with solid tumors	1	Solid tumors	Combination	Docetaxel; paclitaxel + carboplatin; gemcitabine	C	Lockhart et al., 2019
NCT03330106	A study to evaluate the effects of pevonedistat on the QTc interval in participants with advanced solid tumors	1	Advanced solid tumors	Safety (QTc interval)	Docetaxel; paclitaxel + carboplatin	ANR	—
NCT03770260	Ixazomib and pevonedistat in treating patients with MM that has come back or does not respond to treatment	1	Plasma cell myeloma	Combination	Ixazomib	R	—
NCT00677170	Study of MLN4924 in adult patients with nonhematologic malignancies	1	Advanced solid tumors	Single	—	C	Swords et al., 2015; Sarantopoulos et al., 2016
NCT02122770	Effects of fluconazole and itraconazole CYP3A-mediated inhibition on the pharmacokinetics, safety, and tolerability of MLN4924 in participants with advanced solid tumors	1	Advanced solid tumors	Combination (DDI study)	Fluconazole; itraconazole; docetaxel; paclitaxel + carboplatin	C	—
NCT03814005	A study of pevonedistat in combination with azacitidine in participants with higher-risk MDS, CMML, or relapsed/refractory AML with severe renal impairment or mild hepatic impairment	1	MDS CMML AML	Combination	Azacitidine	R	—
NCT03323034	Pevonedistat, irinotecan hydrochloride, and temozolomide in treating patients with recurrent or refractory solid tumors or lymphoma	1	Solid tumors and lymphoma	Combination	Irinotecan; temozolomide	R	—
NCT03486314	A study to evaluate the effects of rifampin on PK of pevonedistat in participants with advanced solid tumors	1	Advanced solid tumors	Combination (DDI study)	Rifampin; docetaxel; paclitaxel + carboplatin	ANR	—
NCT03459859	Pevonedistat and low-dose cytarabine in adult patients with AML and MDS	1	AML and MDS	Combination	Cytarabine	R	—
NCT00722488	Study of MLN4924, a novel inhibitor of Nedd8-activating enzyme, in adult patients with lymphoma or MM	1	Lymphoma and MM	Single	—	C	Shah et al., 2016
NCT00911066	MLN4924 for the treatment of AML, MDS, and ALL	1	AML, MDS, and ALL	Single	—	C	Swords et al., 2015
NCT03745352	Pevonedistat with azacitidine vs. azacitidine alone in treating patients with relapsed or refractory AML	2	AML	Combination	Azacitidine	R	—
NCT03057366	A Study of [¹⁴ C]pevonedistat in participants with advanced solid tumors	1	Advanced solid tumors	Combination (excretion study)	Docetaxel; paclitaxel + carboplatin	C	Lockhart et al., 2019
NCT03386214	Pevonedistat in combination with ruxolitinib for treatment of patients with myelofibrosis	1	Myelofibrosis	Combination	Ruxolitinib	R	—

(continued)

TABLE 4—Continued

Identifier	Tile	Phase	Disease	Single/ Combination	Combined Drug(s)	Status	Reference
NCT03813147	Pevonedistat, azacitidine, fludarabine phosphate, and cytarabine in treating patients with relapsed or refractory AML or relapsed high-risk MDS	1	AML and MDS	Combination	Azacitidine; cytarabine; fludarabine; methotrexate; hydrocortisone	R	—
NCT01415765	MLN4924 compared with MLN4924 plus chemotherapy for LBCL	1/2	Diffuse LBCL	Combination (+EPOCH-R)	Etoposide; prednisone; vincristine; cyclophosphamide; doxorubicin; rituximab; filgrastim	W	—
NCT03479268	Pevonedistat and ibrutinib in treating participants with relapsed or refractory CLL or NHL	1	CLL and NHL	Combination	Ibrutinib	ANR	—
NCT03009240	Pevonedistat and decitabine in treating patients with high-risk AML	1	AML	Combination	Decitabine	R	—
NCT03330821	Pevonedistat, cytarabine, and idarubicin in treating patients with AML	1/2	AML	Combination	Pevonedistat; cytarabine; idarubicin	R	—
NCT03709576	Pevonedistat and azacitidine as maintenance therapy after allogeneic stem cell transplantation for nonremission AML	2	AML	Combination	Azacitidine	T	—
NCT02782468	A study of pevonedistat in adult East Asian participants	1	AML/MDS	Combination	Azacitidine	ANR	—
NCT03268954 (PANTHER)	Pevonedistat plus azacitidine vs. single-agent azacitidine as first-line treatment of participants with higher-risk MDS, CMML, or low-blast AML	3	MDS, CMML, and AML	Combination	Azacitidine	R	—
NCT02610777	An efficacy and safety study of pevonedistat plus azacitidine vs. single-agent azacitidine in participants with HR-MDS, CMML and low-blast AML	2	MDS, CMML, and AML	Combination	Azacitidine	ANR	—
NCT03319537	Pevonedistat with VXLD chemotherapy for adolescent/young adults with relapsed/refractory ALL	1	ALL	Combination	Vincristine; dexamethasone; PEG-asparaginase; doxorubicin	R	—
NCT03228186	Trial of pevonedistat plus docetaxel in patients with previously treated advanced NSCLC	2	NSCLC	Combination	Docetaxel	R	—
NCT03238248	Pevonedistat and azacitidine in patients with MDS or MDS/MPN who fail primary therapy with DNA methyl transferase inhibitors	2	MDS and MDS/MPN	Combination	Azacitidine	R	—
NCT03013998	Study of biomarker-based treatment of AML	1/2	AML	Combination	Azacitidine	R	—
NCT03965689	Testing the combination of pevonedistat, carboplatin, and paclitaxel for NSCLC	2	NSCLC	Combination	Carboplatin; paclitaxel	R	—
NCT03862157	Azacitidine, venetoclax, and pevonedistat in treating patients with newly diagnosed AML	1/2	AML	Combination	Azacitidine; venetoclax	R	—

ALL, acute lymphoblastic leukemia; ANR, active, not recruiting; C, completed; CLL, chronic lymphocytic leukemia; CMML, chronic myelomonocytic leukemia; DDI, drug-drug interaction; EPOCH-R, etoposide, prednisone, vincristine, cyclophosphamide, doxorubicin, and rituximab; LBCL, large B-cell lymphoma; MPN, myeloproliferative neoplasms; N, not yet recruiting; NHL, non-Hodgkin lymphoma; NSCLC, non-small-cell lung cancer; PEG, polyethylene glycol; QTc, corrected QT; R, recruiting; T, terminated; VXLD, vincristine, dexamethasone, pegylated asparaginase, and doxorubicin; W, withdrawn.

TP53 mutations (Swords et al., 2018). These encouraging results with azacitidine prompted several phase 2 clinical trials, and a phase 3 clinical trial, PANTHER, is currently ongoing to evaluate pevonedistat-azacitidine combination versus azacitidine alone as a frontline therapy for specific forms of MDS and AML (clinicaltrials.gov

identifier: NCT03268954). In other hematologic (relapsed/refractory MM and lymphoma) and advanced solid malignancies, pevonedistat has also shown similar tolerability with potential therapeutic benefit in these populations (Bhatia et al., 2016; Sarantopoulos et al., 2016; Shah et al., 2016; Lockhart et al., 2019).

TABLE 5
Properties of ubiquitin-activating enzymes: UBA1 and UBA6 (McGrath et al., 1991; Groettrup et al., 2008; Kulkarni and Smith, 2008; Clague et al., 2015; Hyer et al., 2018)

Enzyme	UBA1	UBA6
Identity ^a	100%	40%
Molecular mass	118 kDa	118 kDa
UBL	Ubiquitin	Ubiquitin; FAT10
Contribution to cellular ubiquitylation	>99%	<1%
E2 enzymes ^c	Ubiquitin E2s except for USE1	Ubiquitin: Few E2s and USE1 is UBA6-specific FAT10: Unknown
Knockout phenotype ^d	Lethality in <i>Caenorhabditis elegans</i> and yeast	Preferential activity toward ubiquitin under basal conditions; cytokines (e.g., IFN-γ) increase activity toward FAT10 Embryonic lethality in mice
Expression ^e	Constitutively expressed in relatively large amounts in all tissues	Expressed in low amounts (compared with UBA1) in all tissues, with specifically higher expression in the testis
Ubiquitin charging in proliferating cells	Fully charged with ubiquitin	50% charged

^aPercent amino acid identity relative to UBA1.
^bFAT10 was reported to serve as a post-translational modification to direct proteins to proteasomal degradation.
^cNine E2s were reported to be charged equally by UBA1 and UBA6.
^dEmbryonic lethality of knockout phenotypes may indicate that UBA1 and UBA6 are not merely redundant versions of each other and that each of them is required for distinct essential biologic functions.
^eUBA1 is ranked among the top 2% of abundant proteins in HeLa cells (>3 × 10⁶ copies per cell), and the relative UBA1:UBA6 abundance ratio is approximately 10:1.

b. **TAS4464.** TAS4464 is an adenosine aminosulfonamide that has structural features similar to adenosine sulfamates but differs in having an aminosulfonamide group in lieu of the sulfamate moiety (Yoshimura et al., 2019). Still, it needs to form a covalent adduct with NEDD8 through this functional group, and this adduct in turn serves as the potent inhibitory species of NAE (Yoshimura et al., 2019). TAS4464 retains the ribose sugar moiety found in AMP, as is the case with several dual and pan E1 inhibitors (see below). Despite this similarity, it exhibits a highly selective activity toward NAE. Accordingly, it induces similar biologic effects on neddylation and downstream signaling, yet with up to 64-fold higher potency and more durable effects compared with pevonedistat. TAS4464 displays a broad-spectrum activity on cancer cell lines of different origin, particularly those of hematologic malignancies. It also displays a superior activity on primary patient samples as well as patient-derived xenografts, including treatment-resistant samples. TAS4464 has been advanced to a phase 1/2 clinical trial in MM and non-Hodgkin lymphoma (clinicaltrials.gov identifier: NCT02978235).

c. **Neural Precursor Cell–Expressed Developmentally Downregulated Protein 8 Adenylate Analogs and Neural Precursor Cell–Expressed Developmentally Downregulated Protein 8–Mimicking Peptides.** Adenosyl-phospho-NEDD8 (APN) has been synthesized as a chemical probe to assist in elucidating the mechanism of NAE inhibition by pevonedistat (Brownell et al., 2010). Similar to the ubiquitin adenylate analog APU (see below), APN acts as a nonhydrolyzable mimetic of NEDD8~AMP. It also resembles pevonedistat-NEDD8 adduct in tightly binding to NAE and inhibiting its activity in an ATP-competitive manner; however, NAE exhibits a slower recovery from pevonedistat-

NEDD8 adduct, which is consistent with tighter binding compared with APN (Brownell et al., 2010).

As the C-terminal sequence of NEDD8 plays a fundamental role in its recognition by NAE, phage display technology has been exploited to screen mutant versions of this sequence for variants with different kinetics and enzyme reactivities (Zhao et al., 2013b). In this respect, pND20 and pND22 have been identified as NEDD8-mimicking heptameric peptides that can still be recognized and activated by NAE, yet with a higher affinity to the enzyme compared with the wild-type NEDD8. Despite successful activation by NAE, these mutant variants cannot be subsequently used in the neddylation cascade, thereby serving as mechanism-based inhibitors that block NEDD8 activation and conjugation. NEDD8 is the closest UBL to ubiquitin in terms of sequence homology, and UBA1 can activate NEDD8 under certain conditions (Leidecker et al., 2012). Therefore, NEDD8-mimicking peptides have also been derived from mutant C-terminal sequences of ubiquitin, resulting in similar inhibitory properties against NAE (Zhao et al., 2013a).

d. **Natural and Semisynthetic Neural Precursor Cell–Expressed Developmentally Downregulated Protein 8–Activating Enzyme Inhibitors.** 6,6''-Biapigenin is a semisynthetic flavonoid derivative whose NAE inhibitory activity was identified through virtual screening of a 20,000-compound library using the crystal structure of the quaternary NAE1-UBA3-NEDD8-ATP complex (Leung et al., 2011). Based on molecular modeling studies, 6,6''-biapigenin is predicted to bind to NAE at a binding site distinct from pevonedistat, resulting in reversible inhibition. Another flavonoid with anti-NAE activity is flavokawain A (Li et al., 2015). Specifically, it belongs to the chalcone subclass of flavonoids and is isolated from the kava extract (Zi and Simoneau, 2005). Flavokawain

A inhibits NAE after binding to the enzyme at the nucleotide-binding pocket, resulting in the deneddylation of cullins and inducing proteasome-dependent degradation of S-phase kinase-associated protein 2 (Skp2). As a result, it shows antitumor activity against Rb-deficient cells that are dependent on Skp2 for survival, as well as antitumor and antimetastatic effects in vivo (Li et al., 2015).

Piperacillin, a Food and Drug Administration–approved semisynthetic β -lactam antibiotic, is also reported to competitively inhibit NAE at the nucleotide-binding pocket, suppressing neddylation of downstream targets and inducing accumulation of p27^{kip1} in cells (Zhong et al., 2014). Another virtual screen of a 90,000-compound library identified a dipeptide-conjugated alkaloid derivative with anti-NAE properties in cell-free and cellular enzymatic assays (Zhong et al., 2012a).

e. Other Natural and Semisynthetic Neural Precursor Cell–Expressed Developmentally Downregulated Protein 8–Activating Enzyme Inhibitors. M22 is a piperidin-4-amine–based NAE inhibitor identified through a structure-based virtual screening of 50,000 compounds (Lu et al., 2016). Molecular modeling studies predict M22 binds to the nucleotide-binding site in a similar conformation to that of ATP. As assessed by biochemical analyses, M22 reversibly inhibits NAE in an ATP-competitive manner, resulting in the stabilization of CRL substrates. These effects are associated with antiproliferative activity in cancer cells of different origin both in vitro and in vivo, as well as synergistic effects with bortezomib. Using zebrafish as an acute toxicity model, M22 is tolerated even at high micromolar concentrations (Lu et al., 2016).

To identify covalent NAE inhibitors, Zhang et al. (2014) leveraged structural information on NAE and pharmacologic data available on pevonedistat's mode of action. Specifically, they applied a ligand- and structure-based pharmacophore modeling approach combined with covalent docking and screened a focused library of free sulfamoyl-containing compounds for such irreversible inhibitors. These endeavors led to the identification of LZ3 as the top hit. LZ3 is a sulfonamide derivative that covalently binds to NAE, inhibiting its activity in cell-free and cell-based assays. It is unclear, however, whether it forms a covalent adduct with NEDD8 to exhibit such irreversible inhibition. On the other hand, target-based virtual screening approaches have been performed using structural data of the noncovalent interaction between pevonedistat and NAE to identify reversible NAE inhibitors. This led to the identification of a series of reversible nonsulfamate-based inhibitors, including ZM223, a benzothiazole with activity in the low micromolar range (Ma et al., 2017).

A rhodium (III)-based complex is reported to possess a similar inhibitory activity against NAE in cell-free assays and Caco-2 cells, with a low micromolar potency

partly attributed to its octahedral coordination geometry (Zhong et al., 2012b).

B. Ubiquitin-Like Modifier–Activating Enzyme 1

UBA1 is the major ubiquitin-activating enzyme that catalyzes the first step in ubiquitin conjugation cascade, and it is involved in the ubiquitylation of most cellular proteins (Hyer et al., 2018). It is an evolutionarily conserved protein and plays an indispensable role in regulating numerous cellular functions (Groen and Gillingwater, 2015). Structurally, UBA1 is a canonical monomeric E1 enzyme with several domains, including the AAD, IAD, UFD, and CCD (Lee and Schindelin, 2008; Schulman and Harper, 2009). Among the eight E1 enzymes identified so far, only UBA1 and UBA6 can activate ubiquitin, but the functions of UBA6-mediated ubiquitylation are not well characterized compared with those mediated by UBA1 (Chiu et al., 2007; Jin et al., 2007; Pelzer et al., 2007; Groettrup et al., 2008). In addition, several differences exist between UBA1 and UBA6 (Table 5).

Ubiquitin itself can be modified on one or more of its seven lysine residues, N terminus, and/or 11 phosphorylation sites by ubiquitin or other post-translational modifications, creating highly diverse ubiquitin signals with multiple topologies (Swatek and Komander, 2016). These distinct forms of ubiquitylation can be simply divided into mono- and polyubiquitylation, and this latter comprises K48- and K63-linked polyubiquitin chains as the most common forms of ubiquitylation (Swatek and Komander, 2016). The diverse ubiquitin-based modifications constitute a “ubiquitin code” that is written by the concerted action of ubiquitin E1–E2–E3 enzymes, erased by a set of DUBs, and read by cellular proteins that specifically recognize this code and translate it into a certain biologic response (Swatek and Komander, 2016). Ubiquitylation is important for protein homeostasis through proteasomal degradation and autophagy and is implicated as a post-translational modification in regulating and fine-tuning a myriad of nondegradative cellular processes, such as cell cycle progression, DNA replication and repair, protein translation, cell death, endocytosis, chromatin architecture and epigenetic regulation, cellular differentiation, inflammation, and immune signaling (Bhoj and Chen, 2009; Ulrich and Walden, 2010; Swatek and Komander, 2016; Dikic, 2017; Varshavsky, 2017; Rape, 2018).

Given its role in almost every aspect of cell biology, ubiquitylation is tightly regulated by multiple mechanisms, including other post-translational modifications and DUBs to specifically conjugate target proteins in a spatiotemporal manner (Grabbe et al., 2011). The ubiquitin system is “pyramidal” in design, as only two E1 enzymes transfer ubiquitin to over 30 E2s, which in turn interact with hundreds of E3 ligases to conjugate the respective protein substrate (Nalepa et al., 2006). The specificity of ubiquitin conjugation is mainly

TABLE 6
UBA1 inhibitors and their pharmacologic properties

Inhibitor	Type	Malignancies	Class	K _i /EC ₅₀ /IC ₅₀ /Dose	Reference
TAK-243 (MLN7243; AOB87172)	Clinical: phase 1	Solid and hematologic malignancies, AML, cutaneous SCC, BCL, and myeloma	Adenosine sulfamate	Cell-free UBCH10 thioester assay: 0.001 μ M Cytotoxicity: 0.006–1.31 μ M	Hyer et al., 2018; McHugh et al., 2018; Barghout et al., 2019; Best et al., 2019a; Zhuang et al., 2019
PYR-41	Experimental	Transformed cells, breast, MM, MCL, and CML cancer cells	Nitropyrazone	Cell-free thioester assay: 10 μ M Cellular thioester formation: 10–25 μ M	Yang et al., 2007; Kapuria et al., 2011
PYZD-4409	Experimental	Leukemia and MM cells in vitro and in vivo	Nitropyrazone	Cell-free thioester formation: 20 μ M Cytotoxicity: 3–20 μ M Dose in mice: 10 mg/kg IP every other day for 8 days	Xu et al., 2010
JS-K	Experimental	Transformed cells, prostate, and MM cancer cells	Nitropiperazine	MM cell lines (48 h): 0.3–1.2 μ M MM primary cells (72 h): 2–2.5 μ M Cellular thioester formation: 2 μ M Dose in mice: 4 μ M/kg 3 times weekly, i.v.	Kitagaki et al., 2009; Kiziltepe et al., 2007; Tan et al., 2017
Largazole	Experimental	Lung epithelial cell line, breast, colon, neuroblastoma, and osteosarcoma cells	Macrocyclic natural compound	Cell-free thioester formation: 29 μ M Cytotoxicity (cancer cells): 7.7–55 nM Cytotoxicity (nontransformed cells): 122–480 nM	Taori et al., 2008; Ungermannova et al., 2012b
APU	Experimental	Cell-free activity	Ubiquitin adenylate	K _i (ATP): 50 nM K _i (Ub): 35 nM K _i (Ub conjugation): 750 nM	Wilkinson et al., 1990
Panepophenanthrin	Experimental	Cell-free activity	Natural compound	Cell-free thioester formation: 40 μ M	Sekizawa et al., 2002
NSC624206	Experimental	Cell-free and cellular activity in lung and HCC cells	Disulfide	Cell-free thioester formation: 9–13 μ M	Ungermannova et al., 2012a
Himeic acid A ^a	Experimental	Cell-free activity	Natural compound	Cell-free thioester formation: 50 μ M	Tsukamoto et al., 2005
Hyr-tireticulins A and B	Experimental	Cell-free activity	Natural compounds	Cell-free thioester formation: 2.4 (A) and 35 μ M (B)	Yamanokuchi et al., 2012
Ub-AMSN	Experimental	Cell-free activity	Semisynthetic ubiquitin adenylate	NA	Lu et al., 2010; Olsen et al., 2010
Ub-AVSN	Experimental	Cell-free activity	Semisynthetic ubiquitin adenylate	NA	Lu et al., 2010; Olsen et al., 2010

BCL, B-cell lymphoma; HCC, hepatocellular carcinoma; NA, not available; SCC, squamous cell carcinoma.

^aThe reported concentration corresponds to the IC₆₅.

controlled downstream at the level of ubiquitin E3 ligases. Therefore, pharmacologic manipulation of the upstream enzyme UBA1 is anticipated to indiscriminately influence numerous biologic processes.

1. Role of Ubiquitin-Like Modifier-Activating Enzyme 1 in Cancer. Although many mutations and copy number variations (CNVs) have been reported to affect *UBA1*, there is no known evidence that these genetic aberrations are functionally important in the pathogenesis or progression of cancer (Tate et al., 2019; UniProt Consortium, 2019). Physiologically, UBA1 is among the most abundant proteins in the cell; however, variations in its expression levels may occur in several cancers (Clague et al., 2015). The analysis of patient survival data has demonstrated that UBA1 expression may

correlate with the prognosis of cancer. Although it is associated with poor prognosis in liver cancer, high expression of UBA1 is associated with a favorable prognosis in prostate cancer (Uhlen et al., 2017). On the other hand, AML cell lines and primary samples exhibited equal levels of UBA1 protein compared with normal hematopoietic cells while having a higher abundance of ubiquitylated proteins (Xu et al., 2010). This increased ubiquitylation is likely attributable to the more active utilization of UBA1 in AML versus normal cells, constituting a potential therapeutic vulnerability in AML and other malignancies (Yang et al., 2007; Hyer et al., 2018). Ubiquitin conjugation is a multienzymatic process, and cancer-associated dysregulation of ubiquitylation may occur at the level of downstream E2 or E3

Fig. 8. Chemical structures of UBA1 inhibitors. Classes of UBA1 inhibitors include adenosine sulfamates (e.g., TAK-243), nitro-based compounds (e.g., PYR-41, PYZD-4409, and JS-K), natural compounds (e.g., panepophenanthrin, himeic acid A, hyrtioreticulins A/B, and largazole), and ubiquitin adenylate analogs (e.g., adenosyl-phospho-ubiquitinol, Ub-AMSN, and Ub-AVSN), and disulfides (e.g., NSC624206).

enzymes with no changes in the ubiquitin E1 enzymes. The role of these other components of the ubiquitin system in cancer has been extensively reviewed (Nakayama and Nakayama, 2006; Fulda et al., 2012; Alpi et al., 2016; Senft et al., 2018).

2. Ubiquitin-Like Modifier-Activating Enzyme 1 Inhibitors. The crystal structure of human UBA1 has not been elucidated until recently (Lv et al., 2018a), and most of the structural information on UBA1 and ubiquitin activation was derived from published structures of yeast, humanized yeast, mouse Uba1, and related bacterial enzymes (Lake et al., 2001; Szczepanowski et al., 2005; Lee and Schindelin, 2008; Olsen and Lima, 2013; Lv et al., 2017a,b; Hyer et al., 2018). Nonetheless, drug discovery endeavors started earlier using several approaches, including structure-based rational design, relying on the close similarity between human UBA1 and related orthologs (Hyer et al., 2018). These efforts have led to the identification of several inhibitors that act by diverse mechanisms (Table 6). The inhibitors reported thus far can be structurally classified into ubiquitin adenylate analogs,

natural compounds, disulfides, nitro-based compounds, and adenosine sulfamates (Fig. 8). Of these, the adenosine sulfamate TAK-243 (MLN7243) is the only drug that has been advanced to phase 1 clinical trials.

a. TAK-243. TAK-243 (MLN7243) is a first-in-class, mechanism-based UBA1 inhibitor (Hyer et al., 2018). It was identified through a screen of more than 700 chemical candidates using an in vitro transthiolation assay that assesses ubiquitin transfer to the ubiquitin-specific E2 enzyme UBCH10 (also known as UBE2C). It is a second-in-class adenosine sulfamate E1 inhibitor developed after the NAE inhibitor pevonedistat, the prototype of this structural class (Soucy et al., 2009a). TAK-243 has been reported to be partially selective for UBA1, as it displays activity against UBA6 and NAE but severalfold higher IC_{50} , as assessed by transthiolation assays (Ciavarrri and Langston, 2017; Hyer et al., 2018). However, it displays minimal or no activity against kinases and carbonic anhydrases. It forms a covalent adduct with ubiquitin (through a covalent linkage between the sulfamate nitrogen of TAK-243 and glycine 76 of ubiquitin), which tightly binds to

TABLE 7
Clinical trials of TAS4464, TAK-243, and TAK-981

Identifier	Title	Phase	Disease	Single/ Combination	Combined Drug	Status	Reference
NCT02045095	A phase 1 dose escalation study of MLN7243 in adult patients with advanced solid tumors	1	Advanced solid tumors	Single	—	T	—
NCT03816319	TAK-243 in treating patients with relapsed or refractory AML or refractory MDS or CMML	1	AML, MDS, and CMML	Single	—	N	—
NCT03648372	A study to evaluate the safety, tolerability, and PK of TAK-981 in adult participants with metastatic solid tumors or lymphomas	1	Metastatic solid tumors and lymphomas	Single	—	R	—
NCT04074330	A study of TAK-981 in combination with rituximab in participants with relapsed/refractory (r/r) CD20 positive (CD20 ⁺) non-Hodgkin lymphoma	1b/2	Non-Hodgkin lymphoma	Combination	Rituximab	R	Assouline et al., 2019
NCT02978235	A dose finding study followed by a safety and efficacy study of TAS4464 for patients with MM or lymphoma	1	MM and lymphoma	Single	—	T	—

CMML, chronic myelomonocytic leukemia; N, not yet recruiting; R, recruiting; T, terminated.

the nucleotide-binding site and inhibits UBA1 by substrate-assisted inhibition (Misra et al., 2017; Hyer et al., 2018).

In cells of different origins, TAK-243 reduces the abundance of mono- and polyubiquitylated proteins and induces the accumulation of ubiquitin-regulated short-lived proteins, including p53, c-Jun, c-Myc, MCL1, and X-linked inhibitor of apoptosis protein (XIAP) (Hyer et al., 2018; Barghout et al., 2019; Best et al., 2019a; Zhuang et al., 2019). These changes are associated with a multitude of ubiquitylation-dependent biologic effects that lead to cell death, including cell cycle arrest, proteotoxic stress, and DNA damage stress (Hyer et al., 2018). In this respect, TAK-243 induces cell cycle arrest at the G2/M phase or both G1 and G2/M phases at high concentrations. Notably, the cell cycle phenotype observed after TAK-243 treatment resembles that observed after proteasome inhibitors and differs from NAE inhibitors, as TAK-243 exhibits no induction of re-replication phenotype (Hyer et al., 2018) or a slight induction of this phenotype at high concentrations (Best et al., 2019a). Given the role of UPS in the degradation of misfolded and unfolded proteins in the ER, TAK-243 induces the accumulation of such proteins, leading to sustained ER stress associated with morphologic ER changes, a terminal UPR, and apoptosis (Wang and Kaufman, 2014; Hyer et al., 2018; Best et al., 2019a; Zhuang et al., 2019). Moreover, ubiquitylation plays a pivotal role in several DNA repair pathways, such as translesion synthesis, Fanconi anemia pathway, double-strand break repair, and nucleotide-excision repair (Milhollen et al., 2015); thus, TAK-243 impairs DNA repair and induces DNA damage stress under irradiated and unirradiated conditions (Hyer et al., 2018; Barghout et al., 2019). Of note, genomic and proteomic analysis of TAK-243 effects demonstrates a pleiotropic impact on diverse signaling pathways, including proliferative signaling, inflammatory responses, apoptosis, hypoxia, oxidative stress, and oxidative phosphorylation (Zhuang et al., 2019).

As expected with these pleiotropic effects, TAK-243 demonstrates a broad antitumor activity in several malignancies, including solid tumors (Hyer et al., 2018), AML (Hyer et al., 2018; Barghout et al., 2019), myeloma (Hyer et al., 2018; Zhuang et al., 2019), B-cell lymphoma (Best et al., 2019a), cutaneous squamous cell carcinoma (McHugh et al., 2018), chronic lymphocytic leukemia (Best et al., 2019b), mantle cell lymphoma (Best et al., 2019b), and small-cell lung cancer (Majeed et al., 2019). Most tested cell lines of these malignancies are sensitive to TAK-243, with low nanomolar IC₅₀ values in vitro and robust responses in vivo, and only a few of them display intrinsic resistance (Hyer et al., 2018; Majeed et al., 2019; Zhuang et al., 2019). In contrast, data with normal human fibroblasts and in vivo studies show a lesser effect on normal versus cancer cells, suggesting TAK-243 possesses a therapeutic window in several malignancies (Hyer et al., 2018; Barghout et al., 2019). Although seemingly narrow, this therapeutic window may lead to beneficial therapeutic outcomes in selected malignancies with dose optimization and careful patient selection, as is the case with proteasome inhibitors (Ruschak et al., 2011; Hyer et al., 2018). Based on preclinical data including efficacy and toxicology studies, TAK-243 has been advanced to phase 1 clinical trials in advanced solid tumors and AML (Table 7).

TAK-243 also exhibits additive and/or synergistic effects with radiotherapy and DNA damaging agents in preclinical models of breast cancer and lung cancer (Milhollen et al., 2015; Hyer et al., 2018; Majeed et al., 2019), as well as histone deacetylase (HDAC) inhibitors in MM (Zhuang et al., 2019). Because of its different mechanism of action and biologic effects that are distinct from proteasome inhibitors, TAK-243 can overcome specific forms of resistance to these drugs (McHugh et al., 2018; Zhuang et al., 2019). Interestingly, B-cell lymphoma cells with Myc overexpression are more sensitive to TAK-243, suggesting higher dependence on ubiquitylation in such cells, resembling the

effects observed with SUMOylation inhibitors (Kessler et al., 2012; Best et al., 2019a).

Although many aspects of TAK-234 action in several malignancies have been studied, the determinants of sensitivity and resistance remain largely unknown (Barghout and Schimmer, 2018). With the drug being tested in clinical trials, it is important to identify patient cohorts that are likely to respond and to maximize the efficacy and minimize unnecessary toxicity from the drug. Such endeavors require a deeper understanding of the drug action at the cellular and molecular levels to identify transporters, metabolic factors, signaling molecules, and/or (epi)genetic mechanisms that may influence response. In this context, unbiased large-scale genetic and chemical screens can serve as a rapid and adaptable tool to achieve this goal and provide, with appropriate validation, useful insights into TAK-243 response (Barghout et al., 2018). Of note, correlating TAK-243 sensitivity with UBA1 expression levels in cell lines has been attempted; however, the data are conflicting, and the number of cell lines tested is not large enough for conclusive findings (Hyer et al., 2018; McHugh et al., 2018). Equally important are the development and optimization of biomarkers to monitor drug response in these clinical trials.

b. Ubiquitin Adenylate Analogs. Adenosyl-phospho-ubiquitinol (APU) was synthesized in 1990 and is reportedly the first experimental inhibitor of UBA1 (Wilkinson et al., 1990). It is a stable analog of the ubiquitin~AMP intermediate, formed during the adenylation step of ubiquitin activation, with a methylene group replacing the carbonyl of the C-terminal glycine residue (Brownell et al., 2010). Biochemically, it inhibits ubiquitin activation in a competitive manner with ATP and a noncompetitive manner with ubiquitin and tightly binds to the nucleotide-binding site of the free rather than ubiquitin-bound form. APU also inhibits protein degradation in cell-free assays; however, its structural features are not suggestive of appreciable cell permeability (Brownell et al., 2010). As such, the biochemical properties of this inhibitor, particularly affinity and ATP competitiveness, are not sufficiently favorable to produce a physiologically relevant UBA1 inhibition (Brownell et al., 2010). However, APU or related analogs could potentially serve as chemical probes to interrogate ubiquitin biology. In this respect, ubiquitin (Ub)-AMSN and Ub-AVSNS have been developed as semisynthetic mechanism-based UBA1 inhibitors by chemical ligation of truncated ubiquitin to a synthetic cysteyleglycylglycyl tripeptide with a C-terminal 5'-sulfonyl-adenosine and an N-terminal cysteine (Lu et al., 2010). Although they are structurally related, Ub-AMSN and Ub-AVSNS possess distinct features that allow them to inhibit different steps of ubiquitin activation. Specifically, Ub-AMSN inhibits the adenylation step because of its nonhydrolyzable sulfamide,

which mimics the phosphate group in the ubiquitin~AMP intermediate. On the other hand, Ub-AVSNS inhibits the ubiquitin thioester formation step because of its electrophilic vinyl sulfonamide, which traps the nucleophilic cysteine of UBA1. In cell-free assays, Ub-AMSN and Ub-AVSNS selectively inhibit Uba1~ubiquitin thioester formation in a concentration-dependent manner. In addition, Ub-AVSNS forms a covalent adduct with Uba1 that is resistant to hydrolysis by dithiothreitol as opposed to the native Uba1~ubiquitin thioester. Importantly, these inhibitors have been exploited to trap ubiquitin activation intermediates to gain insights into the biochemical and structural basis of ubiquitin activation (Olsen et al., 2010).

c. Natural Compounds. This class comprises panepophenanthrin, himeic acid A, hyrtioreticulins A/B, and largazole. The UBA1 inhibitory activities of these compounds were not extensively characterized or verified in other studies. In addition, all of them except largazole were not tested in cellular contexts. Panepophenanthrin is a bioactive compound derived from the fermented broth of the mushroom strain *Panus rudis* Fr. IFO 8994. It is reported to possess UBA1 inhibitory activity in a cell-free thioester formation assay (Sekizawa et al., 2002). Himeic acid A is another natural compound isolated from the marine fungus *Aspergillus* sp. In analogy to panepophenanthrin, it reportedly inhibits cell-free thioester formation at a high micromolar concentration; however, further characterization of its UBA1 inhibitory activity has not been pursued in independent studies (Tsukamoto et al., 2005). Hyrtioreticulins A and B are indole alkaloids isolated from the marine sponge *Hyrtios reticulatus*. When tested in vitro, they showed a more potent UBA1 inhibitory activity compared with panepophenanthrin and himeic acid A; however, they displayed no activity against HeLa cells (Yamanokuchi et al., 2012). Based on these reports, himeic acid A is predicted to inhibit UBA1 at the ubiquitin adenylation step, whereas hyrtioreticulins A and B inhibit UBA1 at the thioester formation step (Yamanokuchi et al., 2012).

Largazole is a macrocyclic compound isolated from the Floridian marine cyanobacterium *Symploca* sp. It demonstrates a potent and preferential antiproliferative activity against breast, osteosarcoma, colon, and neuroblastoma cancer cell lines, with relatively less activity against nontransformed cells (Taori et al., 2008). Although the mechanism of this antiproliferative action has been attributed to HDAC inhibition in several studies (Liu et al., 2010; Ghosh et al., 2012; Hong and Luesch, 2012; Law et al., 2013; Poli et al., 2017), a study has reported that largazole and select analogs also possess UBA1-inhibitory activity (Ungermannova et al., 2012b). Specifically, largazole was identified as a top hit through a cell-based screen for compounds that stabilize GFP-labeled p27 in a lung epithelial cell line with potent activity in the nanomolar range. Further investigation

using cell-free ubiquitylation assays demonstrated that largazole and ketone/ester analogs inhibit UBA1 at the adenylation step, resulting in decreased ubiquitin charging of ubiquitin-conjugating enzyme E2 R1 (CDC34) and reduced ubiquitylation of p27 with no effect on SUMOylation. Nonetheless, the higher micromolar IC_{50} required for in vitro UBA1 inhibition as opposed to the nanomolar cellular activity suggests that largazole-induced cytotoxicity is mediated through HDAC inhibition and subsequent cell cycle arrest rather than UBA1 inhibition (Ungermannova et al., 2012b; Wu et al., 2013).

d. Nitropyrazones and Nitropiperazines. Nitro-based UBA1 inhibitors include pyrazones (PYR-41 and PYZD-4409) and piperazines (JS-K). They share a nitro group that is susceptible to displacement by nucleophilic attack and covalent interaction with the catalytic cysteine of the CCD, resulting in irreversible inhibition (Yang et al., 2007; Xu et al., 2010).

PYR-41, the prototype of this class, was reported in 2007 as the first cell-permeable inhibitor of UBA1. It was discovered through a secondary screen of E3 ligase inhibitors for hits with reactivity to UBA1's catalytic cysteine (Yang et al., 2007). In vitro, PYR-41 demonstrates activity against the E1 enzyme in addition to some activity against homologous to the E6-AP carboxyl terminus (HECT) E3s. Biochemically, it inhibits thioester formation with free UBA1 without impact on ubiquitin transfer to E2s from ubiquitin-charged UBA1. It also inhibits degradative and nondegradative forms of ubiquitylation and cytokine-induced activation of NF- κ B. p53 is an important transcription factor and tumor suppressor protein whose stability is prone to regulation by degradative ubiquitylation (Vucic et al., 2011). PYR-41 induces the stabilization of p53 with a subsequent increase in transcriptional activity and preferential cytotoxicity to transformed cells with wild-type p53. Cellular effects of PYR-41 include reductions in ubiquitin charging of UBA1 and the abundance of ubiquitylated proteins. Interestingly, increases in ubiquitylated proteins with high molecular mass (>250 kDa) as well as total SUMOylation are observed after PYR-41 treatment with no effect on neddylation (Yang et al., 2007; Kapuria et al., 2011). This paradoxical accumulation of ubiquitylated proteins is ascribed to the protein cross-linking activity of PYR-41 resulting in covalent interaction with several DUBs (e.g., USP5), kinases [breakpoint cluster region protein-Abelson tyrosine-protein kinase (BCR-ABL) and Janus kinase 2 (JAK2)], and possibly other proteins, forming adducts with high molecular weight. Therefore, it appears that PYR-41 is a promiscuous inhibitor with multiple targets, and its antineoplastic effects are likely due to the combined inhibition of UBA1 as well as other off-target effects, particularly after prolonged treatment (Kapuria et al., 2011).

PYZD-4409 is a structural analog of PYR-41 that was discovered through a focused screen of a pyrazolidine pharmacophore-based chemical library (Xu et al., 2010).

In addition to cancer cell lines, it has been tested in primary cancer cells and a mouse model of murine leukemia. It is cytotoxic to cell lines of hematologic and solid malignancies, with MM cell lines being particularly sensitive. It also displays preferential cytotoxicity to AML versus normal hematopoietic primary cells (Xu et al., 2010). Mechanistically, it stabilizes the short-lived proteins p53 and cyclin D3 and induces ER stress resulting in cell death. Nonetheless, the structural similarities to PYR-41 suggest that PYZD-4409 may also induce cytotoxicity through off-target effects besides UBA1 inhibition.

JS-K is a nitric oxide (NO $^{\bullet}$)-releasing prodrug that is metabolically activated by glutathione *S*-transferases in the presence of glutathione. Therefore, it is preferentially cytotoxic to glutathione *S*-transferase-overexpressing malignancies, such as MM (Kiziltepe et al., 2007). In this context, it is cytotoxic to MM cell lines and primary cells, with a lesser effect on normal hematopoietic cells. It has also been tested in a mouse xenograft model of MM without significant toxicity. Mechanistically, JS-K induces apoptosis and activates DNA damage response and c-Jun N-terminal kinase (JNK) pathways. The apoptotic effects of JS-K in preclinical models of MM have been attributed to NO $^{\bullet}$ generation without implicating UBA1 (Kiziltepe et al., 2007). A later study, however, showed that JS-K induces *S*-nitrosylation of UBA1 in immortalized human cells, resulting in UBA1 inhibition along with other effects similar to those reported with nitropyrazones in transformed cells (Kitagaki et al., 2009). The effects of JS-K on ubiquitylation and p53 levels have been further corroborated in prostate cancer cells (Tan et al., 2017). Therefore, it appears that the nitro group of JS-K serves as a source of NO $^{\bullet}$ that causes inhibition of multiple targets and is likely a leaving group at which covalent adduct formation with UBA1 occurs in a similar manner to nitropyrazones.

e. Disulfides. As with largazole, the disulfide compound NSC624206 was discovered through a cell-based screen of a National Cancer Institute library for compounds that prevent degradation of p27 (Ungermannova et al., 2012a). As the inhibitory activity is lost with an analog lacking the disulfide group, this group is thought to be essential for activity (Ungermannova et al., 2012a). In a similar manner to nitropyrazones, NSC624206 inhibits UBA1~ubiquitin thioester formation, with little or no effect on the adenylation step. It is speculated that the disulfide group of NSC624206 may undergo a nucleophilic attack by the -SH group of the catalytic cysteine in a thiol-disulfide exchange reaction, forming a covalent adduct coupled with UBA1 inhibition. NSC624206 is a cell-permeable inhibitor, as it stabilizes p27 in lung epithelial and liver cancer cells (Ungermannova et al., 2012a).

C. Small Ubiquitin-Like Modifier-Activating Enzyme

SUMOylation is a post-translational modification that regulates several cellular processes, including nuclear transport, gene expression, chromosome

segregation, quality control and proteasomal degradation of proteins, DNA damage response, and cell cycle progression (Gareau and Lima, 2010; Eifler and Vertegaal, 2015). SAE is the E1 enzyme that catalyzes SUMO activation, which is followed by SUMO conjugation to a variety of cellular proteins (Schulman and Harper, 2009). Structurally, SAE is a canonical heterodimeric enzyme composed of two subunits, SAE1 and SAE2 (UBA2), with several domains analogous to UBA1 and NAE (Fig. 2). Of the four SUMO isoforms known so far, SUMO1–3 are activated by SAE; however, it is not known whether SUMO4 is conjugated to cellular proteins (Gareau and Lima, 2010). Compared with the ubiquitylation cascade that involves tens of E2 enzymes, hundreds of E3 ligases, and over 100 DUBs, the SUMOylation cascade is much less complex, as it involves a single E2 enzyme, UBC9, as well as few E3 ligases and SUMO-specific proteases (SENPs) (Gareau and Lima, 2010). Among the substrates that are regulated by SUMOylation are RanGAP1, I κ B α , PCNA, p53, c-Jun, c-Myc, and promyelocytic leukemia (PML) nuclear bodies (Seeler and Dejean, 2003; Wilkinson and Henley, 2010). Of these, RanGAP1 SUMOylation is commonly used in SUMOylation assays to identify and characterize SUMOylation modulators (Sarge and Park-Sarge, 2009).

1. Role of Small Ubiquitin-Like Modifier-Activating Enzyme in Cancer. Given its critical role in many cellular processes, SUMOylation is tightly coordinated to maintain normal cellular functions (Seeler and Dejean, 2017). In cancer, however, SUMOylation may be dysregulated, and mostly the activation of the SUMOylation pathway is associated with an adverse outcome. The expression, activity, and stability of SAE1/2 are among the key regulators of SUMOylation, and either SAE1, SAE2, or both may be altered in several malignancies. For example, primary mantle cell lymphoma samples and cell lines have been reported to display higher SAE1/2 expression as well as a higher abundance of global SUMOylated proteins compared with normal counterpart cells (Hanel et al., 2019). In addition, gastric and small-cell lung cancer show a high expression of SAE2, which promotes tumor progression and correlates with a poor prognosis (Shao et al., 2014; Liu et al., 2015). SAE1 is upregulated in glioblastoma tumors compared with normal brain samples (Bellail et al., 2014) and is a part of a 75-gene signature associated with lymph node metastasis in lung adenocarcinomas (Inamura et al., 2007). SAE expression is also reported to be important for Myc-driven tumor progression in solid and hematologic malignancies. For example, elevated SAE1/2 expression has been reported in Myc-overexpressing lymphoma, and as assessed by genetic and pharmacologic approaches, SAE is functionally important for Myc-driven tumorigenesis (Hoellein et al., 2014). Consistent with these findings, low SAE1/2 expression in

Myc-overexpressing breast cancers has been associated with favorable survival outcomes in patients (Kessler et al., 2012). In these contexts, targeting SAE is expected to be synthetically lethal with hyperactive Myc, demonstrating SUMOylation is crucial to tolerate aberrant Myc signaling and maintain tumor progression, possibly through alleviating mitotic stress. Similarly, higher dependence on SUMOylation enzymes, including SAE, has been observed in breast cancer cells with Notch1 activation (Licciardello et al., 2015). Moreover, high SAE1 expression is associated with unfavorable prognosis in liver, renal, and thyroid cancers, with similar prognostic outcomes with SAE2 in liver and renal cancers (Uhlen et al., 2017). In preclinical models of colorectal cancer, SAE has been found to play a critical role in cancer stem cell self-renewal and maintenance (Du et al., 2016). The roles of other components of the SUMO pathway in cancer have been reviewed elsewhere (Eifler and Vertegaal, 2015; Seeler and Dejean, 2017). These data provide compelling evidence that SUMOylation may be hyperactive in several cancers compared with normal cells to support cellular stresses associated with the malignant phenotype. This form of nononcogene addiction can be therapeutically targeted at different levels of the SUMOylation machinery including SAE that initiates the cascade (Luo et al., 2009).

2. Role of SUMOylation in Differentiation Therapy of Acute Promyelocytic Leukemia. Acute promyelocytic leukemia (APL) is a subtype of AML characterized by a chromosomal translocation [t(15;17)(q22;q12)] involving the *PML* gene and the retinoic acid receptor α (*RARA*) gene, producing the fusion oncoprotein PML/RARA (de Thé et al., 2017). Biologically, PML/RARA contributes to APL pathogenesis by the deregulation of RAR α -mediated transcriptional control and disruption of PML nuclear bodies (NBs), which are spherical, stress-sensitive nuclear structures involved in regulating many cellular functions and are mainly organized by PML protein (Bernardi and Pandolfi, 2007; de Thé et al., 2017). These effects lead to resistance to apoptosis, differentiation block, and enhancement of self-renewal of myeloid progenitors (de Thé and Chen, 2010). This oncoprotein is the main oncogenic driver in several types of APL, providing a unique opportunity for targeted therapy aimed at inducing differentiation. Currently, differentiation therapy comprising ATO plus ATRA is a frontline therapeutic option that has transformed the outlook of APL in many patients, with cure rates up to 90% (Lallemand-Breitenbach et al., 2012; Lo-Coco et al., 2013; Sanz et al., 2019). Mechanistically, ATO and ATRA bind to PML/RARA to induce its degradation, resulting in the reformation of PML NBs, p53 activation, loss of self-renewal, and differentiation (Jeanne et al., 2010; Zhang et al., 2010; Ablain et al., 2014; de Thé et al., 2017). ATO-induced degradation of PML/RARA is dependent on both SUMOylation and ubiquitylation. Specifically, ATO targets the PML moiety

Fig. 9. Chemical structures of SAE inhibitors. SAE inhibitors can be classified into adenosine sulfamates (e.g., TAK-981 and ML-792), oxabicycloheptadienes (e.g., COH000), natural compounds (e.g., ginkgolic acid, anacardic acid, davidiin, tannic acid, and kerriamycin B), SUMO adenylylate analogs and SUMO-mimicking peptides (e.g., SUMO-AMSN, SUMO-AVSN, and pS50), and urea-based inhibitors (e.g., compound 21).

of the fusion oncoprotein, inducing oxidative and conformational changes that enhance recruitment and tight binding of UBC9, leading to hyperSUMOylation of PML. This form of PML actively recruits the SUMO-dependent ubiquitin E3 ligase RING finger protein 4 (RNF4) followed by K48-linked polyubiquitylation and proteasomal degradation (Lallemand-Breitenbach et al., 2008; Tatham et al., 2008). Of note, genetic and chemical inhibition of SAE in colon cancer cells has been reported to reduce SUMOylation and cause disruption of PML NB organization and dissociation of the PML-interacting protein death-associated protein 6 (DAXX), leading to antitumor effects (He et al., 2015, 2017). Therefore, it appears that the induction of either hyper- or hypoSUMOylation can be leveraged in different contexts to obtain beneficial therapeutic outcomes.

3. Antitumor Immunomodulatory Roles of SUMOylation. Interferons (IFNs) are small secreted proteins that play a crucial role in innate and adaptive immunity. They include three classes: type I (IFN- α , IFN- β , IFN- ϵ , IFN- κ , IFN- ω), type II (IFN- γ), and type III IFNs

(IFN- λ , interleukin-28 [IL-28], and interleukin-29 [IL-29]), which initiate three distinct signaling cascades (Schneider et al., 2014). SUMOylation has been reported to regulate type I IFN signaling (Decque et al., 2016; Crowl and Stetson, 2018). Type I IFN signaling is initiated by binding of IFNs to a heterodimeric cell surface receptor complex composed of IFN- α receptor (IFNAR) 1 and IFNAR2, leading to the activation of Janus kinase-signal transducer and activator of transcription protein (JAK-STAT) signaling and formation of a heterotrimeric complex termed IFN-stimulated gene factor 3 (ISGF3), which is composed of STAT1, STAT2, and IFN-regulatory factor 9 (IRF9) (Schneider et al., 2014). ISGF3 undergoes nuclear translocation, in which it activates the expression of ISGs by modulating IFN-stimulated response elements. Recently, SUMOylation has been reported to negatively regulate the innate inflammatory response by silencing IFN- β expression and attenuating the priming of inflammatory cytokine production by type I IFNs (Decque et al., 2016). Accordingly, inhibiting SUMOylation by *UBC9* knock-down enhanced the innate immune responses. Loss of

TABLE 8
SAE inhibitors and their pharmacologic properties

Inhibitor	Type	Malignancies	Class	EC ₅₀ /IC ₅₀ /Dose	Reference
TAK-981	Clinical: phase 1	BCL and colorectal carcinoma	Adenosine sulfamate	SUMOylation assays: 0.005–0.026 μ M; MCL cell lines: 17–62.5 nM; dose in mice: 5–10 mg/kg, i.v., twice weekly; 15 mg once weekly	Berger et al., 2019; Hanel et al., 2019; Hattori et al., 2019; Khattar et al., 2019; Nakamura et al., 2019
ML-792	Experimental	Cell-free and cellular activity in colorectal, breast, and melanoma cancer cells	Adenosine sulfamate	SUMOylation assays: 0.003–0.051 μ M; cytotoxicity: 0.03–0.45 μ M	He et al., 2017
COH000	Experimental	Cell-free and cellular activity in lymphoma, colorectal, and ovarian cancer cells in vitro and in vivo	Oxabicycloheptadiene	Cell-free RanGAP1 SUMOylation: 0.2 μ M; dose in mice: 10 mg/kg once daily	Lv et al., 2018b; Li et al., 2019
Davidiin	Experimental	Cell-free and cellular activity in gastric, prostate, lung, and hepatocellular cancer cells	Natural compound; ellagitannin	Cell-free RanGAP1-C2 SUMOylation: 0.15 μ M; cytotoxicity: 8.3–16.4 μ M; dose in mice: 10 mg/kg 3 times weekly	Takenoto et al., 2014; Wang et al., 2014
Ginkgolide acid	Experimental	Cell-free and cellular activity	Natural compound; alkyl phenol derivative	Cell-free substrate SUMOylation: 3 μ M	Fukuda et al., 2009a
Kerriamycin B (Urdamycin A)	Experimental	Cell-free and cellular activity	Natural compound; polycyclic antibiotic	Cell-free RanGAP1-C2 SUMOylation: 11.7 μ M	Fukuda et al., 2009b
Tannic acid	Experimental	Cell-free and cellular activity (no cytotoxicity in liver cancer cells)	Natural compound; gallotannin	Cell-free rhLRH-1 SUMOylation: 12.8 μ M	Suzawa et al., 2015
Anacardic acid	Experimental	Cell-free and preclinical activity in AML	Natural compound; alkyl phenol derivative	Cell-free substrate SUMOylation: 2.2 μ M; median IC ₅₀ (primary AML samples): 42 μ M; dose in mice: 2 mg/kg/d	Fukuda et al., 2009a; Bossis et al., 2014
Compound 21 SUMO-AMSN	Experimental	Cell-free activity	Biaryl urea derivative	Cell-free RanGAP1-C2 SUMOylation: 14.4 μ M	Kumar et al., 2013b
SUMO-AVSN	Experimental	Cell-free activity	Semisynthetic SUMO adenylate	NA	Lu et al., 2010; Olsen et al., 2010
pS50 (YSFVSGG)	Experimental	Cell-free activity	Semisynthetic SUMO adenylate	NA	Lu et al., 2010; Olsen et al., 2010
pS90 (YQYVSGG)	Experimental	Cell-free activity	Synthetic heptameric SUMO-mimicking peptides	Cell-free thioester formation: 125 μ M	Zhao et al., 2014
			Synthetic heptameric SUMO-mimicking peptides	Cell-free thioester bond formation: 131 μ M	Zhao et al., 2014

BCL, B-cell lymphoma; NA, not available; rhLRH-1, recombinant human liver receptor homolog-1.

SUMOylation after *SAE* or *UBC9* knockout was also found to induce a type I IFN response, however, through a noncanonical mechanism independent of known IFN-regulating pathways (Crowl and Stetson, 2018). Among the inducers of type I IFN signaling is the cGAS–stimulator of interferon genes protein (cGAS-STING) pathway, which acts as a sensor of aberrant cytosolic DNA and activates the expression of type I IFNs (Schneider et al., 2014; Chen et al., 2016b). SUMOylation has been reported to suppress the activity of cGAS, and this suppression is reversed by the SUMO-specific protease SENP7, which catalyzes deSUMOylation of cGAS, triggering type I IFN-dependent innate immune responses (Cui et al., 2017). The importance of SUMOylation is not confined to regulating innate immunity, as it also plays a key role in adaptive immune responses. In this context, SENP3 has been found to maintain the stability and function of regulatory T cells through deSUMOylation of BTB and CNC homolog 2 (BACH2), a regulatory T-cell-specific transcription factor, facilitating its nuclear retention and function (Yu et al., 2018b). Therefore, it appears that SUMOylation restrains a subset of innate and adaptive immune responses; thus, targeting the SUMOylation machinery including *SAE* can serve, in specific contexts, as a potential therapeutic strategy to obtain useful antitumor and antimicrobial immune effects.

4. Small Ubiquitin-Like Modifier-Activating Enzyme Inhibitors. The structural basis of SUMO activation by human *SAE* and interaction with *UBC9* has been elucidated since 2005, and most *SAE* inhibitors reported later are natural compounds whose *SAE*-inhibiting activity was discovered without regard to the crystal structure of *SAE* (Lois and Lima, 2005; Wang et al., 2007; Olsen et al., 2010). These endeavors started with Gam1, an adenoviral protein that interacts with *SAE* and enhances *SAE*-*UBC9* proteasomal degradation (Boggio et al., 2004). This was followed by the discovery of a series of small-molecule *SAE* inhibitors with diverse structures until the first mechanism-based selective *SAE* inhibitor, ML-792, was developed more than a decade later (He et al., 2017). *SAE* inhibitors reported so far can be classified into SUMO adenylate analogs and SUMO-mimicking peptides, natural compounds, urea-based derivatives, oxabicycloheptadienes, and adenosine sulfamates (Fig. 9; Table 8). A first-in-class *SAE* activator, N106, was reported in 2015 as a potential modulator of heart failure (Kho et al., 2015).

a. ML-792 and TAK-981. ML-792 and TAK-981 are adenosine sulfamate *SAE* inhibitors, and they share similar profiles but differ in terms of their in vivo properties (Huszar, 2019). In pursuit of developing *SAE* inhibitors that are analogous to pevonedistat, ML-792 was discovered through a drug development program that used a pyrazole-carbonylpyrimidine-based scaffold (He et al., 2017). In analogy with other adenosine

sulfamates, ML-792 acts by substrate-assisted inhibition and forms a covalent adduct with SUMO. In cell-free and cell-based assays, ML-792 demonstrates a potent and selective *SAE*-inhibitory activity, with little or no activity against related E1s or other ATP-dependent enzymes. ML-792 is also cytotoxic to colorectal, breast, and melanoma cancer cell lines, with Myc-overexpressing cell lines being particularly sensitive. Despite the importance of the SUMOylation pathway in regulating gene expression, ML-792 only induces modest transcriptional and proteomic changes, with no impact on splicing in different cell lines. As opposed to TAK-243, ML-792 has little or no effect on DNA repair, suggesting that DNA damage plays no role in the observed cytotoxicity (He et al., 2017; Hyer et al., 2018). Instead, ML-792 induces mitotic disruption (multinucleation, cell enlargement, and endoreduplication), senescence, and chromosome segregation defects (decrease in the anaphase/telophase of mitosis and formation of DNA bridges). Cytotoxicity and cell cycle changes are evident only in actively cycling cells and can be rescued by the overexpression of ML-792-resistant *UBA2* S95N M97T mutant, suggesting that mitotic disruption is functionally important for ML-792 cytotoxicity (He et al., 2017).

Despite the excellent profile of ML-792 ex vivo, it shows poor stability in vivo, making it suitable only as a chemical probe for interrogating SUMOylation biology (He et al., 2017; Huszar, 2019). Specifically, high doses are required to maintain in vivo responses, as ML-792-SUMO adducts are rapidly released from *SAE* and degraded after in vivo administration (Huszar, 2019). Conversely, TAK-981—developed thereafter to overcome this limitation—forms SUMO adducts that are stable and causes sustained inhibition of SUMOylation after administration of small doses of the drug (Huszar, 2019). TAK-981 selectively inhibits *SAE*~SUMO and *UBC9*~SUMO thioester formation as well as global SUMOylation, with IC₅₀ values in the low nanomolar range, and without significant effects on neddylation or ubiquitylation (Berger et al., 2019; Hanel et al., 2019; Huszar, 2019). In analogy with ML-792, TAK-981 treatment in mantle cell lymphoma cells induces cell cycle arrest and mitotic disruption, as evidenced by the accumulation of polyploid and aneuploid cells (Hanel et al., 2019).

Interestingly, prolonged effects of TAK-981 in vivo have allowed the observation of several immune responses that are associated with the antitumor effects of the drug, particularly against tumor models with intrinsic resistance to TAK-981 in vitro (Huszar, 2019). The molecular basis of these responses partly derives from the role of SUMOylation in regulating type I IFN signaling (Decque et al., 2016; Crowl and Stetson, 2018). This is consistent with the reduced antitumor effects of TAK-981 with prior administration of anti-IFN- α/β receptor 1 antibody (Berger et al., 2019). The effects of TAK-981 mediated by type I IFN signaling

are pleiotropic and implicate several effectors of innate and adaptive arms of immunity (Berger et al., 2019; Khattar et al., 2019; Lam et al., 2019; Nakamura et al., 2019). Direct microinjection of TAK-981 into tumors is also associated with both local and abscopal immune responses, which are consistent with those observed after systemic dosing (Hatton et al., 2019). In addition, TAK-981 exerts synergistic antitumor effects when administered in combination with other immunotherapies, such as rituximab, an anti-CD20 monoclonal antibody, and immune checkpoint inhibitors (Huszar, 2019; Nakamura et al., 2019). Based on these preclinical data, TAK-981 is currently evaluated in phase 1 clinical trials for patients with metastatic solid tumors and lymphomas (Table 7).

b. COH000. Most SAE inhibitors reported so far target either the nucleotide-binding site or the catalytic cysteine of the enzyme. Through an HTS of a library of ~300,000 small molecules for compounds that inhibit RanGAP1 SUMOylation, Li et al. (2019) discovered COH000, an inhibitor that targets a novel, highly conserved allosteric site in the SAE. COH000 is an oxabicycloheptadiene-based compound that selectively and potently inhibits SUMOylation in cell-free and cell-based assays with no effects on ubiquitylation. In this respect, COH000 inhibits SAE~SUMO thioester formation in a noncompetitive manner with neither ATP nor SUMO, and this is associated with a very slow off-rate, suggesting irreversible inhibition via covalent binding to the SAE (Li et al., 2019). Mass spectrometry data demonstrate covalent adduct formation after the binding of COH000 to a hidden cysteine residue (Cys30), without modifying the other SAE's non-disulfide-bonded cysteines, including the catalytic cysteine (Cys173). Chemically, a Michael addition reaction is involved in the covalent bond formation between the Cys30 and an electrophilic center within the 7-oxabicyclohept-2-ene group of COH000. However, the selectivity observed with COH000 implies that noncovalent interactions precede covalent adduct formation.

Cys30 is located in the AAD at a site proximal to the active adenylation site; however, as per published SAE crystal structures, it is among the inaccessible cysteines, suggesting that COH000 binding is associated with a set of conformational changes that expose this residue (Li et al., 2019). This has been confirmed by a detailed structural analysis of SAE in complex with COH000, in which the drug was shown to target a novel cryptic allosteric site located specifically between the AAD and IAD (Lv et al., 2018b). Although the associated conformational changes involving AAD, IAD, and CCD are an extension of those observed during the normal catalytic mechanism of SAE, they are differently orchestrated in a way that leads ultimately to locking the enzyme in an inactive conformation. Interestingly, COH000 induces ubiquitylation and degradation of

SAE2, suggesting that this inactive conformation is susceptible to proteasomal degradation.

The anti-SAE activity of COH000 is associated with reduced proliferation, apoptosis, increased miR-34b expression, and decreased c-Myc expression in lymphoma and colorectal cancer cell lines and primary cells as well as reduced tumor growth in an esterase-deficient mouse xenograft model of colorectal cancer (Li et al., 2019). Since COH000 contains two ester bonds, it is susceptible to hydrolysis by plasma esterases; however, it is unclear whether cleavage of these bonds will compromise the efficacy of the drug *in vivo*.

In addition to the unique mechanism of action and potent biologic activities, COH000 has enabled a chemically assisted discovery of a novel cryptic pocket that was not previously reported and provided new insights into the structural biology of SAE and possibly other E1 enzymes (Li et al., 2019).

c. Small Ubiquitin-Like Modifier Adenylate Analogs and Small Ubiquitin-Like Modifier-Mimicking Peptides. SUMO-AMSN and SUMO-AVSN are semisynthetic mechanism-based SAE inhibitors that are structurally and biochemically analogous to Ub-AMSN and Ub-AVSN (Lu et al., 2010). They were similarly exploited as chemical probes to interrogate the structural basis and biochemistry of SUMO activation in cell-free assays (Lu et al., 2010; Olsen et al., 2010).

Based on the SUMO C-terminal sequences that can be activated by SAE, as profiled by phage display technology, Zhao et al. (2014) developed pS50 and pS90, two other mechanism-based SAE inhibitors. pS50 and pS90 are heptameric SUMO-mimicking peptides that can be activated by SAE, conjugated to UBC9, and transferred to protein substrates at a higher rate compared with the wild-type SUMO, but with little or no functionality. Therefore, these peptides competitively inhibit SUMOylation at every step of the cascade, starting with the SAE-catalyzed activation step (Zhao et al., 2014).

d. Natural Small Ubiquitin-Like Modifier-Activating Enzyme Inhibitors. This class of SAE inhibitors includes ginkgolic acid, anacardic acid, davidiin, tannic acid, and kerriamycin B, which have diverse chemical structures and molecular mechanisms of SAE inhibition. A cell-based screen of herbal extracts and nutraceuticals for compounds that inhibit SUMOylation identified ginkgolic acid, anacardic acid, and davidiin as SAE inhibitors (Fukuda et al., 2009a; Takemoto et al., 2014). Ginkgolic acid is an alkylphenol derivative and one of the active ingredients of ginkgo (*Ginkgo biloba*) leaf extract, and anacardic acid is a structurally related analog found in cashew nut (*Anacardium occidentale*) shells and immature ginkgo seeds (Fukuda et al., 2009a; Takemoto et al., 2014). Both compounds reduce cell-free protein-specific and cell-based global SUMOylation in a concentration- and

time-dependent manner, with little or no effect on ubiquitin conjugates. Structure-activity relationship (SAR) and molecular docking studies suggest that the carboxylic acid and long aliphatic side chains are essential for activity and that these compounds potentially target the AAD of the enzyme (Hamdoun and Efferth, 2017). Although known to interact with multiple targets, inducing numerous molecular activities, it appears that the effects of ginkgolic and anacardic acid on the SAE are achievable at low concentrations (Fukuda et al., 2009a). A subset of these other targets may themselves be subject to regulation by SUMOylation. For instance, the NF- κ B pathway, regulated partly by SUMOylation of NEMO and I κ B α (Seeler and Dejean, 2017), is modulated by anacardic acid, resulting in antitumor effects in cancer cell lines of solid and hematologic malignancies (Sung et al., 2008). Inhibition of NEMO SUMOylation by ginkgolic acid in cell-free assays has also been reported (Hamdoun and Efferth, 2017). Therefore, the inhibition of SAE and subsequent effects on global SUMOylation may mediate the pleiotropic effects of these two compounds on other targets.

Ginkgolic and anacardic acids have also been reported to display antitumor effects in cancers addicted to oncogenes that promote a higher reliance on SUMOylation. In this respect, anacardic acid selectively induces growth arrest, apoptosis, and aneuploidy in B-cell lymphoma cell lines with Myc activation (Hoellein et al., 2014). Similarly, ginkgolic acid reduces proliferation and induces apoptosis in engineered and patient-derived breast cancer cell lines with Notch1 activation, without cytotoxicity in isogenic cell lines with normal Notch1 activity (Licciardello et al., 2015). Consistent with these findings, ginkgolic acid reduces migration in wound-healing assays of MCF7 and MDA-MB 231 breast cancer cell lines without inducing cytotoxicity (Hamdoun and Efferth, 2017).

Inhibition of SUMOylation by anacardic acid has been reported to overcome the resistance of AML to conventional therapies. In this context, treatment with daunorubicin, cytarabine, or etoposide induced ROS-dependent deSUMOylation that was associated with transcriptional alterations and apoptosis in chemosensitive AML cells (Bossis et al., 2014). Intrinsically chemoresistant AML cells, however, failed to undergo deSUMOylation after chemotherapy. Treatment with anacardic acid in vitro and in vivo induced deSUMOylation and circumvented AML chemoresistance. Most AML subtypes, except for APL, are not responsive to differentiation therapy with ATRA. Baik et al. (2018) reported that this was partly due to the SUMOylation that repressed ATRA-induced expression of genes involved in myeloid differentiation, proliferation, and apoptosis. Suppression of SUMOylation by anacardic acid and other SUMOylation inhibitors sensitized non-APL AML cells to ATRA, suggesting a potential utility of differentiation therapy in other subtypes of AML in

addition to APL (Baik et al., 2018). ATRA is mostly used in APL in combination with ATO that acts partly by inducing hyperSUMOylation and subsequent degradation of PML/RARA. Therefore, SUMOylation inhibitors may be beneficial when combined with ATRA therapy in non-APL AML, although in theory they may cause adverse outcomes when combined with ATO therapy in APL.

Davidiin is an ellagitannin found in the botanical extract of *Davidia involucrata* and *Polygonum capitatum* (Wang et al., 2014). Compared with other known naturally occurring SUMOylation inhibitors, it displays a more potent inhibition of SUMOylation in vitro, with an IC₅₀ in the nanomolar range and antiproliferative activity in several cancer lines in the micromolar range. The antitumor activity of davidiin in vivo is reported to be associated with the downregulation of enhancer of zeste homolog 2 (EZH2), a histone-lysine *N*-methyltransferase that itself is a target for SUMOylation (Riising et al., 2008; Wang et al., 2014).

Suzawa et al. (2015) performed a cellular, gene expression-based, phenotypic screen of a 1600-compound library for drugs that inhibit human liver receptor homolog-1 (hLRH-1) SUMOylation using two genes whose expression undergoes a robust induction upon hLRH-1 deSUMOylation. They identified tannic acid, a polyphenolic gallotannin related to davidiin that is extracted from several botanical sources, as an SAE inhibitor. Compared with ginkgolic acid, tannic acid exhibits much less cytotoxicity in liver cancer cells; however, its effects on the SAE appear to be specific and not due to promiscuous binding or modulation of ROS levels, known with similar polyphenols.

An earlier cell-based screen of samples from microbial cultured broth identified kerriamycin B as a potential inhibitor of SUMOylation (Fukuda et al., 2009b). Kerriamycin B is a polycyclic antibiotic produced by certain strains of actinomycetes and is reported to reduce substrate-specific and global SUMOylation by targeting the SAE and inhibiting the formation of E1~SUMO thioesters.

e. Urea-Based Small Ubiquitin-Like Modifier-Activating Enzyme Inhibitors. The discovery of this class of SAE inhibitors was informed by the crystal structure of the SAE as opposed to natural compounds reported earlier (Kumar et al., 2013b). A structure-based virtual screening of a 78,000-compound library for molecules that docked to the nucleotide-binding site of the SAE identified a series of hits with IC₅₀ values comparable to ginkgolic acid, including compound 21. However, the structural features of these compounds were suggestive of unfavorable solubility properties. Computational interrogation of chemical databases and drug libraries identified structurally related compounds with a better solubility profile and comparable potency (Kumar et al., 2013a, 2016). Notwithstanding the potential of this class, further chemical optimization is still needed

Fig. 10. Chemical structures of UBA6, UBA5, and pan/dual E1 inhibitors. (A) Chemical structure of a selective adenosine sulfamate UBA6 inhibitor with the chemical name ((1S,2S,4R)-4-(4-(1H-indol-2-yl)-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2-hydroxycyclopentyl)methylsulfamate (here referred to as UBA6i) is shown. Moreover, the reported UBA5 inhibitors include the adenosine sulfamate compound 8.5 and the chloroacetamide DKM 2-93. (B) Multi-E1 inhibitors include adenosine sulfamates (e.g., adenosine 5'-sulfamate, compound 1, ABP1, and ABPA3), which are pan-E1 inhibitors, as well as nonnucleoside compounds, e.g., LP0040, that serves as a dual UBA1/NAE inhibitor.

to generate more potent compounds with enhanced drug-like properties.

D. Ubiquitin-Like Modifier-Activating Enzyme 6

UBA6 is a canonical monomeric E1 enzyme with a domain structure similar to UBA1 (Fig. 2). It is 40% identical to UBA1, and compared with other E1 enzymes, it possesses the highest level of similarity to UBA1 at the nucleotide-binding C-terminal domain (Groettrup et al., 2008). The discovery that UBA6 can also activate ubiquitin has challenged the conventional wisdom that UBA1 is the only ubiquitin-activating enzyme, leading to a paradigm shift in the field and serving as the first example in which two distinct E1s can activate the same UBL (Chiu et al., 2007; Jin et al., 2007; Pelzer et al., 2007; Groettrup et al., 2008). Despite having similar ubiquitin affinities and ubiquitin activation kinetics, there exist several differences between the two enzymes, including the spectrum of E2 and E3

enzymes to which they transfer activated ubiquitin (Table 5) (Jin et al., 2007; Schelpe et al., 2016). Although the substrate pools for UBA1 and UBA6 are partially overlapping, many substrates are specifically targeted by only one of the two enzymes, suggesting that they are required for nonredundant biologic functions (Liu et al., 2017c). This notion is consolidated by the lethal knockout phenotypes observed with both enzymes (Groettrup et al., 2008).

As is the case with SAE and ATG7, which can activate more than one UBL, UBA6 can activate both ubiquitin and FAT10 (Chiu et al., 2007). FAT10 is an 18-kDa protein composed of two ubiquitin-like domains, and conjugation of FAT10 to protein substrates (known as FATylation) plays poorly understood roles in immune responses, apoptosis, NF- κ B activation, and mitosis (Gong et al., 2010; Li et al., 2011; Merbl et al., 2013; Schmidtke et al., 2014; Theng et al., 2014). Interestingly, a subset of FATylated proteins can be identified

and degraded by the proteasome; therefore, the role of UBA6 in proteasomal degradation is mediated by both ubiquitylation and FATylation (Schmidtke et al., 2014). There also exist several differences between ubiquitin and FAT10 as signals for proteasomal degradation. As opposed to ubiquitylation, monoFATylation is sufficient to induce effective proteasomal degradation (Schmidtke et al., 2014). Despite having similar mechanisms of UBA6-mediated activation, there is conflicting evidence regarding the preferential affinity of UBA6 toward FAT10 and ubiquitin (Chiu et al., 2007; Gavin et al., 2012). Nevertheless, the expression of FAT10 is known to be induced in response to proinflammatory cytokines such as IFN- γ and TNF- α , which may shift the competition in favor of FAT10 in certain contexts (Raasi et al., 1999; Groettrup et al., 2008). Although *Uba6*-knockout mice are embryonically lethal, *FAT10*-knockout mice are viable, suggesting UBA6-mediated ubiquitylation rather than FATylation is responsible for the lethal phenotype (Canaan et al., 2006, 2014).

1. Role of Ubiquitin-Like Modifier-Activating Enzyme 6 in Cancer. UBA6 is affected by somatic mutations and CNVs, but their role in cancer is uncertain (Tate et al., 2019). Analysis of patient survival data has shown that high UBA6 expression is associated with unfavorable prognosis in a subset of patients with renal cancer (Uhlen et al., 2017). Apart from these, the data on the role of UBA6 in cancer is limited. Nevertheless, other components of the UBA6-initiated cascade have been implicated in the pathogenesis of cancer. For instance, FAT10 has demonstrated protumorigenic functions in hepatic and gastrointestinal malignancies (Lee et al., 2003; Yuan et al., 2014). Similarly, the UBA6-specific E2 enzyme UBA6-specific E2 conjugating enzyme 1 (USE1) is overexpressed in a subset of patients with lung cancer and promotes tumor growth, migration, and invasiveness (Kim et al., 2017). In contrast, UBA6 is inversely related to invasiveness in a subset of breast cancer in which it suppresses mammary carcinogenesis of breast cancer cells by inhibiting epithelial-mesenchymal transition (Liu et al., 2017b). Therefore, UBA6 appears to display context-dependent oncogenic and tumor-suppressive functions in different cancers.

2. Ubiquitin-Like Modifier-Activating Enzyme 6 Inhibitors. So far, there are no reports of the crystal structure of UBA6. However, biochemical studies show the unique ability of the active site of UBA6 to engage both ubiquitin and the larger diubiquitin-like FAT10 (Gavin et al., 2012). A pan-E1 adenosine sulfamate E1 inhibitor, compound 1, has been reported to inhibit several E1s, including UBA6 (Brownell et al., 2010). In a similar manner to other adenosine sulfamates, compound 1 forms an adduct with both ubiquitin and FAT10 in a reaction catalyzed by UBA6 (Gavin et al., 2012). This adduct serves as the active inhibitory species for the enzyme (Brownell et al., 2010). However, the inhibitory effects of compound 1 on UBA6-mediated

FATylation are much weaker compared with its effects on UBA6-mediated ubiquitylation. This weaker potency has been attributed to the slow rate at which the FAT10-compound 1 adduct is formed and the low affinity with which the adduct binds to the enzyme (Gavin et al., 2012). A selective adenosine sulfamate UBA6 inhibitor has been developed, but there are no biochemical or biologic data available for this inhibitor (Fig. 10) (Amidon et al., 2017).

E. Ubiquitin-Like Modifier-Activating Enzyme 7

UBA7 is a canonical monomeric E1 enzyme with a domain structure that resembles both UBA1 and UBA6 (Fig. 2). With a 46% sequence identity, UBA7 possesses the highest overall similarity to UBA1 among all E1 enzymes (Table 1). The function of UBA7 is to activate ISG15, initiating its conjugation to protein substrates, a process termed ISGylation (Krug et al., 2005). ISG15 is a 15-kDa UBL composed of two ubiquitin-like domains and is 29.7% identical to ubiquitin (Narasimhan et al., 2005; Hermann and Bogunovic, 2017). Historically, it was the second member of type 1 UBLs to be identified after ubiquitin (Loeb and Haas, 1992; Schwartz and Hochstrasser, 2003). As opposed to ubiquitin, the sequence of ISG15 is much less conserved among different species (Zhao et al., 2013c). ISG15 is initially produced as a 17-kDa precursor polypeptide that undergoes post-translational processing into the 15-kDa functional form (Blomstrom et al., 1986; Knight et al., 1988). In addition to its function as a post-translational modifier of other proteins, ISG15 can serve as a free intracellular molecule and a secreted cytokine (Knight and Cordova, 1991; Sadler and Williams, 2008; Jeon et al., 2010).

As the name suggests, the expression of ISG15 and ISGylation is induced in response to type I IFN signaling, specifically IFN- α and IFN- β (Hermann and Bogunovic, 2017). In addition, they are induced by other stimuli, including infection, radiation, ischemia, aging, and activation of the NF- κ B and p53 pathways (Villarroya-Beltri et al., 2017). Similarly, the expression of UBA7 is inducible by IFNs, supporting the enhanced ISGylation upon exposure to IFN- α/β (Fan et al., 2015). ISGylated proteins are deconjugated by USP18, an ISG15-specific protease, and a negative regulator of type I IFN signaling by a mechanism that is independent of its protease activity (Malakhov et al., 2002; Ketscher et al., 2015). USP18 is also induced in response to viral infections, IFN signaling, and DNA damage (Malakhov et al., 2002). ISGylation has been implicated in numerous cellular functions, including immune responses against viral and bacterial infections, DNA repair by translesion synthesis, regulation of cytoskeletal dynamics, exosome secretion, protein translation, autophagy, proteasomal degradation, and the hypoxic response (Villarroya-Beltri et al., 2017). In a subset of these processes, ISGylation is associated

Fig. 11. Noncanonical UBL activation by ATG7 and UBA4. (A) ATG7 activates two UBLs, ATG12 and ATG8, which lie at the apex of two cascades that exhibit significant cross talk. After activation, ATG12 is transferred to the E2-like enzyme ATG10, which in turn transfers ATG12 to ATG5. ATG12-ATG5 conjugate then associates with ATG16, forming an E3-like complex, ATG12-ATG5-ATG16, which is catalytically important in the ATG8 conjugation arm. In this other arm, ATG8 is first proteolytically activated by the protease ATG4 and is then activated and transferred by ATG7 to the E2-like enzyme ATG3. ATG8 is then conjugated to PE in a reaction facilitated by the ATG12-ATG5-ATG16 complex. ATG8-PE conjugate is functionally important for the formation of autophagosomes and the identification of cargos during autophagy. The general scheme of UBL activation (UBL→E1→E2→E3) is shown in parallel to the corresponding enzymes in the cascade. (B) UBA4 activates URM1 by a noncanonical mechanism that enables its dual function as a post-translational modifier and a sulfur carrier. After adenylation of URM1 and subsequent thioester formation at the catalytic cysteine (CC), URM1 is transferred by an intramolecular reaction to the other catalytic cysteine located on the RHD of UBA4 to form UBA4~URM1 acyl persulfide intermediate. The formation of the persulfide bond is enabled by the presence of a sulfur donor [S], and the activated URM1 thiocarboxylate species (highlighted in a green box) is released by a reductive cleavage step for subsequent use in sulfur transfer reactions, such as tRNA thiolation. The E2 and E3 enzymes involved in URMylation are unknown.

with context-dependent opposite functions (Hermann and Bogunovic, 2017).

There is substantial evidence of cross talk between the ISG15 and ubiquitin conjugation systems. In this respect, the two ubiquitin E2 enzymes, UBCH6 and UBCH8, have been reported to serve as E2 enzymes that transfer ISG15 from UBA7 to cellular substrates (Kim et al., 2004; Zhao et al., 2004; Takeuchi et al., 2005). A model has also been proposed wherein proteasomal degradation influences the substrate pool available for ISGylation (Liu et al., 2003). In addition, increased expression of ISG15 in tumor cells has been found to antagonize ubiquitylation and subsequent proteasomal degradation of a subset of cellular proteins (Desai et al., 2006). Interestingly,

ISGylation has been reported to act as a signal for proteasomal degradation of misfolded dominant-negative forms of p53 independent of ubiquitylation-regulated degradation (Huang et al., 2014).

1. Role of Ubiquitin-Like Modifier-Activating Enzyme 7 in Cancer. Reduced expression of UBA7 has been reported in lung cancer cell lines compared with normal cells, suggesting that it may possess tumor-suppressive functions in a subset of lung cancers (McLaughlin et al., 2000; Yin et al., 2009). This is corroborated by the observation that frequent genomic lesions affect the chromosomal region 3p21.3, where the *UBA7* gene is located (Hesson et al., 2007). Mechanistic studies of this tumor-suppressive activity have implicated cyclin D1 as a target that is repressed after

ISGylation by UBA7, resulting in antitumor effects (Pitha-Rowe et al., 2004b; Feng et al., 2008). Similar tumor-suppressive activity has been reported in APL, in which UBA7 acts as a retinoid-inducible target that causes PML-RARA degradation and apoptosis (Kitareewan et al., 2002; Pitha-Rowe et al., 2004a). In addition, UBA7 and other components of the ISGylation system exert tumor-suppressive functions in cervical, hepatic, and renal cell cancer cells (Jeon et al., 2012; Yeh et al., 2013; Zhou et al., 2017). In contrast, the upregulation of the ISGylation system has been associated with oncogenic functions in models of breast, pancreatic, and prostate cancer (Kiessling et al., 2009; Burks et al., 2014; Sainz et al., 2014). These contrasting reports support the context-dependent oncogenic and tumor-suppressive functions of UBA7 and ISG15 in cancer (Desai, 2015; Zuo et al., 2016). Consistent with these observations, the analysis of patient survival data has correlated UBA7 expression to unfavorable outcomes in renal cancer and favorable outcomes in urothelial and breast cancer (Uhlen et al., 2017). Of note, the ISG15-specific deconjugase USP18 has also been shown to exert opposite functions in several malignancies (Mustachio et al., 2018).

2. Ubiquitin-Like Modifier-Activating Enzyme 7 Inhibitors. The crystal structure of UBA7 has not been elucidated thus far. However, a docking model of ISG15 onto the published structure of the NAE-NEDD8 complex allowed the prediction of several specific interactions, including residues critical for the binding between the C-terminal domain of ISG15 and the AAD of UBA7 (Narasimhan et al., 2005). With the exception of the pan-E1 inhibitor compound 1, there are no inhibitors reported to target UBA7 so far. As assessed by ATP-PPi exchange assays, compound 1 inhibits UBA7 with a potency that is comparable to UBA1 and SAE but less than NAE and UBA6 (Chen et al., 2011). Even after the development of clinical UBA7 inhibitors/activators, it is reasonable to expect that the use of such agents will be determined by the context-dependent roles of UBA7 and other ISGylation components in different cancers. A model has been proposed in which secreted ISG15 stimulates antitumor immune responses; thus, UBA1-initiated ISGylation may reduce the pool of secreted ISG15 and constitute a mechanism by which tumor cells evade immune surveillance (Desai, 2015). In these contexts, UBA7 inhibitors are anticipated to trigger useful antitumor immune responses.

V. Noncanonical E1 Enzymes

A. Autophagy-Related Protein 7

ATG7 is a noncanonical E1 enzyme that plays a key role in the regulation of autophagy, a process by which cellular proteins and organelles are targeted to the lysosome to be degraded by the lysosomal enzymes (Schulman and Harper, 2009; Choi et al., 2013; Xiong,

2015; Levine and Kroemer, 2019). The UPS and the autophagy-lysosome system are the two major cellular machineries involved in protein degradation and are highly coordinated by ubiquitylation and other UBL modifications (Klaips et al., 2018). ATG7 is essential for the autophagic cell death program, a role that is regulated by signaling molecules involved in apoptosis, such as caspases 8 and 9 (Yu et al., 2004; Han et al., 2014). Moreover, ATG7 is implicated in several autophagy-independent functions, including cellular proliferation, cytokine and exosome secretion, and regulation of immune responses (Galluzzi and Green, 2019). Of note, ATG7 has been reported to bind p53 to regulate cell cycle progression and survival under conditions of metabolic stress, a function that appears to be independent of its E1 enzymatic activity (Lee et al., 2012).

In autophagy, ATG7 lies at the center of two UBL conjugation systems, where it activates two classes of UBLs, ATG12 and ATG8 family proteins, initiating two pathways that are important for the formation of autophagosomes, which are vesicular structures that sequester and transport target proteins to the lysosomes (Nakatogawa, 2013) (Fig. 11A). In humans, ATG8 family proteins (simply referred to as ATG8) comprise two subfamilies: microtubule-associated protein 1 light chain 3 proteins and γ -aminobutyric acid receptor-associated proteins (Abdollahzadeh et al., 2017). In the ATG12 conjugation pathway, ATG7 activates ATG12 and transfers it to the E2-like enzyme ATG10, which in turn conjugates ATG12 to ATG5. The ATG12-ATG5 conjugate forms a complex with ATG16, and the complex serves as an E3-like enzyme that plays a key role in ATG8 conjugation. On the other hand, the ATG8 conjugation pathway requires prior processing of the precursor form of ATG8 into the functional form through proteolytic cleavage mediated by the protease ATG4. After activation by ATG7, ATG4 is transferred to the E2-like enzyme ATG3, which in turn conjugates this UBL to phosphatidylethanolamine (PE) in a reaction facilitated by the E3-like enzyme complex ATG12-ATG5-ATG16, which is formed by the ATG12 conjugation pathway (Geng and Klionsky, 2008; Nakatogawa, 2013). The resultant ATG8-PE conjugate plays an important role in autophagosome formation and serves as a receptor for the selective recognition of cargos during autophagy (Suzuki et al., 2017). In addition to its role in the proteolytic cleavage of pro-ATG8, ATG4 can also serve as a deconjugase for the ATG8-PE conjugate (Maruyama and Noda, 2018). Of note, the ATG12 conjugation pathway has been implicated in the regulation of mitochondrial homeostasis and cell death, a function that appears to be independent of its role in autophagy (Radoshevich et al., 2010).

1. Role of Autophagy-Related Protein 7 in Cancer. Similar to other genes encoding E1 enzymes, ATG7 is affected by a number of mutations and CNVs; however,

they do not appear to play a significant role in cancer development and/or progression (Grossman et al., 2016; Tate et al., 2019). As assessed by the analysis of survival data in a subset of cancer patients, high expression of ATG7 is associated with favorable prognosis in renal and colon cancer and unfavorable prognosis in liver cancer (Luo et al., 2016; Uhlen et al., 2017). ATG7 has been implicated in both oncogenic and tumor-suppressive functions in different malignancies. This derives partly from the role of ATG7 in autophagy, which has dual functions in tumor development and progression in a context-dependent manner (Wen and Klionsky, 2019). In this respect, ATG7 has been reported to promote the tumorigenesis of several malignancies, including prostate cancer (Santanam et al., 2016), chronic myeloid leukemia (Karvela et al., 2016), skin cancer (Xie et al., 2015; Qiang et al., 2017), and bladder cancer (Zhu et al., 2019b), and this protumorigenic role occurs mostly in cooperation with other genetic aberrations, such as *phosphatase and tensin homolog (PTEN)* loss, *Adenomatous polyposis coli gene (APC)* inactivation, *BCR-ABL* fusion, as well as *BRAF* and Kirsten rat sarcoma viral oncogene homolog (*KRAS*) mutations (Guo et al., 2013b). On the other hand, ATG7 has been associated with contextual oncogenic and tumor-suppressive functions in liver cancer (Takamura et al., 2011; Gong et al., 2012; Luo et al., 2016), colon cancer (Zhao et al., 2010; Lévy et al., 2015), and lung cancer (Zhao et al., 2010; Guo and White, 2013; Guo et al., 2013a; Strohecker et al., 2013; Strohecker and White, 2014; Han et al., 2018; Cao et al., 2020). Given the reported role of ATG7 in the maintenance of hematopoietic stem and progenitor cells, genetic inactivation of ATG7 has been found to cause dysregulated myeloid proliferation, suggesting a tumor-suppressive role in the early stages of MDS and AML development (Mortensen et al., 2011a,b; Watson et al., 2011a; Gomez-Puerto et al., 2016). In contrast, ATG7 is required to support tumor progression and resistance to chemotherapy in established AML neoplasms (Piya et al., 2016). Therapeutic targeting of ATG7 has been reported to sensitize several malignancies to cytotoxic chemotherapy and targeted agents, including tyrosine kinase inhibitors, *BRAF* inhibitors, and monoclonal antibodies (Li and Fan, 2010; Han et al., 2011; Xie et al., 2015; Karvela et al., 2016; Piya et al., 2016).

2. Autophagy-Related Protein 7 Inhibitors. The crystal structure of human ATG7 has not been solved to date; however, significant structural and biochemical information has been obtained from Atg7 of other organisms, particularly yeast (Hong et al., 2011; Noda et al., 2011; Taherbhoy et al., 2011; Kaiser et al., 2012; Yamaguchi et al., 2012, 2018). Based on these studies, ATG7 is a homodimeric enzyme with each subunit composed of two domains that are connected by a short flexible linker: 1) N-terminal domain (NTD) and 2) C-terminal domain (CTD). As a homodimer,

ATG7 possesses two nucleotide-binding sites and two catalytic cysteine residues (Noda et al., 2009). Moreover, it assumes a bird-like three-dimensional structure in which the CTD dimer occupies a central position corresponding to the body, and the two NTDs connected to the CTD dimer correspond to the wings (Kaiser et al., 2013). The NTD is structurally unique and does not show significant sequence homology with other E1 enzymes. The CTD is composed of two subdomains: a homodimeric AD that is conserved among all E1s and an extreme CTD (ECTD) that is specific to ATG7 (Noda et al., 2011). The ECTD subdomain comprises a C-terminal tail that identifies ATG7-specific UBLs and subsequently transfers them to the AD subdomain that harbors the adenylation active site and the catalytic cysteine that are important for the UBL-activating function of the enzyme. The NTD is involved in the interaction of ATG7 with E2 enzymes for UBL transfer (Noda et al., 2011; Taherbhoy et al., 2011). There are several unique aspects of the noncanonical mechanisms of ATG7 catalytic activity, including the ability to activate UBLs without undergoing significant active site remodeling (Hong et al., 2011). Given the unique architecture and catalytic mechanism of ATG7, it is not surprising that the pan-E1 inhibitor compound 1 does not display significant activity against this enzyme (Brownell et al., 2010). Apart from genetic ablation approaches, no selective chemical inhibitors have been reported to target ATG7 thus far. It is noteworthy that the utility of ATG7 inhibitors in cancer therapy depends on the specific malignancy and the context (e.g., tumor stage) in which they are used. For example, in contexts in which autophagy is perceived as tumor-suppressive, mainly through induction of autophagic cell death, inhibiting ATG7 is anticipated to cause adverse therapeutic outcomes and vice versa. Moreover, the cytoprotective effects of ATG7-mediated autophagy in normal cells should be considered to minimize the untoward effects of ATG7 inhibitors.

B. Ubiquitin-Like Modifier-Activating Enzyme 4

UBA4 [also known as molybdenum cofactor synthesis 3 (MOCS3)] is a noncanonical E1 enzyme that activates URM1, a UBL that acts as a post-translational modifier conjugated to other proteins (by a process termed URMylation) and a sulfur carrier in the thiolation of cytosolic tRNAs (Schlieker et al., 2008; Leidel et al., 2009; Wang et al., 2011). These two functions of URM1 are highly conserved among eukaryotes (Jüdes et al., 2015). Evolutionarily, URM1 is the most ancient UBL and is closely related—in the structure and mechanism of activation—to the prokaryotic sulfur carrier proteins molybdopterin synthase sulfur carrier subunit (MoaD) and thiamine biosynthesis protein S (ThiS), which play a role in molybdopterin and thiamine biosynthesis, respectively (Furukawa et al., 2000; Xu et al., 2006; Pedrioli et al., 2008; Petroski et al., 2011). Therefore,

TABLE 9
UBA5 inhibitors and their pharmacologic properties

Inhibitor	Type	Malignancies	Class	EC ₅₀	Reference
Compound 8.5	Experimental	Cell-free and cellular activity in UBA5-high cancer cells	Adenosine-based organometallic inhibitor	Cell-free transthiolation assays: 4 μ M	da Silva et al., 2016
DKM 2–93	Experimental	Antitumor activity in pancreatic and glioblastoma cancer cells in vitro and in vivo	Chloroacetamide-based inhibitor	Cell-free thioester formation assays: 430 μ M; dose in mice: 50 mg/kg	Roberts et al., 2017; MacLeod et al., 2019

URM1 is regarded as the evolutionary link between prokaryotic sulfur carriers and UBL eukaryotic modifiers (Iyer et al., 2006; Schlieker et al., 2008; Schmitz et al., 2008). The repertoire of human substrates that are modified by URMylation include proteins implicated in the ubiquitylation pathway (e.g., USP15 and 47), nuclear transport (e.g., RanGAP1), and oxidative stress [e.g., carbonyl reductase (NADPH) 1]; however, the biologic roles of URM1 modifications of these substrates are not well understood (Van der Veen et al., 2011). Nonetheless, few biologic functions have been linked to protein URMylation and tRNA thiolation. For example, genetic ablation of URM1 has been reported to induce cytokinesis defects, suggesting a role of URMylation in cell cycle progression (Schlieker et al., 2008). Moreover, tRNA thiolation plays a regulatory role in translation, metabolic homeostasis, and control of cellular growth (Laxman et al., 2013). Of note, URMylation and tRNA thiolation are both stimulated in response to oxidative stress, suggesting a role for these URM1 modifications in sensing and/or protecting against oxidative damage (Leidel et al., 2009; Van der Veen et al., 2011).

1. Role of Ubiquitin-Like Modifier-Activating Enzyme 4 in Cancer. The expression of UBA4 has been associated with unfavorable prognosis in liver and endometrial cancers and a favorable prognosis in renal cancer (Uhlen et al., 2017). Apart from these data, there are no published reports that directly implicate UBA4 in cancer development and/or progression. However, it has been reported that several enzymes involved in the tRNA modification pathway are upregulated in breast cancer and are important in supporting tumor metastasis, suggesting UBA4 can potentially serve as a therapeutic target in this context (Delaunay et al., 2016; Termathe and Leidel, 2018).

2. Ubiquitin-Like Modifier-Activating Enzyme 4 Inhibitors. As is the case with other noncanonical E1 enzymes, UBA4 is a homodimeric enzyme composed of two symmetric subunits (Schulman and Harper, 2009). Each of these subunits comprises E1-like domains at its N terminus that harbor the active adenylation and catalytic cysteine sites (Schulman and Harper, 2009; Termathe and Leidel, 2018). Different from all other E1 enzymes, UBA4 possesses a characteristic RHD at its C terminus that also harbors a catalytic cysteine and plays an important catalytic role in URM1 activation (<http://www.rcsb.org/structure/3I2V>; Schulman and

Harper, 2009). These structural features enable a non-canonical mechanism of UBL activation that is unique to UBA4 and allow for the dual function of URM1 as a UBL and a sulfur carrier (Wang et al., 2011). As with other UBLs, URM1 undergoes initial adenylation and thioester formation steps to form a UBA4~URM1 thioester intermediate in which URM1 is linked to the catalytic cysteine located on the E1-like domain of UBA4. However, an intramolecular reaction follows whereby URM1 is transferred to the catalytic cysteine located on the RHD, forming a UBA4~URM1 acyl persulfide intermediate. This reaction requires the presence of a sulfur donor, such as cysteine, to form the persulfide linkage (Ternatthe and Leidel, 2018). The final step involves reductive cleavage and release of URM1 thiocarboxylate that serves as the activated species for subsequent sulfur transfer reactions (Fig. 11B). To date, there are no known E2 or E3 enzymes that catalyze URMylation (Pedrioli et al., 2008). UBA4 similarly activates molybdenum cofactor synthesis protein 2A (MOCS2A), a polypeptide structurally related to UBLs, and forms a MOCS2A thiocarboxylate that serves as a sulfur donor in molybdenum cofactor biosynthesis (Chowdhury et al., 2012). No inhibitors have been reported to target UBA4 thus far; however, the possession of two catalytic cysteines and the unique catalytic mechanism of this enzyme may create valuable opportunities to develop inhibitors that selectively target UBA4 at the expense of related E1 enzymes.

C. Ubiquitin-Like Modifier-Activating Enzyme 5

UBA5 (also known as ubiquitin-activating enzyme E1-domain containing 1) is a noncanonical E1 enzyme that catalyzes the activation of UFM1, a metazoan-specific UBL that is conjugated to cellular substrates—by a process termed UFMylation—to modulate their functions (Komatsu et al., 2004; Daniel and Liebau, 2014; Yoo et al., 2015; Wei and Xu, 2016; Gerakis et al., 2019). After activation of UFM1 by UBA5, UFMylation of target substrates is mediated by UFM1-conjugating enzyme 1 (UFC1) and UFM1-specific ligase 1 (UFL1), which serve as UFM1-specific E2 and E3 enzymes, respectively (Komatsu et al., 2004; Dou et al., 2005; Mizushima et al., 2007; Tatsumi et al., 2010). UFMylation is subject to regulation by UFSP1 and 2, UFM1-specific proteases that activate the precursor form of UFM1 and deconjugate UFMylated substrates by

TABLE 10
Nonselective E1 inhibitors and their pharmacologic properties

Inhibitor	Type	Malignancies	Class	EC ₅₀ /IC ₅₀	References
Compound 1	Experimental pan-E1 inhibitor	Cell-free and cellular activity	Adenosine sulfamate	ATP-PPi exchange assays: 0.01–6.4 μ M; UBA1/NAE/SAE-E2 cell-free transthiolation: 0.005 μ M	Brownell et al., 2010; Chen et al., 2011
ADS	Experimental pan-E1 inhibitor	Cell-free and cellular activity in colon cancer cells	Adenosine sulfamate	ATP-PPi exchange assay (UBA5): 13 μ M	Gavin et al., 2014
ABP1	Experimental pan-E1 inhibitor	Cell-free and cellular activity in lung cancer cells	Adenosine sulfamate	NA	An and Statsyuk, 2013
ABPA3	Experimental UBA1/NAE inhibitor	Cell-free and cellular activity in lung cancer cells	Adenosine sulfamate	Cytotoxicity in A549 cells: 2.5 μ M	An and Statsyuk, 2015
LP0040	Experimental UBA1/NAE inhibitor	Cell-free and cellular activity in gastric cancer cells	2H-chromen-2-one derivative	Cytotoxicity: 0.76–3.29 μ M	Lu et al., 2018

NA, not available.

proteolytic cleavage (Ha et al., 2008, 2011; Kim et al., 2018). In humans, it appears that UFSP2 is the only functional UFM1-specific protease (Yoo et al., 2014). UBA5 has also been reported to activate SUMO2 in vitro and in vivo (Zheng et al., 2008). The repertoire of cellular substrates that is modified by UFMylation comprises few known targets, including activating signal cointegrator 1/thyroid receptor-interacting protein 4 (ASC1/TRIP4), which acts as a transcriptional coactivator of nuclear receptors (e.g., estrogen receptor α); ribophorin 1 (RPN1), which serves as an ER quality control factor; and 60S ribosomal protein L26 (RPL26), which is a ribosomal protein and is reportedly the main target of UFMylation in the cell (Lee et al., 1999; Yoo et al., 2014, 2015; Walczak et al., 2019; Liang et al., 2020). Nonetheless, UFMylation is functionally important in regulating several cellular functions, including DNA damage response (Qin et al., 2019; Wang et al., 2019), transcriptional regulation (Yoo et al., 2014), cell cycle control (Yu et al., 2020), autophagy (DeJesus et al., 2016), and ribosomal functions (Simsek et al., 2017; Walczak et al., 2019). Of particular importance are the roles played by UFMylation in regulating hematopoiesis and ER-associated proteostasis (Gerakis et al., 2019). In this respect, UBA5 and other components of the UFMylation cascade have been reported to be essential for hematopoietic stem cell survival, plasma cell development, and erythroid differentiation, as evidenced by the development of anemic phenotype in mice upon genetic ablation of these components (Tatsumi et al., 2011; Cai et al., 2015, 2016; Zhang et al., 2015a; Zhu et al., 2019a). Moreover, UFMylation plays a key regulatory role in ER homeostasis; thus, the disruption of this pathway is associated with the induction of ER stress (DeJesus et al., 2016; Liu et al., 2017a; Walczak et al., 2019; Wang et al., 2020). UFMylation of ER-resident proteins has been implicated in the autophagy of ER (also known as ER-phagy) and regulation of the UPR (Liang et al., 2020). Of note, a subset of UFMylation system components are subject to transcriptional changes in response to ER stress,

establishing a regulatory feedback loop that coordinates UFMylation and ER homeostasis (Zhang et al., 2012).

1. Role of Ubiquitin-Like Modifier-Activating Enzyme 5 in Cancer. The analysis of publicly available datasets has shown that UBA5 is affected by genetic aberrations, particularly amplifications, in several malignancies, including lymphoma, lung, ovarian, and esophageal cancers (Wei and Xu, 2016). As assessed by chemical and genetic approaches, inhibition of UBA5 has revealed that pancreatic and breast cancer cells are dependent on UFMylation for survival and growth in vitro and in vivo (Yoo et al., 2014; Roberts et al., 2017). Besides, a genome-wide CRISPR/Cas9 knockout screen in patient-derived glioblastoma stem cells identified UBA5, among other UFMylation pathway components, as a regulator of glioblastoma stem cell survival (MacLeod et al., 2019). However, in pre-clinical models of lung cancer, inhibition of UFMylation pathway components including UBA5 has been found to induce a prosurvival UPR that promotes resistance to EGFR inhibition by tyrosine kinase inhibitors (Terai et al., 2018). These reports suggest that UBA5 can serve as a potential therapeutic target in glioblastoma, pancreatic, and breast cancer; however, in the context of targeted therapy in lung cancer, UBA5 inhibition may result in adverse therapeutic outcomes by fostering adaptive resistance signaling.

2. Ubiquitin-Like Modifier-Activating Enzyme 5 Inhibitors. As a noncanonical E1 enzyme, UBA5 is predicted to possess a homodimeric organization with two symmetric subunits (Schulman and Harper, 2009; Cappadocia and Lima, 2018). However, compared with other E1s, each of these subunits displays minimalistic structural features that are shared between the more complex canonical E1s and the simpler, ancestral E1-like enzymes. Specifically, UBA5 comprises an AD that harbors both the nucleotide-binding site as well as the catalytic cysteine residue (Cys250), with no dedicated domain for this catalytic cysteine. It also comprises a small UFD-like region at the C terminus—termed UFC1-binding sequence—that facilitates the interaction

Fig. 12. Structural features of adenosine sulfamates. The chemical structures of TAK-243, pevonedistat, ML-792, and TAK-981 are shown in comparison with AMP. Adenosine sulfamates share common chemical features—mainly 1) sulfamate group, 2) cyclopentane/ribose moiety, and 3) nucleobase extension.

with UFC1 and subsequent transfer of activated UFM1 (Bacik et al., 2010; Oweis et al., 2016). UBA5 also possesses characteristic structural motifs, including a UFM1-interacting sequence (UIS) involved in UFM1 interaction, and an N-terminal extension that exists in one of the two known UBA5 isoforms and enhances UFM1 activation (Padala et al., 2017; Soudah et al., 2019). In addition to UFM1, it appears that UBA5 can also interact with γ -aminobutyric acid receptor-associated proteins, and this interaction is required to facilitate the recruitment of UBA5 to the ER membrane (Habisov et al., 2016; Huber et al., 2020).

As opposed to canonical E1s that activate their cognate UBLs via a three-step mechanism, UBA5 activates UFM1 in two steps—adenylation and thioester bond formation—with no second adenylation step, forming a binary rather than a ternary complex from which UFM1 is transferred to UFC1. Besides, UBA5 displays

a slower catalytic activity compared with canonical E1s (Gavin et al., 2014). Of note, UBA5 activates UFM1 by a *trans*-binding mechanism whereby UFM1 interacts with the UIS of one subunit and undergoes adenylation by the AD of the other subunit of the UBA5 homodimer, suggesting that homodimerization is essential for UFM1 activation (Oweis et al., 2016; Mashahreh et al., 2018).

As with other E1 enzymes, UBA5 can be targeted either at the nucleotide-binding site or the catalytic cysteine, taking into consideration the UBA5-specific structural features to maximize the selectivity of UBA5 inhibitors. To date, only two experimental UBA5 inhibitors have been reported: compound 8.5 and DKM 2–93 (Fig. 10; Table 9).

Compound 8.5 is an adenosine-based organometallic UBA5 inhibitor that was identified by a focused screen of compounds that incorporate adenosine and zinc(II) cyclen coordination complex. As assessed by structural and biochemical analyses, compound 8.5 is predicted to target the nucleotide-binding pocket, and with respect to ATP, it serves as a noncompetitive inhibitor that displays selectivity for UBA5 over other E1 enzymes and kinases. Nonetheless, high concentrations are required to induce cytotoxicity in *UBA5*-overexpressing cancer cell lines, suggesting that further optimization is required to enhance the potency of this inhibitor (da Silva et al., 2016).

DKM 2–93 is a UBA5 inhibitor identified through a screen for cysteine-reactive covalent compounds that inhibit pancreatic cancer cell proliferation and subsequent target identification by activity-based protein profiling. DKM 2–93 covalently inhibits UBA5 by targeting the catalytic cysteine, preventing UFM1 charging of UBA5, and resulting in antitumor effects in pancreatic and glioblastoma cancer cells in vitro and in vivo (Roberts et al., 2017; MacLeod et al., 2019). With the high concentrations required to induce cytotoxicity, it is anticipated that DKM 2–93 may have off-target effects; however, it can potentially serve as a useful chemotype for covalent UBA5 inhibitors with higher potency and selectivity (Roberts et al., 2017).

Adenosine 5'-sulfamate (ADS), a pan-E1 inhibitor, has been reported to inhibit UBA5 by targeting the UBA5~UFM1 thioester intermediate and forming an ADS-UFM1 covalent adduct that inhibits UBA5 at the nucleotide-binding site (Table 10). ADS has also displayed cell-based activity in colon cancer cells, as it inhibited UBA5~UFM1 thioester formation with a subsequent reduction in the abundance of UFMylated proteins (Gavin et al., 2014).

VI. Dual and Multi-E1 Inhibitors

E1 inhibitors discussed so far have variable selectivity profiles toward E1 enzymes or other targets of the proteome. Of these, most adenosine sulfamates, particularly those in clinical trials, TAS4464 and COH000

adenine base. Compound 1 is a pan-E1 inhibitor that has been exploited to elucidate important mechanistic aspects of substrate-assisted inhibition (Brownell et al., 2010). While having variable potencies against E1 enzymes in ATP-PPi exchange assays, with the highest potency observed with NAE, compound 1 is equipotent against UBA1, NAE, and SAE in E1-E2 transthioylation assays (Brownell et al., 2010; Chen et al., 2011). Of note, compound 1 displays no appreciable activity against ATG7 in ATP-PPi exchange assays. Replacing the indane moiety of compound 1 with a propargyl moiety generates another nonselective E1 inhibitor, ABP1, with a free alkyne group that has been leveraged for covalent conjugation to fluorescent or biotin tags using click chemistry (An and Statsyuk, 2013). This advantage has enabled the use of ABP1 as a probe to quantify the activity of E1 enzymes, as the adducts are formed by the enzymes themselves and thus the rate of adduct formation is dependent on E1 activity (see below). In addition, it has been used to provide further insights into the biochemistry of UBL conjugation and evaluate

the potency and selectivity of investigational E1 inhibitors (An and Statsyuk, 2013). Experience with ABP1 has been exploited to develop a related analog, ABPA3, which acts as a dual UBA1/NAE inhibitor. ABPA3 is also structurally related to compound 1, with phenylacetylene replacing the indane moiety (An and Statsyuk, 2015). ABPA3 displays anti-UBA1/NAE activity in lung cancer cells, which results in the induction of UPR and prevents cytoprotective aggresome formation leading to cell death. Of note, ABPA3 paradoxically causes increases in SUMOylation and UFMylation, which are likely compensatory responses activated in response to UPR induction (An and Statsyuk, 2015).

B. LP0040

LP0040 is a non-nucleoside dual UBA1/NAE inhibitor developed by rational drug design and molecular modeling based on the structure of the NAE inhibitor M22 (Lu et al., 2018). Replacing the benzyl group of M22 with biphenyl-substituted 2H-chromen-2-one in LP0040 led to an improved antiproliferative activity associated with cell cycle arrest and apoptosis in cancer cell lines of different origin with a low micromolar IC₅₀. LP0040 has also displayed synergistic cytotoxicity with bortezomib (Lu et al., 2018).

VII. Adenosine Sulfamate E1 Inhibitors

This class comprises pevonedistat, TAK-243, ML-792, TAK-981, compound 1, ABPA3, and ABP1, which act as mechanism-based inhibitors of E1 enzymes. They are structurally related to AMP, which tightly binds to E1 enzymes during the catalytic cascade of UBL activation (Soucy et al., 2009a). The structures of adenosine sulfamates share three major chemical features: 1) a ribose or cyclopentane moiety, 2) a sulfamate group analogous to the phosphate group in AMP, and 3) a nucleobase extension analogous to the adenine base in AMP (Misra et al., 2017) (Fig. 12).

A. Discovery and Structure-Activity Relationship

The discovery and SAR of adenosine sulfamates have been reviewed in detail; here, we briefly highlight the main features of these discovery programs (Ciavarri and Langston, 2017). An HTS identified N6-substituted adenosine derivatives as initial chemotypes with weak NAE inhibitory activity. Further chemical optimization was required to enhance potency, cellular activity, selectivity, and stability (Ciavarri and Langston, 2017) (Fig. 13). Although phosphate derivatives demonstrated sub-micromolar potency, they did not permeate into cells and thus lacked cellular activity (Ciavarri and Langston, 2017). Replacement with the neutral sulfamate group retained potency and yielded a series of cell-active E1 inhibitors. The sulfamate functional group is the closest bioisostere to phosphate and does not undergo promiscuous thioester formation in cells as

a result of its low intrinsic reactivity (An and Statsyuk, 2015). Compared with sulfamate, sulfonamide and sulfamide groups are less favorable in many aspects, including cellular activity, potency, and selectivity; therefore, the sulfamate group remained as a common chemical feature of adenosine sulfamates (Ciavarri and Langston, 2017). In this series, certain compounds were prone to an intramolecular cyclization reaction, resulting in the loss of activity. In addition, several compounds lacked selectivity for individual E1 enzymes. A combination of chemical modifications in the ribose sugar, purine base, and stereochemical orientation of the sulfamate group solved these stability and selectivity issues and retained or even improved potency and cellular activity, leading to pevonedistat as the prototype of this class (Soucy et al., 2009a; Ciavarri and Langston, 2017).

The success of the NAE inhibitor program inspired the initiation of another program to develop UBA1 inhibitors by exploring the adenosine sulfamate series and building on the previous SAR and mechanistic studies done with pevonedistat (Brownell et al., 2010). Although chemical modifications of the ribose sugar and aromatic side chain led to improved selectivity to UBA1, the pharmacodynamic profile in vivo was not optimal. The replacement of a purine base with a pyrazolo-pyrimidine scaffold and the placement of the N9-nitrogen in an *exo* position improved the in vivo profile and significantly enhanced selectivity over the SAE; however, the selectivity over the NAE was only partial (Ciavarri and Langston, 2017). Further modifications enhanced selectivity to UBA1, however, at the expense of potency, which was significantly compromised. As NAE regulates the activity of only a subset of ubiquitin E3 ligases downstream of UBA1, it is anticipated that a dual UBA1/NAE inhibitor will demonstrate the same biologic activity and phenotype as a selective UBA1 inhibitor (Ciavarri and Langston, 2017b). Therefore, chemical optimization was pursued with priority given to potency over selectivity, leading ultimately to the selection of TAK-243 as a clinical candidate.

B. Substrate-Assisted Inhibition

The development of pevonedistat involved an extensive characterization of its mode of action using various approaches including crystallographic studies to elucidate the structural basis of NAE inhibition by this drug (Brownell et al., 2010). These studies showcased a continuous electron density between the C terminus of NEDD8 and the sulfamate group of pevonedistat when bound to NAE, suggesting that a pevonedistat-NEDD8 adduct is formed and possibly it is the species that inhibits NAE (Brownell et al., 2010). Given that the UBL activation mechanism is highly conserved among canonical E1 enzymes, this finding has stimulated further studies that revealed this mode of inhibition—termed substrate-assisted inhibition—is a general

mechanism shared by other adenosine sulfamates, including those targeting UBA1 and SAE (Chen et al., 2011; He et al., 2017; Hyer et al., 2018).

Substrate-assisted inhibition is unique in that the UBL substrate of E1 enzymes is involved in the enzyme inhibition through a chemical reaction with adenosine sulfamates to form a covalent adduct that itself serves as the active inhibitory species (Brownell et al., 2010). This mode of inhibition may be pharmacologically beneficial, as it confers selectivity among E1 enzymes because most E1s activate only one or closely related UBLs. Besides, it reduces the likelihood of interaction with other ATP-dependent enzymes, including kinases, which have different catalytic mechanisms and thus are likely less susceptible to these UBL-based inhibitory species (Brownell et al., 2010).

To form these adducts, adenosine sulfamates act as AMP mimetics and exploit the multistep catalytic mechanism of UBL activation by E1 enzymes (detailed in section II. B. *Biochemical and Structural Mechanisms of E1-Catalyzed Ubiquitin-Like Protein Activation*). They specifically target the E1~UBL thioester intermediate, a binary complex in which E1 is loaded with a single UBL molecule at the catalytic cysteine, and bind to the nucleotide-binding site in the AAD. This is followed by a nucleophilic attack of the thioester bond by the sulfamate amino group of adenosine sulfamates to form a covalent adduct with UBL (Fig. 3). Consistent with a mechanism-based inhibitory mode, the adduct formation is catalyzed by the E1 enzyme itself and requires the presence of magnesium, ATP, and an active catalytic cysteine, mimicking the conditions of AMP~UBL formation and implicating most sites of interaction with the E1 enzyme (Brownell et al., 2010; Chen et al., 2011). Although E1~UBL intermediate formation is the rate-limiting step in UBL activation under physiologic conditions, biochemical studies suggest adduct formation is the rate-limiting step in E1 inhibition by adenosine sulfamates (Chen et al., 2011). As opposed to the binary complex, the ternary complex loaded with two ubiquitin molecules is not susceptible to inhibition by adenosine sulfamates, as the nucleotide-binding site, to which these drugs need to bind first, is occupied by AMP~UBL. After formation, adenosine sulfamate adducts are stable and tightly bind—with a picomolar affinity—to E1 enzymes in an ATP-competitive manner. By competing for E1 binding, high ATP concentrations are thus anticipated to slow the binding rate of adenosine sulfamates with a resultant increase in their IC_{50} values (Chen et al., 2011).

Of note, the inhibitory activity of stable AMP~UBL mimetics against E1 enzymes has been known several years before the development of adenosine sulfamates as with APU, and later with UBL-AMSN/AVSN which are preformed semisynthetic UBL phospho- and sulfamoyl-adenosine derivatives (Wilkinson et al., 1990; Olsen et al., 2010). Adenosine sulfamates, however, differ

in that they form adducts in situ after binding to their cognate E1 enzymes, allowing for a preferential activity against cancer cells that display higher activity of these enzymes to support their cellular stresses (Luo et al., 2009; Dobbstein and Moll, 2014). It is also noteworthy that—in cell-free assays—preformed adenosine sulfamate adducts can inhibit E1 enzymes with much higher potency compared with unbound adenosine sulfamates (Chen et al., 2011). This may be ascribed to their different inhibition kinetics, as the adducts rapidly bind to and inhibit unloaded E1 enzymes, halting all subsequent steps, whereas unbound adenosine sulfamates bind at a slower rate to the E1~UBL thioester intermediate with no effect on the first step of UBL activation (Chen et al., 2011).

C. Selectivity and Potency of Adenosine Sulfamate E1 Inhibitors

Biochemical studies suggest that the potency of adenosine sulfamates is a function of the rate of adduct formation and the affinity of this adduct to E1 enzymes, and these biochemical parameters in turn depend on their chemical structure (Chen et al., 2011). Although adenosine sulfamates share general chemical features, they still have several variations that dictate—among other properties—potency and selectivity to E1 enzymes. These include chemically diverse nucleobase extensions, different stereochemical orientation of the sulfamate group, as well as varying numbers and stereochemical orientations of the hydroxyl groups on the cyclopentane/ribose moiety (Misra et al., 2017). For instance, TAK-243 possesses a characteristic thio(trifluoromethyl) group in its nucleobase extension, which is absent in other adenosine sulfamates (Fig. 11). This group serves as a hook that engages TAK-243 in several tight contacts with several amino acid residues within a hydrophobic pocket in the UBA1 enzyme, contributing to the drug selectivity compared with related analogs (Misra et al., 2017). On the other hand, the NAE has an extended hydrophobic pocket, and the presence of a hydrophobic indane cycle in the structure of pevonedistat partly contributes to its selectivity through hydrophobic interactions with the extended pocket of the enzyme (Misra et al., 2017). As they exhibit a wider selectivity profile, dual- or pan-specific E1 inhibitors such as ABPA3 and compound 1 retain the adenine base and the ribose sugar with its two hydroxyl groups that are present in the natural substrate, AMP, thereby increasing their capacity to interact with multiple E1 enzymes with a comparable affinity (Misra et al., 2017). Although the interactions with AMP are generally conserved among different E1 enzymes, there still exist some variations that are reflected in their interactions with panspecific E1 inhibitors with different nucleobase extensions, leading to narrower/wider selectivity profile. For example, ABP1 possesses a small propargyl group linked to the exocyclic amino group of the adenine

TABLE 11
Pharmacokinetic parameters of pevonedistat after 1 h intravenous infusion

Dose (mg/m ²)	C _{max} (ng/ml) ^a	T _{max} (h) ^b	AUC _{0–24 h} (ng.h/ml) ^a	AUC _{0,t} (ng.h/ml) ^a	t _{1/2} (h) ^a	CL _p (l/h) ^a	V _{ss} (l) ^a	Reference	
Monotherapy									
25 (<i>n</i> = 2)	222	1.07	1034	—	—	—	—	Swords et al., 2015	
33 (<i>n</i> = 4)	374	1.08 (1.1–1.2)	1419	—	—	—	—		
44 (<i>n</i> = 3)	431	1.03 (1.0–1.1)	1916	1829 (48 h)	—	—	—		
59 (<i>n</i> = 6)	648	1.08 (0.9–1.6)	2506	2913 (48 h)	—	—	—		
78 (<i>n</i> = 4)	1173	1.04 (1.0–1.2)	3857	3790 (48 h)	—	—	—		
83 (<i>n</i> = 17)	1212	1.05 (0.9–1.3)	3869	—	~12 (<i>n</i> = 1)	—	—	Sarantopoulos et al., 2016	
50 (<i>n</i> = 5)	575	1.08 (1.0–1.1)	2180	—		—	—		—
67 (<i>n</i> = 13)	873	1.08 (1.0–1.2)	3383	3529 (48 h)		—	—		—
110 (<i>n</i> = 13)	1502	1.02 (1.0–1.3)	4685	4685 (48 h)	—	—	—		Shah et al., 2016
196 (<i>n</i> = 7)	4565	1.10 (1.0–1.1)	10,830	12,367 (72 h)	8.5 (<i>n</i> = 5)	28.3	158		
157 (<i>n</i> = 10)	2452	1.00 (0.92–1.7)	7248	8932 (∞)	10.1 (<i>n</i> = 5)	38.8	245	Bhatia et al., 2016	
209 (<i>n</i> = 11)	3591	1.00 (0.98–1.2)	10,716	12,300 (∞)	10.4 (<i>n</i> = 1)	38.1	237		
Combination therapy with azacytidine									
20 (<i>n</i> = 6)	158	1.06 (0.97–2.27)	990	1110 (48 h)	7.80	—	—	Swords et al., 2018	
30 (<i>n</i> = 3)	299	0.98 (0.97–1.00)	1640	1770 (48 h)	7.39	—	—		
20 (<i>n</i> = 26)	155	1.01 (0.65–2.03)	861	976 (48 h)	7.45	—	—		
20 (<i>n</i> = 28)	152	1.00 (0.88–3.00)	890	1000 (48 h)	7.30	—	—		

AUC, area under the concentration-time curve from time zero to 24 h (AUC_{0–24 h}), or to the end of the dosing interval (AUC_{0–t}), or extrapolated to infinity (∞); CL_p, plasma clearance; T_{max}, time to reach C_{max}; V_{ss}, steady-state volume of distribution.

^aData represent geometric means unless otherwise specified.

^bData represent the median.

as opposed to the larger phenylacetylene in the related adenosine sulfamate ABPA3. With its small size, this propargyl can fit into the SAE pocket leading to an inhibitory activity against SAE, which is lacking with ABPA3 (Misra et al., 2017).

D. Resistance to Adenosine Sulfamate E1 Inhibitors

The emergence of acquired anticancer drug resistance is a common problem, particularly with molecularly targeted agents, including drugs targeting the ATP-dependent tyrosine kinases (Holohan et al., 2013; Dobbelsstein and Moll, 2014). Mechanisms of such resistance include the activation of adaptive/secondary signaling pathways that maintain survival by compensating for the inhibitory effects on the targeted pathway, overexpression of multidrug resistance proteins that reduce the intracellular concentrations of the drug by extracellular efflux, and the acquisition of on-target mutations that interfere with binding to or preclude inhibition of the intracellular target (Ellis and Hicklin, 2009). Although the first mechanism is less likely to develop with adenosine sulfamates, especially those targeting UBA1—given their pleiotropic effects, the other two mechanisms are still potential contributors to resistance because of the on-target selectivity and structural similarity of adenosine sulfamates to ATP (Dobbelsstein and Moll, 2014).

Preclinical resistance studies on adenosine sulfamates involved the development of laboratory-evolved resistance models under the selective pressure of exposure to the drug in tissue culture and in vivo. Such models have led to the identification of several on-target mutations in the AADs of UBA1 and NAE that preclude the binding of TAK-243 and pevonedistat, respectively (Milhollen et al., 2012; Toth et al., 2012; Xu et al., 2014; Barghout et al., 2019).

Milhollen et al. (2012) and Toth et al. (2012) independently identified A171T missense mutation as a mechanism of resistance to pevonedistat. Milhollen et al. also identified other mutations, such as A171D, C324Y, G201V, E204K, and N209K. These mutations map to the AAD either within the nucleotide-binding pocket (A171T and A171D) or within/close to the NEDD8-binding cleft (C324Y, G201V, E204K, and N209K), reducing the binding of pevonedistat or NEDD8 to NAE, respectively (Milhollen et al., 2012). Consistent with the heterozygosity of these mutations, pevonedistat-NEDD8 adduct can still be formed by the wild-type copy of the enzyme; however, the mutant form displays a dominant effect, resulting in a resistant phenotype.

Of these mutation sites, A171 is particularly important, as it constitutes a hotspot where two-thirds of the reported mutations are located. A171 serves as a gatekeeper residue that plays a key role in controlling ATP and pevonedistat access to the nucleotide-binding site (Toth et al., 2012). In this respect, the alanine residue with its small side chain contributes to shaping an edge of the nucleotide-binding pocket without engaging in direct contact with ATP (Toth et al., 2012). Structural and computational studies suggest the replacement of alanine with threonine (A171T) or aspartic acid (A171D), which possess larger side chains, leads to conformational changes and steric clashes with the aminoindane group of pevonedistat, particularly with the A171D-mutant NAE, which is completely insensitive to pevonedistat because of the bulkier aspartic acid residue (Milhollen et al., 2012; Toth et al., 2012; Verma et al., 2014).

These mutations alter two fundamental biochemical properties required for potent NAE inhibition (Brownell et al., 2010; Chen et al., 2011). Specifically, they slow

TABLE 12
Pharmacokinetic parameters of TAK-243 after 10 min intravenous infusion^a

Dose (mg)	C _p (ng/ml) ^b	AUC _{0–48 h} (ng.h/ml) ^b	AUC _{0–∞} (ng.h/ml) ^b	t _{1/2} (h) ^b	CL _T (l/h) ^b	V _{ss} (l) ^b	Reference
1	78.7 (48.2)	32.9 (17.4)	—	—	—	—	ClinicalTrials.gov identifier: NCT02045095
2	100.8 (24.1)	52.8 (5.7)	—	—	—	—	
4	473.1 (132.8)	312.7 (193.4)	203 (10.1)	7 (1.4)	19.7 (1)	29.8 (0.9)	
8	921.8 (258.6)	368.1 (8.2)	375.5 (11.3)	9.4 (4.6)	21.3 (0.6)	34 (5.8)	
12	1276.7 (804)	867.4 (374.2)	579.1 (201.2)	5.8 (0.9)	22.7 (8.9)	37.6 (20.8)	
18	1775.5 (463)	922.1 (192.2)	768.7 (175)	8.3 (4.4)	24.3 (5.7)	63.7 (39.6)	

AUC, area under the concentration-time curve from time zero to 48 h (AUC_{0–48 h}) or extrapolated to infinity (AUC_{0–∞}); CL_T, total clearance; C_p, plasma concentration at the end of infusion; V_{ss}, steady-state volume of distribution.

^aNumber of participants = two to eight patients.

^bData represent geometric mean (S.D.).

the rate of pevonedistat-NEDD8 adduct formation and reduce the adduct affinity to the NAE, resulting in the loss of tight binding and faster recovery of enzyme activity with 2- to >2000-fold reduction in pevonedistat potency. Compound 1, which exhibits tighter binding properties to several E1 enzymes, including the NAE, could circumvent pevonedistat-resistant NAE mutations, suggesting that second-generation NAE inhibitors with higher affinities may offer a potential opportunity to overcome acquired pevonedistat resistance. Using a similar approach, Xu et al. (2014) identified two other mutations, I310N and Y352H, that decrease the affinity of NAE to NEDD8 and confer resistance of leukemia cell lines to pevonedistat.

The A171 residue of NAE is highly conserved among E1 enzymes; thus, mutation of the corresponding alanine residues in other E1s is expected to alter the response to adenosine sulfamates selective for these enzymes. In this respect, A580T mutation generated in recombinant UBA1 conferred resistance to TAK-243 in cell-free thioester formation assays (Misra et al., 2017). Consistent with these findings, Barghout et al. (2019) identified A580S mutation in a laboratory-evolved TAK-243-resistant leukemia cell line. They also identified another mutation in a close residue, Y583C. This is a TAK-243-specific mutation that was not reported with pevonedistat as TAK-243 possesses a thio(trifluoromethyl) hook through which it extends in the nucleotide-binding pocket further than pevonedistat and undergoes a favorable interaction with Y583 (Misra et al., 2017; Barghout et al., 2019). Although not highly conserved among E1 enzymes, the 583 position is mostly occupied with amino acid residues that possess bulky side chains. Therefore, replacement with cysteine is anticipated to perturb the hydrophobic core and eliminate hydrogen bonds with neighboring residues, leading to changes in the binding pocket that preclude the efficient binding of TAK-243 to UBA1 (Barghout et al., 2019).

Drug resistance is known to be multifactorial, and the resistance phenotype observed after clonal selection with adenosine sulfamates can result from a combination of several mechanisms. In this context, moderate upregulation of ATP-binding cassette transporter proteins is observed in pevonedistat-resistant cells, suggesting that they may contribute to drug resistance,

particularly at higher expression levels (Milhollen et al., 2012). Recently, pevonedistat has been reported to serve as a substrate for breast cancer resistance protein/ATP-binding cassette sub-family G member 2 (BCRP/ABCG2), and breast cancer resistance protein upregulation confers resistance to pevonedistat in ovarian, colorectal, and non-small-cell lung cancer cells (Kathawala et al., 2020; Wei et al., 2020).

Although A580T-mutant UBA1 exhibits a 6-fold resistance to TAK-243 in cell-free assays, TAK-243-resistant cell lines with A580S and Y583C mutations exhibit 28- to 33-fold resistance, suggesting that other resistance mechanisms may coexist in these cells (Misra et al., 2017; Barghout et al., 2019). Apart from these mechanisms, genetic knockout of the transcriptional regulator BEN domain-containing protein 3 is also reported to confer resistance to TAK-243 in a leukemia cell line; however, it remains unknown how it alters the response to this drug (Barghout et al., 2018).

E. Pharmacodynamic Activity of Adenosine Sulfamate E1 Inhibitors

Adenosine sulfamates targeting UBA1, NAE, and SAE are currently investigated in different phases of clinical trials in various malignancies (Tables 4 and 7). To maximize the benefit from such trials, pharmacodynamic biomarkers need to be assessed either as evidence of mechanism or to inform clinical decision-making in the trial (Schilsky et al., 2012). These biomarkers are usually pathway-specific, and in the case of adenosine sulfamates, adduct formation, E1~UBL thioesters, and/or reduction in the levels of UBL conjugates could serve as potential biomarkers (Bhatia et al., 2016; Sarantopoulos et al., 2016). For preclinical research, qualitative or semiquantitative approaches can be used to assess the pharmacodynamic effects of these drugs as a proof of principle (Soucy et al., 2009a; Hyer et al., 2018). However, for clinical trials, it is important to develop reliable pharmacodynamic assays that can assess and monitor the response to these investigational agents, preferably in a quantitative manner.

The pharmacodynamic evaluation of adenosine sulfamate activity has been performed at different levels pre- and post-E1 inhibition. For example, the cellular

thermal shift assay—a biophysical assay based on the thermostabilization of proteins after binding specific ligands, has been used to assess the binding of TAK-243 to UBA1 in cells and tumors after drug injection into mice (Martinez Molina et al., 2013; Barghout et al., 2019). As the formation of adenosine sulfamate adducts is a prerequisite for potent inhibition of E1 enzymes, antibodies raised against adducts with ubiquitin (e.g., MIL90 antibody), NEDD8 and SUMO have been used to assess drug activity in vitro and in vivo (Yan et al., 2013; He et al., 2017; Hyer et al., 2018; Berger et al., 2019). In addition, levels of E1~UBL thioesters, global protein conjugates, and conjugates of specific proteins such as ubiquitylated H2A, neddylated Cul1, and SUMOylated RanGAP1 have been used (Yan et al., 2013; He et al., 2017; Hyer et al., 2018; Barghout et al., 2019).

In such preclinical studies, semiquantitative assays such as immunoblotting and immunohistochemistry have been used as a proof of principle for E1 engagement and pathway inhibition. To achieve a more quantitative evaluation of these parameters, other platforms have been developed, such as the AlphaScreen format, a nonradioactive amplified luminescent proximity homogeneous assay that provides a faster and more robust assessment as well as higher throughput (Yan et al., 2013). Moreover, a mass spectrometry-based profiling of SDS-PAGE fractionated lysates of biologic samples has been used to provide absolute quantification of the NAE and NEDD8-pevonedistat adducts in these samples (Yang et al., 2013). In addition to gaining quantitative insights into the dynamics of UBL conjugation, the molar ratio of adduct to E1 enzyme can be calculated to provide an indirect pharmacodynamic measure of pathway inhibition (Yang et al., 2013).

Since the adduct is formed by the E1 enzyme itself, the rate of adduct accumulation can provide valuable information on the activity of the enzyme, which in turn can potentially serve as a predictive biomarker of response to these drugs (Brownell et al., 2010). In this context, the availability of specific antibodies against these adducts can be exploited to develop flow cytometry-based quantitative assays that can—with appropriate analytical and clinical validation—be employed to quantitatively measure the rate of adduct formation after administration of a single dose of the drug. This may help with decision-making as to whether patients should continue therapy with these agents, particularly in hematologic malignancies for which blood samples can be easily obtained. It remains challenging, however, to determine the cutoffs of enzyme activity that will be used to make such decisions.

F. Cell Cycle Effects of Adenosine Sulfamate E1 Inhibitors

Post-translational modifications, particularly with ubiquitin, SUMO, and NEDD8, play an important role in regulating cell cycle progression (Teixeira and Reed,

2013; Eifler and Vertegaal, 2015). Therefore, the disruption of these post-translational modifications is anticipated to result in the deregulation of the cell cycle. However, the ultimate phenotype varies among different adenosine sulfamates. Although the inhibition of neddylation with pevonedistat causes cell cycle arrest and accumulation of cells in the S phase, selective inhibition of ubiquitylation with a selective UBA1 inhibitor causes cell cycle arrest in the G2/M phase. Interestingly, dual inhibition of ubiquitylation and neddylation with TAK-243 also results in G2 arrest, suggesting that the effect on ubiquitylation is more dominant (Ciavarri and Langston, 2017). Although TAK-243 resembles pevonedistat in inducing the accumulation of CDT1, it also stabilizes its endogenous inhibitor, geminin, offsetting CDT1 effects (Hyer et al., 2018). As a result, pevonedistat induces CDT1-mediated re-replication, which is not observed with TAK-243. ML-792, on the other hand, causes mitotic disruption with a decrease in the number of cells in the anaphase/telophase subphases of mitosis, without affecting the number of those in the prometaphase and metaphase. These mitotic defects are associated with the induction of endoreduplication and the formation of DNA bridges (He et al., 2017).

G. Pharmacokinetics of Adenosine Sulfamate E1 Inhibitors

Among adenosine sulfamates, pevonedistat is the most extensively studied agent in clinical settings; thus, substantial human pharmacokinetic (PK) data are available on this drug (Table 11). After intravenous infusion, pevonedistat shows biphasic PK with a rapid distribution from the plasma to peripheral tissues, followed by a slower elimination from the body. In addition, it displays dose-dependent increases in plasma concentrations and remains detectable at quantifiable concentrations up to 72 hours after infusion; however, it shows little or no accumulation in the plasma after repeated intermittent or daily infusion (Bhatia et al., 2016; Sarantopoulos et al., 2016; Swords et al., 2015, 2017, 2018; Shah et al., 2016). The terminal elimination half-life time ($t_{1/2}$) of pevonedistat is approximately 7.5–10 hours after intravenous infusion. These PK parameters are not affected when pevonedistat is administered in combination with azacytidine, docetaxel, or gemcitabine, suggesting no significant interactions with these drugs. Conversely, higher plasma concentrations of pevonedistat have been observed after combination with carboplatin plus paclitaxel (Swords et al., 2018; Lockhart et al., 2019).

The PK of the second-in-class adenosine sulfamate, TAK-243, has also been evaluated in immunodeficient mice bearing subcutaneous diffuse, large B-cell lymphoma tumors after intravenous administration of the drug at doses of 12.5–25 mg/kg. As opposed to the short $t_{1/2}$ (0.2–0.4 hour) displayed by TAK-243 in the plasma,

the tumor $t_{1/2}$ is much longer (16.9–20.6 hour), reflecting the high volume of distribution and the higher exposure to the drug in tumors versus plasma (Hyer et al., 2018). The human PK of TAK-243 has also been reported among secondary outcome measures in a dose escalation study of the drug in advanced solid tumors (clinicaltrials.gov identifier: NCT02045095). Based on this study, the plasma $t_{1/2}$ of TAK-243 is 7–9.4 hours after intravenous infusion of the drug at doses of 4–18 mg (Table 12). With respect to the SAE inhibitor TAK-981, there is no available PK data on this drug thus far.

H. Insights into Clinical Trials of E1 Inhibitors

Of the four E1 inhibitors that have entered clinical trials so far (pevonedistat, TAK-243, TAK-981, and TAS4464), pevonedistat is the only drug for which clinical trial data has been published. It appears that pevonedistat shows more promising results when used in combinatorial versus single-agent regimens. In this respect, pevonedistat showed superior clinical benefits in combination with azacytidine in patients with AML and carboplatin plus paclitaxel in patients with advanced solid tumors (Swords et al., 2018; Lockhart et al., 2019). Many of these combinations are supported with preclinical studies showing synergistic antitumor effects with cytotoxic drugs as well as azacytidine (Smith et al., 2011; Jazaeri et al., 2013; Li et al., 2014b; Visconte et al., 2016). This suggests similar results could be obtained with other drug combinations that have shown preclinical evidence of synergism, such as belinostat, olaparib, vemurafenib, pomalidomide, and other conventional anticancer agents (Benamar et al., 2016; Leclerc et al., 2016; Zhou et al., 2016; Guo et al., 2017; Liu et al., 2019).

Better results have been observed with longer treatment durations of pevonedistat, and this could be achieved by dose adjustments and combination with low-toxicity drugs or drugs known to display toxicity profiles distinct from those of pevonedistat, as is the case with azacytidine (Swords et al., 2018). As E1 inhibitors are anticipated to have less favorable toxicity profile compared with other anticancer agents that selectively target aberrant oncogenic molecules, it is important to develop predictive biomarkers that identify patient cohorts most likely to benefit from these drugs and spare potential nonresponders from unnecessary toxicities. Clinical trials of pevonedistat in advanced solid tumors have shown that the expression of excision repair cross-complementation group 1 could potentially serve this role, as high expression levels positively correlated with clinical benefits and continuation of treatment when pevonedistat was combined with paclitaxel plus carboplatin. The need for such biomarkers is even more urgent with TAK-243 and TAK-981, which are expected to have more adverse effects compared with pevonedistat as a result of their broader pharmacologic effects

(Lockhart et al., 2019). Potential biomarkers for E1 inhibitors are highlighted in section VII. *E. Pharmacodynamic Activity of Adenosine Sulfamate E1 Inhibitors.*

VIII. Concluding Remarks

Despite the anticipated toxicity of proteasome inhibitors, they have proved clinically useful in several malignancies, particularly MM. These clinical successes have fueled subsequent research endeavors to explore other components of the UPS including E1 enzymes that initiate the cascade and possess readily druggable active sites, two key attributes that position E1s as attractive therapeutic targets. Although extensively evaluated in preclinical settings and clinical trials, the first-in-class NAE inhibitor pevonedistat has not been approved for clinical use yet. However, it has reached an advanced stage of clinical development as a combinatorial therapy with azacytidine in certain types of AML. The clinical approval of pevonedistat will encourage further exploration of other E1 inhibitors, particularly those that target SAE and UBA1. It must be noted, however, that the adverse drug reactions expected with these inhibitors—given the broad and essential biologic roles of ubiquitylation and SUMOylation—may present several challenges that dictate identifying the patient cohorts who are most likely to benefit from these therapies. Equally encouraging is the unprecedented appreciation of immunotherapy as a promising cancer treatment and the emerging immunomodulatory functions of SUMOylation, ubiquitylation, neddylation, and ISGylation. Identifying these potential immunotherapeutic effects is actively pursued with TAK-981 and pevonedistat, and further investigation may be warranted with other E1 inhibitors. Although possible with UBA1, for which a selective, clinical-grade inhibitor is available, it may be challenging to interrogate these effects with other E1s with no known inhibitors available thus far. Therefore, the discovery of selective inhibitors of UBA6, UBA7, and noncanonical E1s is needed to interrogate many unexplored areas of E1 biology and therapeutic potential in cancer.

Authorship Contributions

Performed data analysis: Barghout.

Wrote or contributed to the writing of the manuscript: Barghout, Schimmer.

References

- Abdollahzadeh I, Schwarten M, Gensch T, Willbold D, and Weiergräber OH (2017) The Atg8 family of proteins-modulating shape and functionality of autophagic membranes. *Front Genet* 8:109.
- Ablain J, Rice K, Soilihi H, de Reynies A, Minucci S, and de Thé H (2014) Activation of a promyelocytic leukemia-tumor protein 53 axis underlies acute promyelocytic leukemia cure. *Nat Med* 20:167–174.
- Agbor TA and Taylor CT (2008) SUMO, hypoxia and the regulation of metabolism. *Biochem Soc Trans* 36:445–448.
- Alontaga AY, Bobkova E, and Chen Y (2012) Biochemical analysis of protein SUMOylation. *Curr Protoc Mol Biol* Chapter 10, p Unit10.29.

- Alpi AF, Chaugule V, and Walden H (2016) Mechanism and disease association of E2-conjugating enzymes: lessons from UBE2T and UBE2L3. *Biochem J* **473**: 3401–3419.
- Amidon BS, Cardin DP, Gould AE, Greenspan PD, and Harrison SJ (2017) inventors, Millennium Pharmaceuticals Inc., assignee. Indole substituted pyrrolopyrimidinyl inhibitors of Uba6. U.S. patent US9593121B2. 2017 Mar 14.
- An H and Statsyuk AV (2013) Development of activity-based probes for ubiquitin and ubiquitin-like protein signaling pathways. *J Am Chem Soc* **135**:16948–16962.
- An H and Statsyuk AV (2015) An inhibitor of ubiquitin conjugation and aggresome formation. *Chem Sci (Camb)* **6**:5235–5245.
- Assouline S, Mehta A, Caimi PF, Wang B, Patel C, Kim M-S, Huszar D, Berger AJ, Friedlander S, Gomez-Pinillos A, et al. (2019) A Phase 1b/2 study of TAK-981, a first-in-class sumoylation inhibitor, in combination with rituximab in patients with relapsed/refractory (r/r) CD20-positive Non-Hodgkin Lymphoma (NHL). *Blood* **134**:1593.
- Bacik JP, Walker JR, Ali M, Schimmer AD, and Dhe-Paganon S (2010) Crystal structure of the human ubiquitin-activating enzyme 5 (UBA5) bound to ATP: mechanistic insights into a minimalistic E1 enzyme. *J Biol Chem* **285**: 20273–20280.
- Baek K, Krist DT, Prabu JR, Hill S, Klügel M, Neumaier LM, von Gronau S, Kleiger G, and Schulman BA (2020) NEDD8 nucleates a multivalent cullin-RING-UBE2D ubiquitin ligation assembly. *Nature* **578**:461–466.
- Baik H, Boulanger M, Hosseini M, Kowalczyk J, Zaghdoudi S, Salem T, Sarry JE, Hicheri Y, Cartron G, Piechaczek M, et al. (2018) Targeting the SUMO pathway primes all-trans retinoic acid-induced differentiation of nonpromyelocytic acute myeloid leukemias. *Cancer Res* **78**:2601–2613.
- Balachandran RS, Heighington CS, Starostina NG, Anderson JW, Owen DL, Vasudevan S, and Kipreos ET (2016) The ubiquitin ligase CRL2ZYG11 targets cyclin B1 for degradation in a conserved pathway that facilitates mitotic slippage. *J Cell Biol* **215**:151–166.
- Barbier-Torres L, Delgado TC, García-Rodríguez JL, Zubiete-Franco I, Fernández-Ramos D, Buqué X, Cano A, Gutiérrez-de Juan V, Fernández-Domínguez I, Lopitz-Otsoa F, et al. (2015) Stabilization of LKB1 and Akt by neddylation regulates energy metabolism in liver cancer. *Oncotarget* **6**:2509–2523.
- Barghout SH (2020) Targeted protein degradation: an emerging therapeutic strategy in cancer. *Anticancer Agents Med Chem* DOI: 10.2174/1871520620666200410082652 [published ahead of print].
- Barghout SH, MacLean N, Xu GW, Thomas G, Blatman Z, Hurren R, and Schimmer AD (2018) A genome-wide CRISPR/Cas9 knockout screen identifies BEND3 as a determinant of sensitivity to UBA1 inhibition in acute myeloid leukemia. *Blood* **132**:1350.
- Barghout SH, Patel PS, Wang X, Xu GW, Kavanagh S, Halgas O, Zarabi SF, Gronda M, Hurren R, Jeyaraju DV, et al. (2019) Preclinical evaluation of the selective small-molecule UBA1 inhibitor, TAK-243, in acute myeloid leukemia. *Leukemia* **33**:37–51.
- Barghout SH and Schimmer AD (2018) The ubiquitin-activating enzyme, UBA1, as a novel therapeutic target for AML. *Oncotarget* **9**:34198–34199.
- Bassi C, Ho J, Srikumar T, Dowling RJ, Gorrini C, Miller SJ, Mak TW, Neel BG, Raught B, and Stambolic V (2013) Nuclear PTEN controls DNA repair and sensitivity to genotoxic stress. *Science* **341**:395–399.
- Bedford L, Lowe J, Dick LR, Mayer RJ, and Brownell JE (2011) Ubiquitin-like protein conjugation and the ubiquitin-proteasome system as drug targets. *Nat Rev Drug Discov* **10**:29–46.
- Bellail AC, Olson JJ, and Hao C (2014) SUMO1 modification stabilizes CDK6 protein and drives the cell cycle and glioblastoma progression. *Nat Commun* **5**:4234.
- Benamar M, Guessous F, Du K, Corbett P, Obeid J, Gioeli D, Slingluff CL Jr., and Abbas T (2016) Inactivation of the CRL4-CDT2-SET8/p21 ubiquitylation and degradation axis underlies the therapeutic efficacy of pevonedistat in melanoma. *EBioMedicine* **10**:85–100.
- Berger AJ, Friedlander S, Ghasemi O, Grossman S, Koenig E, Lightcap E, Milhollen M, Shah P, Shapiro G, Shinde V, et al. (2019) Abstract 3079: pharmacodynamic evaluation of the novel SUMOylation inhibitor TAK-981 in a mouse tumor model. *Cancer Res* **79**.
- Bergink S and Jentsch S (2009) Principles of ubiquitin and SUMO modifications in DNA repair. *Nature* **458**:461–467.
- Bernardi R and Pandolfi PP (2007) Structure, dynamics and functions of promyelocytic leukaemia nuclear bodies. *Nat Rev Mol Cell Biol* **8**:1006–1016.
- Best S, Hashiguchi T, Kittai A, Bruss N, Paiva C, Okada C, Liu T, Berger A, and Danilov AV (2019a) Targeting ubiquitin-activating enzyme induces ER stress-mediated apoptosis in B-cell lymphoma cells. *Blood Adv* **3**:51–62.
- Best S, Lam V, Liu T, Bruss N, Kittai A, Danilova OV, Murray S, Berger A, Pennock ND, Lind EF, and Danilov AV (2020) Immunomodulatory effects of pevonedistat, a NEDD8-activating enzyme inhibitor, in chronic lymphocytic leukemia-derived T cells. *Leukemia* DOI: 10.1038/s41375-020-0794-0 [published ahead of print].
- Best S, Liu T, Bruss N, Kittai A, Berger A, and Danilov AV (2019b) Pharmacologic inhibition of the ubiquitin-activating enzyme induces ER stress and apoptosis in chronic lymphocytic leukemia and ibrutinib-resistant mantle cell lymphoma cells. *Leuk Lymphoma* **60**:2946–2950.
- Bhatia S, Pavlick AC, Boasberg P, Thompson JA, Mulligan G, Pickard MD, Faessel H, Dezube BJ, and Hamid O (2016) A phase I study of the investigational NEDD8-activating enzyme inhibitor pevonedistat (TAK-924/MLN4924) in patients with metastatic melanoma. *Invest New Drugs* **34**:439–449.
- Bhoj VG and Chen ZJ (2009) Ubiquitylation in innate and adaptive immunity. *Nature* **458**:430–437.
- Birkenkamp KU, Geugien M, Schepers H, Westra J, Lemmink HH, and Vellenga E (2004) Constitutive NF-kappaB DNA-binding activity in AML is frequently mediated by a Ras/PI3-K/PKB-dependent pathway. *Leukemia* **18**:103–112.
- Blank JL, Liu XJ, Cosmopoulos K, Bouck DC, Garcia K, Bernard H, Tayber O, Hather G, Liu R, Narayanan U, et al. (2013) Novel DNA damage checkpoints mediating cell death induced by the NEDD8-activating enzyme inhibitor MLN4924. *Cancer Res* **73**:225–234.
- Blomstrom DC, Fahey D, Kutny R, Korant BD, and Knight E Jr. (1986) Molecular characterization of the interferon-induced 15-kDa protein. Molecular cloning and nucleotide and amino acid sequence. *J Biol Chem* **261**:8811–8816.
- Boggio R, Colombo R, Hay RT, Draetta GF, and Chiocca S (2004) A mechanism for inhibiting the SUMO pathway. *Mol Cell* **16**:549–561.
- Bohnsack RN and Haas AL (2003) Conservation in the mechanism of Nedd8 activation by the human AppBp1-Uba3 heterodimer. *J Biol Chem* **278**:26823–26830.
- Bosman MC, Schepers H, Jaques J, Brouwers-Vos AZ, Quax WJ, Schuringa JJ, and Vellenga E (2014) The TAK1-NF-kB axis as therapeutic target for AML. *Blood* **124**:3130–3140.
- Bossis G and Melchior F (2006) Regulation of SUMOylation by reversible oxidation of SUMO conjugating enzymes. *Mol Cell* **21**:349–357.
- Bossis G, Sarry JE, Kifagi C, Ristic M, Saland E, Vergez F, Salem T, Boutzen H, Baik H, Brockly F, et al. (2014) The ROS/SUMO axis contributes to the response of acute myeloid leukemia cells to chemotherapeutic drugs. *Cell Rep* **7**:1815–1823.
- Broemer M and Meier P (2009) Ubiquitin-mediated regulation of apoptosis. *Trends Cell Biol* **19**:130–140.
- Brown JS, Lukashchuk N, Sczaniecka-Clift M, Britton S, le Sage C, Calsou P, Beli P, Galanty Y, and Jackson SP (2015) Neddylation promotes ubiquitylation and release of Ku from DNA-damage sites. *Cell Rep* **11**:704–714.
- Brown P, Richardson CM, Mensah LM, O'Hanlon PJ, Osborne NF, Pope AJ, and Walker G (1999) Molecular recognition of tyrosinyl adenylate analogues by prokaryotic tyrosyl tRNA synthetases. *Bioorg Med Chem* **7**:2473–2485.
- Brownell JE, Sintchak MD, Gavin JM, Liao H, Bruzzese FJ, Bump NJ, Soucy TA, Milhollen MA, Yang X, Burkhardt AL, et al. (2010) Substrate-assisted inhibition of ubiquitin-like protein-activating enzymes: the NEDD8 E1 inhibitor MLN4924 forms a NEDD8-AMP mimetic in situ. *Mol Cell* **37**:102–111.
- Bruzzese FJ, Tsu CA, Ma J, Loke HK, Wu D, Li Z, Tayber O, and Dick LR (2009) Development of a charcoal paper adenosine triphosphate:pyrophosphate exchange assay: kinetic characterization of NEDD8 activating enzyme. *Anal Biochem* **394**:24–29.
- Buac D, Shen M, Schmitt S, Kona FR, Deshmukh R, Zhang Z, Neslund-Dudas C, Mitra B, and Dou QP (2013) From bortezomib to other inhibitors of the proteasome and beyond. *Curr Pharm Des* **19**:4025–4038.
- Burks J, Reed RE, and Desai SD (2014) ISGylation governs the oncogenic function of Ki-Ras in breast cancer. *Oncogene* **33**:794–803.
- Cai Y, Pi W, Sivaprakasam S, Zhu X, Zhang M, Chen J, Makala L, Lu C, Wu J, Teng Y, et al. (2015) UFBP1, a key component of the Ufm1 conjugation system, is essential for ufmylation-mediated regulation of erythroid development. *PLoS Genet* **11**:e1005643.
- Cai Y, Singh N, and Li H (2016) Essential role of Ufm1 conjugation in the hematopoietic system. *Exp Hematol* **44**:442–446.
- Canaan A, DeFuria J, Perelman E, Schultz V, Seay M, Tuck D, Flavell RA, Snyder MP, Obin MS, and Weissman SM (2014) Extended lifespan and reduced adiposity in mice lacking the FAT10 gene. *Proc Natl Acad Sci USA* **111**:5313–5318.
- Canaan A, Yu X, Booth CJ, Lian J, Lazar I, Gamfi SL, Castille K, Kohya N, Nakayama Y, Liu YC, et al. (2006) FAT10/diubiquitin-like protein-deficient mice exhibit minimal phenotypic differences. *Mol Cell Biol* **26**:5180–5189.
- Cao Q, You X, Xu L, Wang L, and Chen Y (2020) PAQR3 suppresses the growth of non-small cell lung cancer cells via modulation of EGFR-mediated autophagy. *Autophagy* **16**:1236–1247.
- Capdeville R, Buchdunger E, Zimmermann J, and Matter A (2002) Glivec (STI571, imatinib), a rationally developed, targeted anticancer drug. *Nat Rev Drug Discov* **1**: 493–502.
- Cappadocia L and Lima CD (2018) Ubiquitin-like protein conjugation: structures, chemistry, and mechanism. *Chem Rev* **118**:889–918.
- Chen D, Li Y, Zhang X, Wu H, Wang Q, Cai J, Cui Y, Liu H, Lan P, Wang J, et al. (2019) Ubiquitin ligase TRIM65 promotes colorectal cancer metastasis by targeting ARHGAP35 for protein degradation. *Oncogene* **38**:6429–6444.
- Chen JJ, Tsu CA, Gavin JM, Milhollen MA, Bruzzese FJ, Mallender WD, Sintchak MD, Bump NJ, Yang X, Ma J, et al. (2011) Mechanistic studies of substrate-assisted inhibition of ubiquitin-activating enzyme by adenosine sulfamate analogues. *J Biol Chem* **286**:40867–40877.
- Chen P, Hu T, Liang Y, Jiang Y, Pan Y, Li C, Zhang P, Wei D, Li P, Jeong LS, et al. (2015) Synergistic inhibition of autophagy and neddylation pathways as a novel therapeutic approach for targeting liver cancer. *Oncotarget* **6**:9002–9017.
- Chen P, Hu T, Liang Y, Li P, Chen X, Zhang J, Ma Y, Hao Q, Wang J, Zhang P, et al. (2016a) Neddylation inhibition activates the extrinsic apoptosis pathway through ATF4-CHOP-DR5 axis in human esophageal cancer cells. *Clin Cancer Res* **22**: 4145–4157.
- Chen Q, Sun L, and Chen ZJ (2016b) Regulation and function of the cGAS-STING pathway of cytosolic DNA sensing. *Nat Immunol* **17**:1142–1149.
- Chène P (2003) Inhibiting the p53-MDM2 interaction: an important target for cancer therapy. *Nat Rev Cancer* **3**:102–109.
- Chiu YH, Sun Q, and Chen ZJ (2007) E1-L2 activates both ubiquitin and FAT10. *Mol Cell* **27**:1014–1023.
- Choi AM, Rytter SW, and Levine B (2013) Autophagy in human health and disease. *N Engl J Med* **368**:651–662.
- Chowdhury MM, Dosche C, Löhmansröben HG, and Leimkühler S (2012) Dual role of the molybdenum cofactor biosynthesis protein MOCS3 in tRNA thiolation and molybdenum cofactor biosynthesis in humans. *J Biol Chem* **287**:17297–17307.
- Ciavarrri J and Langston S (2017) 8.04 - The discovery of first-in-class inhibitors of the Nedd8-Activating Enzyme (NAE) and the Ubiquitin-Activating Enzyme (UAE), in *Comprehensive Medicinal Chemistry III* (Rotella D and Ward SE eds) pp 95–112, Elsevier, Oxford.
- Ciechanover A (2005) Proteolysis: from the lysosome to ubiquitin and the proteasome. *Nat Rev Mol Cell Biol* **6**:79–87.
- Ciechanover A, Elias S, Heller H, and Hershkó A (1982) "Covalent affinity" purification of ubiquitin-activating enzyme. *J Biol Chem* **257**:2537–2542.
- Ciechanover A, Heller H, Katz-Etzion R, and Hershkó A (1981) Activation of the heat-stable polypeptide of the ATP-dependent proteolytic system. *Proc Natl Acad Sci USA* **78**:761–765.

- Clague MJ, Heride C, and Urbé S (2015) The demographics of the ubiquitin system. *Trends Cell Biol* **25**:417–426.
- Cohen P (2002) Protein kinases—the major drug targets of the twenty-first century? *Nat Rev Drug Discov* **1**:309–315.
- Cohen P and Tcherpakov M (2010) Will the ubiquitin system furnish as many drug targets as protein kinases? *Cell* **143**:686–693.
- Cooper J, Xu Q, Zhou L, Pavlovic M, Ojeda V, Moulick K, de Stanchina E, Poirier JT, Zauderer M, Rudin CM, et al. (2017) Combined inhibition of NEDD8-activating enzyme and mTOR suppresses NF2 loss-driven tumorigenesis. *Mol Cancer Ther* **16**:1693–1704.
- Copp LJ (2003) Mechanism-based inhibition, in *Enzyme Kinetics* (Marangoni AG ed) pp 158–173.
- Crowl JT and Stetson DB (2018) SUMO2 and SUMO3 redundantly prevent a noncanonical type I interferon response. *Proc Natl Acad Sci USA* **115**: 6798–6803.
- Cui Y, Yu H, Zheng X, Peng R, Wang Q, Zhou Y, Wang R, Wang J, Qu B, Shen N, et al. (2017) SENP7 potentiates cGAS activation by relieving SUMO-mediated inhibition of cytosolic DNA sensing. *PLoS Pathog* **13**:e1006156.
- Cuijpers SAG and Vertegaal ACO (2018) Guiding mitotic progression by crosstalk between post-translational modifications. *Trends Biochem Sci* **43**:251–268.
- Czuczman NM, Barth MJ, Gu J, Neppalli V, Mavis C, Frys SE, Hu Q, Liu S, Klenner P, Vockova P, et al. (2016) Pevonedistat, a NEDD8-activating enzyme inhibitor, is active in mantle cell lymphoma and enhances rituximab activity in vivo. *Blood* **127**:1128–1137.
- Daniel J and Liebau E (2014) The Ufm1 cascade. *Cells* **3**:627–638.
- Dantuma NP and van Attikum H (2016) Spatiotemporal regulation of post-translational modifications in the DNA damage response. *EMBO J* **35**:6–23.
- da Silva SR, Paiva SL, Bancerz M, Geletu M, Lewis AM, Chen J, Cai Y, Lukkarila JL, Li H, and Gunning PT (2016) A selective inhibitor of the UFM1-activating enzyme, UBA5. *Bioorg Med Chem Lett* **26**:4542–4547.
- Decque A, Joffre O, Magalhaes JG, Cossec JC, Blecher-Gonen R, Lapaquette P, Silvain A, Manel N, Joubert PE, Seeler JS, et al. (2016) Sumoylation coordinates the repression of inflammatory and anti-viral gene-expression programs during innate sensing. *Nat Immunol* **17**:140–149.
- DeJesus R, Moretti F, McAllister G, Wang Z, Bergman P, Liu S, Frias E, Alford J, Reece-Hoyes JS, Lindeman A, et al. (2016) Functional CRISPR screening identifies the ufmylation pathway as a regulator of SQSTM1/p62. *eLife* **5**:e17290.
- Delaunay S, Rapino F, Tharun L, Zhou Z, Heukamp L, Termathe M, Shostak K, Klevernic I, Florin A, Desmecht H, et al. (2016) Elp3 links tRNA modification to IRES-dependent translation of LEF1 to sustain metastasis in breast cancer. *J Exp Med* **213**:2503–2523.
- Dengler MA, Weillbacher A, Gutekunst M, Staiger AM, Vöhringer MC, Horn H, Ott G, Aulitzky WE, and van der Kuip H (2014) Discrepant NOXA (PMAIP1) transcript and NOXA protein levels: a potential Achilles' heel in mantle cell lymphoma. *Cell Death Dis* **5**:e1013.
- Desai SD (2015) ISG15: a double edged sword in cancer. *OncoImmunology* **4**: e1052935.
- Desai SD, Haas AL, Wood LM, Tsai YC, Pestka S, Rubin EH, Saleem A, Nur-E-Kamal A, and Liu LF (2006) Elevated expression of ISG15 in tumor cells interferes with the ubiquitin/26S proteasome pathway. *Cancer Res* **66**:921–928.
- de Thé H and Chen Z (2010) Acute promyelocytic leukaemia: novel insights into the mechanisms of cure. *Nat Rev Cancer* **10**:775–783.
- de Thé H, Pandolfi PP, and Chen Z (2017) Acute promyelocytic leukemia: a paradigm for oncoprotein-targeted cure. *Cancer Cell* **32**:552–560.
- Dhananjayan SC, Ismail A, and Nawaz Z (2005) Ubiquitin and control of transcription. *Essays Biochem* **41**:69–80.
- DiDonato JA, Mercurio F, and Karin M (2012) NF- κ B and the link between inflammation and cancer. *Immunol Rev* **246**:379–400.
- Dikic I (2017) Proteasomal and autophagic degradation systems. *Annu Rev Biochem* **86**:193–224.
- Dobbelstein M and Moll U (2014) Targeting tumour-supportive cellular machineries in anticancer drug development. *Nat Rev Drug Discov* **13**:179–196.
- Dou T, Gu S, Liu J, Chen F, Zeng L, Guo L, Xie Y, and Mao Y (2005) Isolation and characterization of ubiquitin-activating enzyme E1-domain containing 1, UBE1DC1. *Mol Biol Rep* **32**:265–271.
- Du L, Li YJ, Fakih M, Wiarek RL, Duldulao M, Chen Z, Chu P, Garcia-Aguilar J, and Chen Y (2016) Role of SUMO-activating enzyme in cancer stem cell maintenance and self-renewal. *Nat Commun* **7**:12326.
- Dusha I and Dénes G (1977) A simplified assay of enzymes catalyzing ATP-pyrophosphate exchange reactions. *Anal Biochem* **81**:247–250.
- Eifler K and Vertegaal ACO (2015) SUMOylation-mediated regulation of cell cycle progression and cancer. *Trends Biochem Sci* **40**:779–793.
- Ellis LM and Hicklin DJ (2009) Resistance to targeted therapies: refining anticancer therapy in the era of molecular oncology. *Clin Cancer Res* **15**:7471–7478.
- Enchev RI, Schulman BA, and Peter M (2015) Protein neddylation: beyond cullin-RING ligases. *Nat Rev Mol Cell Biol* **16**:30–44.
- Fan JB, Miyauchi-Ishida S, Arimoto K, Liu D, Yan M, Liu CW, Györfy B, and Zhang DE (2015) Type I IFN induces protein ISGylation to enhance cytokine expression and augments colonic inflammation. *Proc Natl Acad Sci USA* **112**:14313–14318.
- Feng Q, Sekula D, Guo Y, Liu X, Black CC, Galimberti F, Shah SJ, Sempere LF, Memoli V, Andersen JB, et al. (2008) UBE1L causes lung cancer growth suppression by targeting cyclin D1. *Mol Cancer Ther* **7**:3780–3788.
- Fenteany G, Gaur P, Sharma G, Pintér L, Kiss E, and Haracska L (2020) Robust high-throughput assays to assess discrete steps in ubiquitination and related cascades. *BMC Mol Cell Biol* **21**:21.
- Flick K and Kaiser P (2012) Protein degradation and the stress response. *Semin Cell Dev Biol* **23**:515–522.
- Forrest AK, Jarvest RL, Mensah LM, O'Hanlon PJ, Pope AJ, and Sheppard RJ (2000) Aminoalkyl adenylate and aminoacyl sulfamate intermediate analogues differing greatly in affinity for their cognate *Staphylococcus aureus* aminoacyl tRNA synthetases. *Bioorg Med Chem Lett* **10**:1871–1874.
- Fujita Y, Tinoco R, Li Y, Senft D, and Ronai ZA (2019) Ubiquitin ligases in cancer immunotherapy - balancing antitumor and autoimmunity. *Trends Mol Med* **25**: 428–443.
- Fukuda I, Ito A, Hirai G, Nishimura S, Kawasaki H, Saitoh H, Kimura K, Sodeoka M, and Yoshida M (2009a) Ginkgolic acid inhibits protein SUMOylation by blocking formation of the E1-SUMO intermediate. *Chem Biol* **16**:133–140.
- Fukuda I, Ito A, Uramoto M, Saitoh H, Kawasaki H, Osada H, and Yoshida M (2009b) Kerriamycin B inhibits protein SUMOylation. *J Antibiot (Tokyo)* **62**: 221–224.
- Fulda S, Rajalingam K, and Dikic I (2012) Ubiquitylation in immune disorders and cancer: from molecular mechanisms to therapeutic implications. *EMBO Mol Med* **4**:545–556.
- Furukawa K, Mizushima N, Noda T, and Ohsumi Y (2000) A protein conjugation system in yeast with homology to biosynthetic enzyme reaction of prokaryotes. *J Biol Chem* **275**:7462–7465.
- Gallo LH, Ko J, and Donoghue DJ (2017) The importance of regulatory ubiquitination in cancer and metastasis. *Cell Cycle* **16**:634–648.
- Galluzzi L and Green DR (2019) Autophagy-independent functions of the autophagy machinery. *Cell* **177**:1682–1699.
- Gao Q, Yu GY, Shi JY, Li LH, Zhang WJ, Wang ZC, Yang LX, Duan M, Zhao H, Wang XY, et al. (2014) Neddylation pathway is up-regulated in human intrahepatic cholangiocarcinoma and serves as a potential therapeutic target. *Oncotarget* **5**: 7820–7832.
- Garcia K, Blank JL, Bouck DC, Liu XJ, Sappal DS, Hather G, Cosmopoulos K, Thomas MP, Kuranda M, Pickard MD, et al. (2014) Nedd8-activating enzyme inhibitor MLN4924 provides synergy with mitomycin C through interactions with ATR, BRCA1/BRCA2, and chromatin dynamics pathways. *Mol Cancer Ther* **13**: 1625–1635.
- Gareau JR and Lima CD (2010) The SUMO pathway: emerging mechanisms that shape specificity, conjugation and recognition. *Nat Rev Mol Cell Biol* **11**:861–871.
- Gavin JM, Chen JJ, Liao H, Rollins N, Yang X, Xu Q, Ma J, Loke HK, Lingaraj T, Brownell JE, et al. (2012) Mechanistic studies on activation of ubiquitin and di-ubiquitin-like protein, FAT10, by ubiquitin-like modifier activating enzyme 6, Uba6. *J Biol Chem* **287**:15512–15522.
- Gavin JM, Hoar K, Xu Q, Ma J, Lin Y, Chen J, Chen W, Bruzzese FJ, Harrison S, Mallender WD, et al. (2014) Mechanistic study of Uba5 enzyme and the Ufm1 conjugation pathway. *J Biol Chem* **289**:22648–22658.
- Geng J and Klionsky DJ (2008) The Atg8 and Atg12 ubiquitin-like conjugation systems in macroautophagy. 'Protein modifications: beyond the usual suspects' review series. *EMBO Rep* **9**:859–864.
- Gerakis Y, Quintero M, Li H, and Hetz C (2019) The UFMylation system in proteostasis and beyond. *Trends Cell Biol* **29**:974–986.
- Ghosh SK, Perrine SP, Williams RM, and Fallor DV (2012) Histone deacetylase inhibitors are potent inducers of gene expression in latent EBV and sensitize lymphoma cells to nucleoside antiviral agents. *Blood* **119**:1008–1017.
- Gilberto S and Peter M (2017) Dynamic ubiquitin signaling in cell cycle regulation. *J Cell Biol* **216**:2259–2271.
- Godberson JC, Humphries LA, Danilova OV, Kebbekus PE, Brown JR, Eastman A, and Danilov AV (2014) The Nedd8-activating enzyme inhibitor MLN4924 thwarts microenvironment-driven NF- κ B activation and induces apoptosis in chronic lymphocytic leukemia B cells. *Clin Cancer Res* **20**:1576–1589.
- Goldstein G, Scheid M, Hammerling U, Schlesinger DH, Niall HD, and Boyse EA (1975) Isolation of a polypeptide that has lymphocyte-differentiating properties and is probably represented universally in living cells. *Proc Natl Acad Sci USA* **72**: 11–15.
- Gómez-Díaz C and Ikeda F (2019) Roles of ubiquitin in autophagy and cell death. *Semin Cell Dev Biol* **93**:125–135.
- Gomez-Puerto MC, Folkerts H, Wierenga AT, Schepers K, Schuringa JJ, Coffey PJ, and Vellenga E (2016) Autophagy proteins ATG5 and ATG7 are essential for the maintenance of human CD34(+) hematopoietic stem-progenitor cells. *Stem Cells* **34**:1651–1663.
- Gong K, Chen C, Zhan Y, Chen Y, Huang Z, and Li W (2012) Autophagy-related gene 7 (ATG7) and reactive oxygen species/extracellular signal-regulated kinase regulate tetrandrine-induced autophagy in human hepatocellular carcinoma. *J Biol Chem* **287**:35576–35588.
- Gong P, Canaan A, Wang B, Leventhal J, Snyder A, Nair V, Cohen CD, Kretzler M, D'Agati V, Weissman S, et al. (2010) The ubiquitin-like protein FAT10 mediates NF- κ B activation. *J Am Soc Nephrol* **21**:316–326.
- Grabbe C, Husnjak K, and Dikic I (2011) The spatial and temporal organization of ubiquitin networks. *Nat Rev Mol Cell Biol* **12**:295–307.
- Graves JD, Lee YJ, Liu K, Li G, Lin FT, and Lin WC (2020) E2F1 sumoylation as a protective cellular mechanism in oxidative stress response. *Proc Natl Acad Sci USA* **117**:14958–14969.
- Groen EJN and Gillingwater TH (2015) UBA1: at the crossroads of ubiquitin homeostasis and neurodegeneration. *Trends Mol Med* **21**:622–632.
- Groettrup M, Pelzer C, Schmidtke G, and Hofmann K (2008) Activating the ubiquitin family: UBA6 challenges the field. *Trends Biochem Sci* **33**:230–237.
- Grossman RL, Heath AP, Ferretti V, Varmus HE, Lowy DR, Kibbe WA, and Staudt LM (2016) Toward a shared vision for cancer genomic data. *N Engl J Med* **375**: 1109–1112.
- Gu Y, Kaufman JL, Bernal L, Torre C, Matulis SM, Harvey RD, Chen J, Sun SY, Boisse LH, and Lonial S (2014) MLN4924, an NAE inhibitor, suppresses AKT and mTOR signaling via upregulation of REDD1 in human myeloma cells. *Blood* **123**: 3269–3276.
- Guo JY, Karshi-Uzunbas G, Mathew R, Aisner SC, Kamphorst JJ, Strohecker AM, Chen G, Price S, Lu W, Teng X, et al. (2013a) Autophagy suppresses progression of K-ras-induced lung tumors to oncocytes and maintains lipid homeostasis. *Genes Dev* **27**:1447–1461.
- Guo JY and White E (2013) Autophagy is required for mitochondrial function, lipid metabolism, growth, and fate of KRAS(G12D)-driven lung tumors. *Autophagy* **9**: 1636–1638.

- Guo JY, Xia B, and White E (2013b) Autophagy-mediated tumor promotion. *Cell* **155**: 1216–1219.
- Guo ZP, Hu YC, Xie Y, Jin F, Song ZQ, Liu XD, Ma T, and Zhou PK (2017) MLN4924 suppresses the BRCA1 complex and synergizes with PARP inhibition in NSCLC cells. *Biochem Biophys Res Commun* **483**:223–229.
- Ha BH, Ahn HC, Kang SH, Tanaka K, Chung CH, and Kim EE (2008) Structural basis for Ufm1 processing by UISP1. *J Biol Chem* **283**:14893–14900.
- Ha BH, Jeon YJ, Shin SC, Tatsumi K, Komatsu M, Tanaka K, Watson CM, Wallis G, Chung CH, and Kim EE (2011) Structure of ubiquitin-fold modifier 1-specific protease UfSP2. *J Biol Chem* **286**:10248–10257.
- Haas AL and Rose IA (1982) The mechanism of ubiquitin activating enzyme. A kinetic and equilibrium analysis. *J Biol Chem* **257**:10329–10337.
- Habisov S, Huber J, Ichimura Y, Akutsu M, Rogova N, Loehr F, McEwan DG, Johansen T, Dikic I, Doetsch V, et al. (2016) Structural and functional analysis of a novel interaction motif within UFM1-activating enzyme 5 (UBA5) required for binding to ubiquitin-like proteins and ufmylation. *J Biol Chem* **291**: 9025–9041.
- Haglund K, Di Fiore PP, and Dikic I (2003) Distinct monoubiquitin signals in receptor endocytosis. *Trends Biochem Sci* **28**:598–603.
- Hamdoun S and Efferth T (2017) Ginkgolic acids inhibit migration in breast cancer cells by inhibition of NEMO sumoylation and NF- κ B activity. *Oncotarget* **8**: 35103–35115.
- Han J, Goldstein LA, Hou W, Chatterjee S, Burns TF, and Rabinowich H (2018) HSP90 inhibition targets autophagy and induces a CASP9-dependent resistance mechanism in NSCLC. *Autophagy* **14**:958–971.
- Han J, Hou W, Goldstein LA, Stolz DB, Watkins SC, and Rabinowich H (2014) A complex between Atg7 and Caspase-9: a novel mechanism of cross-regulation between autophagy and apoptosis. *J Biol Chem* **289**:6485–6497.
- Han K, Wang Q, Cao H, Qiu G, Cao J, Li X, Wang J, Shen B, and Zhang J (2016) The NEDD8-activating enzyme inhibitor MLN4924 induces G2 arrest and apoptosis in T-cell acute lymphoblastic leukemia. *Oncotarget* **7**:23812–23824.
- Han W, Pan H, Chen Y, Sun J, Wang Y, Li J, Ge W, Feng L, Lin X, Wang X, et al. (2011) EGFR tyrosine kinase inhibitors activate autophagy as a cytoprotective response in human lung cancer cells. *PLoS One* **6**:e18691.
- Hanahan D and Weinberg RA (2011) Hallmarks of cancer: the next generation. *Cell* **144**:646–674.
- Hanel W, Tsyba L, Huszar D, Prouty A, Zhang X, Helmig-Mason J, Mundy-Bosse BL, Youssef Y, Parekh S, Maddocks KJ, et al. (2019) Targeting hypersumoylation in mantle cell lymphoma. *Blood* **134**:4060.
- Hatton BA, Grenley M, Garnsey J, Shinde V, Huszar D, Burns C, Ditzler S, Merrell A, Dey J, Beirne E, et al. (2019) Abstract 4136: direct intratumoral microdosing via the CIVO® platform reveals anti-tumor immune responses induced by the SUMO inhibitor TAK-981. *Cancer Res* **79**.
- He X, Riceberg J, Pulkuri SM, Grossman S, Shinde V, Shah P, Brownell JE, Dick L, Newcomb J, and Bence N (2015) Characterization of the loss of SUMO pathway function on cancer cells and tumor proliferation. *PLoS One* **10**:e0123882.
- He X, Riceberg J, Soucy T, Koenig E, Minissale J, Gallery M, Bernard H, Yang X, Liao H, Rabino C, et al. (2017) Probing the roles of SUMOylation in cancer cell biology by using a selective SAE inhibitor. *Nat Chem Biol* **13**:1164–1171.
- Hermann M and Bogunovic D (2017) ISG15 in sickness and in health. *Trends Immunol* **38**:79–93.
- Hesson LB, Cooper WN, and Latif F (2007) Evaluation of the 3p21.3 tumour-suppressor gene cluster. *Oncogene* **26**:7283–7301.
- Ho IL, Kuo KL, Liu SH, Chang HC, Hsieh JT, Wu JT, Chiang CK, Lin WC, Tsai YC, Chou CT, et al. (2015) MLN4924 synergistically enhances cisplatin-induced cytotoxicity via JNK and Bcl-xL pathways in human urothelial carcinoma. *Sci Rep* **5**: 16948.
- Hochstrasser M (2009) Origin and function of ubiquitin-like proteins. *Nature* **458**: 422–429.
- Hoellein A, Fallahi M, Schoeffmann S, Steidle S, Schaub FX, Rudelius M, Laitinen I, Nilsson L, Goga A, Peschel C, et al. (2014) Myc-induced SUMOylation is a therapeutic vulnerability for B-cell lymphoma. *Blood* **124**:2081–2090.
- Hoeller D and Dikic I (2009) Targeting the ubiquitin system in cancer therapy. *Nature* **458**:438–444.
- Hoeller D, Hecker CM, and Dikic I (2006) Ubiquitin and ubiquitin-like proteins in cancer pathogenesis. *Nat Rev Cancer* **6**:776–788.
- Holohan C, Van Schaeckbroeck S, Longley DB, and Johnston PG (2013) Cancer drug resistance: an evolving paradigm. *Nat Rev Cancer* **13**:714–726.
- Hong J and Luesch H (2012) Largazole: from discovery to broad-spectrum therapy. *Nat Prod Rep* **29**:449–456.
- Hong SB, Kim BW, Lee KE, Kim SW, Jeon H, Kim J, and Song HK (2011) Insights into noncanonical E1 enzyme activation from the structure of autophagic E1 Atg7 with Atg8. *Nat Struct Mol Biol* **18**:1323–1330.
- Hong X, Li S, Li W, Xie M, Wei Z, Guo H, Wei W, and Zhang S (2019) Disruption of protein neddylation with MLN4924 attenuates paclitaxel-induced apoptosis and microtubule polymerization in ovarian cancer cells. *Biochem Biophys Res Commun* **508**:986–990.
- Hu H and Sun SC (2016) Ubiquitin signaling in immune responses. *Cell Res* **26**: 457–483.
- Hua W, Li C, Yang Z, Li L, Jiang Y, Yu G, Zhu W, Liu Z, Duan S, Chu Y, et al. (2015) Suppression of glioblastoma by targeting the overactivated protein neddylation pathway. *Neuro Oncol* **17**:1333–1343.
- Huang DT, Hunt HW, Zhuang M, Ohi MD, Holton JM, and Schulman BA (2007) Basis for a ubiquitin-like protein thioester switch toggling E1-E2 affinity. *Nature* **445**:394–398.
- Huang DT, Miller DW, Mathew R, Cassell R, Holton JM, Roussel MF, and Schulman BA (2004a) A unique E1-E2 interaction required for optimal conjugation of the ubiquitin-like protein NEDD8. *Nat Struct Mol Biol* **11**:927–935.
- Huang DT, Walden H, Duda D, and Schulman BA (2004b) Ubiquitin-like protein activation. *Oncogene* **23**:1958–1971.
- Huang J, Zhou Y, Thomas GS, Gu Z, Yang Y, Xu H, Tricot G, and Zhan F (2015) NEDD8 inhibition overcomes CKS1B-induced drug resistance by upregulation of p21 in multiple myeloma. *Clin Cancer Res* **21**:5532–5542.
- Huang TT and D'Andrea AD (2006) Regulation of DNA repair by ubiquitylation. *Nat Rev Mol Cell Biol* **7**:323–334.
- Huang X and Dixit VM (2016) Drugging the undruggables: exploring the ubiquitin system for drug development. *Cell Res* **26**:484–498.
- Huang YF, Wee S, Gunaratne J, Lane DP, and Bulavin DV (2014) Isg15 controls p53 stability and functions. *Cell Cycle* **13**:2200–2210.
- Huangfu WC and Fuchs SY (2010) Ubiquitination-dependent regulation of signaling receptors in cancer. *Genes Cancer* **1**:725–734.
- Hubbard SR and Till JH (2000) Protein tyrosine kinase structure and function. *Annu Rev Biochem* **69**:373–398.
- Huber J, Obata M, Gruber J, Akutsu M, Löhr F, Rogova N, Güntert P, Dikic I, Kirkin V, Komatsu M, et al. (2020) An atypical LIR motif within UBA5 (ubiquitin like modifier activating enzyme 5) interacts with GABARAP proteins and mediates membrane localization of UBA5. *Autophagy* **16**:256–270.
- Huszar D (2019) New drugs on the horizon TAK-981: a first-in-class SUMOylation inhibitor in phase 1 clinical trials promotes a type I interferon response and antitumor immunity in preclinical models, in American Association for Cancer Research Annual Meeting; 2019 March 31; Atlanta, GA.
- Hyer ML, Milhollen MA, Ciavarrri J, Fleming P, Traore T, Sappal D, Huck J, Shi J, Gavin J, Brownell J, et al. (2018) A small-molecule inhibitor of the ubiquitin activating enzyme for cancer treatment. *Nat Med* **24**:186–193.
- Inamura K, Shimoi T, Ninomiya H, Hiramatsu M, Okui M, Satoh Y, Okumura S, Nakagawa K, Noda T, Fukayama M, et al. (2007) A metastatic signature in entire lung adenocarcinomas irrespective of morphological heterogeneity. *Hum Pathol* **38**:702–709.
- Ishikawa Y, Nakayama K, Morimoto M, Mizutani A, Nakayama A, Toyoshima K, Hayashi A, Takagi S, Dairiki R, Miyashita H, et al. (2017) Synergistic anti-AML effects of the LSD1 inhibitor T-3775440 and the NEDD8-activating enzyme inhibitor pevonedistat via transdifferentiation and DNA rereplication. *Oncogenesis* **6**: e377.
- Iyer LM, Burroughs AM, and Aravind L (2006) The prokaryotic antecedents of the ubiquitin-signaling system and the early evolution of ubiquitin-like beta-grasp domains. *Genome Biol* **7**:R60.
- Jackson SP and Durocher D (2013) Regulation of DNA damage responses by ubiquitin and SUMO. *Mol Cell* **49**:795–807.
- Jazaeri AA, Shibata E, Park J, Bryant JL, Conaway MR, Modesitt SC, Smith PG, Milhollen MA, Berger AJ, and Dutta A (2013) Overcoming platinum resistance in preclinical models of ovarian cancer using the neddylation inhibitor MLN4924. *Mol Cancer Ther* **12**:1958–1967.
- Jeanne M, Lallemand-Breitenbach V, Ferhi O, Koken M, Le Bras M, Duffort S, Peres L, Berthier C, Soilihi H, Raught B, et al. (2010) PML/RARA oxidation and arsenic binding initiate the antileukemia response of As2O3. *Cancer Cell* **18**:88–98.
- Jeon YJ, Jo MG, Yoo HM, Hong SH, Park JM, Ka SH, Oh KH, Seol JH, Jung YK, and Chung CH (2012) Chemosensitivity is controlled by p63 modification with ubiquitin-like protein ISG15. *J Clin Invest* **122**:2622–2636.
- Jeon YJ, Yoo HM, and Chung CH (2010) ISG15 and immune diseases. *Biochim Biophys Acta* **1802**:485–496.
- Jia L, Li H, and Sun Y (2011) Induction of p21-dependent senescence by an NAE inhibitor, MLN4924, as a mechanism of growth suppression. *Neoplasia* **13**: 561–569.
- Jin J, Li X, Gygi SP, and Harper JW (2007) Dual E1 activation systems for ubiquitin differentially regulate E2 enzyme charging. *Nature* **447**:1135–1138.
- Jin L, Wang W, and Fang G (2014) Targeting protein-protein interaction by small molecules. *Annu Rev Pharmacol Toxicol* **54**:435–456.
- Jin Y, Zhang P, Wang Y, Jin B, Zhou J, Zhang J, and Pan J (2018) Neddylation blockade diminishes hepatic metastasis by dampening cancer stem-like cells and angiogenesis in uveal melanoma. *Clin Cancer Res* **24**:3741–3754.
- Jüdes A, Ebert F, Bär C, Thüring KL, Harter A, Klassen R, Helm M, Stark MJ, and Schaffrath R (2015) Urm1ylation and tRNA thiolation functions of ubiquitin-like Uba4-Urm1 systems are conserved from yeast to man. *FEBS Lett* **589**: 904–909.
- Jung CR, Hwang KS, Yoo J, Cho WK, Kim JM, Kim WH, and Im DS (2006) E2-EPF UCP targets pVHL for degradation and associates with tumor growth and metastasis. *Nat Med* **12**:809–816.
- Kaiser SE, Mao K, Taherbhoy AM, Yu S, Olszewski JL, Duda DM, Kurinov I, Deng A, Fenn TD, Klionsky DJ, et al. (2012) Noncanonical E2 recruitment by the autophagy E1 revealed by Atg7-Atg3 and Atg7-Atg10 structures. *Nat Struct Mol Biol* **19**: 1242–1249.
- Kaiser SE, Qiu Y, Coats JE, Mao K, Klionsky DJ, and Schulman BA (2013) Structures of Atg7-Atg3 and Atg7-Atg10 reveal noncanonical mechanisms of E2 recruitment by the autophagy E1. *Autophagy* **9**:778–780.
- Kapuria V, Peterson LF, Showalter HD, Kirchhoff PD, Talpaz M, and Donato NJ (2011) Protein cross-linking as a novel mechanism of action of a ubiquitin-activating enzyme inhibitor with anti-tumor activity. *Biochem Pharmacol* **82**: 341–349.
- Karvela M, Baquero P, Kuntz EM, Mukhopadhyay A, Mitchell R, Allan EK, Chan E, Kranc KR, Calabretta B, Salomoni P, et al. (2016) ATG7 regulates energy metabolism, differentiation and survival of Philadelphia-chromosome-positive cells. *Autophagy* **12**:936–948.
- Kathawala RJ, Espitia CM, Jones TM, Islam S, Gupta P, Zhang YK, Chen ZS, Carew JS, and Nawrocki ST (2020) ABCG2 overexpression contributes to pevonedistat resistance. *Cancers (Basel)* **12**:429.
- Kee Y, Huang M, Chang S, Moreau LA, Park E, Smith PG, and D'Andrea AD (2012) Inhibition of the Nedd8 system sensitizes cells to DNA interstrand cross-linking agents. *Mol Cancer Res* **10**:369–377.
- Kessler JD, Kahle KT, Sun T, Meerbreij KL, Schlabach MR, Schmitt EM, Skinner SO, Xu Q, Li MZ, Hartman ZC, et al. (2012) A SUMOylation-dependent transcriptional subprogram is required for Myc-driven tumorigenesis. *Science* **335**:348–353.

- Ketscher L, Hannß R, Morales DJ, Basters A, Guerra S, Goldmann T, Hausmann A, Prinz M, Naumann R, Pekosz A, et al. (2015) Selective inactivation of USP18 isopeptidase activity in vivo enhances ISG15 conjugation and viral resistance. *Proc Natl Acad Sci USA* **112**:1577–1582.
- Khalife J, Radomska HS, Santhanam R, Huang X, Neviani P, Saultz J, Wang H, Wu YZ, Alachkar H, Anghelina M, et al. (2015) Pharmacological targeting of miR-155 via the NEDD8-activating enzyme inhibitor MLN4924 (Pevonedistat) in FLT3-ITD acute myeloid leukemia. *Leukemia* **29**:1981–1992.
- Khattar M, Song K, Grossman S, Xega K, He X, Idamakanti N, and Huszar D (2019) Abstract 3252: TAK-981: a first in class SUMO inhibitor in Phase 1 trials that promotes dendritic cell activation, antigen-presentation, and T cell priming. *Cancer Res* **79**.
- Kho C, Lee A, Jeong D, Oh JG, Gorski PA, Fish K, Sanchez R, DeVita RJ, Christensen G, Dahl R, et al. (2015) Small-molecule activation of SERCA2a SUMOylation for the treatment of heart failure. *Nat Commun* **6**:7229.
- Kiessling A, Hogrefe C, Erb S, Bobach C, Fuessel S, Wessjohann L, and Seliger B (2009) Expression, regulation and function of the ISGylation system in prostate cancer. *Oncogene* **28**:2606–2620.
- Kim KH, Ha BH, and Kim EE (2018) Structural basis for Ufm1 recognition by UfSP. *FEBS Lett* **592**:263–273.
- Kim KI, Giannakopoulos NV, Virgin HW, and Zhang DE (2004) Interferon-inducible ubiquitin E2, Ubc8, is a conjugating enzyme for protein ISGylation. *Mol Cell Biol* **24**:9592–9600.
- Kim SJ, Hyeon Lee T, Hee Nam S, Kim JH, Oh S, Sook Cho Y, Sup Lee M, Choi S, and Lee PC (2017) Association of Uba6-specific-E2 (USE1) with lung tumorigenesis. *J Natl Cancer Inst* **109**:1–11.
- Kitagaki J, Yang Y, Saavedra JE, Colburn NH, Keefer LK, and Perantoni AO (2009) Nitric oxide prodrug JS-K inhibits ubiquitin E1 and kills tumor cells retaining wild-type p53. *Oncogene* **28**:619–624.
- Kitareewan S, Pitha-Rowe I, Sekula D, Lowrey CH, Nemeth MJ, Golub TR, Freemantle SJ, and Dmitrovsky E (2002) UBE1L is a retinoid target that triggers PML/RAR α degradation and apoptosis in acute promyelocytic leukemia. *Proc Natl Acad Sci USA* **99**:3806–3811.
- Kiziltepe T, Hideshima T, Ishitsuka K, Ocio EM, Raju N, Catley L, Li CQ, Trudel LJ, Yasui H, Vallet S, et al. (2007) JS-K, a GST-activated nitric oxide generator, induces DNA double-strand breaks, activates DNA damage response pathways, and induces apoptosis in vitro and in vivo in human multiple myeloma cells. *Blood* **110**:709–718.
- Klaips CL, Jayaraj GG, and Hartl FU (2018) Pathways of cellular proteostasis in aging and disease. *J Cell Biol* **217**:51–63.
- Knight E Jr. and Cordova B (1991) IFN-induced 15-kDa protein is released from human lymphocytes and monocytes. *J Immunol* **146**:2280–2284.
- Knight E Jr., Fahey D, Cordova B, Hillman M, Kutny R, Reich N, and Blomstrom D (1988) A 15-kDa interferon-induced protein is derived by COOH-terminal processing of a 17-kDa precursor. *J Biol Chem* **263**:4520–4522.
- Knorr KL, Schneider PA, Meng XW, Dai H, Smith BD, Hess AD, Karp JE, and Kaufmann SH (2015) MLN4924 induces Noxa upregulation in acute myelogenous leukemia and synergizes with Bcl-2 inhibitors. *Cell Death Differ* **22**:2133–2142.
- Komatsu M, Chiba T, Tatsumi K, Iemura S, Tanida I, Okazaki N, Ueno T, Kominami E, Natsume T, and Tanaka K (2004) A novel protein-conjugating system for Ufm1, a ubiquitin-fold modifier. *EMBO J* **23**:1977–1986.
- Krug RM, Zhao C, and Beaudenon S (2005) Properties of the ISG15 E1 enzyme UBE1L. *Methods Enzymol* **398**:32–40.
- Kulkarni M and Smith HE (2008) E1 ubiquitin-activating enzyme UBA-1 plays multiple roles throughout C. elegans development. *PLoS Genet* **4**:e1000131.
- Kumar A, Ito A, Hirohama M, Yoshida M, and Zhang KY (2013a) Identification of quinoxalinyloxy biaryl urea as a new class of SUMO activating enzyme 1 inhibitors. *Bioorg Med Chem Lett* **23**:5145–5149.
- Kumar A, Ito A, Hirohama M, Yoshida M, and Zhang KY (2013b) Identification of sumoylation activating enzyme 1 inhibitors by structure-based virtual screening. *J Chem Inf Model* **53**:809–820.
- Kumar A, Ito A, Hirohama M, Yoshida M, and Zhang KY (2016) Identification of new SUMO activating enzyme 1 inhibitors using virtual screening and scaffold hopping. *Bioorg Med Chem Lett* **26**:1218–1223.
- Kuo KL, Ho IL, Shi CS, Wu JT, Lin WC, Tsai YC, Chang HC, Chou CT, Hsu CH, Hsieh JT, et al. (2015) MLN4924, a novel protein neddylation inhibitor, suppresses proliferation and migration of human urothelial carcinoma: in vitro and in vivo studies. *Cancer Lett* **363**:127–136.
- Lake MW, Wuebbens MM, Rajagopalan KV, and Schindelin H (2001) Mechanism of ubiquitin activation revealed by the structure of a bacterial MoeB-MoaD complex. *Nature* **414**:325–329.
- Lallemant-Breitenbach V, Jeanne M, Benhenda S, Nasr R, Lei M, Peres L, Zhou J, Zhu J, Raught B, and de Thé H (2008) Arsenic degrades PML or PML-RAR α through a SUMO-triggered RNF4/ubiquitin-mediated pathway. *Nat Cell Biol* **10**:547–555.
- Lallemant-Breitenbach V, Zhu J, Chen Z, and de Thé H (2012) Curing APL through PML/RAR α degradation by As2O3. *Trends Mol Med* **18**:36–42.
- Lam V, Best SR, Bruss N, Liu T, Rowland TH, Huszar D, and Danilov A (2019) Pharmacologic inhibition of SUMO-activating enzyme (SAE) with TAK-981 augments interferon signaling and regulates T cell differentiation in ex vivo studies of chronic lymphocytic leukemia (CLL). *Blood* **134**:1760.
- Lan H, Tang Z, Jin H, and Sun Y (2016) Neddylation inhibitor MLN4924 suppresses growth and migration of human gastric cancer cells. *Sci Rep* **6**:24218.
- Lavie J, De Belvalet H, Sonon S, Ion AM, Dumon E, Melser S, Lacombe D, Dupuy JW, Lalou C, and Bénard G (2018) Ubiquitin-Dependent Degradation of Mitochondrial Proteins Regulates Energy Metabolism. *Cell Rep* **23**:2852–2863.
- Law ME, Corsino PE, Jahn SC, Davis BJ, Chen S, Patel B, Pham K, Lu J, Sheppard B, Nørgaard P, et al. (2013) Glucocorticoids and histone deacetylase inhibitors cooperate to block the invasiveness of basal-like breast cancer cells through novel mechanisms. *Oncogene* **32**:1316–1329.
- Laxman S, Sutter BM, Wu X, Kumar S, Guo X, Trudgian DC, Mirzaei H, and Tu BP (2013) Sulfur amino acids regulate translational capacity and metabolic homeostasis through modulation of tRNA thiolation. *Cell* **154**:416–429.
- Leclerc GM, Zheng S, Leclerc GJ, DeSalvo J, Swords RT, and Barredo JC (2016) The NEDD8-activating enzyme inhibitor pevonedistat activates the eIF2 α and mTOR pathways inducing UPR-mediated cell death in acute lymphoblastic leukemia. *Leuk Res* **50**:1–10.
- Lee CG, Ren J, Cheong IS, Ban KH, Ooi LL, Yong Tan S, Kan A, Nuchprayoon I, Jin R, Lee KH, et al. (2003) Expression of the FAT10 gene is highly upregulated in hepatocellular carcinoma and other gastrointestinal and gynecological cancers. *Oncogene* **22**:2592–2603.
- Lee I and Schindelin H (2008) Structural insights into E1-catalyzed ubiquitin activation and transfer to conjugating enzymes. *Cell* **134**:268–278.
- Lee IH, Kawai Y, Fergusson MM, Rovira II, Bishop AJ, Motoyama N, Cao L, and Finkel T (2012) Atg7 modulates p53 activity to regulate cell cycle and survival during metabolic stress. *Science* **336**:225–228.
- Lee SK, Anzick SL, Choi JE, Bubendorf L, Guan XY, Jung YK, Kallioniemi OP, Kononen J, Trent JM, Azorsa D, et al. (1999) A nuclear factor, ASC-2, as a cancer-amplified transcriptional coactivator essential for ligand-dependent transactivation by nuclear receptors in vivo. *J Biol Chem* **274**:34283–34293.
- Leidecker O, Matic I, Mahata B, Pion E, and Xirodimas DP (2012) The ubiquitin E1 enzyme Ube1 mediates NEDD8 activation under diverse stress conditions. *Cell Cycle* **11**:1142–1150.
- Leidel S, Pedrioli PG, Bucher T, Brost R, Costanzo M, Schmidt A, Aebersold R, Boone C, Hofmann K, and Peter M (2009) Ubiquitin-related modifier Urm1 acts as a sulphur carrier in thiolation of eukaryotic transfer RNA. *Nature* **458**:228–232.
- Leung CH, Chan DS, Yang H, Abagyan R, Lee SM, Zhu GY, Fong WF, and Ma DL (2011) A natural product-like inhibitor of NEDD8-activating enzyme. *Chem Commun (Camb)* **47**:2511–2513.
- Levine B and Kroemer G (2019) Biological functions of autophagy genes: a disease perspective. *Cell* **176**:11–42.
- Lévy J, Cacheux W, Bara MA, L'Hermitte A, Lepage P, Fraudeau M, Trentesaux C, Lemarchand J, Durand A, Crain AM, et al. (2015) Intestinal inhibition of Atg7 prevents tumour initiation through a microbiome-influenced immune response and suppresses tumour growth. *Nat Cell Biol* **17**:1062–1073.
- Li H, Tan M, Jia L, Wei D, Zhao Y, Chen G, Xu J, Zhao L, Thomas D, Beer DG, et al. (2014a) Inactivation of SAG/RBX2 E3 ubiquitin ligase suppresses KrasG12D-driven lung tumorigenesis. *J Clin Invest* **124**:835–846.
- Li JA, Song C, Rong Y, Kuang T, Wang D, Xu X, Yuan J, Luo K, Qin B, Nowsheen S, et al. (2018) Chk1 inhibitor SCH 900776 enhances the antitumor activity of MLN4924 on pancreatic cancer. *Cell Cycle* **17**:191–199.
- Li L, Wang M, Yu G, Chen P, Li H, Wei D, Zhu J, Xie L, Jia H, Shi J, et al. (2014b) Overactivated neddylation pathway as a therapeutic target in lung cancer. *J Natl Cancer Inst* **106**:djy083.
- Li T, Santockyte R, Yu S, Shen RF, Tekle E, Lee CG, Yang DC, and Chock PB (2011) FAT10 modifies p53 and upregulates its transcriptional activity. *Arch Biochem Biophys* **509**:164–169.
- Li X and Fan Z (2010) The epidermal growth factor receptor antibody cetuximab induces autophagy in cancer cells by downregulating HIF-1 α and Bcl-2 and activating the beclin 1/Vps34 complex. *Cancer Res* **70**:5942–5952.
- Li X, Yokoyama NN, Zhang S, Ding L, Liu HM, Lilly MB, Mercola D, and Zi X (2015) Flavokawain A induces deNEDDylation and Skp2 degradation leading to inhibition of tumorigenesis and cancer progression in the TRAMP transgenic mouse model. *Oncotarget* **6**:41809–41824.
- Li YJ, Du L, Wang J, Vega R, Lee TD, Miao Y, Aldana-Masangkay G, Samuels ER, Li B, Ouyang SX, et al. (2019) Allosteric inhibition of ubiquitin-like modifications by a class of inhibitor of SUMO-activating enzyme. *Cell Chem Biol* **26**:278–288.e6.
- Liang JR, Lingeman E, Luong T, Ahmed S, Muhar M, Nguyen T, Olzmann JA, and Corn JE (2020) A genome-wide ER-phagy screen highlights key roles of mitochondrial metabolism and ER-resident UFMylation. *Cell* **180**:1160–1177.e20.
- Licciardello MP, Müller MK, Dürnberger G, Kerzendorfer C, Boidol B, Trefzer C, Sclerli S, Berg T, Penz T, Schuster M, et al. (2015) NOTCH1 activation in breast cancer confers sensitivity to inhibition of SUMOylation. *Oncogene* **34**:3780–3790.
- Lin JJ, Milhollen MA, Smith PG, Narayanan U, and Dutta A (2010) NEDD8-targeting drug MLN4924 elicits DNA rereplication by stabilizing Cdt1 in S phase, triggering checkpoint activation, apoptosis, and senescence in cancer cells. *Cancer Res* **70**:10310–10320.
- Lin WC, Kuo KL, Shi CS, Wu JT, Hsieh JT, Chang HC, Liao SM, Chou CT, Chiang CK, Chiu WS, et al. (2015) MLN4924, a novel NEDD8-activating enzyme inhibitor, exhibits antitumor activity and enhances cisplatin-induced cytotoxicity in human cervical carcinoma: in vitro and in vivo study. *Am J Cancer Res* **5**:3350–3362.
- Liu C, Nie D, Li J, Du X, Lu Y, Li Y, Zhou J, Jin Y, and Pan J (2018) Antitumor effects of blocking protein neddylation in T3151-BCR-ABL leukemia cells and leukemia stem cells. *Cancer Res* **78**:1522–1536.
- Liu J, Song T, Zhou W, Xing L, Wang S, Ho M, Peng Z, Tai YT, Hideshima T, Anderson KC, et al. (2019) A genome-scale CRISPR-Cas9 screening in myeloma cells identifies regulators of immunomodulatory drug sensitivity. *Leukemia* **33**:171–180.
- Liu J, Wang Y, Song L, Zeng L, Yi W, Liu T, Chen H, Wang M, Ju Z, and Cong YS (2017a) A critical role of DDRGK1 in endoplasmic reticulum homeostasis via regulation of IRE1 α stability. *Nat Commun* **8**:14186.
- Liu M, Li XL, and Hassel BA (2003) Proteasomes modulate conjugation to the ubiquitin-like protein, ISG15. *J Biol Chem* **278**:1594–1602.
- Liu X, Sun L, Gursel DB, Cheng C, Huang S, Rademaker AW, Khan SA, Yin J, and Kiyokawa H (2017b) The non-canonical ubiquitin activating enzyme UBA6 suppresses epithelial-mesenchymal transition of mammary epithelial cells. *Oncotarget* **8**:87480–87493.
- Liu X, Xu Y, Pang Z, Guo F, Qin Q, Yin T, Sang Y, Feng C, Li X, Jiang L, et al. (2015) Knockdown of SUMO-activating enzyme subunit 2 (SAE2) suppresses cancer malignancy and enhances chemotherapy sensitivity in small cell lung cancer. *J Hematol Oncol* **8**:67.

- Liu X, Zhao B, Sun L, Bhuripanyo K, Wang Y, Bi Y, Davuluri RV, Duong DM, Nanavati D, Yin J, et al. (2017c) Orthogonal ubiquitin transfer identifies ubiquitination substrates under differential control by the two ubiquitin activating enzymes. *Nat Commun* **8**:14286.
- Liu Y, Salvador LA, Byeon S, Ying Y, Kwan JC, Law BK, Hong J, and Luesch H (2010) Anticancer activity of largazole, a marine-derived tunable histone deacetylase inhibitor. *J Pharmacol Exp Ther* **335**:351–361.
- Liu Z, Xu X, Li X, Liu S, Simoneau AR, He F, Wu XR, and Zi X (2013) Kava chalcone, flavokawain A, inhibits urothelial tumorigenesis in the UPII-SV40T transgenic mouse model. *Cancer Prev Res (Phila)* **6**:1365–1375.
- Lo-Coco F, Avvisati G, Vignetti M, Thiede C, Orlando SM, Iacobelli S, Ferrara F, Fazi P, Cicconi L, Di Bona E, et al.; Gruppo Italiano Malattie Ematologiche dell'Adulto German-Austrian Acute Myeloid Leukemia Study Group Study Alliance Leukemia (2013) Retinoic acid and arsenic trioxide for acute promyelocytic leukemia. *N Engl J Med* **369**:111–121.
- Lockhart AC, Bauer TM, Aggarwal C, Lee CB, Harvey RD, Cohen RB, Sedarati F, Nip TK, Faessel H, Dash AB, et al. (2019) Phase Ib study of pevonedistat, a NEDD8-activating enzyme inhibitor, in combination with docetaxel, carboplatin and paclitaxel, or gemcitabine, in patients with advanced solid tumors. *Invest New Drugs* **37**:87–97.
- Loeb KR and Haas AL (1992) The interferon-inducible 15-kDa ubiquitin homolog conjugates to intracellular proteins. *J Biol Chem* **267**:7806–7813.
- Lois LM and Lima CD (2005) Structures of the SUMO E1 provide mechanistic insights into SUMO activation and E2 recruitment to E1. *EMBO J* **24**:439–451.
- Lu P, Guo Y, Zhu L, Xia Y, Zhong Y, and Wang Y (2018) A novel NAE/UAE dual inhibitor LP0040 blocks neddylation and ubiquitination leading to growth inhibition and apoptosis of cancer cells. *Eur J Med Chem* **154**:294–304.
- Lu P, Liu X, Yuan X, He M, Wang Y, Zhang Q, and Ouyang PK (2016) Discovery of a novel NEDD8 activating enzyme inhibitor with piperidin-4-amine scaffold by structure-based virtual screening. *ACS Chem Biol* **11**:1901–1907.
- Lu X, Olsen SK, Capili AD, Cisar JS, Lima CD, and Tan DS (2010) Designed semi-synthetic protein inhibitors of Ub/Ubl E1 activating enzymes. *J Am Chem Soc* **132**:1748–1749.
- Luo J, Solimini NL, and Elledge SJ (2009) Principles of cancer therapy: oncogene and non-oncogene addiction. *Cell* **136**:823–837.
- Luo T, Fu J, Xu A, Su B, Ren Y, Li N, Zhu J, Zhao X, Dai R, Cao J, et al. (2016) PSMD10/gankyrin induces autophagy to promote tumor progression through cytoplasmic interaction with ATG7 and nuclear transactivation of ATG7 expression. *Autophagy* **12**:1355–1371.
- Luo Z, Yu G, Lee HW, Li L, Wang L, Yang D, Pan Y, Ding C, Qian J, Wu L, et al. (2012) The Nedd8-activating enzyme inhibitor MLN4924 induces autophagy and apoptosis to suppress liver cancer cell growth. *Cancer Res* **72**:3360–3371.
- Lux MC, Standke LC, and Tan DS (2019) Targeting adenylate-forming enzymes with designed sulfonyladenine inhibitors. *J Antibiot (Tokyo)* **72**:325–349.
- Lv Z, Rickman KA, Yuan L, Williams K, Selvam SP, Woosley AN, Howe PH, Ogretmen B, Smogorzewska A, and Olsen SK (2017a) S. pombe Uba1-Ubc15 structure reveals a novel regulatory mechanism of ubiquitin E2 activity. *Mol Cell* **65**:699–714.e6.
- Lv Z, Williams KM, Yuan L, Atkison JH, and Olsen SK (2018a) Crystal structure of a human ubiquitin E1-ubiquitin complex reveals conserved functional elements essential for activity. *J Biol Chem* **293**:18337–18352.
- Lv Z, Yuan L, Atkison JH, Aldana-Masangkay G, Chen Y, and Olsen SK (2017b) Domain alternation and active site remodeling are conserved structural features of ubiquitin E1. *J Biol Chem* **292**:12089–12099.
- Lv Z, Yuan L, Atkison JH, Williams KM, Vega R, Sessions EH, Divlianska DB, Davies C, Chen Y, and Olsen SK (2018b) Molecular mechanism of a covalent allosteric inhibitor of SUMO E1 activating enzyme. *Nat Commun* **9**:5145.
- Lydeard JR, Schulman BA, and Harper JW (2013) Building and remodelling Cullin-RING E3 ubiquitin ligases. *EMBO Rep* **14**:1050–1061.
- Ma H, Zhuang C, Xu X, Li J, Wang J, Min X, Zhang W, Zhang H, and Miao Z (2017) Discovery of benzothiazole derivatives as novel non-sulfamide NEDD8 activating enzyme inhibitors by target-based virtual screening. *Eur J Med Chem* **133**:174–183.
- Ma R, Ma L, Weng W, Wang Y, Liu H, Guo R, Gao Y, Tu J, Xu TL, Cheng J, et al. (2020) DUSP6 SUMOylation protects cells from oxidative damage via direct regulation of Drp1 dephosphorylation. *Sci Adv* **6**:eaaz0361.
- Mackintosh C, García-Domínguez DJ, Ordóñez JL, Ginel-Picardo A, Smith PG, Sacristán MP, and de Alava E (2013) WEE1 accumulation and deregulation of S-phase proteins mediate MLN4924 potent inhibitory effect on Ewing sarcoma cells. *Oncogene* **32**:1441–1451.
- MacLeod G, Bozek DA, Rajakulendran N, Monteiro V, Ahmadi M, Steinhart Z, Kushiada MM, Yu H, Coutinho FJ, Cavalli FMG, et al. (2019) Genome-wide CRISPR-Cas9 screens expose genetic vulnerabilities and mechanisms of temozolomide sensitivity in glioblastoma stem cells. *Cell Rep* **27**:971–986.e9.
- Majeed S, Aparnathi MK, Song L, Schimmer AD, Tsao MS, Liu G, and Lok BH (2019) Abstract 2699: targeting an ubiquitin-activating enzyme in small-cell lung cancer (SCLC). *Cancer Res* **79**:2699.
- Malakhov MP, Malakhova OA, Kim KI, Ritchie KJ, and Zhang DE (2002) UBP43 (USP18) specifically removes ISG15 from conjugated proteins. *J Biol Chem* **277**:9976–9981.
- Malhab LJ, Descamps S, Delaval B, and Xirodimas DP (2016) The use of the NEDD8 inhibitor MLN4924 (Pevonedistat) in a cyclotherapy approach to protect wild-type p53 cells from MLN4924 induced toxicity. *Sci Rep* **6**:37775.
- Manasachin EE and Orlowski RZ (2017) Proteasome inhibitors in cancer therapy. *Nat Rev Clin Oncol* **14**:417–433.
- Martinez Molina D, Jafari R, Ignatushchenko M, Seki T, Larsson EA, Dan C, Sreekumar L, Cao Y, and Nordlund P (2013) Monitoring drug target engagement in cells and tissues using the cellular thermal shift assay. *Science* **341**:84–87.
- Maruyama T and Noda NN (2018) Autophagy-regulating protease Atg4: structure, function, regulation and inhibition. *J Antibiot Res* **71**:72–78.
- Mashahreh B, Hassouna F, Soudah N, Cohen-Kfir E, Strulovich R, Haitin Y, and Wiener R (2018) Trans-binding of UFM1 to UBA5 stimulates UBA5 homodimerization and ATP binding. *FASEB J* **32**:2794–2802.
- Mathewson N, Toubai T, Kapeles S, Sun Y, Oravec-Wilson K, Tamaki H, Wang Y, Hou G, Sun Y, and Reddy P (2013) Neddylation plays an important role in the regulation of murine and human dendritic cell function. *Blood* **122**:2062–2073.
- Mathewson ND, Fujiwara H, Wu SR, Toubai T, Oravec-Wilson K, Sun Y, Rossi C, Zajac C, Sun Y, and Reddy P (2016) SAG/Rbx2-dependent neddylation regulates T-cell responses. *Am J Pathol* **186**:2679–2691.
- Mattern MR, Eddins MJ, Agarwal S, Sterner DE, and Kodrasov MP (2014) In Resistance to Proteasome Inhibitors in Cancer (Dou QP ed), Springer International Publishing, Switzerland.
- McGrath JP, Jentsch S, and Varshavsky A (1991) UBA 1: an essential yeast gene encoding ubiquitin-activating enzyme. *EMBO J* **10**:227–236.
- McHugh A, Fernandes K, South AP, Mellerio JE, Salas-Alanis JC, Proby CM, Leigh IM, and Saville MK (2018) Preclinical comparison of proteasome and ubiquitin E1 enzyme inhibitors in cutaneous squamous cell carcinoma: the identification of mechanisms of differential sensitivity. *Oncotarget* **9**:20265–20281.
- McLaughlin PM, Helfrich W, Kok K, Mulder M, Hu SW, Brinker MG, Ruiter MH, de Leij LF, and Buys CH (2000) The ubiquitin-activating enzyme E1-like protein in lung cancer cell lines. *Int J Cancer* **85**:871–876.
- McMillin DW, Jacobs HM, Delmore JE, Buon L, Hunter ZR, Monrose V, Yu J, Smith PG, Richardson PG, Anderson KC, et al. (2012) Molecular and cellular effects of NEDD8-activating enzyme inhibition in myeloma. *Mol Cancer Ther* **11**:942–951.
- Merbl Y, Refour P, Patel H, Springer M, and Kirschner MW (2013) Profiling of ubiquitin-like modifications reveals features of mitotic control. *Cell* **152**:1160–1172.
- Milholten MA, Narayanan U, Soucy TA, Veiby PO, Smith PG, and Amidon B (2011) Inhibition of NEDD8-activating enzyme induces rereplication and apoptosis in human tumor cells consistent with deregulating CDT1 turnover. *Cancer Res* **71**:3042–3051.
- Milholten MA, Shi J, Traore T, Huck J, Sappal D, Duffy J, Lightcap E, Ishii Y, Ciavarrri J, Fleming P, et al. (2015) Abstract A164: the small molecule UAE inhibitor TAK-243 (MLN7243) prevents DNA damage repair and reduces cell viability/tumor growth when combined with radiation, carboplatin and docetaxel. *Mol Cancer Ther* **14**:A164.
- Milholten MA, Thomas MP, Narayanan U, Traore T, Riceberg J, Amidon BS, Bence NF, Bolen JB, Brownell J, Dick LR, et al. (2012) Treatment-emergent mutations in NAEβ confer resistance to the NEDD8-activating enzyme inhibitor MLN4924. *Cancer Cell* **21**:388–401.
- Milholten MA, Traore T, Adams-Duffy J, Thomas MP, Berger AJ, Dang L, Dick LR, Garnsey JJ, Koenig E, Langston SP, et al. (2010) MLN4924, a NEDD8-activating enzyme inhibitor, is active in diffuse large B-cell lymphoma models: rationale for treatment of NF-kappaB-dependent lymphoma. *Blood* **116**:1515–1523.
- Misra M, Kuhn M, Lobel M, An H, Statsyuk AV, Sottriffer C, and Schindelin H (2017) Dissecting the specificity of adenosyl sulfamate inhibitors targeting the ubiquitin-activating enzyme. *Structure* **25**:1120–1129.e3.
- Mizushima T, Tatsumi K, Ozaki Y, Kawakami T, Suzuki A, Ogasahara K, Komatsu M, Kominami E, Tanaka K, and Yamane T (2007) Crystal structure of Ufc1, the Ufm1-conjugating enzyme. *Biochem Biophys Res Commun* **362**:1079–1084.
- Mortensen M, Soilleux EJ, Djordjevic G, Tripp R, Lutteropp M, Sadighi-Akha E, Stranks AJ, Glanville J, Knight S, Jacobsen SE, et al. (2011a) The autophagy protein Atg7 is essential for hematopoietic stem cell maintenance. *J Exp Med* **208**:455–467.
- Mortensen M, Watson AS, and Simon AK (2011b) Lack of autophagy in the hematopoietic system leads to loss of hematopoietic stem cell function and dysregulated myeloid proliferation. *Autophagy* **7**:1069–1070.
- Mustachio LM, Lu Y, Kawakami M, Roszik J, Freemantle SJ, Liu X, and Dmitrovsky E (2018) Evidence for the ISG15-specific deubiquitinase USP18 as an antineoplastic target. *Cancer Res* **78**:587–592.
- Nakamura A, Grossman S, Song K, Idamakanti N, Shapiro G, and Huszar D (2019) Abstract 1523: inhibition of SUMOylation by TAK-981 induces antitumor innate immune responses by modulating macrophage and NK cell function through Type I IFN pathway activation. *Cancer Res* **79**:1523.
- Nakatogawa H (2013) Two ubiquitin-like conjugation systems that mediate membrane formation during autophagy. *Essays Biochem* **55**:39–50.
- Nakayama KI and Nakayama K (2006) Ubiquitin ligases: cell-cycle control and cancer. *Nat Rev Cancer* **6**:369–381.
- Nalepa G, Rolfe M, and Harper JW (2006) Drug discovery in the ubiquitin-proteasome system. *Nat Rev Drug Discov* **5**:596–613.
- Narasimhan J, Wang M, Fu Z, Klein JM, Haas AL, and Kim JJ (2005) Crystal structure of the interferon-induced ubiquitin-like protein ISG15. *J Biol Chem* **280**:27356–27365.
- Nawrocki ST, Kelly KR, Smith PG, Espitia CM, Possemato A, Beausoleil SA, Milholten M, Blakemore S, Thomas M, Berger A, et al. (2013) Disrupting protein NEDDylation with MLN4924 is a novel strategy to target cisplatin resistance in ovarian cancer. *Clin Cancer Res* **19**:3577–3590.
- Nawrocki ST, Kelly KR, Smith PG, Keaton M, Carraway H, Sekeres MA, Maciejewski JP, and Carew JS (2015) The NEDD8-activating enzyme inhibitor MLN4924 disrupts nucleotide metabolism and augments the efficacy of cytarabine. *Clin Cancer Res* **21**:439–447.
- Noda NN, Ohsumi Y, and Inagaki F (2009) ATG systems from the protein structural point of view. *Chem Rev* **109**:1587–1598.
- Noda NN, Satoo K, Fujioka Y, Kumeta H, Ogura K, Nakatogawa H, Ohsumi Y, and Inagaki F (2011) Structural basis of Atg8 activation by a homodimeric E1, Atg7. *Mol Cell* **44**:462–475.
- Oka Y, Varmark H, Vitting-Seerup K, Beli P, Waage J, Hakobyan A, Mistrik M, Choudhary C, Rohde M, Bekker-Jensen S, et al. (2014) UBL5 is essential for pre-mRNA splicing and sister chromatid cohesion in human cells. *EMBO Rep* **15**:956–964.

- Olsen SK, Capili AD, Lu X, Tan DS, and Lima CD (2010) Active site remodelling accompanies thioester bond formation in the SUMO E1. *Nature* **463**:906–912.
- Olsen SK and Lima CD (2013) Structure of a ubiquitin E1-E2 complex: insights to E1-E2 thioester transfer. *Mol Cell* **49**:884–896.
- Oweis W, Padala P, Hassouna F, Cohen-Kfir E, Gibbs DR, Todd EA, Berndsen CE, and Wiener R (2016) Trans-binding mechanism of ubiquitin-like protein activation revealed by a UBA5-UFM1 complex. *Cell Rep* **16**:3113–3120.
- Padala P, Oweis W, Mashahreh B, Soudah N, Cohen-Kfir E, Todd EA, Berndsen CE, and Wiener R (2017) Novel insights into the interaction of UBA5 with UFM1 via a UFM1-interacting sequence. *Sci Rep* **7**:508.
- Paiva C, Godbersen JC, Berger A, Brown JR, and Danilov AV (2015) Targeting neddylation induces DNA damage and checkpoint activation and sensitizes chronic lymphocytic leukemia B cells to alkylating agents. *Cell Death Dis* **6**:e1807.
- Paiva C, Godbersen JC, Rowland T, Danilova OV, Danes C, Berger A, and Danilov AV (2017) Pevonedistat, a Nedd8-activating enzyme inhibitor, sensitizes neoplastic B-cells to death receptor-mediated apoptosis. *Oncotarget* **8**:21128–21139.
- Pan WW, Zhou JJ, Yu C, Xu Y, Guo LJ, Zhang HY, Zhou D, Song FZ, and Fan HY (2013) Ubiquitin E3 ligase CRL4(CDT2/DCAF2) as a potential chemotherapeutic target for ovarian surface epithelial cancer. *J Biol Chem* **288**:29680–29691.
- Park SY, Park JW, Lee GW, Li L, and Chun YS (2018) Inhibition of neddylation facilitates cell migration through enhanced phosphorylation of caveolin-1 in PC3 and U373MG cells. *BMC Cancer* **18**:30.
- Pedrioli PG, Leidel S, and Hofmann K (2008) Urm1 at the crossroad of modifications. 'Protein Modifications: Beyond the Usual Suspects' Review Series. *EMBO Rep* **9**:1196–1202.
- Pelzer C, Kassner I, Matentzoglou K, Singh RK, Wollscheid HP, Scheffner M, Schmidtke G, and Groettrup M (2007) UBE1L2, a novel E1 enzyme specific for ubiquitin. *J Biol Chem* **282**:23010–23014.
- Pérez-Benavente B, Nasresfahani AF, and Farrás R (2020) Ubiquitin-regulated cell proliferation and cancer. *Adv Exp Med Biol* **1233**:3–28.
- Petropoulos M, Champeris Tsaniras S, Taraviras S, and Lygerou Z (2019) Replication licensing aberrations, replication stress, and genomic instability. *Trends Biochem Sci* **44**:752–764.
- Petroski MD, Salvesen GS, and Wolf DA (2011) Urm1 couples sulfur transfer to ubiquitin-like protein function in oxidative stress. *Proc Natl Acad Sci USA* **108**:1749–1750.
- Picco G, Petti C, Sassi F, Grillone K, Migliardi G, Rossi T, Isella C, Di Nicolaantonio F, Sarotto I, Sapino A, et al. (2016) Efficacy of NEDD8 pathway inhibition in pre-clinical models of poorly differentiated, clinically aggressive colorectal cancer. *J Natl Cancer Inst* **109**:djw209.
- Pitha-Rowe I, Hassel BA, and Dmitrovsky E (2004a) Involvement of UBE1L in ISG15 conjugation during retinoid-induced differentiation of acute promyelocytic leukemia. *J Biol Chem* **279**:18178–18187.
- Pitha-Rowe I, Petty WJ, Feng Q, Koza-Taylor PH, Dimattia DA, Pinder L, Dragnev KH, Memoli N, Memoli V, Turi T, et al. (2004b) Microarray analyses uncover UBE1L as a candidate target gene for lung cancer chemoprevention. *Cancer Res* **64**:8109–8115.
- Piya S, Kornblau SM, Ruvalo VR, Mu H, Ruvalo PP, McQueen T, Davis RE, Hail N Jr., Kantarjian H, Andreeff M, et al. (2016) Atg7 suppression enhances chemotherapeutic agent sensitivity and overcomes stroma-mediated chemoresistance in acute myeloid leukemia. *Blood* **128**:1260–1269.
- Poli G, Di Fabio R, Ferrante L, Summa V, and Botta M (2017) Largazole analogues as histone deacetylase inhibitors and anticancer agents: an overview of structure-activity relationships. *ChemMedChem* **12**:1917–1926.
- Qiang L, Sample A, Shea CR, Soltani K, Macleod KF, and He YY (2017) Autophagy gene ATG7 regulates ultraviolet radiation-induced inflammation and skin tumorigenesis. *Autophagy* **13**:2086–2103.
- Qin B, Yu J, Nowshen S, Wang M, Tu X, Liu T, Li H, Wang L, and Lou Z (2019) UFL1 promotes histone H4 ufmylation and ATM activation. *Nat Commun* **10**:1242.
- Raasi S, Schmidtke G, de Giuli R, and Groettrup M (1999) A ubiquitin-like protein which is synergistically inducible by interferon-gamma and tumor necrosis factor- α . *Eur J Immunol* **29**:4030–4036.
- Rabellino A, Andreani C, and Scaglioni PP (2020) Roles of ubiquitination and SUMOylation in the regulation of angiogenesis. *Curr Issues Mol Biol* **35**:109–126.
- Radoshevič L, Murrow L, Chen N, Fernandez E, Roy S, Fung C, and Debnath J (2010) ATG12 conjugation to ATG3 regulates mitochondrial homeostasis and cell death. *Cell* **142**:590–600.
- Rahimi N (2012) The ubiquitin-proteasome system meets angiogenesis. *Mol Cancer Ther* **11**:538–548.
- Rape M (2018) Ubiquitylation at the crossroads of development and disease. *Nat Rev Mol Cell Biol* **19**:59–70.
- Ravi R, Mookerjee B, Bhujwala ZM, Sutter CH, Artemov D, Zeng Q, Dillehay LE, Madan A, Semenza GL, and Bedi A (2000) Regulation of tumor angiogenesis by p53-induced degradation of hypoxia-inducible factor 1 α . *Genes Dev* **14**:34–44.
- Rising EM, Boggio R, Chiocca S, Helin K, and Pasini D (2008) The polycomb repressive complex 2 is a potential target of SUMO modifications. *PLoS One* **3**:e2704.
- Roberts AM, Miyamoto DK, Huffman TR, Bateman LA, Ives AN, Akopian D, Heslin MJ, Contreras CM, Rape M, Skibola CF, et al. (2017) Chemoproteomic screening of covalent ligands reveals UBA5 as a novel pancreatic cancer target. *ACS Chem Biol* **12**:899–904.
- Ronau JA, Beckmann JF, and Hochstrasser M (2016) Substrate specificity of the ubiquitin and Ubl proteases. *Cell Res* **26**:441–456.
- Rossman TG, Visalli MA, and Komissarova EV (2003) Fub1 and its ubiquitin-like domain (FUBI) transforms human osteogenic sarcoma (HOS) cells to anchorage-independence. *Oncogene* **22**:1817–1821.
- Rulina AV, Mittler F, Obeid P, Gerbaud S, Guyon L, Sulpice E, Kermarrec F, Assard N, Dolega ME, Gidrol X, et al. (2016) Distinct outcomes of CRL-Nedd8 pathway inhibition reveal cancer cell plasticity. *Cell Death Dis* **7**:e2505.
- Ruschak AM, Slassi M, Kay LE, and Schimmer AD (2011) Novel proteasome inhibitors to overcome bortezomib resistance. *J Natl Cancer Inst* **103**:1007–1017.
- Sadler AJ and Williams BR (2008) Interferon-inducible antiviral effectors. *Nat Rev Immunol* **8**:559–568.
- Sainz B Jr., Martin B, Tatari M, Heesch C, and Guerra S (2014) ISG15 is a critical microenvironmental factor for pancreatic cancer stem cells. *Cancer Res* **74**:7309–7320.
- Santanam U, Banach-Petrosky W, Abate-Shen C, Shen MM, White E, and DiPaola RS (2016) Atg7 cooperates with Pten loss to drive prostate cancer tumor growth. *Genes Dev* **30**:399–407.
- Sanz MA, Fenaux P, Tallman MS, Estey EH, Löwenberg B, Naoe T, Lengfelder E, Döhner H, Burnett AK, Chen SJ, et al. (2019) Management of acute promyelocytic leukemia: updated recommendations from an expert panel of the European LeukemiaNet. *Blood* **133**:1630–1643.
- Sarantopoulos J, Shapiro GI, Cohen RB, Clark JW, Kauh JS, Weiss GJ, Cleary JM, Mahalingam D, Pickard MD, Faessel HM, et al. (2016) Phase I study of the investigational NEDD8-activating enzyme inhibitor pevonedistat (TAK-924/MLN4924) in patients with advanced solid tumors. *Clin Cancer Res* **22**:847–857.
- Sarge KD and Park-Sarge OK (2009) Detection of proteins sumoylated in vivo and in vitro. *Methods Mol Biol* **590**:265–277.
- Schapiro M, Calabrese MF, Bullock AN, and Crews CM (2019) Targeted protein degradation: expanding the toolbox. *Nat Rev Drug Discov* **18**:949–963.
- Schelpel J, Monté D, Dewitte F, Sixma TK, and Rucktooa P (2016) Structure of UBE2Z enzyme provides functional insight into specificity in the FAT10 protein conjugation machinery. *J Biol Chem* **291**:630–639.
- Schilsky RL, Doroshov JH, Leblanc M, and Conley BA (2012) Development and use of integral assays in clinical trials. *Clin Cancer Res* **18**:1540–1546.
- Schlieker CD, Van der Veen AG, Damon JR, Spooner E, and Ploegh HL (2008) A functional proteomics approach links the ubiquitin-related modifier Urm1 to a tRNA modification pathway. *Proc Natl Acad Sci USA* **105**:18255–18260.
- Schmidtke G, Aiche M, and Groettrup M (2014) FAT10ylation as a signal for proteasomal degradation. *Biochim Biophys Acta* **1843**:97–102.
- Schmitz J, Chowdhury MM, Hänzelmann P, Mintz M, Lee EY, Schindelin H, and Leimkühler S (2008) The sulfurtransferase activity of Uba4 presents a link between ubiquitin-like protein conjugation and activation of sulfur-carrier proteins. *Biochemistry* **47**:6479–6489.
- Schneider WM, Chevillotte MD, and Rice CM (2014) Interferon-stimulated genes: a complex web of host defenses. *Annu Rev Immunol* **32**:513–545.
- Schulman BA and Harper JW (2009) Ubiquitin-like protein activation by E1 enzymes: the apex for downstream signalling pathways. *Nat Rev Mol Cell Biol* **10**:319–331.
- Schwartz DC and Hochstrasser M (2003) A superfamily of protein tags: ubiquitin, SUMO and related modifiers. *Trends Biochem Sci* **28**:321–328.
- Schwertman P, Bekker-Jensen S, and Mailand N (2016) Regulation of DNA double-strand break repair by ubiquitin and ubiquitin-like modifiers. *Nat Rev Mol Cell Biol* **17**:379–394.
- Seeler JS and Dejean A (2003) Nuclear and unclear functions of SUMO. *Nat Rev Mol Cell Biol* **4**:690–699.
- Seeler JS and Dejean A (2017) SUMO and the robustness of cancer. *Nat Rev Cancer* **17**:184–197.
- Sekizawa R, Ikeno S, Nakamura H, Naganawa H, Matsui S, Iinuma H, and Takeuchi T (2002) Panepophenanthrin, from a mushroom strain, a novel inhibitor of the ubiquitin-activating enzyme. *J Nat Prod* **65**:1491–1493.
- Senft D, Qi J, and Ronai ZA (2018) Ubiquitin ligases in oncogenic transformation and cancer therapy. *Nat Rev Cancer* **18**:69–88.
- Shah JJ, Jakubowski AJ, O'Connor OA, Orlowski RZ, Harvey RD, Smith MR, Lebovic D, Diefenbach C, Kelly K, Hua Z, et al. (2016) Phase I study of the novel investigational NEDD8-activating enzyme inhibitor pevonedistat (MLN4924) in patients with relapsed/refractory multiple myeloma or lymphoma. *Clin Cancer Res* **22**:34–43.
- Shang F and Taylor A (2011) Ubiquitin-proteasome pathway and cellular responses to oxidative stress. *Free Radic Biol Med* **51**:5–16.
- Shao DF, Wang XH, Li ZY, Xing XF, Cheng XJ, Guo T, Du H, Hu Y, Dong B, Ding N, et al. (2014) High-level SAE2 promotes malignant phenotype and predicts outcome in gastric cancer. *Am J Cancer Res* **5**:140–154.
- Simons M, Gordon E, and Claesson-Welsh L (2016) Mechanisms and regulation of endothelial VEGF receptor signalling. *Nat Rev Mol Cell Biol* **17**:611–625.
- Simsek D, Tiu GC, Flynn RA, Byeon GW, Leppke K, Xu AF, Chang HY, and Barna M (2017) The mammalian ribo-interactome reveals ribosome functional diversity and heterogeneity. *Cell* **169**:1051–1065.e18.
- Skaug B, Jiang X, and Chen ZJ (2009) The role of ubiquitin in NF-kappaB regulatory pathways. *Annu Rev Biochem* **78**:769–796.
- Smith MA, Maris JM, Gorlick R, Kolb EA, Lock R, Carol H, Keir ST, Reynolds CP, Kang MH, Morton CL, et al. (2012) Initial testing of the investigational NEDD8-activating enzyme inhibitor MLN4924 by the pediatric preclinical testing program. *Pediatr Blood Cancer* **59**:246–253.
- Smith PG, Traore T, Grossman S, Narayanan U, Carew JS, Lublinksky A, Kuranda M, and Milhollen M (2011) Azacitidine/decitabine synergism with the NEDD8-activating enzyme inhibitor MLN4924 in pre-clinical AML models. *Blood* **118**:578.
- Soucy TA, Dick LR, Smith PG, Milhollen MA, and Brownell JE (2010) The NEDD8 conjugation pathway and its relevance in cancer biology and therapy. *Genes Cancer* **1**:708–716.
- Soucy TA, Smith PG, Milhollen MA, Berger AJ, Gavin JM, Adhikari S, Brownell JE, Burke KE, Cardin DP, Critchley S, et al. (2009a) An inhibitor of NEDD8-activating enzyme as a new approach to treat cancer. *Nature* **458**:732–736.
- Soucy TA, Smith PG, and Rolfe M (2009b) Targeting NEDD8-activated cullin-RING ligases for the treatment of cancer. *Clin Cancer Res* **15**:3912–3916.
- Soudah N, Padala P, Hassouna F, Kumar M, Mashahreh B, Lebedev AA, Isupov MN, Cohen-Kfir E, and Wiener R (2019) An N-terminal extension to UBA5 adenylation

- domain boosts UFM1 activation: isoform-specific differences in ubiquitin-like protein activation. *J Mol Biol* **431**:463–478.
- Stewart MD, Ritterhoff T, Klevit RE, and Brzovic PS (2016) E2 enzymes: more than just middle men. *Cell Res* **26**:423–440.
- Stindt MH, Carter S, Vigneron AM, Ryan KM, and Vousden KH (2011) MDM2 promotes SUMO-2/3 modification of p53 to modulate transcriptional activity. *Cell Cycle* **10**:3176–3188.
- Streich FC Jr. and Lima CD (2014) Structural and functional insights to ubiquitin-like protein conjugation. *Annu Rev Biophys* **43**:357–379.
- Strohecker AM, Guo JY, Karsli-Uzunbas G, Price SM, Chen GJ, Mathew R, McMahon M, and White E (2013) Autophagy sustains mitochondrial glutamine metabolism and growth of Brav600E-driven lung tumors. *Cancer Discov* **3**:1272–1285.
- Strohecker AM and White E (2014) Autophagy promotes Brav600E-driven lung tumorigenesis by preserving mitochondrial metabolism. *Autophagy* **10**:384–385.
- Sumi H, Inazuka M, Morimoto M, Hibino R, Hashimoto K, Ishikawa T, Kuida K, Smith PG, Yoshida S, and Yabuki M (2016) An inhibitor of apoptosis protein antagonist T-3256336 potentiates the antitumor efficacy of the Nedd8-activating enzyme inhibitor pevonedistat (TAK-924/MLN4924). *Biochem Biophys Res Commun* **480**:380–386.
- Sung B, Pandey MK, Ahn KS, Yi T, Chaturvedi MM, Liu M, and Aggarwal BB (2008) Anacardic acid (6-nonadecyl salicylic acid), an inhibitor of histone acetyltransferase, suppresses expression of nuclear factor-kappaB-regulated gene products involved in cell survival, proliferation, invasion, and inflammation through inhibition of the inhibitory subunit of nuclear factor-kappaB kinase, leading to potentiation of apoptosis. *Blood* **111**:4880–4891.
- Suzawa M, Miranda DA, Ramos KA, Ang KK, Faivre EJ, Wilson CG, Caboni L, Arkin MR, Kim YS, Fletterick RJ, et al. (2015) A gene-expression screen identifies a non-toxic sumoylation inhibitor that mimics SUMO-less human LRH-1 in liver. *eLife* **4**:e09003.
- Suzuki H, Osawa T, Fujioka Y, and Noda NN (2017) Structural biology of the core autophagy machinery. *Curr Opin Struct Biol* **43**:10–17.
- Swatek KN and Komander D (2016) Ubiquitin modifications. *Cell Res* **26**:399–422.
- Swords RT, Coutre S, Maris MB, Zeidner JF, Foran JM, Cruz J, Erba HP, Berdeja JG, Tam W, Vardhanabhatu S, et al. (2018) Pevonedistat, a first-in-class NEDD8-activating enzyme inhibitor, combined with azacitidine in patients with AML. *Blood* **131**:1415–1424.
- Swords RT, Erba HP, DeAngelo DJ, Bixby DL, Altman JK, Maris M, Hua Z, Blakemore SJ, Faessel H, Sedarati F, et al. (2015) Pevonedistat (MLN4924), a First-in-Class NEDD8-activating enzyme inhibitor, in patients with acute myeloid leukaemia and myelodysplastic syndromes: a phase 1 study. *Br J Haematol* **169**:534–543.
- Swords RT, Kelly KR, Smith PG, Garnsey JJ, Mahalingam D, Medina E, Oberheu K, Padmanabhan S, O'Dwyer M, Nawrocki ST, et al. (2010) Inhibition of NEDD8-activating enzyme: a novel approach for the treatment of acute myeloid leukemia. *Blood* **115**:3796–3800.
- Swords RT, Watts J, Erba HP, Altman JK, Maris M, Anwer F, Hua Z, Stein H, Faessel H, Sedarati F, et al. (2017) Expanded safety analysis of pevonedistat, a first-in-class NEDD8-activating enzyme inhibitor, in patients with acute myeloid leukemia and myelodysplastic syndromes. *Blood Cancer J* **7**:e520.
- Szczepanowski RH, Filipek R, and Bochtler M (2005) Crystal structure of a fragment of mouse ubiquitin-activating enzyme. *J Biol Chem* **280**:22006–22011.
- Taheribhoy AM, Tait SW, Kaiser SE, Williams AH, Deng A, Nourse A, Hammel M, Kurinov I, Rock CO, Green DR, et al. (2011) Atg8 transfer from Atg7 to Atg3: a distinctive E1-E2 architecture and mechanism in the autophagy pathway. *Mol Cell* **44**:451–461.
- Takamura A, Komatsu M, Hara T, Sakamoto A, Kishi C, Waguri S, Eishi Y, Hino O, Tanaka K, and Mizushima N (2011) Autophagy-deficient mice develop multiple liver tumors. *Genes Dev* **25**:795–800.
- Takemoto M, Kawamura Y, Hirohama M, Yamaguchi Y, Handa H, Saitoh H, Nakao Y, Kawada M, Khalid K, Koshino H, et al. (2014) Inhibition of protein SUMOylation by davidin, an ellagitannin from *Davidia involucrata*. *J Antibiot (Tokyo)* **67**:335–338.
- Takeuchi T, Iwahara S, Saeki Y, Sasajima H, and Yokosawa H (2005) Link between the ubiquitin conjugation system and the ISG15 conjugation system: ISG15 conjugation to the UbcH6 ubiquitin E2 enzyme. *J Biochem* **138**:711–719.
- Tan G, Qiu M, Chen L, Zhang S, Ke L, and Liu J (2017) JS-K, a nitric oxide pro-drug, regulates growth and apoptosis through the ubiquitin-proteasome pathway in prostate cancer cells. *BMC Cancer* **17**:376.
- Tan M, Li H, and Sun Y (2014) Endothelial deletion of Sag/Rbx2/Roc2 E3 ubiquitin ligase causes embryonic lethality and blocks tumor angiogenesis. *Oncogene* **33**:5211–5220.
- Tan M, Li Y, Yang R, Xi N, and Sun Y (2011) Inactivation of SAG E3 ubiquitin ligase blocks embryonic stem cell differentiation and sensitizes leukemia cells to retinoid acid. *PLoS One* **6**:e27726.
- Taori K, Paul VJ, and Luesch H (2008) Structure and activity of largazole, a potent antiproliferative agent from the Floridian marine cyanobacterium *Symploca* sp. *J Am Chem Soc* **130**:1806–1807.
- Tate JG, Bamford S, Jubb HC, Sondka Z, Beare DM, Bindal N, Boutselakis H, Cole CG, Creatore C, Dawson E, et al. (2019) COSMIC: the catalogue of somatic mutations in cancer. *Nucleic Acids Res* **47** (D1):D941–D947.
- Tatham MH, Geoffroy MC, Shen L, Plechanovova A, Hattersley N, Jaffray EG, Palvimo JJ, and Hay RT (2008) RNF4 is a poly-SUMO-specific E3 ubiquitin ligase required for arsenite-induced PML degradation. *Nat Cell Biol* **10**:538–546.
- Tatsumi K, Sou YS, Tada N, Nakamura E, Temura S, Natsume T, Kang SH, Chung CH, Kasahara M, Kominami E, et al. (2010) A novel type of E3 ligase for the Ufm1 conjugation system. *J Biol Chem* **285**:5417–5427.
- Tatsumi K, Yamamoto-Mukai H, Shimizu R, Waguri S, Sou YS, Sakamoto A, Taya C, Shitara H, Hara T, Chung CH, et al. (2011) The Ufm1-activating enzyme Uba5 is indispensable for erythroid differentiation in mice. *Nat Commun* **2**:181.
- Teixeira LK and Reed SI (2013) Ubiquitin ligases and cell cycle control. *Annu Rev Biochem* **82**:387–414.
- Terai H, Kitajima S, Potter DS, Matsui Y, Quiceno LG, Chen T, Kim TJ, Rusan M, Thai TC, Piccioni F, et al. (2018) ER stress signaling promotes the survival of cancer “Persister Cells” tolerant to EGFR tyrosine kinase inhibitors. *Cancer Res* **78**:1044–1057.
- Termahe M and Leidel SA (2018) The Uba4 domain interplay is mediated via a thioester that is critical for tRNA thiolation through Urm1 thioacetylation. *Nucleic Acids Res* **46**:5171–5181.
- Thapa P, Shanmugam N, and Pokrzywa W (2020) Ubiquitin signaling regulates RNA biogenesis, processing, and metabolism. *BioEssays* **42**:e1900171.
- Theng SS, Wang W, Mah WC, Chan C, Zhuo J, Gao Y, Qin H, Lim L, Chong SS, Song J, et al. (2014) Disruption of FAT10-MAD2 binding inhibits tumor progression. *Proc Natl Acad Sci USA* **111**:E5282–E5291.
- Tong S, Si Y, Yu H, Zhang L, Xie P, and Jiang W (2017) MLN4924 (Pevonedistat), a protein neddylation inhibitor, suppresses proliferation and migration of human clear cell renal cell carcinoma. *Sci Rep* **7**:5599.
- Toth JI, Yang L, Dahl R, and Petroski MD (2012) A gatekeeper residue for NEDD8-activating enzyme inhibition by MLN4924. *Cell Rep* **1**:309–316.
- Tsherniak A, Vazquez F, Montgomery PG, Weir BA, Kryukov G, Cowley GS, Gill S, Harrington WF, Pantel S, Krill-Burger JM, et al. (2017) Defining a cancer dependency map. *Cell* **170**:564–576.e16.
- Tsukamoto S, Hirota H, Imachi M, Fujimuro M, Onuki H, Ohta T, and Yokosawa H (2005) Himeic acid A: a new ubiquitin-activating enzyme inhibitor isolated from a marine-derived fungus, *Aspergillus* sp. *Bioorg Med Chem Lett* **15**:191–194.
- Uchida C, Miwa S, Kitagawa K, Hattori T, Isobe T, Otani S, Oda T, Sugimura H, Kamijo T, Okawa K, et al. (2005) Enhanced Mdm2 activity inhibits pRB function via ubiquitin-dependent degradation. *EMBO J* **24**:160–169.
- Uckelmann M and Sixma TK (2017) Histone ubiquitination in the DNA damage response. *DNA Repair (Amst)* **56**:92–101.
- Uhlen M, Zhang C, Lee S, Sjostedt E, Fagerberg L, Bidkhori G, Benfais R, Arif M, Liu Z, Edfors F, et al. (2017) A pathology atlas of the human cancer transcriptome. *Science* **357**:eaan2507.
- Ulrich HD (2014) Two-way communications between ubiquitin-like modifiers and DNA. *Nat Struct Mol Biol* **21**:317–324.
- Ulrich HD and Walden H (2010) Ubiquitin signalling in DNA replication and repair. *Nat Rev Mol Cell Biol* **11**:479–489.
- Ungermannova D, Parker SJ, Nasveschuk CG, Chapnick DA, Phillips AJ, Kuchta RD, and Liu X (2012a) Identification and mechanistic studies of a novel ubiquitin E1 inhibitor. *J Biomol Screen* **17**:421–434.
- Ungermannova D, Parker SJ, Nasveschuk CG, Wang W, Quade B, Zhang G, Kuchta RD, Phillips AJ, and Liu X (2012b) Largazole and its derivatives selectively inhibit ubiquitin activating enzyme (e1). *PLoS One* **7**:e29208.
- UniProt Consortium (2019) UniProt: a worldwide hub of protein knowledge. *Nucleic Acids Res* **47** (D1):D506–D515.
- van der Veen AG and Ploegh HL (2012) Ubiquitin-like proteins. *Annu Rev Biochem* **81**:323–357.
- Van der Veen AG, Schorpp K, Schlieker C, Buti L, Damon JR, Spooner E, Ploegh HL, and Jentsch S (2011) Role of the ubiquitin-like protein Urm1 as a noncanonical lysine-directed protein modifier. *Proc Natl Acad Sci USA* **108**:1763–1770.
- Vanderdys V, Allak A, Guessous F, Benamar M, Read PW, Jameson MJ, and Abbas T (2018) The neddylation inhibitor pevonedistat (MLN4924) suppresses and radiosensitizes head and neck squamous carcinoma cells and tumors. *Mol Cancer Ther* **17**:368–380.
- Varshavsky A (2017) The ubiquitin system, autophagy, and regulated protein degradation. *Annu Rev Biochem* **86**:123–128.
- Verma S, Singh A, and Mishra A (2014) Molecular dynamics investigation on the poor sensitivity of A171T mutant NEDD8-activating enzyme (NAE) for MLN4924. *J Biomol Struct Dyn* **32**:1064–1073.
- Villarroya-Beltri C, Guerra S, and Sánchez-Madrid F (2017) ISGylation - a key to lock the cell gates for preventing the spread of threats. *J Cell Sci* **130**:2961–2969.
- Visconte V, Nawrocki ST, Espitia CM, Kelly KR, Possemato A, Beausoleil SA, Han Y, Carraway HE, Nazha A, Advani AS, et al. (2016) Comprehensive quantitative proteomic profiling of the pharmacodynamic changes induced by MLN4924 in acute myeloid leukemia cells establishes rationale for its combination with azacitidine. *Leukemia* **30**:1190–1194.
- Visscher M, Arkin MR, and Dansen TB (2016) Covalent targeting of acquired cysteines in cancer. *Curr Opin Chem Biol* **30**:61–67.
- Vlieghe P, Lisowski V, Martinez J, and Khrestchatsky M (2010) Synthetic therapeutic peptides: science and market. *Drug Discov Today* **15**:40–56.
- Vucic D, Dixit VM, and Wertz IE (2011) Ubiquitylation in apoptosis: a post-translational modification at the edge of life and death. *Nat Rev Mol Cell Biol* **12**:439–452.
- Walczak CP, Leto DE, Zhang L, Riepe C, Muller RY, DaRosa PA, Ingolia NT, Elias JE, and Kopito RR (2019) Ribosomal protein RPL26 is the principal target of UFMylation. *Proc Natl Acad Sci USA* **116**:1299–1308.
- Walden H, Podgorski MS, Huang DT, Miller DW, Howard RJ, Minor DL Jr., Holton JM, and Schulman BA (2003a) The structure of the APPBP1-UBA3-NEDD8-ATP complex reveals the basis for selective ubiquitin-like protein activation by an E1. *Mol Cell* **12**:1427–1437.
- Walden H, Podgorski MS, and Schulman BA (2003b) Insights into the ubiquitin transfer cascade from the structure of the activating enzyme for NEDD8. *Nature* **422**:330–334.
- Wang F, Liu M, Qiu R, and Ji C (2011) The dual role of ubiquitin-like protein Urm1 as a protein modifier and sulfur carrier. *Protein Cell* **2**:612–619.
- Wang J, Hu W, Cai S, Lee B, Song J, and Chen Y (2007) The intrinsic affinity between E2 and the Cys domain of E1 in ubiquitin-like modifications. *Mol Cell* **27**:228–237.
- Wang J, Lee B, Cai S, Fukui L, Hu W, and Chen Y (2009) Conformational transition associated with E1-E2 interaction in small ubiquitin-like modifications. *J Biol Chem* **284**:20340–20348.
- Wang J, Wang S, Zhang W, Wang X, Liu X, Liu L, Li L, Liang Y, Yu J, Jeong LS, et al. (2017) Targeting neddylation pathway with MLN4924 (Pevonedistat) induces

- NOXA-dependent apoptosis in renal cell carcinoma. *Biochem Biophys Res Commun* **490**:1183–1188.
- Wang L, Xu Y, Rogers H, Saidi L, Noguchi CT, Li H, Yewdell JW, Guydosh NR, and Ye Y (2020) UFMylation of RPL26 links translocation-associated quality control to endoplasmic reticulum protein homeostasis. *Cell Res* **30**:5–20.
- Wang M and Kaufman RJ (2014) The impact of the endoplasmic reticulum protein-folding environment on cancer development. *Nat Rev Cancer* **14**:581–597.
- Wang X, Zhang W, Yan Z, Liang Y, Li L, Yu X, Feng Y, Fu S, Zhang Y, Zhao H, et al. (2016) Radiosensitization by the investigational NEDD8-activating enzyme inhibitor MLN4924 (pevonedistat) in hormone-resistant prostate cancer cells. *Oncotarget* **7**:38380–38391.
- Wang Y, Luo Z, Pan Y, Wang W, Zhou X, Jeong LS, Chu Y, Liu J, and Jia L (2015) Targeting protein neddylation with an NEDD8-activating enzyme inhibitor MLN4924 induced apoptosis or senescence in human lymphoma cells. *Cancer Biol Ther* **16**:420–429.
- Wang Y, Ma J, Chow SC, Li CH, Xiao Z, Feng R, Fu J, and Chen Y (2014) A potential antitumor ellagitannin, davidin, inhibited hepatocellular tumor growth by targeting EZH2. *Tumour Biol* **35**:205–212.
- Wang Z, Gong Y, Peng B, Shi R, Fan D, Zhao H, Zhu M, Zhang H, Lou Z, Zhou J, et al. (2019) MRE11 UFMylation promotes ATM activation. *Nucleic Acids Res* **47**:4124–4135.
- Watson AS, Mortensen M, and Simon AK (2011a) Autophagy in the pathogenesis of myelodysplastic syndrome and acute myeloid leukemia. *Cell Cycle* **10**:1719–1725.
- Watson IR, Irwin MS, and Ohh M (2011b) NEDD8 pathways in cancer, Sine Quibus Non. *Cancer Cell* **19**:168–176.
- Wei D, Li H, Yu J, Sebolt JT, Zhao L, Lawrence TS, Smith PG, Morgan MA, and Sun Y (2012) Radiosensitization of human pancreatic cancer cells by MLN4924, an investigational NEDD8-activating enzyme inhibitor. *Cancer Res* **72**:282–293.
- Wei LY, Wu ZX, Yang Y, Zhao M, Ma XY, Li JS, Yang DH, Chen ZS, and Fan YF (2020) Overexpression of ABCG2 confers resistance to pevonedistat, an NAE inhibitor. *Exp Cell Res* **388**:111858.
- Wei Y and Xu X (2016) UFMylation: a unique & fashionable modification for life. *Genomics Proteomics Bioinformatics* **14**:140–146.
- Weinstein IB and Joe AK (2006) Mechanisms of disease: oncogene addiction—a rationale for molecular targeting in cancer therapy. *Nat Clin Pract Oncol* **3**:448–457.
- Welchman RL, Gordon C, and Mayer RJ (2005) Ubiquitin and ubiquitin-like proteins as multifunctional signals. *Nat Rev Mol Cell Biol* **6**:599–609.
- Wen X and Klionsky DJ (2019) At a glance: a history of autophagy and cancer. *Semin Cancer Biol*.
- Wilkinson KA and Henley JM (2010) Mechanisms, regulation and consequences of protein SUMOylation. *Biochem J* **428**:133–145.
- Wilkinson KD, Smith SE, O'Connor L, Sternberg E, Taggart JJ, Berges DA, and Butt T (1990) A specific inhibitor of the ubiquitin activating enzyme: synthesis and characterization of adenyl-phospho-ubiquitinol, a nonhydrolyzable ubiquitin adenylate analogue. *Biochemistry* **29**:7373–7380.
- Wilson DJ and Aldrich CC (2010) A continuous kinetic assay for adenylation enzyme activity and inhibition. *Anal Biochem* **404**:56–63.
- Wong KM, Micel LN, Selby HM, Tan AC, Pitts TM, Bagby SM, Spreafico A, Klauck PJ, Blakemore SJ, Smith PF, et al. (2017) Targeting the protein ubiquitination machinery in melanoma by the NEDD8-activating enzyme inhibitor pevonedistat (MLN4924). *Invest New Drugs* **35**:11–25.
- Wu HQ, Baker D, and Ovaa H (2020) Small molecules that target the ubiquitin system. *Biochem Soc Trans* **48**:479–497.
- Wu LC, Wen ZS, Qiu YT, Chen XQ, Chen HB, Wei MM, Liu Z, Jiang S, and Zhou GB (2013) Largazole arrests cell cycle at G1 phase and triggers proteasomal degradation of E2F1 in lung cancer cells. *ACS Med Chem Lett* **4**:921–926.
- Xie P, Yang JP, Cao Y, Peng LX, Zheng LS, Sun R, Meng DF, Wang MY, Mei Y, Qiang YY, et al. (2017) Promoting tumorigenesis in nasopharyngeal carcinoma, NEDD8 serves as a potential theranostic target. *Cell Death Dis* **8**:e2834.
- Xie X, Koh JY, Price S, White E, and Mehnert JM (2015) Atg7 overcomes senescence and promotes growth of Brav600E-driven melanoma. *Cancer Discov* **5**:410–423.
- Xiong J (2015) Atg7 in development and disease: panacea or Pandora's Box? *Protein Cell* **6**:722–734.
- Xu B, Deng Y, Bi R, Guo H, Shu C, Shah NK, Chang J, Liu G, Du Y, Wei W, et al. (2018a) A first-in-class inhibitor, MLN4924 (pevonedistat), induces cell-cycle arrest, senescence, and apoptosis in human renal cell carcinoma by suppressing UBE2M-dependent neddylation modification. *Cancer Chemother Pharmacol* **81**:1083–1093.
- Xu GW, Ali M, Wood TE, Wong D, Maclean N, Wang X, Gronda M, Skrtic M, Li X, Hurren R, et al. (2010) The ubiquitin-activating enzyme E1 as a therapeutic target for the treatment of leukemia and multiple myeloma. *Blood* **115**:2251–2259.
- Xu GW, Toth JI, da Silva SR, Paiva SL, Lukkari J, Hurren R, Maclean N, Sukhai MA, Bhattacharjee RN, Goard CA, et al. (2014) Mutations in UBA3 confer resistance to the NEDD8-activating enzyme inhibitor MLN4924 in human leukemic cells [published correction appears in *PLoS One* (2014) **9**:e12004]. *PLoS One* **9**:e93530.
- Xu J, Zhang J, Wang L, Zhou J, Huang H, Wu J, Zhong Y, and Shi Y (2006) Solution structure of Urm1 and its implications for the origin of protein modifiers. *Proc Natl Acad Sci USA* **103**:11625–11630.
- Xu Q, Lin G, Xu H, Hu L, Wang Y, Du S, Deng W, Hu W, Cheng W, and Jiang K (2018b) MLN4924 neddylation inhibitor promotes cell death in paclitaxel-resistant human lung adenocarcinoma cells. *Oncol Lett* **15**:515–521.
- Yamaguchi M, Matoba K, Sawada R, Fujioka Y, Nakatogawa H, Yamamoto H, Kobashigawa Y, Hoshida H, Akada R, Ohsumi Y, et al. (2012) Noncanonical recognition and UBL loading of distinct E2s by autophagy-essential Atg7. *Nat Struct Mol Biol* **19**:1250–1256.
- Yamaguchi M, Satoo K, Suzuki H, Fujioka Y, Ohsumi Y, Inagaki F, and Noda NN (2018) Atg7 activates an autophagy-essential ubiquitin-like protein Atg8 through multi-step recognition. *J Mol Biol* **430**:249–257.
- Yamanokuchi R, Imada K, Miyazaki M, Kato H, Watanabe T, Fujimuro M, Saeki Y, Yoshinaga S, Terasawa H, Iwasaki N, et al. (2012) Hyrtioreticulins A-E, indole alkaloids inhibiting the ubiquitin-activating enzyme, from the marine sponge Hyrtios reticulatus. *Bioorg Med Chem* **20**:4437–4442.
- Yan ZH, Burkhardt A, Loke HK, Chen J, Xu Q, Brauer P, Ma J, Lin Y, Garcia K, Dick LR, et al. (2013) Quantifiable analysis of cellular pathway inhibition of a Nedd8-activating enzyme inhibitor, MLN4924, using AlphaScreen. *Anal Biochem* **439**:109–115.
- Yang D, Tan M, Wang G, and Sun Y (2012) The p21-dependent radiosensitization of human breast cancer cells by MLN4924, an investigational inhibitor of NEDD8 activating enzyme. *PLoS One* **7**:e34079.
- Yang X, Brownell JE, Xu Q, Zhu F, Ma J, Loke HK, Rollins N, Soucy TA, Minissale JJ, Thomas MP, et al. (2013) Absolute quantification of E1, ubiquitin-like proteins and Nedd8-MLN4924 adduct by mass spectrometry. *Cell Biochem Biophys* **67**:139–147.
- Yang Y, Kitagaki J, Dai RM, Tsai YC, Lorick KL, Ludwig RL, Pierre SA, Jensen JP, Davydov IV, Oberoi P, et al. (2007) Inhibitors of ubiquitin-activating enzyme (E1), a new class of potential cancer therapeutics. *Cancer Res* **67**:9472–9481.
- Yao WT, Wu JF, Yu GY, Wang R, Wang K, Li LH, Chen P, Jiang YN, Cheng H, Lee HW, et al. (2014) Suppression of tumor angiogenesis by targeting the protein neddylation pathway. *Cell Death Dis* **5**:e1059.
- Yeh YH, Yang YC, Hsieh MY, Yeh YC, and Li TK (2013) A negative feedback of the HIF-1alpha pathway via interferon-stimulated gene 15 and ISGylation. *Clin Cancer Res* **19**:5927–5939.
- Yin X, Cong X, Yan M, and Zhang DE (2009) Deficiency of a potential 3p21.3 tumor suppressor gene UBE1L (UBA7) does not accelerate lung cancer development in K-rasLA2 mice. *Lung Cancer* **63**:194–200.
- Ying J, Zhang M, Qiu X, and Lu Y (2018) Targeting the neddylation pathway in cells as a potential therapeutic approach for diseases. *Cancer Chemother Pharmacol* **81**:797–808.
- Yoo HM, Kang SH, Kim JY, Lee JE, Seong MW, Lee SW, Ka SH, Sou YS, Komatsu M, Tanaka K, et al. (2014) Modification of ASC1 by UFM1 is crucial for ERα transactivation and breast cancer development. *Mol Cell* **56**:261–274.
- Yoo HM, Park JH, Jeon YJ, and Chung CH (2015) Ubiquitin-fold modifier 1 acts as a positive regulator of breast cancer. *Front Endocrinol (Lausanne)* **6**:36.
- Yoshimura C, Muraoka H, Ochiwa H, Tsuji S, Hashimoto A, Kazuno H, Nakagawa F, Komiya Y, Suzuki S, Takenaka T, et al. (2019) TAS4464, a highly potent and selective inhibitor of NEDD8-activating enzyme, suppresses neddylation and shows antitumor activity in diverse cancer models. *Mol Cancer Ther* **18**:1205–1216.
- Yu J, Huang WL, Xu QG, Zhang L, Sun SH, Zhou WP, and Yang F (2018a) Over-activated neddylation pathway in human hepatocellular carcinoma. *Cancer Med* **7**:3363–3372.
- Yu L, Alva A, Su H, Dutt P, Freundt E, Welsh S, Baehrecke EH, and Lenardo MJ (2004) Regulation of an ATG7-beclin 1 program of autophagic cell death by caspase-8. *Science* **304**:1500–1502.
- Yu L, Li G, Deng J, Jiang X, Xue J, Zhu Y, Huang W, Tang B, and Duan R (2020) The UFM1 cascade times mitosis entry associated with microcephaly. *FASEB J* **34**:1319–1330.
- Yu X, Lao Y, Teng XL, Li S, Zhou Y, Wang F, Guo X, Deng S, Chang Y, Wu X, et al. (2018b) SENP3 maintains the stability and function of regulatory T cells via BACH2 deSUMOylation. *Nat Commun* **9**:3157.
- Yuan R, Wang K, Hu J, Yan C, Li M, Yu X, Liu X, Lei J, Guo W, Wu L, et al. (2014) Ubiquitin-like protein FAT10 promotes the invasion and metastasis of hepatocellular carcinoma by modifying β-catenin degradation. *Cancer Res* **74**:5287–5300.
- Zhang M, Zhu X, Zhang Y, Cai Y, Chen J, Sivaprakasam S, Gurav A, Pi W, Makala L, Wu J, et al. (2015a) RCAD/Ufl1, a Ufm1 E3 ligase, is essential for hematopoietic stem cell function and murine hematopoiesis. *Cell Death Differ* **22**:1922–1934.
- Zhang Q, Hou D, Luo Z, Chen P, Lv B, Wu L, Ma Y, Chu Y, Liu H, Liu F, et al. (2015b) The novel protective role of P27 in MLN4924-treated gastric cancer cells. *Cell Death Dis* **6**:e1867.
- Zhang S, Tan J, Lai Z, Li Y, Pang J, Xiao J, Huang Z, Zhang Y, Ji H, and Lai Y (2014) Effective virtual screening strategy toward covalent ligands: identification of novel NEDD8-activating enzyme inhibitors. *J Chem Inf Model* **54**:1785–1797.
- Zhang XW, Yan XJ, Zhou ZR, Yang FF, Wu ZY, Sun HB, Liang WX, Song AX, Lallemand-Breitenbach V, Jeanne M, et al. (2010) Arsenic trioxide controls the fate of the PML-RARα oncoprotein by directly binding PML. *Science* **328**:240–243.
- Zhang Y, Shi C, Yin L, Zhou W, Wang H, Seng J, and Li W (2017) Inhibition of Mcl-1 enhances Pevonedistat-triggered apoptosis in osteosarcoma cells. *Exp Cell Res* **358**:234–241.
- Zhang Y, Shi CC, Zhang HP, Li GQ, and Li SS (2016) MLN4924 suppresses neddylation and induces cell cycle arrest, senescence, and apoptosis in human osteosarcoma. *Oncotarget* **7**:45263–45274.
- Zhang Y, Zhang M, Wu J, Lei G, and Li H (2012) Transcriptional regulation of the Ufm1 conjugation system in response to disturbance of the endoplasmic reticulum homeostasis and inhibition of vesicle trafficking. *PLoS One* **7**:e48587.
- Zhao B, Villhauer EB, Bhuripanyo K, Kiyokawa H, Schindelin H, and Yin J (2014) SUMO-mimicking peptides inhibiting protein SUMOylation. *ChemBioChem* **15**:2662–2666.
- Zhao B, Zhang K, Bhuripanyo K, Choi CH, Villhauer EB, Li H, Zheng N, Kiyokawa H, Schindelin H, and Yin J (2013a) Profiling the cross reactivity of ubiquitin with the Nedd8 activating enzyme by phage display. *PLoS One* **8**:e70312.
- Zhao B, Zhang K, Villhauer EB, Bhuripanyo K, Kiyokawa H, Schindelin H, and Yin J (2013b) Phage display to identify Nedd8-mimicking peptides as inhibitors of the Nedd8 transfer cascade. *ChemBioChem* **14**:1323–1330.
- Zhao C, Beaudenon SL, Kelley ML, Waddell MB, Yuan W, Schulman BA, Huibregtse JM, and Krug RM (2004) The UbcH8 ubiquitin E2 enzyme is also the E2 enzyme for ISG15, an IFN-α/β-induced ubiquitin-like protein. *Proc Natl Acad Sci USA* **101**:7578–7582.
- Zhao C, Collins MN, Hsiang TY, and Krug RM (2013c) Interferon-induced ISG15 pathway: an ongoing virus-host battle. *Trends Microbiol* **21**:181–186.

- Zhao L, Yue P, Lonial S, Khuri FR, and Sun SY (2011) The NEDD8-activating enzyme inhibitor, MLN4924, cooperates with TRAIL to augment apoptosis through facilitating c-FLIP degradation in head and neck cancer cells. *Mol Cancer Ther* **10**:2415–2425.
- Zhao Y, Xiong X, Jia L, and Sun Y (2012) Targeting Cullin-RING ligases by MLN4924 induces autophagy via modulating the HIF1-REDD1-TSC1-mTORC1-DEPTOR axis. *Cell Death Dis* **3**:e386.
- Zhao Y, Yang J, Liao W, Liu X, Zhang H, Wang S, Wang D, Feng J, Yu L, and Zhu WG (2010) Cytosolic FoxO1 is essential for the induction of autophagy and tumour suppressor activity. *Nat Cell Biol* **12**:665–675.
- Zheng M, Gu X, Zheng D, Yang Z, Li F, Zhao J, Xie Y, Ji C, and Mao Y (2008) UBE1DC1, an ubiquitin-activating enzyme, activates two different ubiquitin-like proteins. *J Cell Biochem* **104**:2324–2334.
- Zheng S, Leclerc GM, Li B, Swords RT, and Barredo JC (2017a) Inhibition of the NEDD8 conjugation pathway induces calcium-dependent compensatory activation of the pro-survival MEK/ERK pathway in acute lymphoblastic leukemia. *Oncotarget* **9**:5529–5544.
- Zheng W, Luo Z, Zhang J, Min P, Li W, Xu D, Zhang Z, Xiong P, Liang H, and Liu J (2017b) Neural precursor cell expressed, developmentally downregulated 8-activating enzyme inhibitor MLN4924 sensitizes colorectal cancer cells to oxaliplatin by inducing DNA damage, G2 cell cycle arrest and apoptosis. *Mol Med Rep* **15**:2795–2801.
- Zhong HJ, Liu LJ, Chan DS, Wang HM, Chan PW, Ma DL, and Leung CH (2014) Structure-based repurposing of FDA-approved drugs as inhibitors of NEDD8-activating enzyme. *Biochimie* **102**:211–215.
- Zhong HJ, Ma VP, Cheng Z, Chan DS, He HZ, Leung KH, Ma DL, and Leung CH (2012a) Discovery of a natural product inhibitor targeting protein neddylation by structure-based virtual screening. *Biochimie* **94**:2457–2460.
- Zhong HJ, Yang H, Chan DS, Leung CH, Wang HM, and Ma DL (2012b) A metal-based inhibitor of NEDD8-activating enzyme. *PLoS One* **7**:e49574.
- Zhou HJ, Xu Z, Wang Z, Zhang H, Zhuang ZW, Simons M, and Min W (2018) SUMOylation of VEGFR2 regulates its intracellular trafficking and pathological angiogenesis. *Nat Commun* **9**:3303.
- Zhou L, Chen S, Zhang Y, Kmiecik M, Leng Y, Li L, Lin H, Rizzo KA, Dumur CI, Ferreira-Gonzalez A, et al. (2016) The NAE inhibitor pevonedistat interacts with the HDAC inhibitor belinostat to target AML cells by disrupting the DDR. *Blood* **127**:2219–2230.
- Zhou L, Jiang Y, Luo Q, Li L, and Jia L (2019a) Neddylation: a novel modulator of the tumor microenvironment. *Mol Cancer* **18**:77.
- Zhou MJ, Chen FZ, Chen HC, Wan XX, Zhou X, Fang Q, and Zhang DZ (2017) ISG15 inhibits cancer cell growth and promotes apoptosis. *Int J Mol Med* **39**:446–452.
- Zhou S, Zhao X, Yang Z, Yang R, Chen C, Zhao K, Wang W, Ma Y, Zhang Q, and Wang X (2019b) Neddylation inhibition upregulates PD-L1 expression and enhances the efficacy of immune checkpoint blockade in glioblastoma. *Int J Cancer* **145**:763–774.
- Zhu H, Bhatt B, Sivaprakasam S, Cai Y, Liu S, Kodeboyina SK, Patel N, Savage NM, Sharma A, Kaufman RJ, et al. (2019a) Ufbp1 promotes plasma cell development and ER expansion by modulating distinct branches of UPR. *Nat Commun* **10**:1084.
- Zhu J, Huang G, Hua X, Li Y, Yan H, Che X, Tian Z, Liufu H, Huang C, Li J, et al. (2019b) CD44s is a crucial ATG7 downstream regulator for stem-like property, invasion, and lung metastasis of human bladder cancer (BC) cells. *Oncogene* **38**:3301–3315.
- Zhuang J, Shirazi F, Singh RK, Kuitse I, Wang H, Lee HC, Berkova Z, Berger A, Hyer M, Chattopadhyay N, et al. (2019) Ubiquitin-activating enzyme inhibition induces an unfolded protein response and overcomes drug resistance in myeloma. *Blood* **133**:1572–1584.
- Zi X and Simoneau AR (2005) Flavokawain A, a novel chalcone from kava extract, induces apoptosis in bladder cancer cells by involvement of Bax protein-dependent and mitochondria-dependent apoptotic pathway and suppresses tumor growth in mice. *Cancer Res* **65**:3479–3486.
- Zuo C, Sheng X, Ma M, Xia M, and Ouyang L (2016) ISG15 in the tumorigenesis and treatment of cancer: an emerging role in malignancies of the digestive system. *Oncotarget* **7**:74393–74409.