

Regulation of GABA_A Receptor Subunit Expression by Pharmacological Agents

MIKKO UUSI-OUKARI AND ESA R. KORPI

Department of Pharmacology, Drug Development and Therapeutics, University of Turku, Turku, Finland (M.U.-O.); and Institute of Biomedicine, Pharmacology, Biomedicum Helsinki, University of Helsinki, Finland (E.R.K.)

Abstract	B
I. Introduction	B
II. Expression of GABA _A receptors	C
A. GABA _A receptor genes	C
B. Brain regional expression of subunit mRNAs	D
C. Brain regional expression of subunit polypeptides	F
D. Brain regional expression of GABA _A receptor subtypes	F
E. In vitro models to study regulation of GABA _A receptor expression	F
1. Mouse cortical neurons	F
2. Cerebellar granule cells	G
3. Primary cultures of hippocampal neurons	G
F. Mechanisms regulating GABA _A receptor subunit expression	G
G. Mechanisms regulating GABA _A receptor cell surface expression	H
III. Regulation of GABA _A receptor expression by pharmacological agents	I
A. Benzodiazepines	I
1. α 1 Subunit	J
2. α 2 Subunit	K
3. α 3 Subunit	M
4. α 4 Subunit	M
5. α 5 Subunit	O
6. α 6 Subunit	P
7. β 1 Subunit	P
8. β 2 Subunit	Q
9. β 3 Subunit	R
10. γ 2 Subunit	S
11. γ 1 and γ 3 Subunits	T
12. δ Subunit	T
13. Conclusions on effects of long-term benzodiazepine administration on receptor subunit expression	T
14. Benzodiazepine-induced changes in the expression of other genes	U
B. Neurosteroids	U
C. Barbiturates	V
D. Ethanol	X
1. α 1 Subunit	Y
2. α 2 Subunit	AB
3. α 3 Subunit	AC
4. α 4 Subunit	AC
5. α 5 Subunit	AD
6. α 6 Subunit	AE
7. β 1 Subunit	AE
8. β 2 Subunit	AE

Address correspondence to: Dr. Mikko Uusi-Oukari, Department of Pharmacology, Drug Development and Therapeutics, University of Turku, Itäinen Pitkakatku 4, 20014 Turku, Finland. E-mail: mikko.uusi-oukari@utu.fi

This article is available online at <http://pharmrev.aspetjournals.org>.

doi:10.1124/pr.109.002063.

9. β 3 Subunit	AE
10. γ 2 Subunit	AE
11. γ 1 and γ 3 Subunits.....	AF
12. δ Subunit.....	AF
13. Studies of continuous intermittent ethanol administration on subunit cell surface expression	AF
14. Conclusions on effects of continuous ethanol administration on GABA _A receptor subunit expression	AF
IV. Novel opportunities to target the GABA _A receptor system with therapeutics	AF
Acknowledgments	AH
References	AH

Abstract—The γ -aminobutyric acid (GABA) type A receptor system, the main fast-acting inhibitory neurotransmitter system in the brain, is the pharmacological target for many drugs used clinically to treat, for example, anxiety disorders and epilepsy, and to induce and maintain sedation, sleep, and anesthesia. These drugs facilitate the function of pentameric GABA_A receptors that exhibit widespread expression in all brain regions and large structural and pharmacological heterogeneity as a result of composition from a repertoire of 19 subunit variants. One of the main problems in clinical use of GABA_A receptor agonists is the development of tolerance. Most drugs, in long-term use and during withdrawal, have been associated with important modulations of the receptor subunit expression in brain-region-specific manner, participating in the mechanisms of tolerance and de-

pendence. In most cases, the molecular mechanisms of regulation of subunit expression are poorly known, partly as a result of neurobiological adaptation to altered neuronal function. More knowledge has been obtained on the mechanisms of GABA_A receptor trafficking and cell surface expression and the processes that may contribute to tolerance, although their possible pharmacological regulation is not known. Drug development for neuropsychiatric disorders, including epilepsy, alcoholism, schizophrenia, and anxiety, has been ongoing for several years. One key step to extend drug development related to GABA_A receptors is likely to require deeper understanding of the adaptational mechanisms of neurons, receptors themselves with interacting proteins, and finally receptor subunits during drug action and in neuropsychiatric disease processes.

I. Introduction

Chemical balance in the brain systems for neuronal excitation, inhibition, rhythmic activity, and scaling of neuronal discharge probabilities is a very important basic mechanism that has been used in pharmacological modulation of behavior, including therapies of emotional and affective disturbances, cognitive impairment, and motor disturbances resulting from neurodegeneration and genetic abnormalities. In simple terms, the chemical balance is primarily set by the activities of the most widely distributed neurotransmitter systems in the CNS,¹ namely the excitatory glutamate system and the inhibitory GABA system, although the balance is also modulated by a large number of other slower acting transmitters, modulators, and ion channels. The GABA system is widely used in the

treatment of anxiety disorders, insomnia, epilepsy, restlessness, and aggressive behaviors, and it is the target for many intravenous and inhalational anesthetics and drugs of abuse, such as alcohol. Several severe problems are related to the long-term therapeutic use of GABA system-affecting drugs, most significantly the loss of efficacy, tolerance development, dependence development, and finally addiction to at least some of these drugs.

The GABA system is a ubiquitous system regulated by a number of different genes, affecting synthesis of various receptor subunits, interacting proteins, and associated transporters and synthetic and catabolic enzymes. Figure 1 illustrates a GABAergic synapse with its main pharmacological components. Pharmacology has focused on the modification of receptors and transmitters of the system: most of the clinically used drugs target the GABA type A receptors (GABA_A), the GABA type B receptors (e.g., agonist baclofen), GABA transporters in neurons and glial cells (e.g., inhibitor tiagabine), and the catabolic enzyme GABA-transaminase (e.g., inhibitor vigabatrin) to increase the level of GABA.

This review will focus on one part of the GABA system: the GABA_A receptors (Fig. 2). GABA_A receptors are pentameric complexes of subunits, and they form an integral anion channel, permeable to chloride and bicarbonate ions. These receptors are also molecular targets for various

¹ Abbreviations: AMPA, α -amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid; BDNF, brain-derived neurotrophic factor; BZ, benzodiazepine; CA, cornu ammonis; CE, continuous ethanol; CGC, cerebellar granule cell; CIE, continuous intermittent ethanol; CNS, central nervous system; CRE, cAMP response element; CREB, cAMP response element binding protein; FG 7142, *N*-methyl- β -carboline-3-carboxamide; GABA, γ -aminobutyric acid; GABARAP, GABA_A receptor-associated protein; ICER, inducible cAMP early repressor; JAK, Janus kinase; L, long splice variant; NMDA, *N*-methyl-D-aspartate; PKA, protein kinase A; Ro 15-4513, 8-azido-5,6-dihydro-5-methyl-6-oxo-4*H*-imidazo[1,5-*a*][1,4]benzodiazepine-3-carboxylic acid ethyl ester; S, short splice variant; STAT, signal transducer and activator of transcription; wdr-CE, withdrawal from CE; wdr-CIE, withdrawal from CIE.




FIG. 1. GABAergic synapse with synthetic and catabolic enzymes, receptors, and transporters. GAD, glutamate decarboxylase, the GABA-synthesizing enzyme that exists in two forms (GAD65 and GAD67; GABA transaminase, GABA metabolizing enzyme; GAT, a GABA transporter existing at least in four subtypes; GABA_B receptor is a G-protein-coupled heptahelix receptor. The most important of the Cl⁻/cation transporters is KCC2, which provides the chloride gradient for the hyperpolarizing action of GABA_A receptor activation in mature neurons.

classes of benzodiazepine-site (BZ-site) ligands, barbiturates, neurosteroids, intravenous anesthetics propofol and etomidate, inhalation anesthetics, and alcohol (Fig. 3). The molecular actions of various drugs on the GABA_A receptors have been well reviewed (Lüddens et al., 1995; Rabow et al., 1995; Sieghart, 1995; Sigel and Buhr, 1997; Korpi et al., 2002; Lambert et al., 2003). In mature neurons, under normal conditions, the activation of GABA_A receptors leads to hyperpolarization of cell membrane potential and inhibition of neuronal activity. However, in immature neurons, in which the ion gradient-forming anion transporters are not yet fully operational, GABA_A receptor activation produces depolarization (Ben-Ari et al., 1997). The opposite process may take place in such pathological conditions as neuropathic pain (see, Kahle et al., 2008). Prolonged activation of the system leads to loss of function, tolerance, the mechanisms of which are still poorly known. This review does not deal with mechanisms of acute tolerance (i.e., the reduced sensitivity of the system due to changes within one use period of a drug) or those of innate tolerance (i.e., inherited mechanisms regulating the acute sensitivity of the system). Drugs acting via GABA_A receptors often affect the regulation of various GABA_A receptor subunits, which might provide the simplest mechanism for tolerance. Therefore, the present review is focused on the data how the expressions of different subunits are affected by specific pharmacological manipulations.

II. Expression of GABA_A Receptors

A. GABA_A Receptor Genes

A total of 19 mammalian genes coding for GABA_A receptor subunits, belonging to eight subunit classes,

have been cloned: α1–α6, β1–β3, γ1–γ3, δ, ε, θ, π, ρ1–ρ3 (Olsen and Sieghart, 2008). Subsequent gene mapping showed that most GABA_A receptor genes are clustered in vertebrate genomes (Wilcox et al., 1992; Russek and Farb, 1994; Bailey et al., 1999; Russek, 1999). Fourteen of the 19 human GABA_A receptor genes are clustered on four chromosomes, 4p12-p13, 5q31-q35, 15q11-q13, and Xq28 (Russek, 1999). Two clusters of four genes each encode two α subunits, one β subunit, and one γ subunit (*GABRA2*, *GABRA4*, *GABRB1*, and *GABRG1* on chromosome 4, and *GABRA1*, *GABRA6*, *GABRB2*, and *GABRG2* on chromosome 5). The two other clusters each contains three genes: the cluster in chromosome 15 comprises one α subunit gene (*GABRA5*), one β subunit gene (*GABRB3*), and one γ subunit gene (*GABRG3*), and the cluster in X chromosome consists of one α subunit gene (*GABRA3*), the θ subunit gene (*GABRQ*), and the ε subunit gene (*GABRE*). The θ and ε polypeptides exhibit about 50% identity to β and γ subunits, respectively. Therefore, the θ and ε subunits are considered “β-like”




FIG. 2. Schematics of GABA_A receptor structure and function. A, topography of a GABA_A receptor subunit partially embedded in the lipid bilayer. 1, N-terminal extracellular domain responsible for transmitter and ligand binding and coupling of the binding sites with ion channel. This part is also important for the allosteric effects and for the assembly of various receptor subunits into functional receptors. 2, four transmembrane regions forming the anion channel are responsible for binding of hydrophobic ligands, ion selectivity, and channel binding sites. 3, intracellular loop between transmembrane helices 3 and 4 forms the motif for regulatory phosphorylation sites and for the intracellular factors anchoring the receptors in appropriate locations (e.g., on the postsynaptic thickening) using interactions with auxiliary and cell structural proteins. B, hypothetical binding sites for GABA and allosteric modulators such as benzodiazepine ligands and a domain essential for the functions of various ligands such as etomidate, loreclezole, and methyl-6,7-dimethoxy-4-ethyl-β-carboline-3-carboxylate in a pentameric receptor complex. C, allosteric activation of GABA_A receptor may increase the peak height (amplitude) of the response and/or prolong the response as compared with the response by GABA alone.




FIG. 3. Schematic illustration of the GABA_A receptor and its associated binding sites. The receptor is pentameric, being composed of two α , two β , and one γ subunit. GABA_A receptors contain recognition sites for a variety of clinically relevant drugs. The binding of GABA in two GABA binding sites at the interface between α and β subunits open the receptor-associated chloride (Cl^-) channel. The benzodiazepine binding site is located at the interface between α and γ subunits. Barbiturates, ethanol, and neurosteroids bind to sites in the membrane-spanning transmembrane regions of the subunits.

and “ γ -like,” respectively. Thus, each cluster contains one or two α , one β (or “ β -like”), and one γ (or “ γ -like”) subunit gene. The rat and mouse GABA_A receptor genes are clustered similarly to human genes (Olsen and Sieghart, 2008).

It has been proposed that the mammalian GABA_A receptor genes evolved from a common ancestral gene cluster, which is presumed to have comprised an “ α -like,” a “ β -like,” and a “ γ -like” subunit gene, as a consequence of two whole-genome duplication events (Russek and Farb, 1994; Bailey et al., 1999; Russek, 1999). It is assumed that a tandem duplication of an α subunit gene took place in one of the clusters after the first but before the second duplication. The clusters on human chromosomes 4 and 5 evolved from a common ancestral cluster, and the clusters on chromosome 15 and X arose from a second common ancestral cluster. The model is supported by the fact that genes having the same location within the clusters on chromosomes 4 and 5 (i.e., α 1 and α 2 or α 4 and α 6, respectively) are more closely related to one another than to any other α subunit genes.

B. Brain Regional Expression of Subunit mRNAs

Each of the GABA_A receptor subunit mRNAs displays a unique distribution (Laurie et al., 1992a; Persohn et al., 1992; Wisden et al., 1992). Some subunits are ubiquitously expressed almost throughout the brain, whereas the localization of most subunits is more narrowly confined. Some neuronal populations coexpress large number of subunit mRNA isoforms, whereas others express only a few GABA_A receptor subunits (Laurie et al., 1992a; Persohn et al., 1992; Wisden et al., 1992). Neocortex, hippocampus, and caudate-putamen display

complex expression patterns of large diversity of subunit combinations. In contrast, essentially one of each isoform of the α , β , and γ subunit classes is expressed in the inferior colliculus, substantia nigra pars reticulata, and cerebellar stellate/basket cells (Laurie et al., 1992a; Wisden et al., 1992). The brain regional expression profiles of GABA_A receptor subunits are presented in Table 1.

The expression of the two major α subunit isoforms, α 1 and α 2, is both widespread in the brain. The most abundant subunit, α 1, is expressed almost ubiquitously in the brain (Wisden et al., 1992). Its expression is especially rich in the olfactory bulb tufted and mitral cells, pyriform cortex, globus pallidus, endopeduncular nucleus, medial septum and diagonal band, parafascicular nucleus of the thalamus, red nucleus, inferior colliculus, cerebellar molecular and granule cell layers, and Purkinje cells (Laurie et al., 1992a; Persohn et al., 1992; Wisden et al., 1992). Despite widespread expression of the α 2 subunit, there is a negative correlation between the brain regional expression patterns of α 1 and α 2 subunits. The expression of α 2 mRNA is high in granule cells of the olfactory bulb, hippocampus, amygdala, lateral septum and in the medial preoptic area of the hypothalamus (Persohn et al., 1992; Wisden et al., 1992). In the cerebellum α 2 mRNA is solely expressed in Bergmann glial cells (Laurie et al., 1992a; Persohn et al., 1992). The expression of α 3 mRNA is localized in the olfactory bulb, cerebral cortex and brain stem nuclei (Persohn et al., 1992; Wisden et al., 1992). The expression of α 4 mRNA is confined to the hippocampus, thalamus, caudate-putamen, nucleus accumbens, and neocortex (Wisden et al., 1992). The α 5 mRNA is highly expressed in the hippocampus. It is also present at lower levels in the olfactory bulb granule cells, neocortex, and hypothalamus (Laurie et al., 1992a; Persohn et al., 1992). The α 6 mRNA expression pattern is the most restricted of the GABA_A receptor subunits: α 6 mRNA is expressed in the cerebellar granule cells and cochlear nucleus granule cells (Laurie et al., 1992a, Varecka et al., 1994).

The expression of β 1 mRNA is strong in the olfactory bulb mitral cells and in the hippocampus. It is also expressed at lower levels in the cerebral cortex, substantia nigra, superior colliculus, and cerebellum (Persohn et al., 1992; Wisden et al., 1992). The expression of β 2 is more widespread and strongly correlates with that of α 1 mRNA. This suggests that the expression of genes coding for α 1 and β 2 subunits, members of the same gene cluster, might be coordinately regulated (Steiger and Russek, 2004). In addition, it suggests that the two subunits assemble within the same receptor complexes with high probability. The mRNA expression pattern of β 3 is also widespread and strongly correlates with that of α 2 mRNA. The genes for these subunits reside in different chromosomes/gene clusters. In any case, their similar expression patterns suggest that the subunits coassemble to form receptors.

TABLE 1
Quantitative estimates for brain regional distribution of GABA_A receptor subunits in adult rat brain

The table is based on the published mRNA in situ hybridization data (Laurie et al., 1992; Persohn et al., 1992; Wisden et al., 1992) and subunit polypeptide immunohistochemical data (Pirker et al., 2000; Schwarzer et al., 2001). The table as such should not be used to compare the absolute levels of various subunits in any brain region. Thus, the brain regional profile for each subunit is given using the same scale, even if the absolute concentrations of subunit mRNAs or peptides are different. The regional profile of each subunit is presented so that 3 denotes the highest expression of the subunit in question; 2, strong expression; 1, low expression; and no grading (empty fields), very low or undetectable expression. The $\gamma 3$ subunit probes have so faint and even staining in many brain regions both in situ hybridization and immunohistochemical assays that we decided to mark its expression with 1.

Brain Region	GABA _A Receptor Subunit mRNA and Polypeptide Expression												
	$\alpha 1$	$\alpha 2$	$\alpha 3$	$\alpha 4$	$\alpha 5$	$\alpha 6$	$\beta 1$	$\beta 2$	$\beta 3$	$\gamma 1$	$\gamma 2$	$\gamma 3$	δ
Olfactory areas													
External plexiform layer of olfactory bulb	3	1	1		1		2	3	3		3	1	
Glomerular layer	3	1	1				1	3	3		3		1
Internal granular layer		3	3	1	2								1
Olfactory tubercle		2		2			1	1	3		1	1	1
Islands of Calleja	3							3				1	
Primary olfactory cortex	3	3	3	2			1	3	3	1	3	1	1
Cerebral cortex													
Layers 1–4	2	2	1	2	1		2	2	3		2	1	2
Layers 5–6	3	1	3	1	2		3	2	3		2	1	1
Anterior cingulate cortex	2	3	1	1			2	1	3		2		
Limbic regions, hippocampus, amygdala													
Entorhinal cortex	2	2	1	1	1		2	2	2		2	1	1
Subiculum	2	2	1	1	1		2	2	2		2	1	1
Hippocampus, CA1	2	3		2	3		3	1	3		3		
Hippocampus, CA3	1	3		1	3		3	1	3		3		
Hippocampus, dentate gyrus	1	3		2	1		2	2	3		3		1
Bed nucleus stria terminalis, medial	1	3	2	1	1		2	1	2	3	1	1	1
Nucleus of horizontal limb of diagonal band	3		1		1			3	1		2	1	1
Septohippocampal nucleus/taenia tecta	1	3	2	2	1		2		3		1		
Lateral septal nuclei	1	3	2	1			2	1	1	3	2	1	
Triangular septal nucleus	1	2						1	1		1		
Bed nucleus of anterior commissural	1	3	1	1						1			
Anterior amygdaloid area	1	1	1		1		1	2	1	1	2	1	
Amygdala	1	3	1	1			1	2	2	3	1	1	
Posteromedial cortical amygdaloid nucleus	1	3	1		1		1	2	2		1	1	1
Basal ganglia/striatum													
Nucleus accumbens	1	3	1	2	1		1	1	3		1	1	2
Caudate/putamen	1	3	1	2	1		1	1	3		1	1	2
Globus pallidus	3	1			1			3		3	1	1	1
Claustrum	2	2	3	1	1		1	2	2		1	1	1
Ventral pallidum/substantia innominata	3		1	1	1		1	3		3	1	1	
Thalamus, epithalamus													
Paraventricular thalamic nucleus	1	2	1	3			2	2	2		2		1
Anterodorsal thalamic nucleus	3			3				3	1		1		3
Centrolateral/medial thalamic nucleus	2	1	1	2			1	3	1	2	1		1
Intermediodorsal thalamic nucleus	2			3				3					1
Lateral posterior/laterodorsal thalamic nucleus	2			3				3					3
Ventroposterior thalamic nucleus	1			3				3					2
Zona incerta/subthalamic nucleus	2		1					2			1		1
Medial habenular nucleus		2							2		2	1	
Medial geniculate nucleus	2			3				3			1	1	3
Hypothalamus													
Medial preoptic area/periventricular nucleus	1	3	1		1		1		2	2	1		1
Lateral preoptic area	1	1	1					1	1		1	1	
Lateral hypothalamic area	1						1	1	1		1		1
Anterior hypothalamic area	1						3	1	1		2		1
Paraventricular hypothalamic nucleus	1	1					2	1	2		1		1
Ventromedial hypothalamic nucleus	1	2	1		1		3		2	1	2		1
Mesencephalon, pons, medulla													
Substantia nigra, pars reticulata	3		1				1	2		1	1	1	
Substantia nigra, pars compacta			2	1			1				1	1	1
Ventral tegmental area	1	1	1				2				2	1	1
Interpeduncular nucleus	2	1		2	1				2		1	1	1
Red nucleus	3							2			2	1	1
Superior colliculus, superior gray layer	2	2		1	2			2	2	1	1		
Superior colliculus, intermediate gray layer	2						1	1	2		1	1	
Central gray	3	1	1				2	1	2		1	1	2
Inferior colliculus	3	1					2	1	2		1	1	1
Raphe nuclei	2	1	2		1		1	2	2		2	1	1
Locus ceruleus			2										
Cerebellum													
Granule cell layer	3					3	1	3	3		1		3
Molecular layer	3	1			1		2	1			2		

The expression of $\gamma 1$ mRNA is almost undetectable in most brain regions. The expression is highest in the hippocampus, globus pallidus, amygdala, septum, and medial preoptic area of the hypothalamus. The expression pattern of $\gamma 1$ mRNA correlates with that of $\alpha 2$ mRNA. The mRNA of the major γ subunit, $\gamma 2$, is expressed throughout the brain. The expression of $\gamma 3$ mRNA, although low, is most clearly expressed in the neocortex and thalamus (Wisden et al., 1992).

The expression of δ mRNA is high in the neocortex, hippocampal dentate granule cells, thalamus, and cerebellar granule cells (Laurie et al., 1992a; Persohn et al., 1992; Wisden et al., 1992). Its likely partners are $\alpha 4$ in the forebrain and $\alpha 6$ in the cerebellum (Jones et al., 1997; Korpi et al., 2002; Peng et al., 2002) and, at least in some cortical interneurons, the $\alpha 1$ subunit as well (Glykys et al., 2007). The ϵ and θ mRNAs are strongly concentrated in the monoaminergic nuclei of the brainstem, such as the noradrenergic locus ceruleus (Sinkkonen et al., 2000; Moragues et al., 2002). The π mRNA has not been detected in the brain, but it is abundant in female peripheral organs such the uterus (Hedblom and Kirkness, 1997). The ρ subunits are mostly expressed in the retina, colliculi, and cerebellum (Boue-Grabot et al., 1998; Wegelius et al., 1998).

C. Brain Regional Expression of Subunit Polypeptides

Brain regional localizations of GABA_A receptor subunit polypeptides strongly correlate with those of the subunit mRNAs. The $\alpha 1$ subunit displays the most widely distributed expression, being present in practically all brain regions (Fritschy and Mohler, 1995; Pirker et al., 2000). Expression of $\alpha 2$ subunit is highest in the accessory olfactory bulb, hippocampus, amygdala, septum, striatum, accumbens, and hypothalamus (Fritschy and Mohler, 1995; Pirker et al., 2000). The $\alpha 3$ subunit is expressed in the olfactory bulb, inner layers of the cerebral cortex, endopiriform cortex, amygdala, lateral septum, claustrum, and superior colliculus (Pirker et al., 2000). It is usually expressed in monoaminergic neurons (Gao et al., 1993, 1995; Gao and Fritschy, 1994). The expression of $\alpha 4$ subunit is strongest in the thalamus, caudate-putamen, nucleus accumbens, olfactory tubercle, and hippocampus (Pirker et al., 2000). The $\alpha 5$ subunit is highly expressed in the olfactory bulb, inner layers of the cerebral cortex, endopiriform nucleus, subiculum, and hippocampus (Fritschy and Mohler, 1995; Pirker et al., 2000). The expression of $\alpha 6$ subunit is restricted to the granule cells of the cerebellum and cochlear nuclei (Gutiérrez et al., 1996; Pirker et al., 2000).

All three β subunits are widely distributed in the brain (Pirker et al., 2000). They are strongly expressed in the cerebral cortex. In many brain regions, their distribution is more or less complementary (Pirker et al., 2000). In the pallidum and thalamus, $\beta 2$ is the main β subunit, whereas in the striatum, $\beta 3$ is the most abun-

dant β subunit (Moreno et al., 1994; Miralles et al., 1999; Pirker et al., 2000). The most abundant γ subunit, $\gamma 2$, is expressed throughout the brain. In the thalamus, however, $\gamma 2$ displays low expression level (Gutiérrez et al., 1994; Fritschy and Mohler, 1995; Pirker et al., 2000). The expression of $\gamma 1$ subunit is located in the pallidum, substantia nigra, septum, and amygdala (Pirker et al., 2000). The $\gamma 3$ subunit is diffusely distributed throughout the brain (Pirker et al., 2000). The δ subunit is expressed in the cerebellar granule cells, thalamus, dentate molecular layer, subiculum, cerebral cortex, and striatum (Fritschy and Mohler, 1995; Pirker et al., 2000).

D. Brain Regional Expression of GABA_A Receptor Subtypes

The most abundant GABA_A receptor subtype, $\alpha 1\beta 2\gamma 2$, is present in most brain regions. In immunostaining, the subunits display striking similarity of location in the internal granular layer of the olfactory bulb, polymorph cell layer, and CA3 region of the hippocampus, cerebral cortical interneurons, the globus pallidus and several thalamic nuclei (Benke et al., 1994; Pirker et al., 2000).

Subunits of the most prominent $\alpha 2$ -containing subtype, $\alpha 2\beta 3\gamma 2$, have been most clearly colocalized in the accessory olfactory bulb, striatum, septum, molecular layer of the dentate gyrus, and hypothalamus (Benke et al., 1994; Pirker et al., 2000). Of the other $\alpha 2$ -containing subtypes, $\alpha 2\beta 1$ is a likely possibility. This subtype is suggested to be expressed in Bergmann glia and nuclei of the limbic systems (McKernan and Whiting, 1996).

The $\alpha 3\beta 2$ subtype is chiefly expressed in several monoaminergic cells in various brain nuclei. The $\alpha 3\theta\epsilon$ subunit combination is also likely, because their mRNAs are all highly enriched in rat locus ceruleus (Sinkkonen et al., 2000). The $\alpha 4\beta 2$ is expressed in thalamus, caudate-putamen, and dentate gyrus (Pirker et al., 2000). The thalamic relay nuclei express $\alpha 4\beta 2\delta$ subtype (Chandra et al., 2006), whereas the dentate granule cells express the $\alpha 4\beta 3\delta$ subtype (Liang et al., 2006). Both $\alpha 4\beta 2$ subtypes are obviously expressed in the cerebral cortex, whereas $\alpha 4\beta 3\delta$ subtype predominates in the striatum. The $\alpha 5\beta 3\gamma 2$ subtype is expressed in CA1 pyramidal neurons (Sur et al., 1998; Caraiscos et al., 2004; Olsen and Sieghart, 2008). The $\alpha 6$ -containing $\alpha 6\beta 1\gamma 2$, $\alpha 6\beta 2\delta$, and $\alpha 6\beta 3\delta$ subtypes are expressed in the cerebellar granule cells and cochlear nuclei.

E. In Vitro Models to Study Regulation of GABA_A Receptor Expression

1. Mouse Cortical Neurons. The procedure of preparation of primary cultures of mouse cortical neurons is based on techniques described in detail by Yavin and Yavin (1980) and Yu et al. (1984). The cultures are highly enriched in well differentiated GABAergic cortical neurons. More than 90% of the cells in the cultures are neurons (Kuriyama et al., 1987). Cerebrum of 15-

day-old mouse embryos is used for the preparation. At that stage of development, large numbers of neurons have just entered their postmitotic stage of differentiation, and only a few proliferative glial precursors are present. The cerebral hemispheres are dissected and minced. The cells are dissociated by trituration and seeded in poly-L-lysine-coated culture dishes (Mehta and Ticku, 1992). The cultures are virtually homogeneous. The cells grow neurite extensions, migrate, and form clumps (Yu et al., 1984). Prenatally and during early postnatal days, cortical neurons express predominantly $\alpha 2$, $\alpha 3$, $\alpha 4$, $\alpha 5$, $\beta 2$, $\beta 3$, and $\gamma 2$ subunits (Laurie et al., 1992b). Cultured cortical neurons express $\alpha 1$ – $\alpha 5$, $\beta 2$, $\beta 3$, $\gamma 2$, and δ subunits (Sheela Rani and Ticku, 2006).

2. Cerebellar Granule Cells. Cerebellar granule cells (CGCs) constitute the vast majority of neurons in the cerebellum. They undergo cell division until postnatal days 7 to 9, at which time their differentiation process is initiated. Because granule cells outnumber the other types of neurons in the cerebellum by a 1000-fold, it is quite easy to establish an *in vitro* cell culture system to study the development of granule cells (Messer, 1977). Cultured CGCs have been widely used as a model system to study regulation of gene expression during development both *in vivo* and *in vitro* (Savill et al., 2005). The expression of GABA_A receptor subunits in CGCs undergoes maturational developmental changes during the first 2 postnatal weeks (Laurie et al., 1992a). Expression of the major GABA_A receptor subunits in adult cerebellum, $\alpha 1$, $\alpha 6$, $\beta 2$, $\beta 3$, $\gamma 2$, and δ , is strongly increased, and that of $\alpha 2$, $\alpha 3$, and $\beta 1$ subunits is decreased during the second postnatal week both *in vivo* and *in vitro* (Laurie et al., 1992a, 1992b; Carlson et al., 1998; Grayson et al., 1998). Cerebella from decapitated 5- to 8-day-old rats or mice are dissected out, cut in small cubes, and subjected to acute trypsin dissociation. After trituration the dissociated cells are suspended in culture medium. The cells are seeded in poly-L-lysine-coated culture dishes. The cultures are homogeneous and consist of at least 90% granule cells (Messer, 1977). Cultured CGCs express $\alpha 1$ – $\alpha 6$, $\beta 1$ – $\beta 3$, $\gamma 1$ – $\gamma 3$, and δ subunits (Bovolin et al., 1992).

3. Primary Cultures of Hippocampal Neurons. Primary cultures of hippocampal neurons are usually prepared from embryonic day 18 rat fetuses (Banker and Cowan, 1977). The cultures are enriched in pyramidal cells and the number of non-neuronal cells present is minimal. Meninges-free hippocampi are microdissected and trypsinized. The cells are then triturated and plated in culture dishes coated with poly-L-lysine and collagen (Banker and Cowan, 1977). Embryonic hippocampal neuronal cultures include a heterogeneous mixture of pyramidal and nonpyramidal neurons of multiple different morphologies (Rothman and Cowan, 1981). The cultured cells express GABA_AR subunits $\alpha 1$ – $\alpha 5$, $\beta 1$ – $\beta 3$, $\gamma 1$ – $\gamma 3$ and δ (Brooks-Kayal et al., 1998; Maric et al., 1999).

F. Mechanisms Regulating GABA_A Receptor Subunit Expression

The large number of GABA_A receptor genes and the various types of neurons and glial cells in the brain with different patterns of subunit expression suggest a complex system regulating their transcription (Laurie et al., 1992a; Wisden et al., 1992; Olsen and Sieghart, 2008). Major changes occur during development in the subunit expression patterns (Laurie et al., 1992b). However, several studies indicate changes in GABA_A receptor subunit expression also in adult brain (Huntsman et al., 1994; Kamphuis et al., 1995; Loup et al., 2000). The changes are often suggested to reflect a change in neuronal activity. In cultured neurons *in vitro*, the expression of several GABA_A receptor subunit mRNAs and polypeptides is up-regulated by depolarization with glutamate receptor agonists (Harris et al., 1995; Gault and Siegel, 1998; Salonen et al., 2006). However, these effects obviously correspond to developmental induction and up-regulation of the expression occurring *in vivo*. It has been shown that stimulation of *N*-methyl-D-aspartate (NMDA) receptors increases the transcription rate of $\alpha 1$ mRNA in cultured rat CGCs (Harris et al., 1995). On the other hand, the presence of GABA_A agonists in medium of cultured neurons down-regulates GABA_A receptor subunits (Montpied et al., 1991a; Mehta and Ticku, 1992; Baumgartner et al., 1994). Prolonged presence of GABA has been shown to reduce the transcription rate of $\alpha 1$ mRNA (Lyons et al., 2000). This use-dependent regulation of GABA_A receptors has been suggested to include (in putative temporal order): 1) desensitization, 2) endocytosis of subunit polypeptides, 3) subunit polypeptide degradation, and 4) repression of subunit gene expression (Barnes, 1996).

Activity-dependent signaling pathways modulate the function of both transcriptional activators and repressors (West et al., 2002). Calcium is a crucial second messenger in the transduction of synaptic activity into gene expression (Carafoli et al., 2001), and it is involved in the mechanisms of GABA_A receptor up- and down-regulation (Gault and Siegel, 1998; Lyons et al., 2001). The transcription factor cAMP response element (CRE) binding protein (CREB) is induced in response to neurotransmitters, neuromodulators, and neurotrophic factors (Lonze and Ginty, 2002). It was recently shown that the activation of protein kinase C in primary rat neocortical cultures increases transcription of $\alpha 1$ mRNA via phosphorylation of CREB that is bound to the *GABRA1* promoter (Hu et al., 2008). In contrast, activation of protein kinase A (PKA) represses $\alpha 1$ mRNA transcription via inducible cAMP early repressor (ICER) that forms inactive heterodimers with CREB (Hu et al., 2008). Brain-derived neurotrophic factor (BDNF) decreases $\alpha 1$ transcription via activation of the Janus kinase/signal transducer and activator of transcription (STAT) pathway (Lund et al., 2008). BDNF-dependent

phosphorylation of STAT3 induces the synthesis of ICER that binds with phosphorylated CREB at the *GABRA1* promoter CRE site, thereby repressing transcription (Lund et al., 2008).

Expression of GABA_A receptor subunits is regulated in cultured CGCs by both cAMP- and BDNF-mediated signaling mechanisms. Activation of adenylate cyclase or the presence of cAMP analogs in cultured CGCs up-regulate $\beta 2$ and down-regulate $\alpha 6$ mRNA expression but have no effect on $\alpha 1$ and $\beta 3$ mRNA (Thompson et al., 2000). In addition, $\alpha 6$ and $\beta 3$ polypeptides are down-regulated, whereas $\alpha 1$ and $\beta 2$ are up-regulated, indicating subunit-specific regulation of expression. The transcriptional and/or translational mechanisms mediating these effects are only partially mediated by PKA (Thompson et al., 1996, 2000). Addition of BDNF in cultured rat visual cortex cells induces a rapid increase in the total number of functional cell surface GABA_A receptors (Mizoguchi et al., 2003). In cultured CGCs, BDNF induces $\alpha 6$ mRNA expression and enhances the expression of $\alpha 1$ and $\gamma 2$ mRNA (Bulleit and Hsieh, 2000). These enhancements are mediated via mitogen-activated protein kinase pathway (Bulleit and Hsieh, 2000). In contrast, in cultured hippocampal pyramidal cells, BDNF reduced cell surface expression of $\alpha 2$, $\beta 2/3$, and $\gamma 2$ subunits (Brünig et al., 2001). The results suggest that BDNF affects GABA_A receptor expression in a brain region- and cell-specific manner.

The transcription factors responsible for developmental and brain region/cell-specific expression of GABA_A receptor subunits are presently unknown. Both coordinated and independent expression of the genes in a GABA_A receptor gene cluster is suggested by the $\beta 2$ - $\alpha 6$ - $\alpha 1$ - $\gamma 2$ cluster: the expression patterns of $\alpha 1$ and $\beta 2$ are almost identical, whereas $\gamma 2$ is expressed virtually throughout the brain, and $\alpha 6$ expression is restricted to cerebellar granule cells (Laurie et al., 1992a; Wisden et al., 1992). Coordinated expression is strongly suggested for the subunits in the θ - $\alpha 3$ - ϵ cluster: all subunits colocalize in monoaminergic neurons (Fritschy et al., 1992; Sinkkonen et al., 2000; Moragues et al., 2002). But it is clear nonetheless that although coordinated expression of genes in a cluster seems to apply to some subunits, the brain regional and temporal expression of genes in most clusters is independent of other genes of the cluster.

The findings of protein kinase C/CREB mediated initiation of $\alpha 1$ mRNA transcription that is repressed by PKA/ICER, and BDNF/Janus kinase/STAT/ICER-mediated repression of $\alpha 1$ mRNA transcription are new promising examples of approaches that produce information on how expression of GABA_A receptor genes is regulated.

G. Mechanisms Regulating GABA_A Receptor Cell Surface Expression

More than 20 intracellular or transmembrane proteins can be considered GABA_A receptor accessory pro-

teins because they intimately interact with various sites of the large TM3–TM4 intracellular loops to regulate the surface expression of receptors. They regulate receptor trafficking from intracellular machinery to cell membranes and back, modify vesicular trafficking of receptors along the neurites, link receptors to cytoskeleton, affect inhibitory postsynaptic structures, and phosphorylate/dephosphorylate specific subunits (for review, see Lüscher and Keller, 2004; Birnir and Korpi, 2007; Chen and Olsen, 2007; Kneussel and Loebrich, 2007; Jacob et al., 2008). Most of the accessory proteins interact with the β or γ subunits, often very specifically with a certain subunit. This means that the interactions may participate in the regulation of functional activities of selected receptor subtypes rather than that of the whole GABA_A receptor population. All these processes contribute to the dynamic nature of GABA_A receptor-mediated inhibition, but in most cases the significance of the interactions on pharmacology of intact nervous systems remains to be assessed.

One of the interacting proteins is GABA_A receptor-associated protein (GABARAP) (Wang et al., 1999), which interacts with the $\gamma 2$ subunit intracellularly, for the most part in the Golgi apparatus. GABARAP affects the receptor function in a heterologous expression system (Everitt et al., 2004), but it has not been found in synapses (Kittler et al., 2001). The actions of GABARAP are regulated by its post-translational lipid modifications (Chen et al., 2007).

Another interesting interacting protein is gephyrin (Prior et al., 1992) found originally to associate with strychnine-sensitive glycine receptors. It is needed for synaptic clustering and anchoring of $\alpha 2$ or $\gamma 2$ subunit-containing GABA_A receptors (Essrich et al., 1998; Jacob et al., 2005), thus stabilizing the inhibitory receptors. A 10-amino acid part of the $\alpha 2$ subunit has been identified as the critical domain for gephyrin interaction (Tretter et al., 2008).

Various kinases (e.g., protein kinase A, protein kinase C, Ca²⁺/calmodulin-dependent protein kinase II) and phosphatases (e.g., protein phosphatases PP1 α and PP2A) target mostly the GABA_A receptor β subunits (Jacob et al., 2008; Houston et al., 2009; Vithlani and Moss, 2009) and affect, for example, pharmacological sensitivity of the receptors to neurosteroids in a subunit-dependent manner (Koksma et al., 2003; Harney et al., 2003). Some ancillary proteins interact only with a selected GABA_A receptor subtype. For example, radixin, a member of the ezrin, radixin, and moesin family of proteins that link the actin cytoskeleton to the cell plasma membrane, binds to $\alpha 5$ subunits that form extrasynaptic $\alpha 5\beta 3\gamma 2/3$ receptor clusters on dendrites of hippocampal principal neurons (Loebrich et al., 2006). Quantum dot labeling of single receptor molecules have shown that individual receptor molecules can move laterally in the surface membrane (Bouzigués and Dahan, 2007) and

move in and out of synapses apparently independently of endocytosis (Bogdanov et al., 2006).

It is a major challenge to translate the novel findings on protein-protein interactions and processes of GABA_A receptors to possible mechanisms affecting pharmacological properties, such as drug sensitivity and tolerance development.

III. Regulation of GABA_A Receptor Expression By Pharmacological Agents

A. Benzodiazepines

BZs were developed in the mid-1950s and 1960s in response to a need for safe and effective anxiolytics (Sternbach, 1978; Shader and Greenblatt, 1993). Classic 1,4-benzodiazepines such as diazepam display a wide variety of behavioral effects, and they are clinically used as anticonvulsants, sedatives/hypnotics, anxiolytics, muscle relaxants, and preanesthetics (Ashton, 1994). They are especially effective in short-term treatment. However, therapy with BZs is often prolonged, which is associated with the most serious problems of BZ usage: development of tolerance and dependence (Ashton, 1991; Pétersson, 1994). Tolerance to sedative and ataxic effects of BZs usually develops at a faster rate than tolerance to their anxiolytic effects (File, 1985; Hutchinson et al., 1996). Physical dependence in preclinical experiments can be measured as the emergence of characteristic withdrawal signs upon cessation of the drug and/or administration of a BZ antagonist such as flumazenil (Licata and Rowlett, 2008). BZs are also addictive drugs; i.e., their use is associated with acute rewarding effects in animals (Heikkinen et al., 2009; Straub et al., 2010). In some humans, these effects may turn into BZ abuse and finally to compulsive drug-seeking behavior. The treatment of BZ addiction is very difficult, especially in patients with multiple, complicated drug addictions, and even a reduction of BZ dosing/usage is often a good treatment outcome (Vorma et al., 2002, 2004).

BZs exert their action by interacting with several GABA_A receptor subtypes with different pharmacological characteristics (Olsen and Sieghart, 2008). The majority of the pentameric GABA_A receptors are believed to be composed of α , β , and γ subunits in the ratio of 2:2:1, respectively (Sieghart and Sperk, 2002; Ernst et al., 2003). The BZ binding site is located at the interface between an α and a γ subunit, and its pharmacology is thus influenced by both α and γ subunits (Fig. 4) (Ernst et al., 2003; Ogris et al., 2004). Most classic BZs bind to $\alpha\beta\gamma 2$ receptors containing $\alpha 1$, $\alpha 2$, $\alpha 3$, or $\alpha 5$ subunits with approximately the same affinity. In contrast, several non-BZs, such as zolpidem, zaleplon, and abecarnil, have high affinity (low nanomolar) to $\alpha 1\beta\gamma 2$ receptors and intermediate affinity (high nanomolar) to $\alpha 2$ - and $\alpha 3$ -containing receptors, the affinity of zolpidem to $\alpha 5\beta\gamma 2$ receptors being very low (Korpi et al., 2002; Olsen and Sieghart, 2008).

Insensitivity of $\alpha 4$ - and $\alpha 6$ subunit-containing receptors to BZs is based on the presence of an arginine residue instead of a histidine at a conserved position in BZ binding site (residue 101 in Fig. 1) (Wieland et al., 1992). The need for the particular His residue has been used to generate knockin mutant mouse lines [$\alpha 1$ (H101R), $\alpha 2$ (H101R), $\alpha 3$ (H126R), $\alpha 5$ (H105R)] in which the particular arginine-containing receptor subtype is insensitive to classic BZs (for review, see Rudolph and Möhler, 2004). These knockin mouse lines have been used to demonstrate which subtypes of GABA_A receptors mediate specific behavioral actions of diazepam. The $\alpha 1$ -containing receptors seem to mediate sedative, anterograde amnesic, and antimyoclonic actions of diazepam (Rudolph et al., 1999). Anxiolytic activity of BZs is mediated by $\alpha 2$ -containing $\alpha\beta\gamma 2$ receptors, especially in the amygdala and hippocampus, whereas some anxiolytic activity is probably mediated by $\alpha 3$ -containing receptors (Löw et al., 2000; Crestani et al., 2001). Muscle relaxant activity of BZs is mediated partially by each of the $\alpha\beta\gamma 2$ receptor subtypes containing $\alpha 1$, $\alpha 2$, $\alpha 3$, or $\alpha 5$ subunits (Löw et al., 2000; Crestani et al., 2001, 2002). In addition, hippocampal extrasynaptic $\alpha 5$ -containing receptors are involved in learning and memory processes, such as trace fear conditioning (Crestani et al., 2002). It remains to be seen whether the roles of various GABA_A receptor subtypes in humans are similar to the roles the above rodent models suggest. For example, the BZ-site ligand ocinaplon exhibits $\alpha 1$ -preferring full agonist profile in human recombinant GABA_A receptors and anxiolytic activity without sedative effects in patients suffering from generalized anxiety disorder (Lippa et al., 2005). Anxioreselective partial agonists in rodents may have adverse sedative effects in man arising from the differences in $\alpha 1$ subunit-containing BZ binding sites. This may explain the difficulties found in translating the promising preclinical rodent-based results into clinical benefit in man (for example, see Atack, 2003).

Development of tolerance and dependence to BZs upon long-term administration suggests that continuous exposure of GABA_A receptors to BZs might affect receptor function by desensitizing the receptors and/or affecting receptor subunit expression and thereby changing the number and/or subtype of cell surface receptors. Decreased number of BZ binding sites after long-term BZ treatment has been observed in some studies (Rosenberg and Chiu, 1979; Miller et al., 1988, 1989), whereas no change has been found in most studies (Gallager et al., 1984; Heninger and Gallager, 1988; Ramsey-Williams et al., 1994). Long-term treatment of rats with BZs results in so-called "uncoupling," a decrease in the ability of BZs to potentiate the action of GABA on GABA_A receptors and in a decrease in the ability of GABA to potentiate BZ binding (Gallager et al., 1984; Marley and Gallager, 1989; Tietz et al., 1989). This uncoupling might be due to changes from BZ-sensitive to -insensitive receptor subtypes (changes in receptor subunit com-




FIG. 4. Three-dimensional homology model of the extracellular domains of the $\alpha 1$ and $\gamma 2$ subunits, indicating the interface between the subunits where the BZ binding pocket is located. The view is approximately perpendicular to the pore mouth, the bottom end of the figure corresponding to the C-terminal ends of the extracellular domains. The $\alpha 1$ subunit is depicted in yellow and the $\gamma 2$ subunit in blue. Residues $\alpha 1$ His101, $\alpha 1$ Tyr209, and $\gamma 2$ Phe77, indicated with gray arrows, reside in the so-called loops A, C, and D, respectively. The residues line the BZ binding pocket and interact with the binding BZ ligands. [Reproduced from Ogris W, Pörtl A, Hauer B, Ernst M, Oberto A, Wulff P, Höger H, Wisden W, and Sieghart W (2004) Affinity of various benzodiazepine site ligands in mice with a point mutation in the GABA_A receptor $\gamma 2$ subunit. *Biochem Pharmacol* 68:1621–1629. Copyright © 2004 Elsevier Science. Used with permission.]

bination) and/or changes in receptor function without changes in receptor subtype. Administration of the BZ antagonist flumazenil blocks the uncoupling, indicating that the changes are mediated via GABA_A receptor-associated BZ binding sites (Roca et al., 1990; Primus et al., 1996). Uncoupling, however, occurs too rapidly to be the mechanism responsible for BZ tolerance and dependence (for review, see Bateson, 2002).

The effects of long-term BZ administration on GABA_A receptor subunit expression have been studied using various BZ site compounds from several structural classes and with different pharmacological efficacy and receptor subtype selectivity. Treatment durations (7–32 days) and doses (0.25–150 mg/kg/day, the dose usually inversely correlating with efficacy)

used have often differed, which is apparently reflected as variability in the results between the studies. Most of the studies have been performed using Sprague-Dawley rats. Mice have been used in only a few studies. The use of mice instead of rats has been noted in the text and tables of this review. There are only a few studies using cultured neuronal cells. Cultured rat CGCs and hippocampal cells have been used to study the effects of long-term BZ treatment and withdrawal from the treatment on the expression of GABA_A receptor subunit mRNAs and polypeptides. Details for animals, treatment times, and doses used in the studies are listed in Tables 2 to 11.

1. $\alpha 1$ Subunit. The expression of the most abundant α subunit, the $\alpha 1$ subunit, has been studied most often.

Sedative doses of the compounds have been used in most in vivo studies. In the cerebral cortex, $\alpha 1$ mRNA expression was significantly down-regulated by diazepam, the general agonist for $\alpha 1/2/3/5\beta\gamma 2$ receptors in several (Heninger et al., 1990; Impagnatiello et al., 1996; Longone et al., 1996) but not all (Wu et al., 1994; Holt et al., 1996) studies (Table 2). Long-term treatment of rats with the general agonist lorazepam and the $\alpha 1$ -preferring agonist zolpidem down-regulated cerebral cortical $\alpha 1$ mRNA by 50 and 27%, respectively (Kang and Miller, 1991; Holt et al., 1997a). In contrast, long-term treatment with other full or partial agonists (e.g., abecarnil, alprazolam, flurazepam, imidazenil, and triazolam) were ineffective (Zhao et al., 1994a,b; Holt et al., 1996; Impagnatiello et al., 1996; Ramsey-Williams and Carter, 1996; Fahey et al., 1999; Tietz et al., 1999a). Long-term treatment with the inverse agonist FG 7142 slightly up-regulated $\alpha 1$ mRNA (Primus and Gallager, 1992). Withdrawal (2 days) from long-term flurazepam treatment reduced $\alpha 1$ mRNA in the cerebral cortical layers II to III and IV (Tietz et al., 1993), whereas on withdrawal day 7, $\alpha 1$ mRNA had returned to the control level (Tietz et al., 1999a) (Table 4). Long-term treatment with diazepam and flurazepam (but not imidazenil) reduced $\alpha 1$ polypeptide in the cerebral cortex (Impagnatiello et al., 1996; Pesold et al., 1997; Chen et al., 1999) (Table 6). This reduction was not detected on day 2 of withdrawal from long-term flurazepam treatment (Tietz et al., 1999b) (Table 7).

In the cerebellum, none of the above-mentioned BZ-site compounds affected $\alpha 1$ mRNA expression except for a 21% up-regulation by diazepam (Holt et al., 1999). Treatment of cultured rat CGCs with diazepam reduced $\alpha 1$ mRNA expression slightly (Follesa et al., 2001a, 2002), but treatments with imidazenil, zaleplon, or zolpidem were ineffective (Table 8). Withdrawal (6 h) of CGCs from diazepam, imidazenil, zaleplon, and zolpidem decreased $\alpha 1$ mRNA by 20 to 37% (Table 9). Long-term treatment of CGCs with diazepam, flunitrazepam, and bretazenil reduced the level of $\alpha 1$ polypeptide (Brown and Bristow, 1996; Brown et al., 1998; Johnston and Bristow, 1998; Follesa et al., 2001a), whereas imidazenil had no effect on it (Johnston and Bristow, 1998) (Table 10). Withdrawal (6 h) from long-term diazepam treatment down-regulated $\alpha 1$ polypeptide in CGCs by 75% (Follesa et al., 2001a) (Table 11).

Effects of long-term BZ treatments on GABA_A receptor subunit expression have been extensively studied in the hippocampus. Long-term treatment with diazepam either down-regulated (Wu et al., 1994; Impagnatiello et al., 1996) or had no effect (Heninger et al., 1990) on hippocampal $\alpha 1$ mRNA expression. Long-term treatment with flurazepam, imidazenil, or lorazepam had no effect on the expression (Kang and Miller, 1991; Zhao et al., 1994a,b; Impagnatiello et al., 1996), whereas FG 7142 up-regulated it (Primus and Gallager, 1992). The use of in situ hybridization revealed down-regulation of

$\alpha 1$ mRNA in the hippocampal CA1 region and in dentate gyrus granule cells in brain sections from rats continuously treated with flurazepam (Tietz et al., 1999a). Long-term treatment of mice with alprazolam or rats with triazolam had no effect on $\alpha 1$ mRNA expression in any hippocampal subregion (Ramsey-Williams and Carter, 1996; Fahey et al., 1999). Withdrawal (2 days) from long-term flurazepam treatment slightly reduced $\alpha 1$ mRNA in the CA1 region, the expression returning to control level in 7 days (Tietz et al., 1993, 1999a). Long-term treatment with flurazepam down-regulated the $\alpha 1$ polypeptide in hippocampal CA1 and CA3 regions and dentate gyrus (Chen et al., 1999). On day 2 of withdrawal from long-term treatment with flurazepam, down-regulation of $\alpha 1$ polypeptide persisted in the stratum oriens region of the CA1 (Tietz et al., 1999b). Long-term treatment of cultured rat hippocampal cells with lorazepam decreased $\alpha 1$ mRNA by 19%, which returned to control level during a withdrawal of 6 h. Long-term treatment of cultured primary hippocampal cells with lorazepam reduced $\alpha 1$ polypeptide by 36% (Sanna et al., 2005), remaining reduced by 26% after 6-h withdrawal. Treatment of cultured hippocampal cells with etizolam or withdrawal from it had no effect on $\alpha 1$ mRNA or polypeptide (Sanna et al., 2005). Triazolam reduced $\alpha 1$ mRNA in the diagonal band and up-regulated it in the nucleus basalis, substantia nigra pars reticulata, and inferior colliculus (Ramsey-Williams and Carter, 1996).

Overall, there are controversies in results of studies on long-term BZ treatments on expression of $\alpha 1$ subunit. Many of the studies indicate down-regulation of $\alpha 1$ mRNA and polypeptide after long-term treatment and especially on withdrawal in the cerebral cortex and in hippocampus. There is a tight control between $\alpha 1$ mRNA and $\alpha 1$ polypeptide expression that was particularly clearly manifested in the GABA_A receptor $\alpha 6$ subunit knockout mice, where an alteration in the structure of $\alpha 6$ gene within the $\beta 2$ - $\alpha 6$ - $\alpha 1$ - $\gamma 2$ gene cluster altered the expression of the other genes in the cluster (Uusi-Oukari et al., 2000). Down-regulation of $\alpha 1$ mRNA transcription in the forebrain of $\alpha 6$ knockout mice was accompanied with similar down-regulation of forebrain $\alpha 1$ polypeptide (Uusi-Oukari et al., 2000).

2. $\alpha 2$ Subunit. Long-term treatment with abecarnil, diazepam, imidazenil, and zolpidem in vivo had no effect on $\alpha 2$ mRNA expression in any brain region quantified (Wu et al., 1994; Holt et al., 1996; Impagnatiello et al., 1996) (Table 2). Long-term treatment with alprazolam up-regulated $\alpha 2$ mRNA in the brainstem by 76% (Tanay et al., 2001). Of all brain regions studied, the caudate-putamen was the only one in which flurazepam down-regulated $\alpha 2$ mRNA (Tietz et al., 1999a). Triazolam down-regulated $\alpha 2$ mRNA in the prefrontal, olfactory, and cingulate cortices and in several other brain regions, up-regulated it in the hippocampal CA3 region, dentate gyrus, and some other brain regions, but had no effect in many of the regions studied (Ramsey-Williams and

TABLE 2
Effect of long-term benzodiazepine treatment in vivo on GABA_A receptor subunit mRNA expression

Compound, Species, Dose, Duration, Brain Region	Subunit mRNA						Reference
	$\alpha 1$	$\alpha 2$	$\alpha 3$	$\alpha 4$	$\alpha 5$	$\alpha 6$	
	% change						
Abecarnil							
Rat, 6 mg/kg/d, 14 d							Holt et al., 1996
Cerebral cortex	N.S.	N.S.	N.S.	N.S.	N.S.	—	
Alprazolam							
Mouse, 28 d							Fahey et al., 1999
Cerebral cortex							
Layers II–IV	N.S.						
Layer V	N.S.						
Layer VI	N.S.						
Hippocampus							
CA1	N.S.						
CA3	N.S.						
Dentate gyrus	N.S.						
Rat, 10 mg/kg/d, 21 d							Tanay et al., 2001
Brainstem	N.S.	+76	N.S.	N.S.	N.S.	—	
Diazepam							
Rat, 0.25 μ g/g tissue, ^a 21 d							Heninger et al., 1990
Cerebral cortex	–25						
Cerebellum	N.S.						
Hippocampus	N.S.						
Rat, 0.25 μ g/g tissue, ^a 21 d							Wu et al., 1994
Cerebral cortex	N.S.	N.S.	N.S.	N.S.	–28		
Cerebellum	N.S.						
Hippocampus	–40	N.S.		N.S.	–15		
Rat, 15 mg/kg/d, 14 d							
Cerebral cortex	N.S.	N.S.	+36	+50	+42	—	Holt et al., 1996
Cerebellum	+21						Holt et al., 1999
Rat, 3 \times 5 to 3 \times 20 mg/kg/d, 14 d							Impagnatiello et al., 1996
Frontoparietal motor cortex	–42	N.S.	N.S.		+30	—	
Frontoparietal somatosensory cortex	N.S.	N.S.	N.S.		N.S.	—	
Cerebellum	N.S.	N.S.	N.S.		N.S.	—	
Hippocampus	–20	N.S.	N.S.		N.S.	—	
Olfactory bulb	N.S.						
Striatum	N.S.						
Rat, 3 \times 5 to 3 \times 20 mg/kg/d, 14 d							Longone et al., 1996
Frontoparietal motor cortex	–30				+37		
Rat, 15 mg/kg/d, 14 d							Arnot et al., 2001
Cerebral cortex							
Minipump infusion	N.S.	N.S.	N.S.	–18	N.S.		
One daily injection	N.S.	N.S.	N.S.	+19	N.S.		
FG 7142							
Rat, 0.25 g/g, 8 d							Primus and Gallager, 1992
Cerebral cortex	+17						
Cerebellum	N.S.						
Hippocampus	+34						
Flurazepam							
Rat, 40 mg/kg/d, 32 d							O'Donovan et al., 1992b
Whole brain	N.S.	N.S.	+50		–37	+50	
Rat, 150 mg/kg/d, 14d							Zhao et al., 1994b
Cerebral cortex	N.S.				–50		
Hippocampus	N.S.				–50		
Rat, 150 mg/kg/d, 28 d							Zhao et al., 1994b
Cerebral cortex	N.S.				N.S.		
Cerebellum	N.S.						
Hippocampus	N.S.				N.S.		
Rat, 100 mg/kg/day, 7 d							Tietz et al., 1999a
Hippocampus							
CA1	–14	N.S.		N.S.	N.S.	—	
CA2	N.S.	N.S.		N.S.	N.S.	—	
CA3	N.S.	N.S.		N.S.	N.S.	—	
Dentate polymorphic cells	N.S.	N.S.		N.S.	N.S.	—	
Granule cells	–20	N.S.		N.S.	N.S.	—	
Cerebral cortex							
Frontal	N.S.	N.S.		N.S.	N.S.	—	
Parieto-occipital	N.S.	N.S.		N.S.	—	—	
Caudate-putamen	—	–32		—	—	—	
Thalamus	N.S.	N.S.		N.S.	—	—	
Cerebellum, granule cell layer	N.S.	N.S.		N.S.	—	N.S.	
Imidazenil							
Rat, 3 \times 1 to 3 \times 4 mg/kg/d, 14 d							Impagnatiello et al., 1996
Frontoparietal motor cortex	N.S.	N.S.	N.S.		N.S.	—	

Continued

TABLE 2—Continued

Compound, Species, Dose, Duration, Brain Region	Subunit mRNA						Reference
	$\alpha 1$	$\alpha 2$	$\alpha 3$	$\alpha 4$	$\alpha 5$	$\alpha 6$	
Frontoparietal somatosensory cortex	N.S.	N.S.	N.S.		N.S.	—	
Cerebellum	N.S.	N.S.	N.S.		N.S.	—	
Hippocampus	N.S.	N.S.	N.S.		N.S.	—	
Striatum	N.S.						
Lorazepam							
Mouse, 2 mg/kg/d, 14–28 d							Kang and Miller, 1991
Cerebral cortex	–50						
Cerebellum	N.S.						
Hippocampus	N.S.						
Triazolam							
Rats, 6 mg/kg/d, 28 d							Ramsey-Williams and Carter, 1996
Prefrontal cortex	N.S.	–41	N.S.				
Olfactory cortex	N.S.	–28	N.S.				
Diagonal band	–20	N.S.	N.S.				
Cingulate cortex	N.S.	–24	N.S.				
Sensory cortex	N.S.	N.S.	N.S.				
Fornix	N.S.	N.S.	+43				
Nucleus basalis	+32	N.S.	N.S.				
Dorsal raphe	N.S.	N.S.	N.S.				
Anteroventral thalamus	N.S.	N.S.	N.S.	+30			
Central amygdala	N.S.	N.S.	N.S.				
Hippocampus							
CA1	N.S.	N.S.	N.S.		N.S.		
CA3	N.S.	+30	N.S.		N.S.		
Dentate gyrus	N.S.	+20	N.S.		N.S.		
Subthalamus	N.S.						
Lateral geniculate	N.S.	–21	–39				
Mammillary body	N.S.	–46	N.S.				
Substantia nigra reticulata	+32	N.S.	N.S.				
Substantia nigra compacta	N.S.	–28	N.S.				
Entorhinal cortex	N.S.	N.S.	–30				
Inferior colliculus	+26	–34	N.S.				
Periaqueductal grey	N.S.	–27	N.S.				
Zolpidem							
Rats, 15 mg/kg/d, 7 and 14 d							Holt et al., 1997a
Cerebral cortex	–27 ^{14d}	N.S.	N.S.	+71 ^{7d}	N.S.	—	

d, day(s); N.S., not significantly different from the control value; —, not detected.

^a Continuous concentration in brain tissue.

Carter, 1996). Flurazepam withdrawal had no effect on $\alpha 2$ mRNA expression (Tietz et al., 1999a). Treatment of cultured rat hippocampal cells with 10 μ M lorazepam or etizolam had no effect on the expression of $\alpha 2$ mRNA, but withdrawal (6 h) from them slightly increased the expression (Sanna et al., 2005). The $\alpha 2$ polypeptide was not affected by long-term treatment with diazepam or flurazepam, but imidazenil down-regulated it in the frontoparietal cortex (Pesold et al., 1997; Chen et al., 1999).

In contrast to $\alpha 1$, studies with most BZs suggest no effect of long-term BZ treatment or withdrawal from the treatment on $\alpha 2$ subunit expression. Triazolam might be an exception in having heterogeneous and brain region-specific effects on $\alpha 2$ expression.

3. $\alpha 3$ Subunit. Holt et al. (1996) found that long-term treatment with diazepam up-regulated $\alpha 3$ mRNA expression in the cerebral cortex, but other studies (Wu et al., 1994; Impagnatiello et al., 1996) found no such effect. Long-term treatment with abecarnil, imidazenil, triazolam, and zolpidem had no effect on cerebral cortical $\alpha 3$ mRNA expression (Wu et al., 1994; Holt et al., 1996, 1997a; Impagnatiello et al., 1996; Ramsey-Williams and Carter, 1996). Triazolam up-regulated $\alpha 3$

mRNA in the fornix and down-regulated it in the lateral geniculate nucleus and entorhinal cortex (Ramsey-Williams and Carter, 1996). Treatment of cultured rat hippocampal cells with lorazepam increased $\alpha 3$ mRNA by 59%, whereas a similar treatment with etizolam had no effect on the expression (Sanna et al., 2005). Withdrawal from lorazepam and etizolam (6 h) increased $\alpha 3$ mRNA by 38 and 30%, respectively. The $\alpha 3$ polypeptide was up-regulated in frontoparietal somatosensory cortex by long-term treatment with diazepam but not imidazenil (Pesold et al., 1997).

In conclusion, studies in vivo indicate that $\alpha 3$ expression, although not affected by most BZs, is regulated by long-term treatment with some BZs. The effects are brain region-specific and, in contrast to $\alpha 1$, $\alpha 3$ expression is usually up-regulated.

4. $\alpha 4$ Subunit. Expression of $\alpha 4$ mRNA was up-regulated in the cerebral cortex of rats by long-term treatment with diazepam or zolpidem (Holt et al., 1996, 1997a) and in the anteroventral thalamus by long-term treatment with triazolam (Ramsey-Williams and Carter, 1996). Long-term treatment of rats with diazepam continuously infused with minipumps down-regulated cortical $\alpha 4$ mRNA, whereas the same dose given as subcu-

TABLE 3
Effect of long-term benzodiazepine treatment in vivo on GABA_A receptor subunit β, γ and δ mRNA expression

Compound, Species, Dose, Duration, Brain Region	Subunit mRNA							Reference
	β1	β2	β3	γ1	γ2	γ3	δ	
	% change							
Abecarnil								
Rat, 6 mg/kg/d, 14 d								Holt et al., 1996
Cerebral cortex	N.S.	-27	N.S.	N.S.	-41		N.S.	
Alprazolam								
Rat, 10 mg/kg/d, 21 d								Tanay et al., 2001
Brainstem	N.S.	-24	N.S.	N.S.	N.S.		N.S.	
Mouse, 2 mg/kg/d, 28 d								Fahey et al., 1999
Cerebral cortex								
Layers II-IV					N.S.			
Layer V					N.S.			
Layer VI					N.S.			
Hippocampus								
CA1					N.S.			
CA3					N.S.			
Dentate gyrus					+27			
Diazepam								
Rat, 0.25 μg/g tissue, ^a 21 d								Heninger et al., 1990
Cerebral cortex	N.S.							
Cerebellum	N.S.							
Hippocampus	N.S.							
Rat, 0.25 μg/g tissue, ^a 21 d								Primus and Gallager, 1992
Cerebral cortex					-10			
Cerebellum					N.S.			
Hippocampus					N.S.			
Rat, 0.25 μg/g tissue, ^a 21 d								Wu et al., 1994
Cerebral cortex		N.S.	N.S.		-40			
Cerebellum		N.S.	N.S.		N.S.			
Hippocampus		N.S.	N.S.		N.S.			
Rat, 15 mg/kg/d, 14 d								Holt et al., 1996
Cerebral cortex	+32	N.S.	N.S.	N.S.	-26	+57		
Cerebellum		N.S.			+58			Holt et al., 1999
Rat, 3×5 to 3×20 mg/kg/d, 14 d								Impagnatiello et al., 1996
Frontoparietal motor cortex		N.S.		N.S.	-30,-50 ^b		N.S.	
Frontoparietal somatosensory cortex		N.S.		N.S.	N.S.		N.S.	
Cerebellum		N.S.		N.S.	N.S.		N.S.	
Hippocampus		N.S.		N.S.	N.S.		N.S.	
Rat, 3×5 to 3×20 mg/kg/d, 14 d								Longone et al., 1996
Frontoparietal motor cortex					-30,-50 ^b			
Rat, 15 mg/kg/d, 14 d								Arnot et al., 2001
Cerebral cortex,								
Minipump infusion	N.S.	N.S.	N.S.	+37	N.S.		N.S.	
One daily injection	N.S.	N.S.	+21	N.S.	N.S.		N.S.	
FG 7142								
Rat, 0.25 mg/kg tissue, ^a 8 d								Primus and Gallager, 1992
Cerebral cortex	N.S.				+27			
Cerebellum	N.S.				N.S.			
Hippocampus	—				N.S.			
Flurazepam								
Rat, 40 mg/kg/d, 32 d								O'Donovan et al., 1992a
Whole brain	N.S.	N.S.	N.S.		N.S.			
Rat, 150 mg/kg/d, 14 d								Zhao et al., 1994b
Cerebral cortex					N.S.			
Hippocampus					N.S.			
Rat, 150 mg/kg/d, 28 d								Zhao et al., 1994b
Cerebral cortex					-31			
Cerebellum					N.S.			
Hippocampus					-39			
Rat, 150 mg/kg/d, 28 d								Zhao et al., 1994a
Cerebral cortex		-27	N.S.					
Cerebellum		-49	-35					
Hippocampus		-48	-30					
Rat, 100 mg/kg/day, 7 d								Tietz et al., 1999a
Hippocampus								
CA1	N.S.	N.S.	-24		N.S.			
CA2	N.S.	N.S.	-15		N.S.			
CA3	N.S.	N.S.	-18		N.S.			
Dentate cells	N.S.	N.S.	N.S.		N.S.			
Granule cells	N.S.	N.S.	-21		N.S.			
Cerebral cortex frontal	N.S.	N.S.	-37		N.S.			
Parieto-occipital	N.S.	N.S.	N.S.		N.S.			
Caudate-putamen	—	—	-29		N.S.			

Continued

TABLE 3—Continued

Compound, Species, Dose, Duration, Brain Region	Subunit mRNA							Reference
	$\beta 1$	$\beta 2$	$\beta 3$	$\gamma 1$	$\gamma 2$	$\gamma 3$	δ	
Thalamus	—	N.S.	—		N.S.			
Cerebellum, granule cell layer	N.S.	N.S.	N.S.		N.S.			
Imidazenil								
Rat, 3×1 to 3×4 mg/kg/d, 14 d								Impagnatiello et al., 1996
Frontoparietal motor cortex		N.S.		N.S.	N.S.		N.S.	
Frontoparietal somatosensory cortex		N.S.		N.S.	N.S.		N.S.	
Cerebellum		N.S.		N.S.	N.S.		N.S.	
Hippocampus		N.S.		N.S.	N.S.		N.S.	
Lorazepam								
Mouse, 2 mg/kg/d, 14–28 d								Kang and Miller, 1991
Cerebral cortex					–50			
Cerebellum					N.S.			
Hippocampus					N.S.			
Triazolam								
Rat, 6 mg/kg/d, 28 d								Ramsey-Williams and Carter, 1996
Olfactory cortex			–43					
Cingulate cortex	+62							
Sensory cortex			–46					
Dorsal raphe							–46	
Anteroventral thalamus							–47	
Central amygdala			–42					
Hippocampus CA1	+32							
Subthalamus							–58	
Lateral geniculate			–48				–39	
Substantia nigra compacta							–51	
Interpeduncular nucleus	–45	–49						
Periaqueductal grey							–51	
Zolpidem								
Rat, 15 mg/kg/d, 7 and 14 d								Holt et al., 1997a
Cerebral cortex	+49 ^{7d}	N.S.	N.S.	N.S.	N.S.		N.S.	

d, day(s); N.S., not significantly different from the control value; —, not detected.

^a Continuous concentration in brain tissue.

^b Two values represent S and L splice variants, respectively.

taneous injections up-regulated it (Arnot et al., 2001). No effect on $\alpha 4$ mRNA expression was found for long-term treatment with abecarnil, alprazolam, or flurazepam (Table 2). Flurazepam withdrawal did not affect $\alpha 4$ subunit expression (Tietz et al., 1999a). Treatment of cultured rat CGCs with diazepam, imidazenil, zaleplon, or zolpidem had no effect on $\alpha 4$ mRNA expression (Follesa et al., 2001a, 2002), but 6-h withdrawal from them increased the expression by 32 to 59%. Long-term treatment of rat CGC cultures with diazepam had no effect on $\alpha 4$ polypeptide (Follesa et al., 2001a), but the withdrawal from the treatment increased it by 45%. Long-term treatment of cultured rat hippocampal cells with lorazepam or etizolam had no effect on $\alpha 4$ mRNA or polypeptide expression (Sanna et al., 2005), but the withdrawal from lorazepam (6 h) increased the mRNA and polypeptide expression by 23 and 81%, respectively. Etizolam withdrawal had no effects on $\alpha 4$ mRNA or polypeptide expression (Sanna et al., 2005).

Overall, studies on the effects of long-term treatment with BZ on $\alpha 4$ expression suggest strong up-regulation of the subunit by some BZs. In cultured CGCs and hippocampal neurons, $\alpha 4$ was up-regulated on BZ withdrawal. These findings are especially interesting because the ligands tested do not bind to $\alpha 4$ subunit-containing GABA_A receptors.

5. $\alpha 5$ Subunit. Long-term treatment of rats with diazepam either increased (Holt et al., 1996; Impagna-

tello et al., 1996; Longone et al., 1996) or reduced (Wu et al., 1994) cerebral cortical $\alpha 5$ mRNA expression. A 2-week treatment with flurazepam down-regulated cerebral cortical and hippocampal $\alpha 5$ mRNA expression for 50%, whereas after a 4-week treatment, the expressions returned to same level as in control rat cerebral cortex (Zhao et al., 1994b). Long-term treatment with abecarnil, imidazenil, triazolam and zolpidem in vivo had no effect on cortical $\alpha 5$ mRNA expression (Holt et al., 1996, 1997a; Impagnatiello et al., 1996; Ramsey-Williams and Carter, 1996). Hippocampal $\alpha 5$ mRNA was slightly down-regulated by long-term treatment with diazepam in the study by Wu et al. (1994) but not in that of Impagnatiello et al. (1996). Long-term treatment with imidazenil and triazolam did not affect hippocampal $\alpha 5$ mRNA expression (Impagnatiello et al., 1996; Ramsey-Williams and Carter, 1996). Treatment with diazepam, but not imidazenil, produced a drastic up-regulation of $\alpha 5$ polypeptide in frontoparietal motor and somatosensory cortices (Impagnatiello et al., 1996; Pesold et al., 1997). Treatment of cultured rat hippocampal cells with etizolam slightly reduced $\alpha 5$ mRNA expression, whereas lorazepam had no effect (Sanna et al., 2005). Etizolam or lorazepam withdrawal had no effect on $\alpha 5$ mRNA expression (Sanna et al., 2005).

In summary, long-term BZ treatment and withdrawal do not much influence $\alpha 5$ mRNA expression. Up-regulation of $\alpha 5$ polypeptide in the frontoparietal motor and

TABLE 4

Effect of withdrawal from long-term benzodiazepine treatment in vivo on GABA_A receptor α subunit mRNA expression

Compound, Species, Dose, Duration, Withdrawal Time, Brain Region	Subunit mRNA					Reference
	α 1	α 2	α 4	α 5	α 6	
% change						
Diazepam						
Rat, 0.25 μ g/g tissue, 21 d, 2 d						
Cerebral cortex				N.S.		Wu et al., 1994
Hippocampus	N.S.			N.S.		
Rat, 3 \times 5 to 3 \times 20 mg/kg/d, 14 d, 18 h						
Frontoparietal motor cortex	-31			+31		Longone et al., 1996
Hippocampus	-24			N.S.		
Frontoparietal somatosensory cortex	N.S.					
Olfactory bulb	N.S.					
Striatum	N.S.					
Cerebellum	N.S.					
Rat, 3 \times 5 to 3 \times 20 mg/kg/d, 14 d, 3 d						
Frontoparietal motor cortex	N.S.			N.S.		Longone et al., 1996
Hippocampus	N.S.			N.S.		
Flurazepam						
Rat, 150 mg/kg/d, 7 d, 2 d						
Cerebral cortex	-50 ^{Layers II-III, IV}					Tietz et al., 1993
Hippocampus	-35 ^{CA1 region}			N.S.		
Rat, 100 mg/kg/d, 7 d, 2 d						
Hippocampus						Tietz et al., 1999a
CA1	-11	N.S.	N.S.	N.S.	—	
CA2	N.S.	N.S.	N.S.	N.S.	—	
CA3	N.S.	N.S.	N.S.	N.S.	—	
Dentate polymorphic cells	N.S.	N.S.	N.S.	N.S.	—	
Granule cells	N.S.	N.S.	N.S.	N.S.	—	
Cerebral cortex						
Frontal	N.S.	N.S.	N.S.	N.S.	—	
Parieto-occipital	N.S.	N.S.	N.S.	—	—	
Caudate-putamen	—	N.S.	—	—	—	
Thalamus	N.S.	N.S.	N.S.	—	—	
Cerebellum, granule cell layer	N.S.	N.S.	N.S.	—	N.S.	
Rat, 100 mg/kg/d, 7 d, 7 d						
Hippocampus						Tietz et al., 1999a
CA1	N.S.	N.S.	N.S.	N.S.	—	
CA2	N.S.	N.S.	N.S.	N.S.	—	
CA3	N.S.	N.S.	N.S.	N.S.	—	
Dentate polymorphic cells	N.S.	N.S.	N.S.	N.S.	—	
Granule cells	N.S.	N.S.	N.S.	N.S.	—	
Cerebral cortex						
Frontal	N.S.	N.S.	N.S.	N.S.	—	
Parieto-occipital	N.S.	N.S.	N.S.	—	—	
Caudate-putamen	—	N.S.	—	—	—	
Thalamus	N.S.	N.S.	N.S.	—	—	
Cerebellum, granule cell layer	N.S.	N.S.	N.S.	—	N.S.	
Imidazenil						
Rat, 3 \times 5 to 3 \times 20 mg/kg/d, 14 d, 18 h						
Frontoparietal motor cortex	N.S.			N.S.		Longone et al., 1996

d, day(s); N.S., not significantly different from the control value; —, not detected.
^a Continuous concentration in brain tissue.

somatosensory cortices by diazepam seems to be an exception (Impagnatiello et al., 1996). However, using mutant α 5(H105R) mice possessing diazepam-insensitive α 5 β 2 receptors, it has been suggested that α 5-containing receptors are required for development of tolerance to the sedative action of diazepam (van Rijnsoever et al., 2004). After long-term diazepam treatment, the mutant α 5(H105R) mice developed no sedative tolerance to diazepam (van Rijnsoever et al., 2004).

6. α 6 Subunit. Increased α 6 mRNA expression in whole brain has been found after long-term flurazepam treatment by O'Donovan et al. (1992a), whereas no effect of flurazepam was found in cerebellar α 6 mRNA (Tietz et al., 1999a). Withdrawal from flurazepam did not affect α 6 mRNA expression (Tietz et al., 1999a). Long-

term treatment with flunitrazepam had no effect on α 6 mRNA expression in cultured GCGs (Brown and Bristow, 1996). The results do not suggest significant regulation of cerebellar α 6 subunit expression by long-term BZ administration.

7. β 1 Subunit. Holt et al. (1996) found that long-term treatment with diazepam up-regulated cerebral cortical β 1 mRNA expression, but Heninger et al. (1990) found it to be ineffective. Abecarnil, FG 7142, and flurazepam had no effect on cortical β 1 mRNA (Primus and Gallager, 1992; Holt et al., 1996; Tietz et al., 1999a), whereas zolpidem up-regulated it (Holt et al., 1997a). Long-term treatment with diazepam, FG 7142, and flurazepam had no effect on cerebellar β 1 mRNA (Heninger et al., 1990; Primus and Gallager, 1992; Tietz et al., 1999a). Hippocampal β 1

TABLE 5
Effect of withdrawal from long-term benzodiazepine treatment in vivo on GABA_A receptor subunit β and γ mRNA expression

Compound, Species, Dose, Duration, Withdrawal Time, Brain Region	Subunit mRNA				Reference
	β1	β2	β3	γ2	
% change					
Diazepam					
Rat, 0.25 μg/g tissue, 21 d, 2 d Cerebral cortex				N.S.	Wu et al., 1994
Rat, 3×5 to 3×20 mg/kg/d, 14 d, 18 h Frontoparietal motor cortex				-38, -52 ^a	Longone et al., 1996
Hippocampus				N.S., N.S.	
Rat, 3×5 to 3×20 mg/kg/d, 14 d, 3 d Frontoparietal motor cortex				N.S., N.S. ^a	Longone et al., 1996
Hippocampus				N.S., N.S. ^a	
Flurazepam					
Rat, 150 mg/kg/d, 7 d, 2 d Hippocampus				N.S.	Tietz et al., 1993
Rat, 150 mg/kg/d, 28 d, 2 d Cerebral cortex				N.S.	Zhao et al., 1994b
Cerebellum				N.S.	
Hippocampus				N.S.	
Rat, 150 mg/kg/d, 28 d, 2 d Cerebral cortex		N.S.	N.S.		Zhao et al., 1994a
Cerebellum		N.S.	N.S.		
Hippocampus		N.S.	-11		
Rat, 100 mg/kg/d, 7 d, 2 d Hippocampus					Tietz et al., 1999a
CA1	N.S.	+31	-17	N.S.	
CA2	N.S.	N.S.	-18	N.S.	
CA3	N.S.	+37	-14	N.S.	
Dentate polymorphic cells	N.S.	N.S.	-17	N.S.	
Granule cells	N.S.	+29	-15	N.S.	
Cerebral cortex					
Frontal	N.S.	N.S.	N.S.	N.S.	
Parieto-occipital	N.S.	N.S.	N.S.	N.S.	
Caudate-putamen	—	—	N.S.	N.S.	
Thalamus	—	N.S.	—	N.S.	
Cerebellum, granule cell layer	N.S.	N.S.	-20	N.S.	
Rat, 100 mg/kg/d, 7 d, 7 d Hippocampus					Tietz et al., 1999a
CA1	N.S.	N.S.	N.S.	N.S.	
CA2	N.S.	N.S.	N.S.	N.S.	
CA3	N.S.	N.S.	+18	N.S.	
Dentate polymorphic cells	N.S.	N.S.	N.S.	N.S.	
Granule cells	N.S.	N.S.	N.S.	N.S.	
Cerebral cortex					
Frontal	N.S.	N.S.	+40	+34	
Parieto-occipital	N.S.	N.S.	N.S.	N.S.	
Caudate-putamen	—	—	N.S.	N.S.	
Thalamus	—	N.S.	—	N.S.	
Cerebellum, granule cell layer	N.S.	N.S.	N.S.	N.S.	
Imidazenil					
Rat, 3×5 to 3×20 mg/kg/d, 14 d, 18 h Frontoparietal motor cortex				N.S., N.S. ^a	Longone et al., 1996

d, day(s); N.S., not significantly different from the control value; —, not detected.

^a Two values represent S and L splice variants, respectively.

mRNA was not affected by long-term treatment with diazepam (Heninger et al., 1990) or flurazepam (Tietz et al., 1999a). Long-term treatment with flurazepam did not affect β1 polypeptide expression in the hippocampus (Chen et al., 1999). Triazolam up-regulated β1 mRNA in the cingulate cortex and hippocampal CA1 region and down-regulated it in the interpeduncular nucleus (Ramsey-Williams and Carter, 1996). Flurazepam withdrawal for 2 or 7 days had no effect on β1 mRNA expression (Tietz et al., 1999a). Treatment of rat cultured CGCs with diazepam increased β1 expression by 47% (Follesa et al., 2002). Similar treatment with zaleplon or zolpidem had no effect on β1 expression (Follesa et al., 2002). Withdrawal from diazepam, zaleplon, and zolpidem increased β1 mRNA expres-

sion (31–57%) (Follesa et al., 2002). Most in vivo studies suggest that β1 subunit is not affected by long-term BZ treatment.

8. β2 Subunit. A reduction in β2 mRNA expression in vivo was found after long-term treatment with abecarnil in the cerebral cortex (Holt et al., 1996), with alprazolam in the brainstem (Tanay et al., 2001), with flurazepam in the cerebral cortex, cerebellum, and hippocampus (Zhao et al., 1994a), and with triazolam in the interpeduncular nucleus (Ramsey-Williams and Carter, 1996) (Table 3). No effects on β2 mRNA expression in any brain region studied were found after long-term treatment with diazepam, flurazepam (Tietz et al., 1999a), imidazenil, or zolpidem (Wu et al., 1994; Holt et

TABLE 6
Effect of long-term benzodiazepine treatment in vivo on GABA_A receptor subunit polypeptide expression

Compound, Species, Dose, Duration, Brain Region	Subunit mRNA					Reference
	$\alpha 1$	$\alpha 2$	$\alpha 5$	$\beta 2$	$\beta 3$	
% change						
Diazepam						
Rat, 3×5 to 3×20 mg/kg/d, 14 d						
Frontoparietal motor cortex	-37		+158	+47 ^{$\beta 2/3$}		+50
Frontoparietal somatosensory cortex	N.S.		+209	N.S. ^{$\beta 2/3$}		N.S.
Rat, 3×5 to 3×20 mg/kg/d, 14 d						
Frontoparietal motor cortex	-37 ^{LIII-IV}	N.S.	+150	+48 ^{$\beta 2/3$}		+48
Frontoparietal somatosensory cortex	N.S.	N.S.	+221	N.S. ^{$\beta 2/3$}		N.S.
Flurazepam						
Rat, 100 mg/kg/day, 7 d						Chen et al., 1999
Hippocampus						
CA1	-20	N.S.		N.S.	-19	N.S.
CA2	N.S.	N.S.		N.S.	-25 ^{SR}	N.S.
CA3	-38	N.S.		N.S.	-14 ^{SL}	N.S.
Dentate gyrus	-34	N.S.		N.S.	-18	-8 ^{ML}
Cerebral cortex						
Layer I	N.S.				N.S.	N.S.
Layer II/III	-24 ^{Fr}				-23 ^{Fr,Par}	N.S.
Layer IV	-26 ^{Par}				-20 ^{Par}	N.S.
Layer V	-19 ^{Fr,Occ}				-42 ^{Fr}	N.S.
Layer VI	-23				-38 ^{Fr}	N.S.
Caudate-putamen	—	N.S.		—	N.S.	N.S.
Thalamus	N.S.	—		N.S.	—	N.S.
Substantia nigra reticulata	N.S.	—		N.S.	—	N.S.
Inferior colliculus	N.S.	—		N.S.	—	N.S.
Superior colliculus	—	N.S.		—	N.S.	N.S.
Cerebellum	N.S.	—		N.S.	N.S.	N.S.
Imidazenil						
Rat, 3×1 to 3×4 mg/kg/d, 14 d						Impagnatiello et al., 1996
Frontoparietal motor cortex	N.S.		N.S.	N.S. ^{$\beta 2/3$}		N.S.
Frontoparietal somatosensory cortex	N.S.		N.S.	N.S. ^{$\beta 2/3$}		N.S.
Rat, 3×1 to 3×4 mg/kg/d, 14 d						Pesold et al., 1997
Frontoparietal motor cortex	N.S.	N.S.	N.S.	N.S. ^{$\beta 2/3$}		N.S.
Frontoparietal somatosensory cortex	N.S.	-22 ^{LIII-IV}	N.S.	N.S. ^{$\beta 2/3$}		N.S.

d, day(s); N.S., not significantly different from the control value; —, not detected; $\beta 2/3$, detected using an antibody specific to $\beta 2$ and $\beta 3$ subunits; LIII-IV, layers III and IV; SR, stratum radiatum; SL, stratum lacunosum; ML, molecular layer; Fr, frontal; Par, parietal; Occ, occipital.

al., 1996, 1997a, 1999; Impagnatiello et al., 1996) (Table 3). Flurazepam withdrawal for 2 days up-regulated $\beta 2$ mRNA expression in the hippocampal CA1 and CA3 regions and dentate gyrus (Tietz et al., 1999a). On day 7 after discontinuation of flurazepam treatment, these changes returned to control level (Tietz et al., 1999a) (Table 5). Flurazepam-induced $\beta 2$ mRNA down-regulation observed by Zhao et al. (1994a) was reversed after a 2-day withdrawal. Long-term treatment with diazepam up-regulated $\beta 2/3$ subunit immunoreactivity in the fron-

toparietal motor cortex but not in somatosensory cortex, whereas imidazenil had no effects (Impagnatiello et al., 1996; Pesold et al., 1997). Long-term flurazepam treatment had no effect on $\beta 2$ polypeptide (Chen et al., 1999). Treatment of cultured rat CGCs with diazepam decreased $\beta 2$ mRNA (Follesa et al., 2002), whereas zaleplon or zolpidem had no effect (Follesa et al., 2002). Long-term treatment with flunitrazepam down-regulated $\beta 2/3$ immunoreactivity in CGCs (Brown et al., 1998). Withdrawal from diazepam, zaleplon, and zolpidem treatments reduced $\beta 2$ mRNA by 27 to 48% in CGCs (Follesa et al., 2002). In summary, studies on in vivo effects of long-term BZ treatment on $\beta 2$ subunit expression suggest brain region-specific down-regulation of the subunit with some BZs.

9. $\beta 3$ Subunit. Expression of $\beta 3$ mRNA in vivo was reduced by long-term flurazepam treatment in the cerebellum and hippocampus (Zhao et al., 1994a) and in the frontal cortex, hippocampus, and caudate-putamen (Tietz et al., 1999a). Long-term treatment with triazolam strongly down-regulated $\beta 3$ mRNA in the olfactory and sensory cortices, central amygdala, and lateral geniculate nucleus (Ramsey-Williams and Carter, 1996). Long-term treatment of rats with abecarnil, diazepam, or zolpidem had no effect on $\beta 3$ mRNA expression in the

TABLE 7

Effect of withdrawal from long-term benzodiazepine treatment in vivo on GABA_A receptor subunit polypeptide expression

Compound, Species, Dose Duration, Withdrawal Time Brain Region	Subunit mRNA		Reference
	$\alpha 1$	$\beta 3$	
% change			
Flurazepam			
Rat, 100 mg/kg/d, 7 d, 2 d			Tietz et al., 1999b
Hippocampus			
CA1	-14 ^{SO}	N.S.	
CA3	N.S.	+10 ^{SO} , 9 ^{SR}	
Dentate	N.S.	N.S.	
Cerebral cortex	N.S.		
Inferior colliculus	N.S.		
Cerebellum	N.S.		

d, day(s); N.S., not significantly different from the control value; SO, stratum oriens; SR, stratum radiatum.

TABLE 8
Effect of long-term benzodiazepine treatment on GABA_A receptor subunit mRNA expression in cultured cells in vitro

Cells, Compound, Concentration, Duration	Subunit mRNA								Reference
	α1	α3	α4	α5	β1	β2	β3	γ2	
	% change								
Rat hippocampal cells									
Etizolam, 10 μM, 5 d	N.S.	N.S.	N.S.	-17				-26	Sanna et al., 2005
Lorazepam, 10 μM, 5 d	-19	+59	N.S.	N.S.				-21	Sanna et al., 2005
Rat cerebellar granule cells									
Diazepam, 10 μM, 5 d	-20		N.S.					-24, -26 ^a	Follesa et al., 2001a
Diazepam, 10 μM, 5 d	-23		N.S.		+47	-27	N.S.	-25, -27 ^a	Follesa et al., 2002
Imidazenil, 10 μM, 5 d	N.S.		N.S.					-21, N.S. ^a	Follesa et al., 2001a
Zaleplon, 10 μM, 5 d	N.S.		N.S.		N.S.	N.S.	N.S.	N.S., N.S. ^a	Follesa et al., 2002
Zolpidem, 10 μM, 5 d	N.S.		N.S.		N.S.	N.S.	N.S.	N.S., N.S. ^a	Follesa et al., 2002

d, day(s); N.S., not significantly different from the control value.
^a Two values represent S and L splice variants, respectively.

cerebral cortex (Wu et al., 1994; Holt et al., 1996, 1997a), with the exception of diazepam given as daily injections (Arnot et al., 2001). Long-term alprazolam had no effect on β3 mRNA expression in the brainstem (Tanay et al., 2001). Withdrawal from flurazepam for 2 days down-regulated β3 mRNA expression in all hippocampal subregions (Tietz et al., 1993, 1999a) (Table 5). In addition, β3 was down-regulated in the cerebellar granule cell layer (Tietz et al., 1999a). On day 7 after discontinuation of flurazepam treatment these changes returned to control level (Tietz et al., 1999a). Long-term flurazepam treatment down-regulated β3 polypeptide in the cerebral cortex and hippocampus (Chen et al., 1999; Tietz et al., 1999b). However, after 2-day withdrawal β3 polypeptide was returned to control levels and even slightly up-regulated in hippocampal CA3 region (Tietz et al., 1999b). Treatment of CGCs with diazepam, zaleplon, and zolpidem or withdrawal had no effect on β3 mRNA expression (Follesa et al., 2002). In conclusion, in vivo studies suggest brain region-specific down-regulation of β3 mRNA and polypeptide in several hippocampal and cortical subregions after long-term flurazepam treatment.

10. γ2 Subunit. Long-term treatment of rats with abecarnil, diazepam, and lorazepam down-regulated γ2 mRNA expression in the cerebral cortex (Kang and Miller, 1991; Primus and Gallager, 1992; Wu et al., 1994; Holt et al., 1996). A reduction in γ2 mRNA was

seen in the frontoparietal motor cortex by diazepam but not by imidazenil treatment (Impagnatiello et al., 1996; Longone et al., 1996). Long-term treatment with flurazepam for 7 days (Tietz et al., 1999a) or 14 days (Zhao et al., 1994b) did not affect cerebral cortical γ2 mRNA expression, but a 4-week treatment down-regulated it for 31% (Zhao et al., 1994b). The same applies to the hippocampus, where flurazepam-induced down-regulation was seen only after the long, 4-week treatment (Zhao et al., 1994b). Long-term treatment with zolpidem had no effect on cerebral cortical γ2 mRNA expression (Holt et al., 1997a). After discontinuation of long-term treatment with diazepam, cortical γ2 mRNA level of the treated rats returned to control rat level (Wu et al., 1994; Longone et al., 1996). The expression of γ2 mRNA was up-regulated on withdrawal day 7 from long-term treatment with flurazepam (34%) while being at control level on day 2 (Tietz et al., 1993, 1999a). Diazepam up-regulated γ2 polypeptide measured with immunogold labeling in the frontoparietal motor cortex (50%), but not in the somatosensory cortex, whereas imidazenil had no effects (Impagnatiello et al., 1996; Pesold et al., 1997). No effect of long-term treatment with flurazepam treatment was found on cortical γ2 polypeptide (Chen et al., 1999). Long-term diazepam treatment increased γ2 mRNA expression in the cerebellum in one (Holt et al., 1999) of three studies (Wu et al., 1994; Impagnatiello et al., 1996), whereas no effect of long-term treatment with

TABLE 9
Effect of withdrawal from long-term benzodiazepine treatment on GABA_A receptor subunit mRNA expression in cultured cells in vitro

Cells, Compound, Concentration, Duration, Withdrawal Time	Subunit mRNA								Reference
	α1	α2	α3	α4	β1	β2	β3	γ2	
	% change								
Rat hippocampal neurons									
Etizolam, 10 μM, 5 d, 6 h	N.S.	+24	+30	N.S.				-21(S)	Sanna et al., 2005
Lorazepam, 10 μM, 5 d, 6 h	N.S.	+17	+38	+23				-18(S)	Sanna et al., 2005
Rat cerebellar granule cells									
Diazepam, 10 μM, 5 d, 6 h	-37			+40				-18, -23 ^a	Follesa et al., 2001a
Diazepam, 10 μM, 5 d, 6 h	-32			+38	+57	-48	N.S.	-19, -25 ^a	Follesa et al., 2002
Imidazenil, 10 μM, 5 d, 6 h	-20			+42				-37, -21 ^a	Follesa et al., 2001a
Zaleplon, 10 μM, 5 d, 6 h	-23			+59	+42	-27	N.S.	-20, -32 ^a	Follesa et al., 2002
Zolpidem, 10 μM, 5 d, 6 h	-29			+32	+31	-31	N.S.	-19, -27 ^a	Follesa et al., 2002

d, day(s); N.S., not significantly different from the control value.
^a Two values represent S and L splice variants, respectively.

TABLE 10
Effect of long-term benzodiazepine treatment on GABA_A receptor subunit polypeptide expression in cultured cells in vitro

Cells, Compound, Concentration, Duration	Subunit mRNA					Reference
	α1	α4	α6	β2/3	γ2	
	% change					
Rat hippocampal neurons						
Etizolam, 10 μM, 5 d	N.S.	N.S.				Sanna et al., 2005
Lorazepam, 10 μM, 5 d	-36	N.S.				Sanna et al., 2005
Rat cerebellar granule cells						
Bretazenil, 1 μM, 2 d	-11					Johnston and Bristow, 1998
Diazepam, 1 μM, 2 d	-19					Johnston and Bristow, 1998
Diazepam, 10 μM, 5 d	-37	N.S.			-50	Follesa et al., 2001a
Flunitrazepam, 1 μM, 2 d	-20					Johnston and Bristow, 1998
Flunitrazepam, 1 μM, 2 d	-41		N.S.			Brown and Bristow, 1996
Flunitrazepam, 1 μM, 2 d	-40			-67		Brown et al., 1998
Imidazenil, 1 μM, 2 d	N.S.					Johnston and Bristow, 1998

d, day(s); N.S., not significantly different from the control value.

diazepam, lorazepam, or imidazenil was found on hippocampal γ2 mRNA expression (Kang and Miller 1991; Wu et al., 1994; Impagnatiello et al., 1996; Tietz et al., 1999). In accordance, long-term treatment with flurazepam treatment had no effect on γ2 polypeptide in the cerebellum or hippocampus (Chen et al., 1999). Long-term treatment of cultured CGCs with diazepam reduced the γ2 mRNA expression by 24 to 27% and γ2 polypeptide by 50%, whereas zaleplon and zolpidem had no effect (Follesa et al., 2001a, 2002). Withdrawal from long-term treatment with diazepam, zaleplon, zolpidem, and imidazenil reduced CGC γ2 mRNA expression by 18 to 37%, and withdrawal from diazepam treatment reduced the γ2 polypeptide by 63% (Follesa et al., 2001a, 2002). Long-term treatments of cultured hippocampal cells with etizolam and lorazepam reduced γ2S mRNA expression by 21 and 26%, respectively (Sanna et al., 2005). The reduced expression of γ2S mRNA was sustained during 6-h withdrawal from lorazepam and etizolam (Sanna et al., 2005). Studies on the effects of in vivo long-term BZ agonist treatment and withdrawal suggest down-regulation of γ2 expression in the cerebral cortex by some, but not all agonists, but had little effect on γ2 expression in the cerebellum or hippocampus.

11. γ1 and γ3 Subunits. Long-term treatment of rats with continuous diazepam infusion using minipumps up-regulated γ1 mRNA expression (Arnot et al., 2001). No other significant effects of long-term BZ treatment on γ1 mRNA or polypeptide expression have been found thus far (Table 3). Long-term in vivo treatment with

diazepam, but not abecarnil, increased γ3 mRNA expression in the cerebral cortex (Holt et al., 1996).

12. δ Subunit. Long-term treatment with diazepam, imidazenil, or zolpidem did not affect δ mRNA expression in several brain regions studied (Impagnatiello et al., 1996; Holt et al., 1997a). In contrast, triazolam treatment strongly down-regulated δ expression in several brain regions with a strong basal expression, such as the thalamus (Ramsey-Williams and Carter, 1996) (Table 3). These data do not allow any clear conclusion on the effects of long-term BZ administration on δ subunit expression in the brain. This would be important to clarify because the δ subunit-containing receptors are regarded to form BZ-insensitive receptor populations (but see Hanchar et al., 2006) that function especially at extrasynaptic and perisynaptic areas.

13. Conclusions on Effects of Long-Term Benzodiazepine Administration on Receptor Subunit Expression. Studies on regulation of GABA_A receptor subunit expression in vivo by long-term BZ treatment have shown that the regulation is subunit-specific, is brain region-specific, and occurs at subunit-specific time scales. In addition, the inverse agonist studied (FG 7142) often had opposite effects on the expression as compared with BZ agonists. The partial agonist imidazenil was mostly without an effect on expressions of receptor subunits (Tables 2 and 3, Impagnatiello et al., 1996; Longone et al., 1996; Pesold et al., 1997; Johnston and Bristow, 1998). In vitro studies with cultured cells treated with long-term BZs produced surprisingly controversial results compared with studies in

TABLE 11
Effect of withdrawal from long-term benzodiazepine treatment on GABA_A receptor subunit polypeptide expression in cultured cells in vitro

Cells, Compound, Concentration, Duration, Withdrawal Time	Subunit Polypeptide			Reference
	α1	α4	γ2	
	% change			
Rat hippocampal neurons				
Etizolam, 10 μM, 5 d/6 h	N.S.	N.S.		Sanna et al., 2005
Lorazepam, 10 μM, 5 d/6 h	-26	+81		Sanna et al., 2005
Rat cerebellar granule cells				
Diazepam, 10 μM, 5 d/6 h	-75	+45	-63	Follesa et al., 2001a

d, day(s); N.S., not significantly different from the control value.

vivo. The in vitro results from rat CGCs and hippocampal neurons are often opposite those received from hippocampus or cerebellum of rats treated in vivo. Thus, the in vitro results should be interpreted very cautiously.

Quantitative RT-PCR studies usually monitor the steady-state mRNA levels, not the rates of mRNA synthesis or degradation. Therefore, it has not been possible to deduce whether down-regulation of a subunit mRNA results from reduced transcription rate or increased degradation rate. However, using a nuclear run-off assay, Holt et al. (1997b) showed a 65% decrease in the $\gamma 2$ mRNA synthesis rate in the cerebral cortex of rats continuously treated with diazepam, whereas in the cerebellum, the rate was increased by 42% (Holt et al., 1999). The changes in $\gamma 2$ mRNA transcription rate paralleled the changes in $\gamma 2$ mRNA steady-state levels indicating that diazepam predominantly regulates $\gamma 2$ mRNA at the level of transcription (Holt et al., 1997a, 1999). The results suggest a brain region-specific regulation of $\gamma 2$ mRNA transcription by long-term treatment with diazepam. Lorazepam-induced reduction in transcriptional activity is suggested in $\alpha 1$ subunit down-regulation (Kang et al., 1994). The group isolated human $\alpha 1$ gene promoter and showed that long-term lorazepam treatment down-regulates transcriptional activity of $\alpha 1$ promoter in neurons transiently transfected with a $\alpha 1$ promoter-firefly luciferase construct (Kang et al., 1994). Lorazepam treatment dose-dependently attenuated expression of luciferase activity in the cells (Kang et al., 1994). The signaling mechanism by which lorazepam represses $\alpha 1$ gene promoter activity is unknown. However, it has been shown that GABA agonist-induced reduction of cell surface GABA_A receptors occurs before down-regulation of receptor subunit mRNA expression (Baumgartner et al., 1994; Miranda and Barnes, 1997). It was subsequently found that long-term BZ treatment also induces GABA_A receptor internalization (Tehrani and Barnes, 1997). How this receptor internalization is signaled into cell nucleus to suppress receptor subunit gene transcription is currently not known.

Studies on long-term BZ treatment in vivo indicate alterations in GABA_A receptor subunit expression. The clearest effects are the down-regulation of $\alpha 1$ and $\beta 3$ in several cortical and hippocampal subregions and the down-regulation of $\gamma 2$ in the cerebral cortex. In addition, $\alpha 4$ subunit is strongly up-regulated in BZ withdrawal. There is also an indication of up-regulation of $\gamma 3$ subunit. BZ-induced alterations in GABA_A receptor subunits may produce receptors with lower sensitivity or insensitivity to BZs (e.g., formation of $\alpha 4\beta\gamma 2$ and $\alpha\beta\gamma 3$ -containing receptors). The changes, however, are quantitatively rather small and short-lasting. Therefore, the changes cannot solely explain the development of BZ tolerance, dependence or withdrawal syndrome after the discontinuation, but they may partially contribute to mechanisms underlying these phenomena.

14. Benzodiazepine-Induced Changes in the Expression of Other Genes. BZ administration induces a wide variety of other changes in neuronal gene expression that may participate to the development of tolerance and dependence. In a microarray study using wild-type and $\alpha 1$ (H101R) knockin mouse lines, it was shown that even a single dose of diazepam significantly changed the expression of 54 transcripts (0.43% of the transcripts on the array), 34 transcripts being down-regulated and 20 transcripts being up-regulated (Huopaniemi et al., 2004). Changes in the expression of six transcripts, CaMKII α , BDNF, MKP-1, GIF, *c-fos*, and NGFI-A, were mediated via action of diazepam on $\alpha 1$ subunit-containing GABA_A receptors (Huopaniemi et al., 2004).

According to glutamate hypothesis of BZ tolerance and dependence, excitatory mechanisms become up-regulated to compensate for BZ-induced enhancement of inhibition (Stephens, 1995). Expression of *N*-methyl-D-aspartate (NMDA) and α -amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid (AMPA) type glutamate receptors has been shown to be regulated after long-term BZ treatment. Cerebral cortical GluN1 and GluN2B, but not GluN2A subunits of NMDA receptors (Tsuda et al., 1998), glutamic acid decarboxylase (GAD₆₇), and GluA1 subunit of AMPA receptor were increased in diazepam-withdrawn rats (Izzo et al., 2001). In rats withdrawn from flurazepam, amplitudes of AMPA receptor-mediated miniature excitatory postsynaptic currents were increased in hippocampal CA1 neurons (Van Sickle et al., 2004; Xiang and Tietz, 2007). The 50% enhancement in AMPA receptor function was attributed to an increase in GluA1 polypeptide trafficking from the endoplasmic reticulum and its subsequent incorporation into membranes (Song et al., 2007; Das et al., 2008), whereas NMDA receptor-mediated currents were reduced in this brain region (Van Sickle et al., 2004; Xiang and Tietz, 2007). Mice deficient in GluA1 subunits have reduced short-term tolerance (i.e., prolonged short-term impairment to high doses of flurazepam) and develop less tolerance but show increased withdrawal signs after a 7-day flurazepam treatment and challenge with flumazenil (Aitta-Aho et al., 2009). Even a single injection of diazepam to young mice increases the AMPA receptor function over that of NMDA receptors in dopamine neurons of the ventral tegmental area, measured in vitro 24 h after the drug injection in the ventral tegmental area dopamine neurons in vitro (Heikkinen et al., 2009).

Long-term BZ administration up-regulates L-type high voltage-gated calcium channels in the cerebral cortex (Katsura et al., 2007) and potentiates high voltage-gated calcium channel currents in the hippocampal CA1 neurons (Xiang et al., 2008).

B. Neurosteroids

Endogenous neurosteroids 3 α -OH-5 α -pregnan-20-one (3 α ,5 α -THP) and its active 5 β isomer 3 α -OH-5 β -pregnan-20-one are metabolites of the ovarian/adrenal steroid

progesterone (Barbaccia, 2004). Allopregnanolone can also be synthesized de novo in the brain from cholesterol (Compagnone and Mellon, 2000). Other endogenous neurosteroids include 5α -pregnane- $3\alpha,21$ -diol- 20 -one, which is a metabolite of the adrenal steroid corticosterone (Barbaccia, 2004). These neurosteroids are potent positive allosteric modulators of GABA_A receptors (Lambert et al., 2009), the δ -containing receptors ($\alpha4\beta\delta$, $\alpha6\beta\delta$) being especially sensitive to them (Wohlfarth et al., 2002; Spigelman et al., 2003). In addition to modulation of GABA_A receptor function, fluctuations in brain neurosteroid concentrations (e.g., during estrous cycle, pregnancy, and stress) affect GABA_A receptor subunit expression via an action that does not need the activation of steroid hormone receptors (Maguire and Mody, 2007). During estrous cycle, the hippocampal δ polypeptide level decreases 30% from the late diestrus (high-progesterone phase) to estrus (low-progesterone phase), whereas the pattern of $\gamma2$ subunit during the cycle is complementary to that of δ (Maguire et al., 2005). The expression of $\alpha4$ does not change, suggesting an up-regulation of $\alpha4\beta\delta$ receptors, down-regulation of $\alpha4\beta\gamma2$, and increase in tonic inhibition in late diestrus and the reverse effects at estrus (Maguire et al., 2005).

In vivo treatment of rats with progesterone or allopregnanolone up-regulated $\alpha4$ and δ polypeptide expression in the hippocampus (Gulinello et al., 2001; Hsu et al., 2003; Shen et al., 2005; Maguire et al., 2005; Maguire and Mody, 2007), whereas the $\gamma2$ polypeptide was down-regulated by 40 to 50% (Shen et al., 2005). Therefore, exogenous neurosteroids also up-regulate $\alpha4\beta\delta$ receptors and increase tonic inhibition. Discontinuation of administration of a neurosteroid agonist or the parent compound progesterone (i.e., withdrawal) leads to a 3-fold increase in hippocampal $\alpha4$ and δ (but no change in $\gamma2$ expression) (Smith et al., 1998; Sundstrom-Poromaa et al., 2002; Gangisetty and Reddy, 2009) and in premenstrual syndrome-like symptoms such as anxiety and increased seizure susceptibility (Dennerstein et al., 1985; Herzog, 2009).

Although studies on the effects of neurosteroids on GABA_A receptor subunit expression in vivo have been focused on the hippocampus, studies in cultured cells in vitro have been performed using mouse cortical neurons and rat cerebellar granule cells. Progesterone treatment (1 μ M, 5 days) had no effect on $\alpha4$ mRNA in cortical neurons or CGCs, whereas 6-h progesterone withdrawal up-regulated the expression in both cultures (Follesa et al., 2000, 2001b). In contrast, in differentiated P19 cells, allopregnanolone (1 μ M, 4 days) down-regulated $\alpha4$ mRNA expression, but 1-day withdrawal from the treatment did not affect the expression (Grobin and Morrow, 2000). Long-term treatment with allopregnanolone (0.1 μ M, 2 days) up-regulated $\alpha4$ polypeptide by 146% in IMR-32 neuroblastoma cells (Zhou and Smith, 2007, 2009).

Long-term treatment with progesterone and allopregnanolone had no effect on $\gamma2$ mRNA in cultured cortical

neurons (Yu et al., 1996; Follesa et al., 2001b), whereas both compounds down-regulated it in CGCs (Follesa et al., 2000). Progesterone withdrawal down-regulated $\gamma2$ mRNA in both types of cell cultures (Follesa et al., 2000, 2001b). The δ polypeptide tends to be down-regulated in CGCs by long-term treatment with progesterone and allopregnanolone (Biggio et al., 2006), which effect is increased at withdrawal (Biggio et al., 2006).

C. Barbiturates

Barbiturates were introduced in early 20th century as relatively nonselective general depressants (Estes, 1995). Although also affecting several other ionotropic receptors, GABA_A receptors are the primary target of barbiturates (Smith and Riskin, 1991). Barbiturates have been used in clinical practice as sedatives, anxiolytics, hypnotics, anesthetics, and anticonvulsants (Ito et al., 1996b).

There are a number of studies on the effects of long-term treatment with pentobarbital on GABA_A receptor subunit mRNA expression (Tables 12 and 13). Treatment durations have been 6 to 14 days and daily pentobarbital doses 30 to 120 mg/kg/day (intraperitoneal injection) or 24 to 225 mg/day when using osmotic minipumps. Long-term treatment with pentobarbital had no effect on $\alpha1$ mRNA in the cerebral cortex, cerebellum, and several other brain regions, although it slightly down-regulated $\alpha1$ mRNA in the hippocampus and inferior and superior colliculi (Morrow et al., 1990, 1991; Tseng et al., 1994) (Table 12). During pentobarbital withdrawal (1 day), $\alpha1$ mRNA was up-regulated in the neocortex and cerebellar granule and Purkinje cells (Tseng et al., 1994) (Table 13). The expression of cerebellar $\alpha6$ mRNA was up-regulated by long-term treatment with pentobarbital (Ito et al., 1996a) (Table 12), and during withdrawal (1 day), the expression returned to control levels (Ito et al., 1996). Long-term treatment with pentobarbital drastically up-regulated $\beta1$ mRNA 600% over basal expression in the CA1 and CA2 regions of the hippocampus, although the expression was not affected in other hippocampal areas (Yin and Lee, 1998). On 1-day pentobarbital withdrawal, the $\beta1$ expression was still elevated by 130 to 180% but returned to control levels in 7-day withdrawal (Yin and Lee, 1998). Long-term treatment with pentobarbital had no effect on $\beta3$ mRNA expression (Tseng et al., 1994) (Table 12). During withdrawal $\beta3$ expression was up-regulated in the neocortex, whereas no changes in $\beta3$ expression were found in other brain regions studied (Tseng et al., 1994) (Table 13). The expression of $\gamma2$ mRNA was slightly reduced by long-term treatment with pentobarbital in the superior and inferior colliculi (Tseng et al., 1993a). Long-term treatment with pentobarbital up-regulated δ mRNA expression in the cerebellum but not in the frontal cortex (Lin and Wang, 1996) (Table 12). Withdrawal from pentobarbital down-regulated δ expression in the cerebellum, whereas no effect was found in the frontal cortex (Lin and Wang, 1996) (Table 13).

TABLE 12
Effect of long-term pentobarbital treatment in vivo on GABA_A receptor subunit mRNA expression

Species, Dose, Duration, Regimen, Brain Region	Subunit mRNA						Reference
	α1	α6	β1	β3	γ2	δ	
	% change						
Rat, 30 mg/kg, 14 d Intraperitoneal injection Cerebral cortex	N.S.						Morrow et al., 1990
Rat, increasing dose 30 mg->120 mg/kg, 14 d Intraperitoneal injection Cerebral cortex	N.S.						Morrow et al., 1991
Rat, 7.2 mg/d, ^a 6 d Infusion, osmotic minipump Neocortex					N.S.		Tseng et al., 1993a
Piriform cortex					N.S.		
Hippocampus					N.S.		
Caudate putamen					N.S.		
Medial habenular nucleus					N.S.		
Thalamus					N.S.		
Superior colliculus					-12		
Inferior colliculus					-8		
Central gray					N.S.		
Cerebellum					N.S.		
Rat, 7.2 mg/d, ^a 6 d Infusion, osmotic minipump Neocortex	N.S.			N.S.			Tseng et al., 1994
Piriform cortex	N.S.			N.S.			
Hippocampus							
CA1	-19			N.S.			
CA3	-10			N.S.			
Dentate gyrus	-10			N.S.			
Caudate putamen	N.S.			N.S.			
Medial habenular nucleus	N.S.			N.S.			
Thalamus	N.S.			—			
Superior colliculus	-25			—			
Inferior colliculus	-16			—			
Central gray	N.S.			—			
Cerebellum							
Molecular cells	N.S.			—			
Granule cells	N.S.			N.S.			
Purkinje cells	N.S.			—			
Rat, 7.2 mg/d, ^a 6 d Infusion, osmotic minipump Cerebellum							Ito et al., 1996a
Granule cells		+47					
Mouse, 3×75 mg/d, 7 d Infusion, osmotic minipump Frontal cortex					N.S.		Lin and Wang, 1996
Cerebellum					+44		
Rat, increasing dose 30 mg->120 mg/kg, 9 d Intraperitoneal injection Hippocampus							Yin and Lee, 1998
CA1			+636				
CA2			+623				
CA3			N.S.				
CA4			N.S.				
Dentate gyrus			N.S.				

d, day(s); N.S., not significantly different from the control value; —, not detected.
^a Continuous infusion.

The effect of long-term pentobarbital treatment on GABA_A receptors has also been studied at polypeptide level using ligand binding assays. Consistent with slight or no effects of long-term treatment with pentobarbital on subunit mRNA expressions, the maximal number of binding sites (B_{max}) for [³H]flunitrazepam, [³H]muscimol, and [³⁵S]*t*-butylbicyclophosphorothionate binding to brain sections of rats chronically treated with pentobarbital did not differ from the control levels (Tseng et

al., 1993b; Miyaoka et al., 1994). One-day pentobarbital withdrawal increased the [³H]flunitrazepam B_{max} values in the frontal cortex, cerebellum and striatum (Tseng et al., 1993b; Miyaoka et al., 1994). The B_{max} value of [³H]muscimol was increased in the frontal cortex and that of [³⁵S]*t*-butylbicyclophosphorothionate in the frontal cortex and striatum (Tseng et al., 1993b). Consistent with increased α6 mRNA expression in the study of Ito et al. (1996a), cerebellar [³H]Ro 15-4513 binding was

TABLE 13
Effect of withdrawal from long-term pentobarbital treatment in vivo on GABA_A receptor subunit mRNA expression

Species, Dose, Duration, Withdrawal Time, Regimen, Brain Region	Subunit mRNA					Reference
	α1	α6	β1	β3	δ	
	% change					
Rat, 7.2 mg/d, ^a 6 d, 1 d Infusion, osmotic minipump						Tseng et al., 1994
Neocortex						
Layer II/III	+52			+21		
Layer IV	+32			+14		
Layer V/VI	+39			+16		
Piriform cortex	+14			+14		
Hippocampus						
CA1	N.S.			N.S.		
CA3	N.S.			N.S.		
Dentate gyrus	N.S.			N.S.		
Caudate putamen	N.S.			N.S.		
Medial habenular nucleus	N.S.			N.S.		
Thalamus	N.S.			—		
Superior colliculus	N.S.			—		
Inferior colliculus	N.S.			—		
Central gray	N.S.			—		
Cerebellum						
Molecular layer	N.S.			—		
Granule cells	+18			N.S.		
Purkinje cells	+45			—		
Rat, 7.2 mg/d, ^a 6 d, 1 d Infusion, osmotic minipump						Ito et al., 1996a
Cerebellum						
Granule cells		N.S.				
Mouse, 3×75 mg/d, 7 d, 1 d Infusion, osmotic minipump						Lin and Wang, 1996
Frontal cortex					N.S.	
Cerebellum					-41	
Rat, increasing dose 60→90 mg/kg/d, 9 d, 1 d Intraperitoneal injection						Yin and Lee, 1998
Hippocampus						
CA1			+128			
CA2			+176			
CA3			N.S.			
CA4			N.S.			
Dentate gyrus			N.S.			
Rat, increasing dose 60→90 mg/kg/d, 9 d, 7 d Intraperitoneal injection						Yin and Lee, 1998
Hippocampus						
CA1			N.S.			
CA2			N.S.			
CA3			N.S.			
CA4			N.S.			
Dentate gyrus			N.S.			

d, day(s); N.S., not significantly different from the control value; —, not detected.
^a Continuous infusion.

increased by long-term pentobarbital treatment and on pentobarbital withdrawal (Ito et al., 1996a).

In conclusion, there are rather few studies on the effects of barbiturates on the expression of GABA_A receptors/subunits. Long-term treatment studies with pentobarbital suggest that receptor subunit expressions are usually slightly down-regulated, whereas pentobarbital withdrawal leads in receptor up-regulation.

D. Ethanol

The pharmacological actions of ethanol include anxiolysis, sedation, motor incoordination, impairment of judgment, and, at high concentrations, anesthesia. According to current view, the primary targets mediating ethanol intoxication are GABA_A (Wallner

et al., 2003; Lovinger and Homanics, 2007), NMDA (Ron, 2004), glycine (Crawford et al., 2007), 5-hydroxytryptamine 3 (serotonin) (Lovinger, 1999), and nicotinic acetylcholine receptors (Cardoso et al., 1999) as well as L-type Ca²⁺ channels (Wang et al., 1994), G-protein-activated inwardly rectifying K⁺ channels (Kobayashi et al., 1999), and Ca²⁺-activated K⁺ channels (Brodie et al., 2007) (for review, see Vengeliene et al., 2008). Long-term administration of ethanol produces tolerance, dependence, and withdrawal signs upon cessation. Desensitization of ethanol-sensitive mechanisms obviously plays a role in the development of tolerance (Dopico and Lovinger, 2009), and the activity of the glutamatergic system is enhanced during withdrawal (De Witte et al., 2003; Nagy, 2008).

TABLE 15
Effect of long-term ethanol treatment in vivo on GABA_A receptor β , γ and δ subunit mRNA expression

Species, Dose, Duration, Regimen, Brain Region	Subunit mRNA							Reference
	β 1	β 2	β 3	γ 1	γ 2	γ 3	δ	
	% change							
Rat, 5 g/kg/d, 6 d Intragastric intubation Cerebral cortex	+29	+55	+72					Mhatre and Ticku, 1994
Rat, 10–12 g/kg/d, 14 d Forced ethanol drinking Cerebral cortex	N.S.	N.S.	N.S.	70	+32, N.S. ^a	N.S.	N.S.	Devaud et al., 1995
Mouse, 5 g/kg/d, 14 d Intragastric intubation Cerebellum								Wu et al., 1995
					+82			
		N.S.			+46			
		N.S.	N.S.		+62			
Rat, 5 g/kg, 18 months Forced ethanol drinking Frontoparietal cortex	N.S.	N.S.	-20	N.S.	N.S.	N.S.	N.S.	Sarviharju et al., 2006
Olfactory bulb	N.S.	N.S.	-18	N.S.	N.S.	N.S.	N.S.	
Olfactory tubercle	N.S.	N.S.	N.S.	N.S.	N.S.	N.S.	N.S.	

d, day(s); N.S., not significantly different from the control value.

^a Two values represent S and L splice variants, respectively.

drawal from the treatment (Follesa et al., 2003; Biggio et al., 2007). CE down-regulated α 1 polypeptide expression in the cerebellum (Mhatre et al., 1993; Charlton et al., 1997; Marutha Ravindran et al., 2007a). After a 2-day withdrawal, the α 1 polypeptide returned to control level (Marutha Ravindran et al., 2007a).

In the hippocampus, CE down-regulated α 1 mRNA expression (Charlton et al., 1997), whereas the expression of α 1 polypeptide was not affected (Charlton et al., 1997; Matthews et al., 1998; Marutha Ravindran et al., 2007a). A 2-day withdrawal from 6-day CE treatment had no effect on α 1 polypeptide expression in the hippocampus (Marutha Ravindran et al., 2007a). In con-

trast, a 2-day withdrawal from 60-day continuous intermittent ethanol (wdr-CIE) down-regulated it by 48% (Cagetti et al., 2003). In cultured hippocampal neurons, α 1 mRNA was down-regulated by CE and on withdrawal from CE (Sanna et al., 2003). Expression of α 1 mRNA was down-regulated by CE in the amygdala (Papadeas et al., 2001) but not in the nucleus accumbens or ventral tegmental area (Charlton et al., 1997; Papadeas et al., 2001). CE decreased the α 1 mRNA content in whole-brain samples of withdrawal seizure-prone mice but not in withdrawal seizure-resistant mice (Buck et al., 1991).

The effects of CE have also been studied in GABA_AR α 1 knockin mice harboring S270H and L277A mutations in

TABLE 16
Effect of withdrawal from long-term ethanol treatment in vivo on GABA_A receptor α subunit mRNA expression

Species, Dose, Duration, Withdrawal Time Regimen, Brain Region	Subunit mRNA						Reference
	α 1	α 2	α 3	α 4	α 5	α 6	
	% change						
Rat, 5 g/kg/d, 6 d, 1d Intragastric intubation Cerebral cortex	-71	-61, -43	N.S.		-45		Mhatre and Ticku, 1992
	-34					+150	
Rat, 6 g/kg/d, 60 d, 2 d Intragastric intubation, CIE Hippocampus							Mahmoudi et al., 1997
				+26	N.S.		
				+46	N.S.		
				+30	N.S.		
				+12			
				+18			
				+15			
Rat, 6 g/kg/d, 30 d, 2 d Intragastric intubation, CIE Cerebellum	N.S.					N.S.	Petrie et al., 2001
Rat, 6 g/kg/d, 60 d, 2 d Intragastric intubation, CIE Hippocampus					N.S.	N.S.	Petrie et al., 2001
Rat, 6 g/kg/d, 60 d, 2 d Intragastric intubation, CIE Hippocampus							Cagetti et al., 2003
		N.S.					

d, day(s); N.S., not significantly different from the control value.

TABLE 17
Effect of withdrawal from long-term ethanol treatment in vivo on GABA_A receptor β , γ , and δ subunit mRNA expression

Species, Dose, Duration, Withdrawal Time Regimen, Brain Region	Subunit mRNA						Reference
	β 1	β 2	β 3	γ 1	γ 2	δ	
	% change						
Rat, 5 g/kg/d, 6 d, 1 d Intragastric intubation Cerebral cortex	N.S.	+47	+28				Mhatre and Ticku, 1994
Rat, 6 g/kg/d, 30 d, 2 d Intragastric intubation, CIE Cerebellum		N.S.				N.S.	Petrie et al., 2001
Rat, 6 g/kg/d, 60 d, 2 d Intragastric intubation, CIE Hippocampus				N.S.	N.S., N.S. ^a	N.S.	Petrie et al., 2001
Rat, 6 g/kg/d, 60 d, 2 d Intragastric intubation, CIE Hippocampus				+80	+48, N.S. ^a		Cagetti et al., 2003

d, day(s); N.S., not significantly different from the control value.

^a Two values represent S and L splice variants, respectively.

α 1 subunit (Borghese et al., 2006). Recombinant receptors containing these mutations are insensitive to ethanol (Ueno et al., 2000; Borghese et al., 2006). CE treatment did not affect α 1 polypeptide level in α 1 knockin mice, although it reduced the level in wild-type mice (Werner et al., 2009). The results strongly suggest that CE down-

regulates wild-type α 1 mRNA and polypeptide. Furthermore, the study suggests that potentiation of α 1-containing receptors by ethanol would be needed for CE-induced α 1-receptor down-regulation.

The only study where an increase of α 1 mRNA was observed is that of Hirouchi et al. (1993). They found a

TABLE 18
Effect of long-term ethanol treatment in vivo on GABA_A receptor α subunit polypeptide expression

Species, Dose, Duration, Regimen, Brain Region	Subunit Polypeptide						Reference
	α 1	α 2	α 3	α 4	α 5	α 6	
	% change						
Rat, 5 g/kg/d, 6 d Intragastric intubation Cerebral cortex	-61	-47	-30				Mhatre et al., 1993
	-56						
Rat, 10 g/kg/d, 28 d Ethanol cont. liquid diet Frontoparietal cortex	-49				N.S.		Charlton et al., 1997
	-30				-36		
Hippocampus	N.S.				N.S.		
	N.S.						
Rat, 10–12 g/kg/d, 14 d Forced ethanol administration Cerebral cortex	-33			+26			Devaud et al., 1997
Rat, 10 g/kg/d, 14 d Forced ethanol administration Hippocampus	N.S.	N.S.	N.S.	N.S.			Matthews et al., 1998
Rat, 10 g/kg/d, 40 d Forced ethanol administration Cerebral cortex	-18	N.S.	N.S.	+28			Matthews et al., 1998
	N.S.	N.S.	N.S.	+43			
Rat, 15 g/kg/d, 14 d Forced ethanol administration Prefrontal cortex	-18			+24			Grobin et al., 2000
	N.S.			+110			
Cingulate cortex	N.S.			+40			
Motor cortex	N.S.			N.S.			
Parietal cortex	-45			+60			
	-21						
Rat, 15 g/kg/d, 14 d Forced ethanol administration Amygdala	-21			-22			Papadeas et al., 2001
	N.S.			-28			
Nucleus accumbens	N.S.			N.S.			
	N.S.						
Rat, 9 g/kg/d, 6 d Intragastric intubation Cerebral cortex	-41	-41		+29			Marutha Ravindran et al., 2007a
	-28	-27			+66		
Hippocampus	N.S.	N.S.		+34			

d, day(s); N.S., not significantly different from the control value.

TABLE 19
Effect of long-term ethanol treatment in vivo on GABA_A receptor β , γ , and δ subunit polypeptide expression

Species, Dose, Duration, Regimen, Brain Region	Subunit Polypeptide						Reference
	β 1	β 2	β 3	γ 1	γ 2	δ	
	% change						
Rat, 5g/kg/d, 6 d Intragastric intubation Cerebral cortex		+23	+23				Mhatre and Ticku, 1994
Rat, 10–12 g/kg/d, 14 d Forced ethanol administration Cerebral cortex		+36	+36	+30	N.S.		Devaud et al., 1997
Rat, 10 g/kg/d, 14 d Forced ethanol administration Hippocampus		N.S.	N.S.	N.S.			Matthews et al., 1998
Rat, 10 g/kg/d, 40 d Forced ethanol administration Cerebral cortex	N.S.	N.S.	N.S.				Matthews et al., 1998
Rat, 10 g/kg/d, 40 d Forced ethanol administration Hippocampus	N.S.	N.S.	N.S.				Matthews et al., 1998
Rat, 9 g/kg/d, 6 d Intragastric intubation Cerebral cortex		+34			N.S.		Marutha Ravindran et al., 2007a
Cerebellum		+31			N.S.		
Hippocampus		N.S.			+32		
Rat, 9 g/kg/d, 6 d Intragastric intubation Cerebral cortex						N.S.	Marutha Ravindran et al., 2007b
Cerebellum						–35	
Hippocampus						–25	

d, day(s); N.S., not significantly different from the control value.

37% increase in α 1 mRNA in samples purified from total brain of mice receiving a daily 1 mmol/kg injection of the alcohol dehydrogenase inhibitor pyrazole and kept 7 days at continuous inhalation of ethanol vapor. The presence of pyrazole and the absence of acetaldehyde on α 1 mRNA expression were not studied, but these factors may be responsible for the opposite result received.

2. α 2 Subunit. Long-term ethanol treatment down-regulated α 2 mRNA (Montpied et al., 1991b; Mhatre and Ticku, 1992; Morrow et al., 1992) and α 2 polypeptide (Mhatre et al., 1993; Marutha Ravindran et al., 2007a) in the cerebral cortex (Tables 14 and 18). This down-regulation was not seen by Matthews et al. (1998). Down-regulation of cortical α 2 mRNA was also detected

after a 1-day wdr-CE (Mhatre and Ticku, 1992). However, during 2-day wdr-CE, cortical α 2 polypeptide expression was reversed to control levels (Marutha Ravindran et al., 2007a). CE and CIE down-regulated α 2 mRNA in cultured mouse cortical neurons (Sheela Rani and Ticku, 2006). The expression of α 2 polypeptide was down-regulated in cortical neurons with CE, whereas CIE had no effect on expression (Sheela Rani and Ticku, 2006). Withdrawal from CE or CIE had no effect on α 2 mRNA or polypeptide expression in cultured mouse cortical neurons (Sheela Rani and Ticku, 2006).

Long-term ethanol treatment down-regulated α 2 polypeptide in the cerebellum (Marutha Ravindran et al., 2007a), and the expression returned to control levels

TABLE 20
Effect of withdrawal from long-term ethanol treatment in vivo on GABA_A receptor subunit polypeptide expression

Species, Dose, Duration, Withdrawal Time Regimen, Brain Region	Subunit Polypeptide								Reference
	α 1	α 2	α 4	β 2	β 3	γ 1	γ 2	δ	
	% change								
Rat, 10–12 g/kg/d, 14 d, 6–8 h Forced ethanol administration Cerebral cortex	–34		+30	+32	+32	+54	N.S.		Devaud et al., 1997
Rat, 6 g/kg/d, 60 d, 2 d Intragastric intubation, CIE Hippocampus	–48		+50				+38	–52	Cagetti et al., 2003
Rat, 9 g/kg/d, 6 d, 2 d Intragastric intubation Cerebral cortex	N.S.	N.S.	+29	N.S.			N.S.		Marutha Ravindran et al., 2007a
Cerebellum	N.S.	N.S.		N.S.			N.S.		
Hippocampus	N.S.	N.S.	+36	N.S.			N.S.		
Rat, 9 g/kg/d, 6 d, 2 d Intragastric intubation Cerebral cortex								N.S.	Marutha Ravindran et al., 2007b
Cerebellum								N.S.	
Hippocampus								N.S.	

d, day(s); N.S., not significantly different from the control value.

TABLE 21
Effect of long-term ethanol treatment on GABA_A receptor subunit mRNA expression in cultured cells in vitro

Cells, [EtOH], Treatment Time	Subunit mRNA										Reference	
	$\alpha 1$	$\alpha 2$	$\alpha 3$	$\alpha 4$	$\alpha 5$	$\alpha 6$	$\beta 2$	$\beta 3$	$\gamma 2$	δ		
	% change											
Rat hippocampal neurons												
100 mM, 5 d	-24	N.S.	-23	N.S.	N.S.				-22, -34 ^a			Sanna et al., 2003
100 mM, 5 d										+120		Follesa et al., 2005
Rat cerebellar granule cells												
100 mM, 5 d	N.S.			N.S.		N.S.			-23, -32 ^a			Follesa et al., 2003
100 mM, 5 d	N.S.											Follesa et al., 2004
100 mM, 5 d										N.S.		Follesa et al., 2005
Mouse cortical neurons												
75 mM, 5 d	-24	-24		N.S.			N.S.	N.S.	-32			Sheela Rani and Ticku, 2006
75 mM, 5 d, CIE	-55	-52		48			N.S.	N.S.	-29			Sheela Rani and Ticku, 2006

d, day(s); N.S., not significantly different from the control value.

^a Two values represent S and L splice variants, respectively.

after a 2-day withdrawal from the treatment (Marutha Ravindran et al., 2007a). In cultured rat CGCs, CE had no effect on $\alpha 2$ mRNA (Follesa et al., 2004), whereas withdrawal from CE drastically up-regulated $\alpha 2$ mRNA and polypeptide expression (Follesa et al., 2004; Biggio et al., 2007).

In the hippocampus, wdr-CIE had no effect on $\alpha 2$ mRNA expression (Cagetti et al., 2003). Hippocampal $\alpha 2$ polypeptide level was not affected by CE treatment or withdrawal from the treatment (Matthews et al., 1998; Marutha Ravindran et al., 2007a). In cultured hippocampal neurons $\alpha 2$ mRNA expression was not affected by CE, whereas it was up-regulated by withdrawal from the treatment (Sanna et al., 2003). Studies on $\alpha 2$ subunit in vivo suggest that $\alpha 2$ is down-regulated brain region-specifically in the cerebral cortex and cerebellum by CE.

3. $\alpha 3$ Subunit. The expression of $\alpha 3$ mRNA in the cerebral cortex is generally not affected by CE (Montpied et al., 1991b; Morrow et al., 1991; Mhatre and Ticku, 1992) or wdr-CE (Mhatre and Ticku, 1992) (Tables 14 and 18). Down-regulation of cortical $\alpha 3$ polypeptide expression has been observed (Mhatre et al., 1993). CE had no effect on hippocampal $\alpha 3$ polypeptide expression (Matthews et al., 1998). In cultured hippocampal neu-

rons, CE down-regulated $\alpha 3$ mRNA expression, whereas withdrawal from the treatment up-regulated its expression (Sanna et al., 2003). The results on ethanol regulation of $\alpha 3$ expression are controversial but suggest mostly that ethanol does not regulate the $\alpha 3$ expression.

4. $\alpha 4$ Subunit. Cerebral cortical $\alpha 4$ mRNA expression was up-regulated after CE (Devaud et al., 1995) or withdrawal from the treatment (Mahmoudi et al., 1997) (Tables 14 and 18). Likewise, CE up-regulated cortical $\alpha 4$ polypeptide expression (Devaud et al., 1997; Matthews et al., 1998; Grobin et al., 2000; Marutha Ravindran et al., 2007a). There was heterogeneity in the up-regulation, the increase in expression being greatest (110%) in the cingulate cortex, although no effect was found in the parietal cortex (Grobin et al., 2000). The expression of $\alpha 4$ mRNA was up-regulated by CIE and $\alpha 4$ polypeptide by CE in cultured mouse cortical neurons (Sheela Rani and Ticku, 2006). Withdrawal from CE also up-regulated cortical $\alpha 4$ in vivo in rats (Devaud et al., 1997; Marutha Ravindran et al., 2007a). In contrast, wdr-CE and wdr-CIE had no effect on $\alpha 4$ mRNA in cultured mouse cortical neurons (Sheela Rani and Ticku, 2006). CE did not affect $\alpha 4$ mRNA expression in cultured rat cerebellar granule cells, whereas up-regulation of $\alpha 4$ mRNA and polypeptide was detected after withdrawal

TABLE 22
Effect of withdrawal from long-term ethanol treatment on GABA_A receptor subunit mRNA expression in cultured cells in vitro

Cells, [EtOH], Duration, Withdrawal Time	Subunit mRNA										Reference	
	$\alpha 1$	$\alpha 2$	$\alpha 3$	$\alpha 4$	$\alpha 5$	$\alpha 6$	$\beta 2$	$\beta 3$	$\gamma 2$	δ		
	% change											
Rat hippocampal neurons												
100 mM, 5 d, 3 h	-15	+30	+40	+30	N.S.				-21, -40 ^a			Sanna et al., 2003
100 mM, 5 d, 9 h											N.S.	Follesa et al., 2005
Rat cerebellar granule cells												
100 mM, 5 d, 3-12 h	-29			+46		-27			-76, -64 ^a			Follesa et al., 2003
100 mM, 5 d, 3 h		+159										Follesa et al., 2004
100 mM, 5 d, 6 h											-27	Follesa et al., 2005
100 mM, 5 d, 6 h	-22	+160		+43		-28			-33, -22 ^a		-23	Biggio et al., 2007
Mouse cortical neurons												
75 mM, 5 d, 5d	N.S.	N.S.		N.S.			N.S.	N.S.	N.S.			Sheela Rani and Ticku, 2006
75 mM, 5 d, 5d, CIE	N.S.	N.S.		N.S.			N.S.	N.S.	N.S.			Sheela Rani and Ticku, 2006

d, day(s); N.S., not significantly different from the control value.

^a Two values represent S and L splice variants, respectively.

TABLE 23
Effect of long-term ethanol treatment on GABA_A receptor subunit polypeptide expression in cultured cells in vitro

Cells, [EtOH], Treatment Time	Subunit Polypeptide						Reference
	$\alpha 1$	$\alpha 2$	$\alpha 4$	$\beta 2$	$\gamma 2$	δ	
% change							
Rat hippocampal neurons 100 mM, 5 d						+85	Follesa et al., 2005
Rat cerebellar granule cells 100 mM, 5 d						N.S.	Follesa et al., 2005
Mouse cortical neurons 75 mM, 5 d	-30			+35	N.S.		Marutha Ravindran and Ticku, 2006
75 mM, 5 d, CIE	-25			+29	N.S.		Marutha Ravindran and Ticku, 2006
75 mM, 5 d	-26	-30	+74	+24	N.S.		Sheela Rani and Ticku, 2006
75 mM, 5 d, CIE	-15	N.S.	N.S.	+158	N.S.		Sheela Rani and Ticku, 2006

d, day(s); N.S., not significantly different from the control value.

from the treatment (Follesa et al., 2003; Biggio et al., 2007). CE up-regulated hippocampal $\alpha 4$ mRNA and polypeptide expression (Devaud et al., 1995, 1997; Matthews et al., 1998; Marutha Ravindran et al., 2007a). Withdrawal from CE up-regulated hippocampal $\alpha 4$ mRNA in the study by Mahmoudi et al. (1997) [but not in that of Petrie et al. (2001)] and hippocampal $\alpha 4$ polypeptide in studies of Cagetti et al. (2003) and Marutha Ravindran et al. (2007a) (Table 16). CE did not affect $\alpha 4$ mRNA expression in cultured rat hippocampal neurons (Sanna et al., 2003), whereas withdrawal from the treatment up-regulated $\alpha 4$ mRNA and especially polypeptide expression (Sanna et al., 2003). CE down-regulated $\alpha 4$ polypeptide in the amygdala and nucleus accumbens (Papadeas et al., 2001). Life-long ethanol drinking down-regulated $\alpha 4$ mRNA in the olfactory tubercle (Sarviharju et al., 2006). Withdrawal from long-term ethanol treatment slightly up-regulated $\alpha 4$ mRNA in the thalamus (Mahmoudi et al., 1997).

Studies on $\alpha 4$ expression suggest that $\alpha 4$ subunit is strongly up-regulated by long-term ethanol treatment and especially on withdrawal. The effect is brain region-specific. More recently, Pignataro et al. (2007) have described a number of genes that are up-regulated by a short-term alcohol treatment of mouse cultured cortical

neurons. It is noteworthy that one the genes was *GABRA4*, the expression of which can be up-regulated by relevant concentrations of ethanol (10–60 mM) via activation of heat shock factor 1 that binds to “alcohol response element” of the $\alpha 4$ subunit gene promoter. The detailed mechanisms of other GABA_A receptor subunit gene regulation by alcohol are not known.

5. $\alpha 5$ Subunit. The expression of $\alpha 5$ mRNA was down-regulated by CE in the cerebral cortex (Mhatre and Ticku, 1992), whereas no effect was found by Devaud et al. (1995) (Table 14). CE did not affect cerebral cortical $\alpha 5$ polypeptide expression (Charlton et al., 1997). Withdrawal from CE down-regulated cortical $\alpha 5$ mRNA expression (Mhatre and Ticku, 1992). CE down-regulated $\alpha 5$ polypeptide expression in the cerebellum (Charlton et al., 1997). In the hippocampus, CE up-regulated $\alpha 5$ mRNA expression, although it had no effect on $\alpha 5$ polypeptide expression (Charlton et al., 1997). Withdrawal from CE did not affect hippocampal $\alpha 5$ mRNA expression (Mahmoudi et al., 1997; Petrie et al., 2001). Long-term ethanol treatment or withdrawal from it did not affect $\alpha 5$ mRNA expression in cultured rat hippocampal neurons (Sanna et al., 2003). The results of studies on the CE effect on $\alpha 5$ subunit suggest brain region-specific modulation of the expression.

TABLE 24
Effect of withdrawal from long-term ethanol treatment on GABA_A receptor subunit polypeptide expression in cultured cells in vitro

Cells, [EtOH], Duration, Withdrawal Time	Subunit Polypeptide						Reference
	$\alpha 1$	$\alpha 2$	$\alpha 4$	$\beta 2$	$\gamma 2$	δ	
% change							
Rat hippocampal neurons 100 mM, 5 d, 6 h			+120				Sanna et al., 2003
100 mM, 5 d, 6 h						-27	Follesa et al., 2005
100 mM, 5 d, 12 h						N.S.	Follesa et al., 2005
Rat cerebellar granule cells 100 mM, 5 d, 3 h		+159					Follesa et al., 2004
100 mM, 5 d, 6 h						+82	Follesa et al., 2005
100 mM, 5 d, 12 h						N.S.	Follesa et al., 2005
100 mM, 5 d, 6 h			+140				Biggio et al., 2007
Mouse cortical neurons 75 mM, 5 d, 2d	N.S.			N.S.	N.S.		Marutha Ravindran and Ticku, 2006
75 mM, 5 d, 7d, CIE	-20			+26	N.S.		Marutha Ravindran and Ticku, 2006
75 mM, 5 d, 5d	N.S.	N.S.	N.S.	N.S.	-61		Sheela Rani and Ticku, 2006
75 mM 5 d, 5d, CIE	N.S.	N.S.	N.S.	N.S.	-17		Sheela Rani and Ticku, 2006

N.S., not significantly different from the control value.

6. $\alpha 6$ Subunit. Cerebellar $\alpha 6$ mRNA expression was up-regulated by CE in rat (Mhatre and Ticku, 1992; Morrow et al., 1992) and mouse (Wu et al., 1995) cerebellum (Table 14). Withdrawal from CE resulted in a drastic increase in $\alpha 6$ mRNA expression in the study by Mhatre and Ticku (1992), but not in that of Petrie et al. (2001). Cerebellar $\alpha 6$ polypeptide was up-regulated by CE (Marutha Ravindran et al., 2007a), and after a 2-day wdr-CE, it was reversed to expression level of the control animals (Marutha Ravindran et al., 2007a). CE had no effect on $\alpha 6$ mRNA expression in cultured rat CGCs (Follesa et al., 2003), whereas wdr-CE resulted in $\alpha 6$ mRNA down-regulation (Follesa et al., 2003; Biggio et al., 2007). Long-term ethanol treatment of the $\alpha 6$ knockout mice expressing *Escherichia coli* β -galactosidase under the control of $\alpha 6$ gene promoter (Jones et al., 1997) did not alter β -galactosidase activity, suggesting that $\alpha 6$ up-regulation requires functional $\alpha 6$ subunits (Vekovisheva et al., 2000). The results suggest that CE up-regulates $\alpha 6$ expression in the cerebellum.

7. $\beta 1$ Subunit. Long-term ethanol up-regulated $\beta 1$ mRNA in the cerebral cortex (Mhatre and Ticku (1994), whereas no effect was found by Devaud et al. (1995) (Table 15). Expression of $\beta 1$ mRNA returned to control level after a 1-day wdr-CE (Mhatre and Ticku., 1994). CE had no effect on cortical or hippocampal $\beta 1$ polypeptide expression (Matthews et al., 1998). CE does not seem to consistently regulate $\beta 1$ expression.

8. $\beta 2$ Subunit. Cerebral cortical $\beta 2$ mRNA was up-regulated by CE (Mhatre and Ticku (1994), whereas no effect was found by Devaud et al. (1995) (Table 15). In cultured mouse cortical neurons, CE, CIE, wdr-CE, and wdr-CIE had no effect on $\beta 2$ mRNA expression (Sheela Rani and Ticku, 2006). Cerebral cortical $\beta 2$ polypeptide was up-regulated by CE in studies by Mhatre and Ticku (1994), Devaud et al. (1997), and Marutha Ravindran et al. (2007a), whereas no effect was found by Matthews et al. (1998). Expression of cortical $\beta 2$ mRNA was slightly up-regulated after a 1-day wdr-CE (Mhatre and Ticku, 1994), but it was not affected after a 2-day wdr-CIE (Petrie et al., 2001). Cortical $\beta 2$ polypeptide was found to be up-regulated by wdr-CE Devaud et al. (1997) but not by Marutha Ravindran et al. (2007a). In cultured mouse cortical neurons, $\beta 2$ polypeptide was up-regulated by CE and CIE (Marutha Ravindran and Ticku, 2006; Ravindran and Ticku, 2006; Sheela Rani and Ticku, 2006). The up-regulation persisted for at least 7 days after wdr-CIE (Marutha Ravindran and Ticku, 2006; Ravindran and Ticku, 2006) but not wdr-CE (Marutha Ravindran and Ticku, 2006; Ravindran and Ticku, 2006; Sheela Rani and Ticku, 2006). Cerebellar $\beta 2$ polypeptide was up-regulated by CE in rats in vivo (Marutha Ravindran et al., 2007), whereas it had no effect on cerebellar $\beta 2$ mRNA in mice (Wu et al., 1995). After 2-day wdr-CE (Marutha Ravindran et al., 2007) and 2-day wdr-CIE (Petrie et al., 2001), the expressions of cerebellar $\beta 2$ mRNA and polypeptide, respectively, were not differ-

ent from control values. CE or wdr-CE did not affect $\beta 2$ polypeptide expression in the hippocampus (Matthews et al., 1998; Marutha Ravindran et al., 2007a). Studies on $\beta 2$ subunit in vivo suggest that $\beta 2$ is up-regulated brain region-specifically in the cerebral cortex and cerebellum.

9. $\beta 3$ Subunit. Treatment of rats with CE up-regulated cerebral cortical $\beta 3$ mRNA in the study of Mhatre and Ticku (1994), but not in that of Devaud et al. (1995) (Table 15). After 1-day wdr-CE, $\beta 3$ mRNA value was still greater than in the corresponding control animals (Mhatre and Ticku, 1994). In cultured mouse cortical neurons CE, CIE, wdr-CE, and wdr-CIE had no effect on $\beta 3$ mRNA (Sheela Rani and Ticku, 2006). Cortical $\beta 3$ polypeptide was up-regulated by CE in the studies of Mhatre and Ticku (1994) and Devaud et al. (1997), but not in that of Matthews et al. (1998). The up-regulation in $\beta 3$ polypeptide persisted 6 to 8 h after wdr-CE (Devaud et al., 1997). In contrast to short CE exposure, a life-long ethanol consumption in alcohol-preferring rats down-regulated $\beta 3$ mRNA expression in the frontoparietal cortex and olfactory bulb (Sarviharju et al., 2006). The results suggest up-regulation of $\beta 3$ by short-term treatment, but down-regulation in some brain regions after life-long consumption.

10. $\gamma 2$ Subunit. Long-term ethanol treatment up-regulated the short (S) but not the long (L) splice variant of $\gamma 2$ mRNA in the cerebral cortex (Devaud et al., 1995) (Table 15). CE or wdr-CE had no effect on cortical $\gamma 2$ polypeptide expression (Devaud et al., 1997; Marutha Ravindran et al., 2007a). In cultured mouse cortical neurons, CE, CIE, wdr-CE, or wdr-CIE had no effect on $\gamma 2$ polypeptide expression in the studies of Marutha Ravindran and Ticku (2006) and Ravindran and Ticku (2006). In contrast, CE down-regulated $\gamma 2$ mRNA but not polypeptide expression (Sheela Rani and Ticku, 2006). In that study, wdr-CE and wdr-CIE had no effect on $\gamma 2$ mRNA, but $\gamma 2$ polypeptide expression was down-regulated after withdrawal from both types of long-term ethanol administration (Sheela Rani and Ticku, 2006). CE or wdr-CE had no effect on cerebellar $\gamma 2$ polypeptide expression in rats in vivo (Marutha Ravindran et al., 2007a). The S and L variants of $\gamma 2$ mRNA were down-regulated by CE and wdr-CE in cultured rat CGCs (Follesa et al., 2003; Biggio et al., 2007). Expressions of $\gamma 2$ mRNA splice variants in the hippocampus were not affected by wdr-CIE according to Petrie et al. (2001), whereas an increase in $\gamma 2S$ but not in $\gamma 2L$ mRNA after wdr-CIE was found by Cagetti et al. (2003). The $\gamma 2$ polypeptide was up-regulated by a 6-day CE in the hippocampus (Marutha Ravindran et al., 2007a). Expression of hippocampal $\gamma 2$ polypeptide was up-regulated after wdr-CIE (Cagetti et al., 2003), but not wdr-CE (Marutha Ravindran et al., 2007a). In cultured rat hippocampal neurons, CE and wdr-CE down-regulated $\gamma 2S$ and $\gamma 2L$ mRNA expression (Sanna et al., 2003). The

results suggest brain region-specific up-regulation of $\gamma 2$ expression by CE in the hippocampus.

11. $\gamma 1$ and $\gamma 3$ Subunits. Long-term ethanol up-regulated $\gamma 1$ mRNA and polypeptide expression in the cerebral cortex (Devaud et al., 1995, 1997) (Tables 15 and 19). Increased expression of cerebral cortical $\gamma 1$ polypeptide persisted on wdr-CE (Devaud et al., 1997). Hippocampal $\gamma 1$ mRNA expression was found to be up-regulated on wdr-CIE by Cagetti et al. (2003) but not by Petrie et al. (2001). CE did not up-regulate hippocampal $\gamma 1$ polypeptide (Matthews et al., 1998). Long-term ethanol had no effect on $\gamma 3$ mRNA expression in the cerebral cortex (Devaud et al., 1995). The results suggest up-regulation of $\gamma 1$ expression in ethanol withdrawal and unaltered expression of $\gamma 3$ by ethanol.

12. δ Subunit. The expression of δ mRNA and polypeptide was not affected by CE or wdr-CE in the cerebral cortex (Devaud et al., 1995; Marutha Ravindran et al., 2007b) (Tables 15, 17, 19, and 20). Expression of δ polypeptide in the cerebellum and hippocampus was down-regulated by CE (Marutha Ravindran et al., 2007b). After wdr-CE (Marutha Ravindran et al., 2007b) or wdr-CIE (Petrie et al., 2001), the expression of cerebellar and hippocampal δ mRNA and polypeptide did not differ from the expression in control animals, whereas a down-regulation of hippocampal δ polypeptide after wdr-CIE was detected (Cagetti et al., 2003). In cultured rat CGCs, CE had no effect on δ mRNA and polypeptide expression, whereas wdr-CE slightly down-regulated both δ mRNA and polypeptide (Follesa et al., 2005; Biggio et al., 2007). In cultured rat hippocampal neurons, CE strongly up-regulated δ mRNA and polypeptide expression (Follesa et al., 2005). The δ mRNA and polypeptide levels elevated by CE in hippocampal neurons returned to control levels by 12 h (Follesa et al., 2005). The results suggest brain region-specific down-regulation of δ subunit.

13. Studies of Continuous Intermittent Ethanol Administration on Subunit Cell Surface Expression. The group of Olsen and Spigelman have further investigated their CIE model in the rat. Cagetti et al. (2003) found that CIE strongly down-regulates hippocampal $\alpha 1$ and δ subunits but up-regulates synaptic $\alpha 4$ and $\gamma 2$ subunits. They dissected the GABA_A receptor-related molecular events in hippocampal CA1 slices after 1-h ethanol intoxication (Liang et al., 2007). The cell surface fractions of $\alpha 4$ and δ subunits (extrasynaptic), but not those of $\alpha 1$, $\alpha 5$, or $\gamma 2$ subunits, were decreased. This was accompanied by decreased magnitude of tonic GABA_A current, the enhancement of which was reduced by ethanol. At 48 h, the cell surface subunit content of $\alpha 4$ (80%) and $\gamma 2$ (82%) increased, whereas that of $\alpha 1$ (−50%) and δ (−79%) decreased (Liang et al., 2007). These changes were fully reversible, but persisted long after withdrawal from CIE treatment. The authors hypothesized that early ethanol tolerance might result from activation and subsequent internalization of extrasynaptic $\alpha 4\beta\delta$

receptors, subsequently leading to transcriptionally regulated increases in $\alpha 4$ and $\gamma 2$ subunits, resulting in insertion of the newly formed $\alpha 4\beta\gamma 2$ receptors at synapses.

14. Conclusions on Effects of Continuous Ethanol Administration on GABA_A Receptor Subunit Expression. CE and CIE treatments of rats in vivo down-regulate $\alpha 1$ expression in most brain regions studied and $\alpha 2$ expression specifically in the cerebral cortex and cerebellum. These treatments and especially withdrawal from them up-regulate $\alpha 4$ expression in the cerebral cortex and hippocampus. CE up-regulates $\alpha 6$ expression in the cerebellum. $\alpha 1$ and $\alpha 6$ subunits are located in $\beta 2$ - $\alpha 6$ - $\alpha 1$ - $\gamma 2$ GABA_A receptor subunit gene cluster in the same positions as $\alpha 2$ and $\alpha 4$ in the homologous $\beta 1$ - $\alpha 4$ - $\alpha 2$ - $\gamma 1$ gene cluster (Russek, 1999). These two clusters have obviously evolved from a common ancestral cluster. The $\alpha 1/\alpha 2$ genes, homologous genes from $\beta 2$ - $\alpha 6$ - $\alpha 1$ - $\gamma 2$ and $\beta 1$ - $\alpha 4$ - $\alpha 2$ - $\gamma 1$ GABA_A receptor subunit gene clusters may contain similar regulatory elements responsible for their down-regulation by CE; likewise, the $\alpha 6/\alpha 4$ genes may contain similar regulatory elements responsible for the up-regulation by CE. Up-regulation of $\beta 2$ and $\beta 3$ is suggested by most studies. CE and CIE brain region-specific up-regulate $\gamma 2$ in the hippocampus. The δ subunit is down-regulated brain region-specifically in the cerebellum and in hippocampus. These changes suggest down-regulation of $\alpha 1\beta\gamma 2$, $\alpha 2\beta\gamma 2$, $\alpha 4\beta\delta$, and $\alpha 6\beta\delta$ receptor subtypes and up-regulation of $\alpha 4\beta\gamma 2$ and $\alpha 6\beta\gamma 2$ receptor subtypes. The CE-induced changes in GABA_A receptor expression are short-lived (usually 1–2 days), whereas CIE-induced changes take much longer to revert.

In vitro studies with cultured cells treated with CE or CIE have produced quite controversial results compared with studies in vivo. The in vitro results from rat hippocampal neurons or cerebellar granule cells are often opposite the results received from hippocampus or cerebellum of rats treated in vivo. Therefore, the in vitro results should be interpreted very cautiously.

IV. Novel Opportunities to Target the GABA_A Receptor System with Therapeutics

The drugs now used clinically that target the GABA_A receptor system are often very efficacious in the short term, but they lose their pharmacological effects to a great extent during repeated administration. The most important disease group to find more efficient pharmacological treatment is the anxiety disorders, but the treatment of many other indications also would benefit from compounds with more selective action either at BZ-site of synaptic GABA_A receptors or at other sites of, for example, extrasynaptic receptors in the hippocampus, cortex, and thalamus or at selective cell types, such as cortical interneurons. We give here some examples.

Medical genetics has progressed to the level that we know several GABA_A receptor subunit genes that are

somehow associated with neuropsychiatric illnesses (Korpi and Sinkkonen, 2006). Unfortunately, in most cases, it is premature to make detailed hypotheses regarding the mechanisms by which these genetic associations might affect the development or progress of a particular set of symptoms or a disease. However, there is one very good example for rational drug development work, namely schizophrenia, in which deficient function in cortical interneurons has been established (Benes et al., 1991; Lewis et al., 2005; Benes, 2009) and specific GABA_A receptor $\alpha 2$ subunit changes have been documented in principal neurons (Cruz et al., 2009), provoking a simple idea of targeting the $\alpha 2$ subunits with subtype-selective compounds, apparently with agonists to counteract the reduced interneuron activity. Similar compounds should be tested also for generalized anxiety and other anxiety disorders, the idea being based on results from a study on mouse model with targeted inactivation of $\alpha 2$ BZ-sites [so called $\alpha 2$ (H101R) mice]. In that model, diazepam is inactive in inducing short-term anxiolysis (Löw et al., 2000). This idea was corroborated in experiments with $\alpha 2$ knockout mice (Dixon et al., 2008). It is noteworthy that work with $\alpha 3$ knockout mice suggests a role for reduced function of $\alpha 3$ subunit-containing GABA_A receptors in schizophrenia-like sensorimotor deficits and excessive dopamine function (Yee et al., 2005); also, in this mouse model, diazepam had little anxiolytic effect. Experimental compounds having $\alpha 2$ and/or $\alpha 3$ agonist selectivity already exist (e.g., Atack et al., 2006; Jennings et al., 2006; Van Laere et al., 2008; Taliani et al., 2009), but they have not been well tested in patients with anxiety (Atack, 2008). Furthermore, neurobiological findings suggest that these same receptor subtypes at the spinal level should be tested as targets for alleviating neuropathic pain (Knabl et al., 2008).

As discussed previously (Korpi and Sinkkonen, 2006), the arsenal for hypnotic drugs acting on the BZ-site is already satisfactory; triazolam, midazolam, zolpidem, zopiclone, and zaleplon demonstrate acceptable short-term efficacies with a selection of half-lives and pharmacokinetic variations to fit needs of different patients. It is also possible that these compounds retain their hypnotic activity in long-term treatment (up to 12 months) as has been recently shown for the *S* isomer of zopiclone (eszopiclone) (Krystal et al., 2003; Roth et al., 2005). The GABA-site agonist gaboxadol acts differently from BZ-site ligands to alleviate insomnia (Wafford and Ebert, 2006), but its phase III trial as a hypnotic was stopped as a result of adverse effects in some patient populations. The role of direct GABA-site agonists should still be examined, because gaboxadol seems to target preferentially a clearly different receptor population than BZ compounds. Gaboxadol targets the extrasynaptic GABA_A receptors (Chandra et al., 2006), its high-affinity binding occurs in different brain regions from BZ binding (Friemel et al., 2007), and it does not

show cross-tolerance with BZs in rodent motor function tests (Voss et al., 2003). It is interesting that neuroprotective efficacy (e.g., ischemia models) has been stronger with direct GABA agonists such as muscimol than with BZs (Green et al., 2000). In a number of knockout and transgenic mouse models, the sedative actions of gaboxadol and muscimol correlate with the mouse forebrain density of high-affinity muscimol binding, presumably reflecting an extrasynaptic population of GABA_A receptors containing $\alpha 4$ and δ subunits (Chandra et al., 2010).

Neurosteroids and neuroactive steroids have actions on GABA_A receptors (Schumacher et al., 2003; Lambert et al., 2003), especially on δ subunit-containing extrasynaptic receptors (Chandra et al., 2006), and mediate the fast nongenomic effects of these compounds either by enhancing or inhibiting the receptor activity (Majewska, 1992). There have been several attempts to develop neurosteroid compounds as anesthetics; since the pioneering studies of Hans Selye, they have been long known to possess powerful, rapidly acting and quickly residing (quick metabolism) effects in rats. A synthetic steroid ganaxolone (3 α -hydroxy-3 β -methyl-5 α -pregnan-20-one) is a potent GABA_A receptor modulator without efficacy on nuclear hormone receptors (Carter et al., 1997). It holds promise in some forms of epilepsy, such as infantile spasms and catamenial epilepsy (Rogawski and Reddy, 2002), when there is a probable deficiency in brain concentrations of neurosteroids.

Inverse agonists of the GABA_A receptors might be used to treat alcoholism (and eating disorders/obesity?) and cognitive impairment (Korpi and Sinkkonen, 2006). In this context, it seems to be clear that the new compounds should have subtype selectivity rather than being general nonselective inverse agonists. Nonselective BZ-site inverse agonists are strongly anxiogenic and proconvulsant or convulsant, and in long-term treatment, these actions might get sensitized and abolish any initial treatment effect. Subtype-selective inverse agonists might be safer in this regard, but it remains to be fully established. Inverse agonists for the $\alpha 5$ subunit-containing receptors should be tested for cognitive improvement, because both genetically and pharmacologically rendered impairment of $\alpha 5$ subunits seems to improve learning and memory in mouse and monkey models (Crestani et al., 2002; Collinson et al., 2006; Ballard et al., 2009).

Regarding alcoholism, the situation is still unclear. Ethanol actions at extrasynaptic GABA_A receptors containing δ and $\alpha 4$ or $\alpha 6$ subunits have been shown to be antagonized by the BZ ligand Ro 15-4513 in some experiments (Wallner et al., 2006; Hanchar et al., 2006) but not in others (Borghese et al., 2007; Korpi et al., 2007). The excitement on alcohol-antagonistic effects of Ro 15-4513 and several other inverse agonists started already in the 1980s (Suzdak et al., 1986; Bonetti et al., 1988) without any therapeutic breakthroughs up to now. Still needed are more selective molecules and/or better ani-

mal models to study the mechanisms of the putative alcohol antagonistic action.

In addition to the drugs targeting the main GABA_A receptor subtypes, one might also attempt to target the minor ones (e.g., ϵ and θ subunit-containing receptors) that are highly enriched (e.g., in monoaminergic nuclei) (Sinkkonen et al., 2000; Moragues et al., 2002) and might serve as selective targets for non-BZ site compounds to regulate neuronal activity in ascending monoamine pathways. The ϵ and θ subunits are most likely assembled with the $\alpha 3$ subunits, but hardly anything has been published in terms of selective pharmacology and functional activity of these receptor subtypes (Ranna et al., 2006). Neuronal cell population-specific modulation may eventually be needed (e.g., for the pharmacological regulation of behaviors such as feeding), because brain regionally discrete effects are often different from systemic drug effects.

Finally, it should be remembered that the GABA_A receptor is a large molecular complex having surprisingly many different binding sites and interactions with other proteins. Therefore, it is possible that the future will bring more compounds having novel target sites at the receptor complex or at the receptor-associated proteins, which might then provide different pharmacological profiles and effects in human.

Acknowledgments. This work was supported by the Finnish Society of Sciences and Letters (to M.U.-O.), Oskar Öflund Foundation (to M.U.-O.), the Turku University Foundation (to M.U.-O.), the Academy of Finland (to E.R.K.), the Finnish Foundation for Alcohol Studies (to E.R.K.), and the Sigrid Jusélius Foundation (to E.R.K.). In addition, we thank Petri Vainio-Ketola for Fig. 3 and Prof. Werner Sieghart and Elsevier Ltd. for the use of Fig. 4.

REFERENCES

Aitta-Aho T, Vekovischeva OY, Neuvonen PJ, and Korpi ER (2009) Reduced benzodiazepine tolerance, but increased flumazenil-precipitated withdrawal in AMPA-receptor GluR-A subunit-deficient mice. *Pharmacol Biochem Behav* **92**:283–290.

Arnot MI, Davies M, Martin IL, and Bateson AN (2001) GABA_A receptor gene expression in rat cortex: differential effects of two chronic diazepam treatment regimes. *J Neurosci Res* **64**:617–625.

Ashton H (1991) Protracted withdrawal syndromes from benzodiazepines. *J Subst Abuse Treat* **8**:19–28.

Ashton H (1994) Guidelines for the rational use of benzodiazepines. When and what to use. *Drugs* **48**:25–40.

Atack JR (2003) Anxiolytic compounds acting at the GABA_A receptor benzodiazepine binding site. *Curr Drug Targets CNS Neurol Disord* **2**:213–232.

Atack JR (2008) GABA_A receptor subtype-selective efficacy: TPA023, an $\alpha 2/\alpha 3$ selective non-sedating anxiolytic and $\alpha 5$ selective cognition enhancer. *CNS Neurosci Ther* **14**:25–35.

Atack JR, Wafford KA, Tye SJ, Cook SM, Sohal B, Pike A, Sur C, Melillo D, Bristow L, Bromidge F, et al. (2006) TPA023 [7-(1,1-Dimethylethyl)-6-(2-ethyl-2H-1,2,4-triazol-3-ylmethoxy)-3-(2-fluorophenyl)-1,2,4-triazolo[4,3-b]pyridazine], an agonist selective for $\alpha 2$ - and $\alpha 3$ -containing GABA_A receptors, is a non-sedating anxiolytic in rodents and primates. *J Pharmacol Exp Ther* **316**:410–422.

Bailey ME, Albrecht BE, Johnson KJ, and Darlison MG (1999) Genetic linkage and radiation hybrid mapping of the three human GABA_C receptor ρ subunit genes: GABRR1, GABRR2 and GABRR3. *Biochim Biophys Acta* **1447**:307–312.

Ballard TM, Knoflach F, Prinszen E, Borroni E, Vivian JA, Basile J, Gasser R, Moreau JL, Wettstein JG, Buettelmann B, et al. (2009) RO4938581, a novel cognitive enhancer acting at GABA_A $\alpha 5$ subunit-containing receptors. *Psychopharmacology (Berl)* **202**:207–223.

Banker GA and Cowan WM (1977) Rat hippocampal neurons in dispersed cell culture. *Brain Res* **126**:397–442.

Barbaccia ML (2004) Neurosteroidogenesis: relevance to neurosteroid actions in brain and modulation by psychotropic drugs. *Crit Rev Neurobiol* **16**:67–74.

Barnes EM Jr (1996) Use-dependent regulation of GABA_A receptors. *Int Rev Neurobiol* **39**:53–76.

Bateson AN (2002) Basic pharmacologic mechanisms involved in benzodiazepine tolerance and withdrawal. *Curr Pharm Des* **8**:5–21.

Baumgartner BJ, Harvey RJ, Darlison MG, and Barnes EM Jr (1994) Developmental

up-regulation and agonist-dependent down-regulation of GABA_A receptor subunit mRNAs in chick cortical neurons. *Mol Brain Res* **26**:9–17.

Ben-Ari Y, Khazipour R, Leinekugel X, Caillard O, and Gaiarsa JL (1997) GABA_A, NMDA and AMPA receptors: a developmentally regulated 'menage a trois'. *Trends Neurosci* **20**:523–529.

Benes FM (2009) Neural circuitry models of schizophrenia: is it dopamine, GABA, glutamate, or something else? *Biol Psychiatry* **65**:1003–1005.

Benes FM, McSparren J, Bird ED, SanGiovanni JP, and Vincent SL (1991) Deficits in small interneurons in prefrontal and cingulate cortices of schizophrenic and schizoaffective patients. *Arch Gen Psychiatry* **48**:996–1001.

Benke D, Fritschy JM, Trzeciak A, Bannwarth W, and Mohler H (1994) Distribution, prevalence, and drug binding profile of γ -aminobutyric acid type A receptor subtypes differing in the β -subunit variant. *J Biol Chem* **269**:27100–27107.

Biggio F, Gorini G, Caria S, Murru L, Mostallino MC, Sanna E, and Follesa P (2006) Plastic neuronal changes in GABA_A receptor gene expression induced by progesterone metabolites: in vitro molecular and functional studies. *Pharmacol Biochem Behav* **84**:545–554.

Biggio F, Gorini G, Caria S, Murru L, Sanna E, and Follesa P (2007) Flumazenil selectively prevents the increase in $\alpha 4$ -subunit gene expression and an associated change in GABA_A receptor function induced by ethanol withdrawal. *J Neurochem* **102**:657–666.

Birnir B and Korpi ER (2007) The impact of sub-cellular location and intracellular neuronal proteins on properties of GABA_A receptors. *Curr Pharm Des* **13**:3169–3177.

Bogdanov Y, Michels G, Armstrong-Gold C, Haydon PG, Lindstrom J, Pangalos M, and Moss SJ (2006) Synaptic GABA_A receptors are directly recruited from their extrasynaptic counterparts. *EMBO J* **25**:4381–4389.

Bonetti EP, Burkard WP, Gabl M, Hunkeler W, Lorez HP, Martin JR, Moehler H, Osterrieder W, Pieri L, and Polc P (1988) Ro 15-4513: partial inverse agonism at the BZR and interaction with ethanol. *Pharmacol Biochem Behav* **31**:733–749.

Borghese CM, Störustovu S, Ebert B, Herd MB, Belelli D, Lambert JJ, Marshall G, Wafford KA, and Harris RA (2007) The δ subunit of γ -aminobutyric acid type A receptors does not confer sensitivity to low concentrations of ethanol. *J Pharmacol Exp Ther* **316**:1360–1368.

Borghese CM, Werner DF, Topf N, Baron NV, Henderson LA, Boehm SL 2nd, Blednov YA, Saad A, Dai S, Pearce RA, et al. (2006) An isoflurane- and alcohol-insensitive mutant GABA_A receptor $\alpha 1$ subunit with near-normal apparent affinity for GABA: characterization in heterologous systems and production of knockin mice. *J Pharmacol Exp Ther* **319**:208–218.

Bovolenta P, Santi MR, Puia G, Costa E, and Grayson D (1992) Expression patterns of gamma-aminobutyric acid type A receptor subunit mRNAs in primary cultures of granule neurons and astrocytes from neonatal rat cerebellum. *Proc Natl Acad Sci U S A* **89**:9344–9348.

Boue-Grabot E, Roudbaraki M, Bascles L, Tramu G, Bloch B, and Garret M (1998) Expression of GABA receptor ρ subunits in rat brain. *J Neurochem* **70**:899–907.

Bouzigues C and Dahan M (2007) Transient directed motions of GABA_A receptors in growth cones detected by a speed correlation index. *Biophys J* **92**:654–660.

Brodie MS, Scholz A, Weiger TM, and Dopic AM (2007) Ethanol interactions with calcium-dependent potassium channels. *Alcohol Clin Exp Res* **31**:1625–1632.

Brooks-Kayal AR, Jin H, Price M, and Dichter MA (1998) Developmental expression of GABA_A receptor subunit mRNAs in individual hippocampal neurons in vitro and in vivo. *J Neurochem* **70**:1017–1028.

Brown MJ and Bristow DR (1996) Molecular mechanisms of benzodiazepine-induced down-regulation of GABA_A receptor $\alpha 1$ subunit protein in rat cerebellar granule cells. *Br J Pharmacol* **118**:1103–1110.

Brown MJ, Wood MD, Coldwell MC, and Bristow DR (1998) γ -Aminobutyric acid_A receptor function is desensitized in rat cultured cerebellar granule cells following chronic flunitrazepam treatment. *J Neurochem* **71**:1232–1240.

Brüning I, Penschuck S, Berninger B, Benson J, and Fritschy JM (2001) BDNF reduces miniature inhibitory postsynaptic currents by rapid downregulation of GABA_A receptor surface expression. *Eur J Neurosci* **13**:1320–1328.

Buck KJ, Hahner L, Sikela J, and Harris RA (1991) Chronic ethanol treatment alters brain levels of γ -aminobutyric acid_A receptor subunit mRNAs: relationship to genetic differences in ethanol withdrawal seizure severity. *J Neurochem* **57**:1452–1455.

Bulleit RF and Hsieh T (2000) MEK inhibitors block BDNF-dependent and independent expression of GABA_A receptor subunit mRNAs in cultured mouse cerebellar granule neurons. *Brain Res Dev Brain Res* **119**:1–10.

Cagetti E, Liang J, Spigelman I, and Olsen RW (2003) Withdrawal from chronic intermittent ethanol treatment changes subunit composition, reduces synaptic function, and decreases behavioral responses to positive allosteric modulators of GABA_A receptors. *Mol Pharmacol* **63**:53–64.

Carafoli E, Santella L, Branca D, and Brint M (2001) Generation, control, and processing of cellular calcium signals. *Crit Rev Biochem Mol Biol* **36**:107–260.

Caraiscos VB, Elliott EM, You-Ten KE, Cheng VY, Belelli D, Newell JG, Jackson MF, Lambert JJ, Rosahl TW, Wafford KA, et al. (2004) Tonic inhibition in mouse hippocampal CA1 pyramidal neurons is mediated by $\alpha 5$ subunit-containing γ -aminobutyric acid type A receptors. *Proc Natl Acad Sci U S A* **101**:3662–3667.

Cardoso RA, Brozowski SJ, Chavez-Noriega LE, Harpold M, Valenzuela CF, and Harris RA (1999) Effects of ethanol on recombinant human neuronal nicotinic acetylcholine receptors expressed in *Xenopus* oocytes. *J Pharmacol Exp Ther* **289**:774–780.

Carlson BX, Elster L, and Schousboe A (1998) Pharmacological and functional implications of developmentally-regulated changes in GABA_A receptor subunit expression in the cerebellum. *Eur J Pharmacol* **352**:1–14.

Carter RB, Wood PL, Wieland S, Hawkinson JE, Belelli D, Lambert JJ, White HS, Wolf HH, Mirsadeghi S, Tahir SH, et al. (1997) Characterization of the anticonvulsant properties of ganaxolone (CCD 1042; 3 α -hydroxy-3 β -methyl-5 α -pregnan-20-one), a selective, high-affinity, steroid modulator of the γ -aminobutyric acid_A receptor. *J Pharmacol Exp Ther* **280**:1284–1295.

Chandra D, Halonen LM, Linden A-M, Proccaccini C, Hellsten K, Homanics GE, and

- Korpi ER (2010) Prototypic GABA_A receptor agonist muscimol acts preferentially via forebrain high-affinity binding sites. *Neuropsychopharmacology*. doi: 10.1038/npp.2009.203.
- Chandra D, Jia F, Liang J, Peng Z, Suryanarayanan A, Werner DF, Spigelman I, Houser CR, Olsen RW, Harrison NL, et al. (2006) GABA_A receptor $\alpha 4$ subunits mediate extrasynaptic inhibition in thalamus and dentate gyrus and the action of gaboxadol. *Proc Natl Acad Sci U S A* **103**:15230–15235.
- Charlton ME, Sweetnam PM, Fitzgerald LW, Terwilliger RZ, Nestler EJ, and Duman RS (1997) Chronic ethanol administration regulates the expression of GABA_A receptor $\alpha 1$ and $\alpha 2$ subunits in the ventral tegmental area and hippocampus. *J Neurochem* **68**:121–127.
- Chen S, Huang X, Zeng XJ, Sieghart W, and Tietz EI (1999) Benzodiazepine-mediated regulation of $\alpha 1$, $\alpha 2$, $\beta 1$ –3 and $\gamma 2$ GABA_A receptor subunit proteins in the rat brain hippocampus and cortex. *Neuroscience* **93**:33–44.
- Chen ZW, Chang CS, Leil TA, and Olsen RW (2007) C-terminal modification is required for GABARAP-mediated GABA_A receptor trafficking. *J Neurosci* **27**:6655–6663.
- Chen ZW and Olsen RW (2007) GABA_A receptor associated proteins: a key factor regulating GABA_A receptor function. *J Neurochem* **100**:279–294.
- Collinson N, Attack JR, Laughton P, Dawson GR, and Stephens DN (2006) An inverse agonist selective for $\alpha 5$ subunit-containing GABA_A receptors improves encoding and recall but not consolidation in the Morris water maze. *Psychopharmacology (Berl)* **188**:619–628.
- Compagnone NA and Mellon SH (2000) Neurosteroids: biosynthesis and function of these novel neuromodulators. *Front Neuroendocrinol* **21**:1–56.
- Crawford DK, Trudell JR, Bertaccini EJ, Li K, Davies DL, and Alkana RL (2007) Evidence that ethanol acts on a target in loop 2 of the extracellular domain of $\alpha 1$ glycine receptors. *J Neurochem* **102**:2097–2109.
- Crestani F, Keist R, Fritschy JM, Benke D, Vogt K, Prut L, Blüthmann H, Möhler H, and Rudolph U (2002) Trace fear conditioning involves hippocampal $\alpha 5$ GABA_A receptors. *Proc Natl Acad Sci U S A* **99**:8980–8985.
- Crestani F, Löw K, Keist R, Mandelli M, Möhler H, and Rudolph U (2001) Molecular targets for the myorelaxant action of diazepam. *Mol Pharmacol* **59**:442–445.
- Cruz DA, Weaver CL, Lovallo EM, Melchitzky DS, and Lewis DA (2009) Selective alterations in postsynaptic markers of chandelier cell inputs to cortical pyramidal neurons in subjects with schizophrenia. *Neuropsychopharmacology* **34**:2112–2124.
- Das P, Lilly SM, Zerda R, Gunning WT 3rd, Alvarez FJ, and Tietz EI (2008) Increased AMPA receptor GluR1 subunit incorporation in rat hippocampal CA1 synapses during benzodiazepine withdrawal. *J Comp Neurol* **511**:832–846.
- Dennerstein L, Spencer-Gardner C, Gotts G, Brown JB, Smith MA, and Burrows GD (1985) Progesterone and the premenstrual syndrome: a double blind crossover trial. *Br Med J (Clin Res Ed)* **290**:1617–1621.
- Devaud LL, Fritschy JM, Sieghart W, and Morrow AL (1997) Bidirectional alterations of GABA_A receptor subunit peptide levels in rat cortex during chronic ethanol consumption and withdrawal. *J Neurochem* **69**:126–130.
- Devaud LL, Smith PD, Grayson DR, and Morrow AL (1995) Chronic ethanol consumption differentially alters the expression of γ -aminobutyric acid_A receptor subunit mRNAs in rat cerebral cortex: competitive, quantitative reverse transcriptase-polymerase chain reaction analysis. *Mol Pharmacol* **48**:861–868.
- De Witte P, Pinto E, Anseau M, and Verbanck P (2003) Alcohol and withdrawal: from animal research to clinical issues. *Neurosci Biobehav Rev* **27**:189–197.
- Dixon CI, Rosahl TW, and Stephens DN (2008) Targeted deletion of the GABRA2 gene encoding $\alpha 2$ -subunits of GABA_A receptors facilitates performance of a conditioned emotional response, and abolishes anxiolytic effects of benzodiazepines and barbiturates. *Pharmacol Biochem Behav* **90**:1–8.
- Dopico AM and Lovinger DM (2009) Acute alcohol action and desensitization of ligand-gated ion channels. *Pharmacol Rev* **61**:98–114.
- Ernst M, Brauchart D, Boesch S, and Sieghart W (2003) Comparative modeling of GABA_A receptors: limits, insights, future developments. *Neuroscience* **119**:933–943.
- Essrich C, Lorez M, Benson JA, Fritschy JM, and Lüscher B (1998) Postsynaptic clustering of major GABA_A receptor subtypes requires the $\gamma 2$ subunit and gephyrin. *Nat Neurosci* **1**:563–571.
- Estes JW (1995) The road to tranquility: the search for selective anti-anxiety agents. *Synapse* **21**:10–20.
- Everitt AB, Luu T, Cromer B, Tierney ML, Birnir B, Olsen RW, and Gage PW (2004) Conductance of recombinant GABA_A channels is increased in cells co-expressing GABA_A receptor-associated protein. *J Biol Chem* **279**:21701–21706.
- Fahey JM, Pritchard GA, Grassi JM, Pratt JS, Shader RI, and Greenblatt DJ (1999) In situ hybridization histochemistry as a method to assess GABA_A receptor subunit mRNA expression following chronic alprazolam administration. *J Psychopharmacol* **13**:211–218.
- File SE (1985) Tolerance to the behavioral actions of benzodiazepines. *Neurosci Biobehav Rev* **9**:113–121.
- Follesa P, Biggio F, Mancuso L, Cabras S, Caria S, Gorini G, Manca A, Orru A, and Biggio G (2004) Ethanol withdrawal-induced up-regulation of the $\alpha 2$ subunit of the GABA_A receptor and its prevention by diazepam or γ -hydroxybutyric acid. *Mol Brain Res* **120**:130–137.
- Follesa P, Cagett E, Mancuso L, Biggio F, Manca A, Maciocco E, Massa F, Desole MS, Carta M, Busonero F, et al. (2001a) Increase in expression of the GABA_A receptor $\alpha 4$ subunit gene induced by withdrawal of, but not by long-term treatment with, benzodiazepine full or partial agonists. *Mol Brain Res* **92**:138–148.
- Follesa P, Concas A, Porcu P, Sanna E, Serra M, Mostallino MC, Purdy RH, and Biggio G (2001b) Role of allopregnanolone in regulation of GABA_A receptor plasticity during long-term exposure to and withdrawal from progesterone. *Brain Res Rev* **37**:81–90.
- Follesa P, Mancuso L, Biggio F, Cagett E, Franco M, Trapani G, and Biggio G (2002) Changes in GABA_A receptor gene expression induced by withdrawal of, but not by long-term exposure to, zaleplon or zolpidem. *Neuropharmacology* **42**:191–198.
- Follesa P, Mancuso L, Biggio F, Mostallino MC, Manca A, Mascia MP, Busonero F, Talani G, Sanna E, and Biggio G (2003) γ -Hydroxybutyric acid and diazepam antagonize a rapid increase in GABA_A receptors $\alpha 4$ subunit mRNA abundance induced by ethanol withdrawal in cerebellar granule cells. *Mol Pharmacol* **63**:896–907.
- Follesa P, Mostallino MC, Biggio F, Gorini G, Caria S, Busonero F, Murru L, Mura ML, Sanna E, and Biggio G (2005) Distinct patterns of expression and regulation of GABA receptors containing the delta subunit in cerebellar granule and hippocampal neurons. *J Neurochem* **94**:659–671.
- Follesa P, Serra M, Cagett E, Pisu MG, Porta S, Floris S, Massa F, Sanna E, and Biggio G (2000) Allopregnanolone synthesis in cerebellar granule cells: roles in regulation of GABA_A receptor expression and function during progesterone treatment and withdrawal. *Mol Pharmacol* **57**:1262–1270.
- Friemel A, Ebert B, Hutson PH, Brust P, Nieber K, and Deuther-Conrad W (2007) Postnatal development and kinetics of [³H]gaboxadol binding in rat brain: in vitro homogenate binding and quantitative autoradiography. *Brain Res* **1170**:39–47.
- Fritschy JM, Benke D, Mertens S, Oertel WH, Bachi T, and Möhler H (1992) Five subtypes of type A γ -aminobutyric acid receptors identified in neurons by double and triple immunofluorescence staining with subunit-specific antibodies. *Proc Natl Acad Sci U S A* **89**:6726–6730.
- Fritschy JM and Möhler H (1995) GABA_A-receptor heterogeneity in the adult rat brain: differential regional and cellular distribution of seven major subunits. *J Comp Neurol* **359**:154–194.
- Gallager DW, Lakoski JM, Gonsalves SF, and Rauch SL (1984) Chronic benzodiazepine treatment decreases postsynaptic GABA sensitivity. *Nature* **308**:74–77.
- Gangisetty O and Reddy DS (2009) The optimization of TaqMan real-time RT-PCR assay for transcriptional profiling of GABA_A receptor subunit plasticity. *J Neurosci Methods* **181**:58–66.
- Gao B and Fritschy JM (1994) Selective allocation of GABA_A receptors containing the $\alpha 1$ subunit to neurochemically distinct subpopulations of rat hippocampal interneurons. *Eur J Neurosci* **6**:837–853.
- Gao B, Fritschy JM, Benke D, and Möhler H (1993) Neuron-specific expression of GABA_A-receptor subtypes: differential association of the $\alpha 1$ - and $\alpha 3$ -subunits with serotonergic and GABAergic neurons. *Neuroscience* **54**:881–892.
- Gao B, Hornung JP, and Fritschy JM (1995) Identification of distinct GABA_A-receptor subtypes in cholinergic and parvalbumin-positive neurons of the rat and marmoset medial septum-diagonal band complex. *Neuroscience* **65**:101–117.
- Gault LM and Siegel RE (1998) NMDA receptor stimulation selectively initiates GABA_A receptor δ subunit mRNA expression in cultured rat cerebellar granule neurons. *J Neurochem* **70**:1907–1915.
- Glykys J, Peng Z, Chandra D, Homanics GE, Houser CR, and Mody I (2007) A new naturally occurring GABA(A) receptor subunit partnership with high sensitivity to ethanol. *Nat Neurosci* **10**:40–48.
- Grayson DR, Zhu W, Harris BT, Vicini S, and Zheng T (1998) Differentially expressed GABA_A-receptor subunits result in structurally and functionally receptor assemblies following excitatory afferent synaptic transmission. *Perspect Dev Neurobiol* **5**:193–205.
- Green AR, Hainsworth AH, and Jackson DM (2000) GABA potentiation: a logical pharmacological approach for the treatment of acute ischaemic stroke. *Neuropharmacology* **39**:1483–1494.
- Grobin AC, Fritschy JM, and Morrow AL (2000) Chronic ethanol administration alters immunoreactivity for GABA_A receptor subunits in rat cortex in a region-specific manner. *Alcohol Clin Exp Res* **24**:1137–1144.
- Grobin AC and Morrow AL (2000) 3 α -hydroxy-5 α -pregnan-20-one exposure reduces GABA_A receptor $\alpha 4$ subunit mRNA levels. *Eur J Pharmacol* **409**:R1–2.
- Gulinello M, Gong QH, Li X, and Smith SS (2001) Short-term exposure to a neuroactive steroid increases $\alpha 4$ GABA_A receptor subunit levels in association with increased anxiety in the female rat. *Brain Res* **910**:55–66.
- Gutiérrez A, Khan ZU, and De Blas AL (1994) Immunocytochemical localization of $\gamma 2$ short and $\gamma 2$ long subunits of the GABA_A receptor in the rat brain. *J Neurosci* **14**:7168–7179.
- Gutiérrez A, Khan ZU, and De Blas AL (1996) Immunocytochemical localization of the $\alpha 6$ subunit of the γ -aminobutyric acid_A receptor in the rat nervous system. *J Comp Neurol* **365**:504–510.
- Hancher HJ, Chutrinopkun P, Meera P, Supavilai P, Sieghart W, Wallner M, and Olsen RW (2006) Ethanol potently and competitively inhibits binding of the alcohol antagonist Ro15-4513 to $\alpha 4/\beta 3\delta$ GABA_A receptors. *Proc Natl Acad Sci U S A* **103**:8546–8551.
- Harney SC, Freguelli BG, and Lambert JJ (2003) Phosphorylation influences neurosteroid modulation of synaptic GABA_A receptors in rat CA1 and dentate gyrus neurons. *Neuropharmacology* **45**:873–883.
- Harris BT, Costa E, and Grayson DR (1995) Exposure of neuronal cultures to K⁺ depolarization or to N-methyl-D-aspartate increases the transcription of genes encoding the $\alpha 1$ and $\alpha 5$ GABA_A receptor subunits. *Mol Brain Res* **28**:338–342.
- Hedblom E and Kirkness EF (1997) A novel class of GABA_A receptor subunit in tissues of the reproductive system. *J Biol Chem* **272**:15346–15350.
- Heikkinen AE, Möykkynen TP, and Korpi ER (2009) Long-lasting modulation of glutamatergic transmission in VTA dopamine neurons after a single dose of benzodiazepine agonists. *Neuropsychopharmacology* **34**:290–298.
- Heninger C and Gallager DW (1988) Altered γ -aminobutyric acid/benzodiazepine interaction after chronic diazepam exposure. *Neuropharmacology* **27**:1073–1076.
- Heninger C, Saito N, Tallman JF, Garrett KM, Vitek MP, Duman RS, and Gallager DW (1990) Effects of continuous diazepam administration on GABA_A subunit mRNA in rat brain. *J Mol Neurosci* **2**:101–107.
- Herzog AG (2009) Hormonal therapies: progesterone. *Neurotherapeutics* **6**:383–391.
- Hirouchi M, Hashimoto T, and Kuriyama K (1993) Alteration of GABA_A receptor $\alpha 1$ -subunit mRNA in mouse brain following continuous ethanol inhalation. *Eur J Pharmacol* **247**:127–130.
- Holt RA, Bateson AN, and Martin IL (1996) Chronic treatment with diazepam or abecarnil differentially affects the expression of GABA_A receptor subunit mRNAs in the rat cortex. *Neuropharmacology* **35**:1457–1463.
- Holt RA, Bateson AN, and Martin IL (1997a) Chronic zolpidem treatment alters GABA_A receptor mRNA levels in the rat cortex. *Eur J Pharmacol* **329**:129–132.

- Holt RA, Bateson AN, and Martin IL (1999) Decreased GABA enhancement of benzodiazepine binding after a single dose of diazepam. *J Neurochem* **72**:2219–2222.
- Holt RA, Martin IL, and Bateson AN (1997b) Chronic diazepam exposure decreases transcription of the rat GABA_A receptor γ 2-subunit gene. *Mol Brain Res* **48**:164–166.
- Hsu FC, Waldeck R, Faber DS, and Smith SS (2003) Neurosteroid effects on GABAergic synaptic plasticity in hippocampus. *J Neurophysiol* **89**:1929–1940.
- Houston CM, He Q, and Smart TG (2009) CaMKII phosphorylation of the GABA_A receptor: receptor subtype- and synapse-specific modulation. *J Physiol* **587**:2115–2125.
- Hu Y, Lund IV, Gravielle MC, Farb DH, Brooks-Kayal AR, and Russek SJ (2008) Surface expression of GABA_A receptors is transcriptionally controlled by the interplay of cAMP-response element-binding protein and its binding partner inducible cAMP early repressor. *J Biol Chem* **283**:9328–9340.
- Huntsman MM, Isackson PJ, and Jones EG (1994) Lamina-specific expression and activity-dependent regulation of seven GABA_A receptor subunit mRNAs in monkey visual cortex. *J Neurosci* **14**:2236–2259.
- Huopaniemi L, Keist R, Randolph A, Certa U, and Rudolph U (2004) Diazepam-induced adaptive plasticity revealed by α 1 GABA_A receptor-specific expression profiling. *J Neurochem* **88**:1059–1067.
- Hutchinson MA, Smith PF, and Darlington CL (1996) The behavioural and neuronal effects of the chronic administration of benzodiazepine anxiolytic and hypnotic drugs. *Prog Neurobiol* **49**:73–97.
- Impagnatiello F, Pesold C, Longone P, Caruncho H, Fritschy JM, Costa E, and Guidotti A (1996) Modifications of γ -aminobutyric acid_A receptor subunit expression in rat neocortex during tolerance to diazepam. *Mol Pharmacol* **49**:822–831.
- Ito T, Suzuki T, Wellman SE, and Ho IK (1996a) Chronic pentobarbital administration alters γ -aminobutyric acid_A receptor α 6-subunit mRNA levels and diazepam-insensitive [³H]Ro15-4513 binding. *Synapse* **22**:106–113.
- Ito T, Suzuki T, Wellman SE, and Ho IK (1996b) Pharmacology of barbiturate tolerance/dependence: GABA_A receptors and molecular aspects. *Life Sci* **59**:169–195.
- Izzo E, Auta J, Impagnatiello F, Pesold C, Guidotti A, and Costa E (2001) Glutamic acid decarboxylase and glutamate receptor changes during tolerance and dependence to benzodiazepines. *Proc Natl Acad Sci U S A* **98**:3483–3488.
- Jacob TC, Bogdanov YD, Magnus C, Saliba RS, Kittler JT, Haydon PG, and Moss SJ (2005) Gephyrin regulates the cell surface dynamics of synaptic GABA_A receptors. *J Neurosci* **25**:10469–10478.
- Jacob TC, Moss SJ, and Jurd R (2008) GABA_A receptor trafficking and its role in the dynamic modulation of neuronal inhibition. *Nat Rev Neurosci* **9**:331–343.
- Jennings AS, Lewis RT, Russell MG, Hallett DJ, Street LJ, Castro JL, Atack JR, Cook SM, Lincoln R, Stanley J, et al. (2006) Imidazo[1,2-b][1,2,4]triazines as α 2/ α 3 subtype selective GABA A agonists for the treatment of anxiety. *Bioorg Med Chem Lett* **16**:1477–1480.
- Johnston JD and Bristow DR (1998) Regulation of GABA_A receptor α 1 protein is a sensitive indicator of benzodiazepine agonist efficacy. *Eur J Pharmacol* **348**:321–324.
- Jones A, Korpi ER, McKernan RM, Pelz R, Nusser Z, Mäkelä R, Mellor JR, Pollard S, Bahn S, Stephenson FA, et al. (1997) Ligand-gated ion channel subunit partnerships: GABA_A receptor α 6 subunit gene inactivation inhibits δ subunit expression. *J Neurosci* **17**:1350–1362.
- Kahle KT, Staley KJ, Nahed BV, Gamba G, Hebert SC, Lifton RP, and Mount DB (2008) Roles of the cation-chloride cotransporters in neurological disease. *Nat Clin Pract Neurol* **4**:490–503.
- Kamphuis W, De Rijk TC, and Lopes da Silva FH (1995) Expression of GABA_A receptor subunit mRNAs in hippocampal pyramidal and granular neurons in the kindling model of epileptogenesis: an in situ hybridization study. *Mol Brain Res* **31**:33–47.
- Kang I, Lindquist DG, Kinane TB, Ercolani L, Pritchard GA, and Miller LG (1994) Isolation and characterization of the promoter of the human GABA_A receptor α 1 subunit gene. *J Neurochem* **62**:1643–1646.
- Kang I and Miller LG (1991) Decreased GABA_A receptor subunit mRNA concentrations following chronic lorazepam administration. *Br J Pharmacol* **103**:1285–1287.
- Katsura M, Shibasaki M, Kurokawa K, Tsujimura A, and Ohkuma S (2007) Up-regulation of L-type high voltage-gated calcium channel subunits by sustained exposure to 1,4- and 1,5-benzodiazepines in cerebrotical neurons. *J Neurochem* **103**:2518–2528.
- Kittler JT, Rostaing P, Schiavo G, Fritschy JM, Olsen R, Triller A, and Moss SJ (2001) The subcellular distribution of GABARAP and its ability to interact with NSF suggest a role for this protein in the intracellular transport of GABA_A receptors. *Mol Cell Neurosci* **18**:13–25.
- Knabl J, Witschi R, Hösl K, Reinold H, Zeilhofer UB, Ahmadi S, Brockhaus J, Sergejeva M, Hess A, Brune K, et al. (2008) Reversal of pathological pain through specific spinal GABA_A receptor subtypes. *Nature* **451**:330–334.
- Kneussel M and Loebrich S (2007) Trafficking and synaptic anchoring of ionotropic inhibitory neurotransmitter receptors. *Biol Cell* **99**:297–309.
- Kobayashi T, Ikeda K, Kojima H, Niki H, Yano R, Yoshioka T, and Kumanishi T (1999) Ethanol opens G-protein-activated inwardly rectifying K⁺ channels. *Nat Neurosci* **2**:1091–1097.
- Koksma JJ, van Kesteren RE, Rosahl TW, Zwart R, Smit AB, Lüddens H, and Brussaard AB (2003) Oxytocin regulates neurosteroid modulation of GABA_A receptors in supraoptic nucleus around parturition. *J Neurosci* **23**:788–797.
- Korpi ER, Debus F, Linden AM, Malécot C, Leppä E, Vekovischeva O, Rabe H, Böhme I, Aller MI, Wisden W, and Lüddens H (2007) Does ethanol act preferentially via selected brain GABA_A receptor subtypes? The current evidence is ambiguous. *Alcohol* **41**:163–176.
- Korpi ER, Gründer G, and Lüddens H (2002) Drug interactions at GABA_A receptors. *Prog Neurobiol* **67**:113–159.
- Korpi ER and Sinkkonen ST (2006) GABA_A receptor subtypes as targets for neuropsychiatric drug development. *Pharmacol Ther* **109**:12–32.
- Krystal AD, Walsh JK, Laska E, Caron J, Amato DA, Wessel TC, and Roth T (2003) Sustained efficacy of eszopiclone over 6 months of nightly treatment: results of a randomized, double-blind, placebo-controlled study in adults with chronic insomnia. *Sleep* **26**:793–799.
- Kuriyama K, Tomono S, Kishi M, Mukainaka T, and Ohkuma S (1987) Development of γ -aminobutyric acid (GABA)ergic neurons in cerebral cortical neurons in primary culture. *Brain Res* **416**:7–21.
- Lambert JJ, Belelli D, Peden DR, Vardy AW, and Peters JA (2003) Neurosteroid modulation of GABA_A receptors. *Prog Neurobiol* **71**:67–80.
- Lambert JJ, Cooper MA, Simmons RD, Weir CJ, and Belelli D (2009) Neurosteroids: Endogenous allosteric modulators of GABA_A receptors. *Psychoneuroendocrinology* doi: 10.1016/j.psyneuen.2009.08.009.
- Laurie DJ, Seeburg PH, and Wisden W (1992a) The distribution of 13 GABA_A receptor subunit mRNAs in the rat brain. II. Olfactory bulb and cerebellum. *J Neurosci* **12**:1063–1076.
- Laurie DJ, Wisden W, and Seeburg PH (1992b) The distribution of thirteen GABA_A receptor subunit mRNAs in the rat brain. III. Embryonic and postnatal development. *J Neurosci* **12**:4151–4172.
- Lewis DA, Hashimoto T, and Volk DW (2005) Cortical inhibitory neurons and schizophrenia. *Nat Rev Neurosci* **6**:312–324.
- Liang J, Suryanarayanan A, Abriam A, Snyder B, Olsen RW, and Spigelman I (2007) Mechanisms of reversible GABA_A receptor plasticity after ethanol intoxication. *J Neurosci* **27**:12367–12377.
- Liang J, Zhang N, Cagetti E, Houser CR, Olsen RW, and Spigelman I (2006) Chronic intermittent ethanol-induced switch of ethanol actions from extrasynaptic to synaptic hippocampal GABA_A receptors. *J Neurosci* **26**:1749–1758.
- Licata SC and Rowlett JK (2008) Abuse and dependence liability of benzodiazepine-type drugs: GABA_A receptor modulation and beyond. *Pharmacol Biochem Behav* **90**:74–89.
- Lin LH and Wang LH (1996) Region-specific changes in GABA_A receptor δ subunit mRNA level by tolerance to and withdrawal from pentobarbital. *Neurosci Lett* **202**:149–152.
- Lippa A, Czobor P, Stark J, Beer B, Kostakis E, Gravielle M, Bandyopadhyay S, Russek SJ, Gibbs TT, Farb DH, et al. (2005) Selective anxiolysis produced by ocinaplon, a GABA_A receptor modulator. *Proc Natl Acad Sci U S A* **102**:7380–7385.
- Loebrich S, Bähring R, Katsuno T, Tsukita S, and Kneussel M (2006) Activated radixin is essential for GABA_A receptor α 5 subunit anchoring at the actin cytoskeleton. *EMBO J* **25**:987–999.
- Longone P, Impagnatiello F, Guidotti A, and Costa E (1996) Reversible modification of GABA_A receptor subunit mRNA expression during tolerance to diazepam-induced cognition dysfunction. *Neuropharmacology* **35**:1465–1473.
- Lonze BE and Ginty DD (2002) Function and regulation of CREB family transcription factors in the nervous system. *Neuron* **35**:605–623.
- Loup F, Wieser HG, Yonekawa Y, Aguzzi A, and Fritschy JM (2000) Selective alterations in GABA_A receptor subtypes in human temporal lobe epilepsy. *J Neurosci* **20**:5401–5419.
- Lovinger DM (1999) 5-HT₃ receptors and the neural actions of alcohols: an increasingly exciting topic. *Neurochem Int* **35**:125–130.
- Lovinger DM and Homanics GE (2007) Tonic for what ails us? high-affinity GABA_A receptors and alcohol. *Alcohol* **41**:139–143.
- Löw K, Crestani F, Keist R, Benke D, Brünig I, Benson JA, Fritschy JM, Rüllicke T, Bluethmann H, Möhler H, et al. (2000) Molecular and neuronal substrate for the selective attenuation of anxiety. *Science* **290**:131–134.
- Lüddens H, Korpi ER, and Seeburg PH (1995) GABA_A/benzodiazepine receptor heterogeneity: neurophysiological implications. *Neuropharmacology* **34**:245–254.
- Lüscher B and Keller CA (2004) Regulation of GABA_A receptor trafficking, channel activity, and functional plasticity of inhibitory synapses. *Pharmacol Ther* **102**:195–221.
- Lund IV, Hu Y, Raol YH, Benham RS, Faris R, Russek SJ, and Brooks-Kayal AR (2008) BDNF selectively regulates GABA_A receptor transcription by activation of the JAK/STAT pathway. *Sci Signal* **1**:ra9.
- Lyons HR, Gibbs TT, and Farb DH (2000) Turnover and down-regulation of GABA_A receptor α 1, β 2S, and γ 1 subunit mRNAs by neurons in culture. *J Neurochem* **74**:1041–1048.
- Lyons HR, Land MB, Gibbs TT, and Farb DH (2001) Distinct signal transduction pathways for GABA-induced GABA_A receptor down-regulation and uncoupling in neuronal culture: a role for voltage-gated calcium channels. *J Neurochem* **78**:1114–1126.
- Maguire JL, Stell BM, Rafizadeh M, and Mody I (2005) Ovarian cycle-linked changes in GABA_A receptors mediating tonic inhibition alter seizure susceptibility and anxiety. *Nat Neurosci* **8**:797–804.
- Maguire J and Mody I (2007) Neurosteroid synthesis-mediated regulation of GABA_A receptors: relevance to the ovarian cycle and stress. *J Neurosci* **27**:2155–2162.
- Mahmoudi M, Kang MH, Tillakaratne N, Tobin AJ, and Olsen RW (1997) Chronic intermittent ethanol treatment in rats increases GABA_A receptor α 4-subunit expression: possible relevance to alcohol dependence. *J Neurochem* **68**:2485–2492.
- Majewska MD (1992) Neurosteroids: endogenous bimodal modulators of the GABA_A receptor. Mechanism of action and physiological significance. *Prog Neurobiol* **38**:379–395.
- Maric D, Maric I, Wen X, Fritschy JM, Sieghart W, Barker JL, and Serafini R (1999) GABA_A receptor subunit composition and functional properties of Cl⁻ channels with differential sensitivity to zolpidem in embryonic rat hippocampal cells. *J Neurosci* **19**:4921–4937.
- Marley RJ and Gallager DW (1989) Chronic diazepam treatment produces regionally specific changes in GABA-stimulated chloride influx. *Eur J Pharmacol* **159**:217–223.
- Marutha Ravindran CR and Ticku MK (2006) Tyrosine kinase phosphorylation of GABA_A receptor subunits following chronic ethanol exposure of cultured cortical neurons of mice. *Brain Res* **1086**:35–41.
- Marutha Ravindran CR, Mehta AK, and Ticku MK (2007a) Effect of chronic admin-

- istration of ethanol on the regulation of tyrosine kinase phosphorylation of the GABA_A receptor subunits in the rat brain. *Neurochem Res* **32**:1179–1187.
- Marutha Ravindran CR, Mehta AK, and Ticku MK (2007b) Effect of chronic administration of ethanol on the regulation of the δ -subunit of GABA_A receptors in the rat brain. *Brain Res* **1174**:47–52.
- Matthews DB, Devaud LL, Fritschy JM, Sieghart W, and Morrow AL (1998) Differential regulation of GABA_A receptor gene expression by ethanol in the rat hippocampus versus cerebral cortex. *J Neurochem* **70**:1160–1166.
- McKernan RM and Whiting PJ (1996) Which GABA_A-receptor subtypes really occur in the brain? *Trends Neurosci* **19**:139–143.
- Mehta AK and Ticku MK (1992) Chronic GABA exposure down-regulates GABA-benzodiazepine receptor-ionophore complex in cultured cerebral cortical neurons. *Mol Brain Res* **16**:29–36.
- Messer A (1977) The maintenance and identification of mouse cerebellar granule cells in monolayer culture. *Brain Res* **130**:1–12.
- Mhatre MC, Pena G, Sieghart W, and Ticku MK (1993) Antibodies specific for GABA_A receptor α subunits reveal that chronic alcohol treatment down-regulates α -subunit expression in rat brain regions. *J Neurochem* **61**:1620–1625.
- Mhatre MC and Ticku MK (1992) Chronic ethanol administration alters γ -aminobutyric acid_A receptor gene expression. *Mol Pharmacol* **42**:415–422.
- Mhatre M and Ticku MK (1994) Chronic ethanol treatment upregulates the GABA receptor β subunit expression. *Mol Brain Res* **23**:246–252.
- Miller LG, Greenblatt DJ, Barnhill JG, and Shader RI (1988) Chronic benzodiazepine administration. I. Tolerance is associated with benzodiazepine receptor downregulation and decreased γ -aminobutyric acid_A receptor function. *J Pharmacol Exp Ther* **246**:170–176.
- Miller LG, Roy RB, and Weill CL (1989) Chronic clonazepam administration decreases γ -aminobutyric acid_A receptor function in cultured cortical neurons. *Mol Pharmacol* **36**:796–802.
- Miralles CP, Li M, Mehta AK, Khan ZU, and De Blas AL (1999) Immunocytochemical localization of the β 3 subunit of the γ -aminobutyric acid_A receptor in the rat brain. *J Comp Neurol* **413**:535–548.
- Miranda JD and Barnes EM Jr (1997) Repression of γ -aminobutyric acid type A receptor α 1 polypeptide biosynthesis requires chronic agonist exposure. *J Biol Chem* **272**:16288–16294.
- Miyaoka T, Kimura T, Saunders PA, Tseng YT, and Ho IK (1994) Binding characteristics of [³H]flunitrazepam in pentobarbital-withdrawal rats. *Neurochem Res* **19**:37–42.
- Mizoguchi Y, Kanematsu T, Hirata M, and Nabekura J (2003) A rapid increase in the total number of cell surface functional GABA_A receptors induced by brain-derived neurotrophic factor in rat visual cortex. *J Biol Chem* **278**:44097–44102.
- Montpied P, Ginns EI, Martin BM, Roca D, Farb DH, and Paul SM (1991a) γ -Aminobutyric acid (GABA) induces a receptor-mediated reduction in GABA_A receptor α subunit messenger RNAs in embryonic chick neurons in culture. *J Biol Chem* **266**:6011–6014.
- Montpied P, Morrow AL, Karanian JW, Ginns EI, Martin BM, and Paul SM (1991b) Prolonged ethanol inhalation decreases γ -aminobutyric acid_A receptor α subunit mRNAs in the rat cerebral cortex. *Mol Pharmacol* **39**:157–163.
- Moragues N, Ciofi P, Tramu G, and Garret M (2002) Localisation of GABA_A receptor ϵ -subunit in cholinergic and aminergic neurones and evidence for co-distribution with the θ -subunit in rat brain. *Neuroscience* **111**:657–669.
- Moreno JI, Piva MA, Miralles CP, and De Blas AL (1994) Immunocytochemical localization of the β 2 subunit of the γ -aminobutyric acid_A receptor in the rat brain. *J Comp Neurol* **350**:260–271.
- Morrow AL, Herbert JS, and Montpied P (1992) Differential effects of chronic ethanol administration on GABA_A receptor α 1 and α 6 subunit mRNA levels in rat cerebellum. *Mol Cell Neurosci* **3**:251–258.
- Morrow AL, Montpied P, Lingford-Hughes A, and Paul SM (1990) Chronic ethanol and pentobarbital administration in the rat: effects on GABA_A receptor function and expression in brain. *Alcohol* **7**:237–244.
- Morrow AL, Montpied P, and Paul SM (1991) GABA_A receptor function and expression following chronic ethanol and barbiturate administration. *Ann N Y Acad Sci* **625**:496–507.
- Nagy J (2008) Alcohol related changes in regulation of NMDA receptor functions. *Curr Neuropharmacol* **6**:39–54.
- O'Donovan MC, Buckland PR, and McGuffin P (1992a) Levels of GABA_A receptor subunit mRNA in rat brain following flurazepam treatment. *J Psychopharmacol* **6**:364–369.
- O'Donovan MC, Buckland PR, Spurlock G, and McGuffin P (1992b) Bi-directional changes in the levels of messenger RNAs encoding γ -aminobutyric acid_A receptor α subunits after flurazepam treatment. *Eur J Pharmacol* **226**:335–341.
- Ogris W, Pörtl A, Hauer B, Ernst M, Oberto A, Wulff P, Höger H, Wisden W, and Sieghart W (2004) Affinity of various benzodiazepine site ligands in mice with a point mutation in the GABA_A receptor γ 2 subunit. *Biochem Pharmacol* **68**:1621–1629.
- Olsen RW and Sieghart W (2008) International Union of Pharmacology. LXX. Subtypes of γ -aminobutyric acid_A receptors: classification on the basis of subunit composition, pharmacology, and function. Update. *Pharmacol Rev* **60**:243–260.
- Papadeas S, Grobin AC, and Morrow AL (2001) Chronic ethanol consumption differentially alters GABA_A receptor α 1 and α 4 subunit peptide expression and GABA_A receptor-mediated ³⁶Cl⁻ uptake in mesocorticolimbic regions of rat brain. *Alcohol Clin Exp Res* **25**:1270–1275.
- Peng Z, Hauer B, Mihalek RM, Homanics GE, Sieghart W, Olsen RW, and Houser CR (2002) GABA_A receptor changes in δ subunit-deficient mice: altered expression of α 4 and γ 2 subunits in the forebrain. *J Comp Neurol* **446**:179–197.
- Persohn E, Malherbe P, and Richards JG (1992) Comparative molecular neuroanatomy of cloned GABA_A receptor subunits in the rat CNS. *J Comp Neurol* **326**:193–216.
- Pesold C, Caruncho HJ, Impagnatiello F, Berg MJ, Fritschy JM, Guidotti A, and Costa E (1997) Tolerance to diazepam and changes in GABA_A receptor subunit expression in rat neocortical areas. *Neuroscience* **79**:477–487.
- Petrie J, Sapp DW, Tyndale RF, Park MK, Fanselow M, and Olsen RW (2001) Altered GABA_A receptor subunit and splice variant expression in rats treated with chronic intermittent ethanol. *Alcohol Clin Exp Res* **25**:819–828.
- Pétursson H (1994) The benzodiazepine withdrawal syndrome. *Addiction* **89**:1455–1459.
- Pignataro L, Miller AN, Ma L, Midha S, Protiva P, Herrera DG, and Harrison NL (2007) Alcohol regulates gene expression in neurons via activation of heat shock factor 1. *J Neurosci* **27**:12957–12966.
- Pirker S, Schwarzer C, Wieselthaler A, Sieghart W, and Sperk G (2000) GABA_A receptors: immunocytochemical distribution of 13 subunits in the adult rat brain. *Neuroscience* **101**:815–850.
- Primus RJ and Gallager DW (1992) GABA_A receptor subunit mRNA levels are differentially influenced by chronic FG 7142 and diazepam exposure. *Eur J Pharmacol* **226**:21–28.
- Primus RJ, Yu J, Xu J, Hartnett C, Meyyappan M, Kostas C, Ramabhadran TV, and Gallager DW (1996) Allosteric uncoupling after chronic benzodiazepine exposure of recombinant γ -aminobutyric acid_A receptors expressed in Sf9 cells: ligand efficacy and subtype selectivity. *J Pharmacol Exp Ther* **276**:882–890.
- Prior P, Schmitt B, Grenningloh G, Pribilla I, Multhaup G, Beyreuther K, Maulet Y, Werner P, Langosch D, and Kirsch J (1992) Primary structure and alternative splice variants of gephyrin, a putative glycine receptor-tubulin linker protein. *Neuron* **8**:1161–1170.
- Rabow LE, Russek SJ, and Farb DH (1995) From ion currents to genomic analysis: recent advances in GABA_A receptor research. *Synapse* **21**:189–274.
- Ramsey-Williams VA and Carter DB (1996) Chronic triazolam and its withdrawal alters GABA_A receptor subunit mRNA levels: an in situ hybridization study. *Mol Brain Res* **43**:132–140.
- Ramsey-Williams VA, Wu Y, and Rosenberg HC (1994) Comparison of anticonvulsant tolerance, cross-tolerance, and benzodiazepine receptor binding following chronic treatment with diazepam or midazolam. *Pharmacol Biochem Behav* **48**:765–772.
- Ranna M, Sinkkonen ST, Möykkynen T, Uusi-Oukari M, and Korpi ER (2006) Impact of ϵ and θ subunits on pharmacological properties of α 3 β 1 GABA_A receptors expressed in *Xenopus* oocytes. *BMC Pharmacol* **6**:1.
- Ravindran CR and Ticku MK (2006) Tyrosine kinase phosphorylation of GABA_A receptor α 1, β 2 and γ 2 subunits following chronic intermittent ethanol (CIE) exposure of cultured cortical neurons of mice. *Neurochem Res* **31**:1111–1118.
- Roca DJ, Schiller GD, Friedman L, Rozenberg I, Gibbs TT, and Farb DH (1990) γ -Aminobutyric acid_A receptor regulation in culture: altered allosteric interactions following prolonged exposure to benzodiazepines, barbiturates, and methylxanthines. *Mol Pharmacol* **37**:710–719.
- Rogawski MA and Reddy DS (2002) Neurosteroids and infantile spasms: the deoxycorticosterone hypothesis. *Int Rev Neurobiol* **49**:199–219.
- Ron D (2004) Signaling cascades regulating NMDA receptor sensitivity to ethanol. *Neuroscientist* **10**:325–336.
- Rosenberg HC and Chiu TH (1979) Decreased ³H-diazepam binding is a specific response to chronic benzodiazepine treatment. *Life Sci* **24**:803–807.
- Roth T, Walsh JK, Krystal A, Wessel T, and Roehrs TA (2005) An evaluation of the efficacy and safety of eszopiclone over 12 months in patients with chronic primary insomnia. *Sleep Med* **6**:487–495.
- Rothman S and Cowan WM (1981) A scanning electron microscope study of the in vitro development of dissociated hippocampal cells. *J Comp Neurol* **195**:141–155.
- Rudolph U, Crestani F, Benke D, Brünig I, Benson JA, Fritschy JM, Martin JR, Bluethmann H, and Möhler H (1999) Benzodiazepine actions mediated by specific γ -aminobutyric acid(A) receptor subtypes. *Nature* **401**:796–800.
- Rudolph U and Möhler H (2004) Analysis of GABA_A receptor function and dissection of the pharmacology of benzodiazepines and general anesthetics through mouse genetics. *Annu Rev Pharmacol Toxicol* **44**:475–498.
- Russek SJ (1999) Evolution of GABA_A receptor diversity in the human genome. *Gene* **227**:213–222.
- Russek SJ and Farb DH (1994) Mapping of the β 2 subunit gene (GABRB2) to microdissected human chromosome 5q34–q35 defines a gene cluster for the most abundant GABA_A receptor isoform. *Genomics* **23**:528–533.
- Salonen V, Kallinen S, Lopez-Picon FR, Korpi ER, Holopainen IE, and Uusi-Oukari M (2006) AMPA/kainate receptor-mediated up-regulation of GABA_A receptor δ subunit mRNA expression in cultured rat cerebellar granule cells is dependent on NMDA receptor activation. *Brain Res* **1087**:33–40.
- Sanna E, Busonero F, Talani G, Mostallino MC, Mura ML, Pisu MG, Maciocco E, Serra M, and Biggio G (2005) Low tolerance and dependence liabilities of etizolam: molecular, functional, and pharmacological correlates. *Eur J Pharmacol* **519**:31–42.
- Sanna E, Mostallino MC, Busonero F, Talani G, Tranquilli S, Mamei M, Spiga S, Follasa P, and Biggio G (2003) Changes in GABA_A receptor gene expression associated with selective alterations in receptor function and pharmacology after ethanol withdrawal. *J Neurosci* **23**:11711–11724.
- Sarviharju M, Hyytiä P, Hervonen A, Jaatinen P, Kiianmaa K, and Korpi ER (2006) Lifelong ethanol consumption and brain regional GABA_A receptor subunit mRNA expression in alcohol-preferring rats. *Alcohol* **40**:159–166.
- Savill RM, Scotting PJ, and Coyle B (2005) Strategies to investigate gene expression and function in granule cells. *Cerebellum* **4**:271–278.
- Schumacher M, Weill-Engerer S, Liere P, Robert F, Franklin RJ, Garcia-Segura LM, Lambert JJ, Mayo W, Melcangi RC, Parducz A, et al. (2003) Steroid hormones and neurosteroids in normal and pathological aging of the nervous system. *Prog Neurobiol* **71**:3–29.
- Schwarzer C, Berresheim U, Pirker S, Wieselthaler A, Fuchs K, Sieghart W, and Sperk G (2001) Distribution of the major γ -aminobutyric acid_A receptor subunits in the basal ganglia and associated limbic brain areas of the adult rat. *J Comp Neurol* **433**:526–549.
- Shader RI and Greenblatt DJ (1993) Use of benzodiazepines in anxiety disorders. *N Engl J Med* **328**:1398–1405.
- Sheela Rani CS and Ticku MK (2006) Comparison of chronic ethanol and chronic

- intermittent ethanol treatments on the expression of GABA_A and NMDA receptor subunits. *Alcohol* **38**:89–97.
- Shen H, Gong QH, Yuan M, and Smith SS (2005) Short-term steroid treatment increases δ GABA_A receptor subunit expression in rat CA1 hippocampus: pharmacological and behavioral effects. *Neuropharmacology* **49**:573–586.
- Sieghart W (1995) Structure and pharmacology of γ -aminobutyric acid_A receptor subtypes. *Pharmacol Rev* **47**:181–234.
- Sieghart W and Sperk G (2002) Subunit composition, distribution and function of GABA_A receptors. *Curr Top Med Chem* **2**:795–816.
- Sigel E and Buhr A (1997) The benzodiazepine binding site of GABA_A receptors. *Trends Pharmacol Sci* **18**:425–429.
- Sinkkonen ST, Hanna MC, Kirkness EF, and Korpi ER (2000) GABA_A receptor ϵ and θ subunits display unusual structural variation between species and are enriched in the rat locus ceruleus. *J Neurosci* **20**:3588–3595.
- Smith MC and Riskin BJ (1991) The clinical use of barbiturates in neurological disorders. *Drugs* **42**:365–378.
- Smith SS, Gong QH, Hsu FC, Markowitz RS, French-Mullen JM, and Li X (1998) GABA_A receptor $\alpha 4$ subunit suppression prevents withdrawal properties of an endogenous steroid. *Nature* **392**:926–930.
- Song J, Shen G, Greenfield LJ Jr., and Tietz EI (2007) Benzodiazepine withdrawal-induced glutamatergic plasticity involves up-regulation of GluR1-containing α -amino-3-hydroxy-5-methylisoxazole-4-propionic acid receptors in hippocampal CA1 neurons. *J Pharmacol Exp Ther* **322**:569–581.
- Spigelman I, Li Z, Liang J, Cagetti E, Samzadeh S, Mihalek RM, Homanics GE, and Olsen RW (2003) Reduced inhibition and sensitivity to neurosteroids in hippocampus of mice lacking the GABA_A receptor δ subunit. *J Neurophysiol* **90**:903–910.
- Steiger JL and Russek SJ (2004) GABA_A receptors: building the bridge between subunit mRNAs, their promoters, and cognate transcription factors. *Pharmacol Ther* **101**:259–281.
- Stephens DN (1995) A glutamatergic hypothesis of drug dependence: extrapolations from benzodiazepine receptor ligands. *Behav Pharmacol* **6**:425–446.
- Sternbach LH (1978) The benzodiazepine story. *Prog Drug Res* **22**:229–266.
- Straub CJ, Carlezon WA Jr., and Rudolph U (2010) Diazepam and cocaine potentiate brain stimulation reward in C57BL/6J mice. *Behav Brain Res* **206**:17–20.
- Sundstrom-Poromaa I, Smith DH, Gong QH, Sabado TN, Li X, Light A, Wiedmann M, Williams K, and Smith SS (2002) Hormonally regulated $\alpha 4\beta 2\delta$ GABA_A receptors are a target for alcohol. *Nat Neurosci* **5**:721–722.
- Sur C, Quirk K, Dewar D, Atack J, and McKernan R (1998) Rat and human hippocampal $\alpha 5$ subunit-containing γ -aminobutyric acid_A receptors have $\alpha 5\beta 2\gamma 2$ pharmacological characteristics. *Mol Pharmacol* **54**:928–933.
- Suzdak PD, Glowa JR, Crawley JN, Schwartz RD, Skolnick P, and Paul SM (1986) A selective imidazobenzodiazepine antagonist of ethanol in the rat. *Science* **234**:1243–1247.
- Taliani S, Cosimelli B, Da Settimo F, Marini AM, La Motta C, Simorini F, Salerno S, Novellino E, Greco G, Cosconati S, et al. (2009) Identification of anxiolytic/nosedative agents among indol-3-ylglyoxylamides acting as functionally selective agonists at the γ -aminobutyric acid_A (GABA_A) $\alpha 2$ benzodiazepine receptor. *J Med Chem* **52**:3723–3734.
- Tanay VM, Greenshaw AJ, Baker GB, and Bateson AN (2001) Common effects of chronically administered antipanic drugs on brainstem GABA_A receptor subunit gene expression. *Mol Psychiatry* **6**:404–412.
- Tehrani MH and Barnes EM Jr. (1997) Sequestration of γ -aminobutyric acid_A receptors on clathrin-coated vesicles during chronic benzodiazepine administration in vivo. *J Pharmacol Exp Ther* **283**:384–390.
- Thompson CL, Pollard S, and Stephenson FA (1996) Bidirectional regulation of GABA_A receptor $\alpha 1$ and $\alpha 6$ subunit expression by a cyclic AMP-mediated signaling mechanism in cerebellar granule cells in primary culture. *J Neurochem* **67**:434–437.
- Thompson CL, Razzini G, Pollard S, and Stephenson FA (2000) Cyclic AMP-mediated regulation of GABA_A receptor subunit expression in mature rat cerebellar granule cells: evidence for transcriptional and translational control. *J Neurochem* **74**:920–931.
- Tietz EI, Chiu TH, and Rosenberg HC (1989) Regional GABA/benzodiazepine receptor/chloride channel coupling after acute and chronic benzodiazepine treatment. *Eur J Pharmacol* **167**:57–65.
- Tietz EI, Huang X, Chen S, and Ferenak WF 3rd (1999a) Temporal and regional regulation of $\alpha 1$, $\beta 2$ and $\beta 3$, but not $\alpha 2$, $\alpha 4$, $\alpha 5$, $\alpha 6$, $\beta 1$ or $\gamma 2$ GABA_A receptor subunit messenger RNAs following one-week oral flurazepam administration. *Neuroscience* **91**:327–341.
- Tietz EI, Huang X, Weng X, Rosenberg HC, and Chiu TH (1993) Expression of $\alpha 1$, $\alpha 5$, and $\gamma 2$ GABA_A receptor subunit mRNAs measured in situ in rat hippocampus and cortex following chronic flurazepam administration. *J Mol Neurosci* **4**:277–292.
- Tietz EI, Zeng XJ, Chen S, Lilly SM, Rosenberg HC, and Kometiani P (1999b) Antagonist-induced reversal of functional and structural measures of hippocampal benzodiazepine tolerance. *J Pharmacol Exp Ther* **291**:932–942.
- Tretter V, Jacob TC, Mukherjee J, Fritschy JM, Pangalos MN, and Moss SJ (2008) The clustering of GABA_A receptor subtypes at inhibitory synapses is facilitated via the direct binding of receptor $\alpha 2$ subunits to gephyrin. *J Neurosci* **28**:1356–1365.
- Tseng YT, Wellman SE, and Ho IK (1993a) Differential effects on GABA_A receptor $\gamma 2$ -subunit messenger RNA by tolerance to and withdrawal from pentobarbital - an in situ hybridization study. *Life Sci* **53**:PL321–326.
- Tseng YT, Miyaoka T, and Ho IK (1993b) Region-specific changes of GABA_A receptors by tolerance to and dependence upon pentobarbital. *Eur J Pharmacol* **236**:23–30.
- Tseng YT, Wellman SE, and Ho IK (1994) In situ hybridization evidence of differential modulation by pentobarbital of GABA_A receptor $\alpha 1$ - and $\beta 3$ -subunit mRNAs. *J Neurochem* **63**:301–309.
- Tsuda M, Chiba Y, Suzuki T, and Misawa M (1998) Upregulation of NMDA receptor subunit proteins in the cerebral cortex during diazepam withdrawal. *Eur J Pharmacol* **341**:R1–R2.
- Ueno S, Lin A, Nikolaeva N, Trudell JR, Mihic SJ, Harris RA, and Harrison NL (2000) Tryptophan scanning mutagenesis in TM2 of the GABA_A receptor α subunit: effects on channel gating and regulation by ethanol. *Br J Pharmacol* **131**:296–302.
- Uusi-Oukari M, Heikkilä J, Sinkkonen ST, Mäkelä R, Hauer B, Homanics GE, Sieghart W, Wisden W, and Korpi ER (2000) Long-range interactions in neuronal gene expression: evidence from gene targeting in the GABA_A receptor $\beta 2$ - $\alpha 6$ - $\alpha 1$ - $\gamma 2$ subunit gene cluster. *Mol Cell Neurosci* **16**:34–41.
- Van Laere K, Bormans G, Sanabria-Bohórquez SM, de Groot T, Dupont P, De Lepeleire I, de Hoon J, Mortelmans L, Hargreaves RJ, Atack JR, et al. (2008) In vivo characterization and dynamic receptor occupancy imaging of TPA023B, an $\alpha 2/\alpha 3/\alpha 5$ subtype selective γ -aminobutyric acid_A partial agonist. *Biol Psychiatry* **64**:153–161.
- van Rijnsoever C, Täuber M, Choulli MK, Keist R, Rudolph U, Mohler H, Fritschy JM, and Crestani F (2004) Requirement of $\alpha 5$ -GABA_A receptors for the development of tolerance to the sedative action of diazepam in mice. *J Neurosci* **24**:6785–6789.
- Van Sickle BJ, Xiang K, and Tietz EI (2004) Transient plasticity of hippocampal CA1 neuron glutamate receptors contributes to benzodiazepine withdrawal-anxiety. *Neuropsychopharmacology* **29**:1994–2006.
- Varecka L, Wu CH, Rotter A, and Frostholm A (1994) GABA_A/benzodiazepine receptor $\alpha 6$ subunit mRNA in granule cells of the cerebellar cortex and cochlear nuclei: expression in developing and mutant mice. *J Comp Neurol* **339**:341–352.
- Vekovischeva OY, Uusi-Oukari M, and Korpi ER (2000) Chronic ethanol treatment and GABA_A receptor $\alpha 6$ subunit gene expression: a study using $\alpha 6$ subunit-deficient mice. *Addict Biol* **5**:463–467.
- Vengelien V, Bilbao A, Molander A, and Spanagel R (2008) Neuropharmacology of alcohol addiction. *Br J Pharmacol* **154**:299–315.
- Vithlani M and Moss SJ (2009) The role of GABA_AR phosphorylation in the construction of inhibitory synapses and the efficacy of neuronal inhibition. *Biochem Soc Trans* **37**:1355–1358.
- Vorma H, Naukkarinen H, Sarna S, and Kuoppasalmi K (2002) Treatment of outpatients with complicated benzodiazepine dependence: comparison of two approaches. *Addiction* **97**:851–859.
- Vorma H, Naukkarinen H, Sarna S, and Kuoppasalmi K (2004) Symptom severity and quality of life after benzodiazepine withdrawal treatment in participants with complicated dependence. *Addict Behav* **29**:1059–1065.
- Voss J, Sanchez C, Michelsen S, and Ebert B (2003) Rotarod studies in the rat of the GABA_A receptor agonist gaboxadol: lack of ethanol potentiation and benzodiazepine cross-tolerance. *Eur J Pharmacol* **482**:215–222.
- Wafford KA and Ebert B (2006) Gaboxadol—a new awakening in sleep. *Curr Opin Pharmacol* **6**:30–36.
- Wallner M, Hancher HJ, and Olsen RW (2003) Ethanol enhances $\alpha 4\beta 3\delta$ and $\alpha 6\beta 3\delta$ γ -aminobutyric acid type A receptors at low concentrations known to affect humans. *Proc Natl Acad Sci U S A* **100**:15218–15223.
- Wallner M, Hancher HJ, and Olsen RW (2006) Low-dose alcohol actions on $\alpha 4\beta 3\delta$ GABA_A receptors are reversed by the behavioral alcohol antagonist Ro15-4513. *Proc Natl Acad Sci U S A* **103**:8540–8545.
- Wang H, Bedford FK, Brandon NJ, Moss SJ, and Olsen RW (1999) GABA_A-receptor-associated protein links GABA_A receptors and the cytoskeleton. *Nature* **397**:69–72.
- Wang X, Wang G, Lemos JR, and Treisman SN (1994) Ethanol directly modulates gating of a dihydropyridine-sensitive Ca²⁺ channel in neurohypophysial terminals. *J Neurosci* **14**:5453–5460.
- Wegele K, Pasternack M, Hiltunen JO, Rivera C, Kaila K, Saarma M, and Reeben M (1998) Distribution of GABA receptor ρ subunit transcripts in the rat brain. *Eur J Neurosci* **10**:350–357.
- Werner DF, Swihart AR, Ferguson C, Lariviere WR, Harrison NL, and Homanics GE (2009) Alcohol-induced tolerance and physical dependence in mice with ethanol insensitive $\alpha 1$ GABA_A receptors. *Alcohol Clin Exp Res* **33**:289–299.
- West AE, Griffith EC, and Greenberg ME (2002) Regulation of transcription factors by neuronal activity. *Nat Rev Neurosci* **3**:921–931.
- Wieland HA, Lüddens H, and Seeburg PH (1992) A single histidine in GABA_A receptors is essential for benzodiazepine agonist binding. *J Biol Chem* **267**:1426–1429.
- Wilcox AS, Warrington JA, Gardiner K, Berger R, Whiting P, Altherr MR, Wasmuth JJ, Patterson D, and Sikela JM (1992) Human chromosomal localization of genes encoding the $\gamma 1$ and $\gamma 2$ subunits of the γ -aminobutyric acid receptor indicates that members of this gene family are often clustered in the genome. *Proc Natl Acad Sci U S A* **89**:5857–5861.
- Wisden W, Laurie DJ, Monyer H, and Seeburg PH (1992) The distribution of 13 GABA_A receptor subunit mRNAs in the rat brain. I. Telencephalon, diencephalon, mesencephalon. *J Neurosci* **12**:1040–1062.
- Wohlfarth KM, Bianchi MT, and Macdonald RL (2002) Enhanced neurosteroid potentiation of ternary GABA_A receptors containing the δ subunit. *J Neurosci* **22**:1541–1549.
- Wu CH, Frostholm A, De Blas AL, and Rotter A (1995) A Differential expression of GABA_A/benzodiazepine receptor subunit mRNAs and ligand binding sites in mouse cerebellar neurons following in vivo ethanol administration: an autoradiographic analysis. *J Neurochem* **65**:1229–1239.
- Wu Y, Rosenberg HC, Chiu TH, and Zhao TJ (1994) Subunit- and brain region-specific reduction of GABA_A receptor subunit mRNAs during chronic treatment of rats with diazepam. *J Mol Neurosci* **5**:105–120.
- Yavin Z and Yavin E (1980) Survival and maturation of cerebral neurons on poly-L-lysine surfaces in the absence of serum. *Dev Biol* **75**:454–459.
- Yee BK, Keist R, von Boehmer L, Studer R, Benke D, Hagenbuch N, Dong Y, Malenka RC, Fritschy JM, Bluethmann H, et al. (2005) A schizophrenia-related sensorimotor deficit links $\alpha 3$ -containing GABA_A receptors to a dopamine hyperfunction. *Proc Natl Acad Sci U S A* **102**:17154–17159.
- Yin HS and Lee YP (1998) Effects of pentobarbital on the expression of GABA_A receptor $\beta 1$ mRNA in the hippocampus: differential responses of CA1 and CA3. *Synapse* **29**:371–378.
- Yu AC, Hertz E, and Hertz L (1984) Alterations in uptake and release rates for

- GABA, glutamate, and glutamine during biochemical maturation of highly purified cultures of cerebral cortical neurons, a GABAergic preparation. *J Neurochem* **42**:951–960.
- Yu R, Follesa P, and Ticku MK (1996) Down-regulation of the GABA receptor subunits mRNA levels in mammalian cultured cortical neurons following chronic neurosteroid treatment. *Mol Brain Res* **41**:163–168.
- Xiang K, Earl DE, Davis KM, Giovannucci DR, Greenfield LJ Jr., and Tietz EI (2008) Chronic benzodiazepine administration potentiates high voltage-activated calcium currents in hippocampal CA1 neurons. *J Pharmacol Exp Ther* **327**:872–883.
- Xiang K and Tietz EI (2007) Benzodiazepine-induced hippocampal CA1 neuron α -amino-3-hydroxy-5-methylisoxasole-4-propionic acid (AMPA) receptor plasticity linked to severity of withdrawal anxiety: differential role of voltage-gated calcium channels and *N*-methyl-D-aspartic acid receptors. *Behav Pharmacol* **18**:447–460.
- Zhao TJ, Chiu TH, and Rosenberg HC (1994a) Decreased expression of γ -aminobutyric acid type A/benzodiazepine receptor β subunit mRNAs in brain of flurazepam-tolerant rats. *J Mol Neurosci* **5**:181–192.
- Zhao TJ, Chiu TH, and Rosenberg HC (1994b) Reduced expression of γ -aminobutyric acid type A/benzodiazepine receptor γ 2 and α 5 subunit mRNAs in brain regions of flurazepam-treated rats. *Mol Pharmacol* **45**:657–663.
- Zhou X and Smith SS (2009) Expression levels of the α 4 subunit of the GABA_A receptor in differentiated neuroblastoma cells are correlated with GABA-gated current. *Neuropharmacology* **56**:1041–1053.
- Zhou X and Smith SS (2007) Steroid requirements for regulation of the α 4 subunit of the GABA_A receptor in an in vitro model. *Neurosci Lett* **411**:61–66.