

Intestinal barrier, immunity and microbiome: partners in the depression crime

Eva M. Medina-Rodríguez^{1,2}, José Martínez-Raga¹, Yolanda Sanz².

¹ Psychiatry Service, Doctor Peset University Hospital. FISABIO, Valencia, Spain.

² Institute of Agrochemistry and Food Technology, Spanish National Research Council (IATA-CSIC), Valencia, Spain.

Details of financial support are included in the footnotes section at the end of the manuscript.

Running Title: Mechanisms controlling depression; potential interventions.

Corresponding author:

Eva M. Medina-Rodríguez, PhD

Psychiatry Service

Doctor Peset University Hospital, FISABIO

Av. Gaspar Aguilar, 90

46017, Valencia, Spain

phone: +34 963 900 022

e-mail: eva.medina@iata.csic.es

Number of text pages: 22

Number of figures: 2

Number of references: 132

Number of words in the Abstract: 179

Non-standard abbreviations: AhR: aryl hydrocarbon receptor; EGFR: epidermal growth factor receptor; EPA: eicosapentaenoic acid; fMRI: functional magnetic resonance imaging; FXR: farnesoid X receptor; GPR: G-protein-coupled receptor; HEPE: hydroxyeicosapentaenoic acid; HPA: hypothalamic-pituitary-adrenal; HPLC: high-performance liquid chromatography; ILC: innate lymphoid cell; MDD: major depressive disorder; PPAR: peroxisome proliferator-activated receptor; PUFA: polyunsaturated fatty acid; RCT: randomized controlled trial; ROR: RAR-related orphan receptor; ROS: reactive oxygen species; SCFA: short-chain fatty acid; SFB: Segmented filamentous bacteria; SPM: specialized pro-resolving lipid mediators; S1PR2: sphingosine 1 phosphate receptor 2; TAMP: tight junction-associated-MARVEL protein; TGR5: G-protein-coupled receptor 5; TNF: tumor necrosis factor; UHPLC-MS/MS: ultra-high performance liquid chromatography-tandem mass spectrometry.

ABSTRACT

Depression is a highly prevalent disorder and a leading cause of disability worldwide. It has a major impact on the affected individual and on society as a whole. Regrettably, current available treatments for this condition are insufficient in many patients. In recent years, the gut microbiome has emerged as a promising alternative target for treating and preventing depressive disorders. However, the microbes that form this ecosystem do not act alone, but are part of a complicated network connecting the gut and the brain that influences our mood. Host cells that are in intimate contact with gut microbes, such as the epithelial cells forming the gut barrier and the immune cells in their vicinity, play a key role in the process. These cells continuously shape immune responses to maintain healthy communication between gut microbes and the host. In this article, we review how the interplay between epithelial cells, the immune system, and gut microbes mediates gut-brain communication to influence mood. We also discuss how advances in our knowledge of the mechanisms underlying the gut-brain axis could contribute to addressing depression.

SIGNIFICANCE STATEMENT

This review does not aim to systematically describe intestinal microbes that might be beneficial or detrimental for depression. We have adopted a novel point of view by focusing on potential mechanisms underlying the crosstalk between gut microbes and their intestinal environment to control mood. These pathways could be targeted by well-defined and individually tailored dietary interventions, microbes, or microbial metabolites to ameliorate depression and decrease its important social and economic impact.

TABLE OF CONTENTS

1. Introduction
2. Structure and function of the gut epithelial barrier
3. Intestinal immune system
4. The gut microbiome and manipulation of the gut microbes and their metabolites as a strategy to treat depression
 - 4.1 Probiotic supplementation
 - 4.2 Engineered probiotics
 - 4.3 Fecal transplants
 - 4.4 Probiotic food
 - 4.5 Dietary changes
 - a) Short-chain fatty acids (SCFAs) and lactate
 - b) Secondary bile acids
 - c) Tryptophan metabolites
 - d) Vitamins
 - e) Polyphenols
 - f) Polyunsaturated fatty acids (PUFAs)
5. Diet-mediated pathways involving the microbiome, the host intestinal barrier, and the immune system as potential targets to treat depression
 - a) SCFAs and lactate-linked pathways
 - b) Secondary bile acids routes
 - c) Tryptophan-linked pathways
 - d) Vitamin-mediated pathways
 - e) Polyphenol-induced routes
 - f) PUFA-related pathways
 - g) Pathways induced by microbial components
6. Conclusion

1. Introduction

Depression is a highly prevalent and heterogeneous disorder characterized by low mood and loss of interest or pleasure in all or nearly all activities, in addition to symptoms such as changes in appetite or weight, sleep, and psychomotor activity; decreased energy; feelings of worthlessness or guilt; difficulty concentrating or making decisions; and suicidal thoughts or behaviors. More than 280 million people worldwide suffer from depressive disorders (Institute of Health Metrics and Evaluation. Global Health Data Exchange (GHDx)). Furthermore, depression is a leading cause of disability and premature death, making it a significant contributor to the global burden of disease with an elevated economic impact (Disease et al., 2018). Despite the ongoing development of new pharmacological and psychological therapies, there are many unmet needs in the treatment of depressive disorders. More than 50% of people treated with existing therapies show an inadequate response or fail to achieve remission (Ormel et al., 2019). Consequently, new therapeutic options are needed to reduce the economic burden and alleviate the suffering of patients, as depression is a debilitating and often fatal disease.

Microbes that populate our gut, also known as the gut microbiome, communicate with the central nervous system in a bidirectional manner and influence our mood (Margolis et al., 2021). For this reason, researchers are beginning to consider targeting or using these microbes as alternative therapeutic agents for mood disorders such as depression. Bacteria that exert a health benefit on the host are called probiotics. They can be obtained through supplements, naturally fermented food, or artificially enriched food. The specific name for probiotics that benefit mental health is “psychobiotics” (Dinan et al., 2013; Sarkar et al., 2016).

In recent years, an increasing number of studies have focused on the gut microbiome in depressed people to identify potential beneficial microbes. Interest in the microbiome’s influence on mental health has transcended the realm of the scientific

community to the wider public, becoming the subject of news stories, wellness forums, social networks, and advertising campaigns. However, most published studies on the gut microbiome in the context of depression are merely descriptive, reporting differences in depressed people compared with healthy controls (Alli et al., 2022). In addition, the great variability of the individual microbiome resulting from the many factors that influence its composition and function (e.g. microbes inherited at birth, diet, use of antibiotics or other medications) makes it very difficult to reach a consensus on a microbial signature associated with depression. Nonetheless, preclinical studies that have replicated depression-like behaviors in animals following the transfer of fecal microbiota from depressed people point to a causal role of the gut microbiota (Kelly et al., 2016; Knudsen et al., 2021). Researchers have confirmed this hypothesis by comparing the behavior of germ-free or microbiota-depleted rodents to that of rodents with normal gut colonization (Kunugi, 2021).

The effects of intestinal microbes depend, first, on communication with the host's intestinal epithelial cells and the surrounding environment, which can initiate the cascade of signals that eventually reach the central nervous system and influence mental health. Thus, the epithelial cell barrier, which is in close communication with the immune, endocrine, and nervous systems, serves as a first meeting point between the microbes and the host. Later, signals can be transmitted to the central nervous system by: 1) the physical gut-brain connection through the autonomic nervous system, the vagus nerve, and the enteric nervous system; 2) neuroendocrine mediators from the hypothalamic-pituitary-adrenal (HPA) axis; and 3) the immune system, whose immune cells or their immune mediators, such as cytokines, travel to the brain where they influence mood.

The third mechanism is one focus of this review. Researchers have used various approaches to investigate how gut microbes interact with the host epithelial and immune cells in the intestine to mediate depression, but our understanding of these mechanisms is far from complete.

In the present review, we summarize the current evidence on the interactions between intestinal microbes, epithelial cells, and the intestinal immune system, in the context of wellbeing and depression. We also discuss the gaps and future perspectives of microbiome-based approaches in the field of psychiatric medicine. This review does not provide a systematic description of intestinal microbes that might be beneficial for treating depression; rather, we have adopted a novel perspective by highlighting potential mechanisms of the crosstalk between gut microbes and their intestinal environment that have significant implications for brain function and behavior. These pathways could be targeted by dietary interventions, microbes, or microbial metabolites to ameliorate depression. Therefore, despite our incomplete understanding of the gut-immune-brain axis, this review provides insights into the potential translation of research from bench to bedside.

2. Structure and function of the gut epithelial barrier

Trillions of microorganisms, including bacteria, archaea, fungi, and viruses, can live in the digestive tracts of animals without upsetting their health. This implies a very well-organized physical, molecular, and immune defense that confines the microorganisms to the gut lumen and protects the host.

The first line of defense in the gut is a mucus layer composed of mucins (highly glycosylated proteins); it provides a crucial barrier between the microbiome and the host cells (Johansson et al., 2013). Lying beneath this mucus layer is the intestinal epithelium: a monolayer of specialized cells such as enterocytes or intraepithelial cells, Paneth cells, goblet cells, and microfold cells. Each cell type has a different function. The most abundant cells are intraepithelial cells or enterocytes, whose main role is to maintain epithelial barrier integrity while also producing cytokines, chemokines and immunological factors against luminal bacteria (Goto, 2019). Paneth cells are secretory cells residing in small clusters at the base of crypts of Lieberkühn. They have large

secretory granules containing antimicrobials, which are discharged into the crypt lumen and diffuse into the mucous layer overlying the mucosal epithelium, thus preventing microbial invasion (Bevins and Salzman, 2011). Goblet cells secrete the mucus that forms the mucus layer while also secreting antimicrobial proteins, chemokines, and cytokines (Knoop and Newberry, 2018). Microfold cells transfer microbial antigens from the lumen that are essential for the induction of efficient immune responses (Gutzeit et al., 2014). Therefore, the epithelial cells have immune functions beyond physical barrier maintenance.

The epithelial cells are linked to each other by junctional complexes consisting of tight junctions, adherens junctions, desmosomes, and gap junctions (Schoultz and Keita, 2020). Tight junctions, also called zonula occludens, are multiprotein complexes embedded in the plasma membrane that interact with the adjacent cells. They include the tight junction-associated-MARVEL proteins (TAMP), claudins, and members of the junctional adhesion molecule protein family. Adherens junctions are molecules pertaining to the cadherin family. Adherence junctions and tight junctions form the apical junctional complex. The desmosomes comprise the desmoglein and desmocollin families of desmosomal cadherins, as well as connecting proteins such as desmoglein and keratin. The gap junctions link the interior of adjacent cells (Schoultz and Keita, 2020). All these junctions maintain the barrier composed by the epithelial cells and prevent microbes leaking from the gut. Gut epithelial barrier integrity is crucial for maintaining a balanced bacteria-host interaction.

Underneath the epithelial monolayer is a thin layer of connective tissue called the lamina propria. Epithelial cells communicate with the microbiota and with immune cells of the lamina propria. This communication between the microbes, the epithelium, and the immune system of the host plays a crucial role in health and disease, including mental health and mental disorders such as depression.

3. Intestinal immune system

As commented above, the epithelial cells help to control the microbiome by secreting antimicrobial molecules. In addition, the intestinal mucosa is in close contact with cells of the innate and adaptative immune systems to control the microbes and trigger a suitable immune response when homeostasis is disrupted (Zheng et al., 2020). The interplay between the intestinal epithelium and microbiota maintains a basal state of inflammation, which prevents excess inflammatory responses under physiological conditions.

The immune cells in the gut are located in the intestinal lamina propria as well as interleaved in the epithelial layer. The gut contains cells of the innate immune system, such as macrophages, dendritic cells, and innate lymphoid cells (ILCs); and of the adaptative immune system, such as T and B cells (Yuan and Walker, 2004). Intestinal macrophages are highly heterogeneous in terms of their function and localization within the gut. There are three main types of macrophages: monocyte-derived mature macrophages, monocyte-derived inflammatory macrophages, and self-maintaining macrophages. Monocyte-derived mature macrophages (Ly6C^+ , CD64^+ , MHCII^+ , $\text{CX3CR1}^{\text{hi}}$) are located mainly in the lamina propria and secrete anti-inflammatory cytokines. Monocyte-derived inflammatory macrophages (CD64^+ , MHCII^+ , $\text{CX3CR1}^{\text{int}}$) also reside in the lamina propria, but they secrete proinflammatory cytokines. Self-maintaining macrophages (Tim-4^+ , CD4^+) arise from embryonic precursors and adult bone marrow-derived monocytes; they have a relatively low replenishment rate from blood monocytes compared with the other two types of macrophages, which suggests they are maintained locally. They are located predominantly within the muscularis externa layer, although they are also present in the lamina propria (Yip et al., 2021). Macrophages play a key role in the gut immune system and the regulation of gastrointestinal physiology, including gut motility and secretion. Although the gut is in constant contact with foreign antigens, macrophages protect it from chronic inflammation and thus contribute to intestinal homeostasis. They also contribute to neuroimmune interactions.

Dendritic cells are bone marrow-derived antigen-presenting cells that can be divided into two major subsets: conventional (or classical) and plasmacytoid. Although they are developmentally distinct from macrophages, both cell types share many phenotypic markers (CD11c, MHCII, and CX3CR1). However, research suggests that CD64 expression can differentiate intestinal macrophages from dendritic cells (Tamoutounour et al., 2012). Dendritic cells constantly sample luminal content to monitor for pathogens while also mediating tolerance to food antigens and controlling reactivity to the gut microbiota. Collectively, they generate both regulatory and effector T-cell responses (Stagg, 2018).

Innate lymphoid cells are an important family of immune cells that share characteristics with T cells but do not express antigen receptors or undergo clonal selection and expansion when stimulated. Instead, ILCs react promptly to damage signals and produce cytokines. Thus, natural killer cells can be considered the innate counterparts of cytotoxic CD8⁺ T cells, whereas ILC1s, ILC2s, and ILC3s may represent the innate counterparts of CD4⁺ T helper 1, T helper 2, and T helper 17 cells (Eberl et al., 2015; Vivier et al., 2018). In addition to all these innate immune cells, T lymphocytes, as cells of the adaptive immune system, have adapted to intestinal environments, constantly discriminating between natural stimuli (from commensal microbiota and food) and pathogens that need to be eliminated, and helping to maintain the integrity of the intestinal barrier and immune homeostasis. T lymphocytes are interleaved in the epithelial layer between the enterocytes (intraepithelial lymphocytes). Due to this specific location, intraepithelial lymphocytes are in direct contact with enterocytes and are in the immediate proximity of antigens in the gut lumen, representing an important part of the gut lymphoid tissue (Vitale et al., 2016). They have regulatory and effector capabilities, so they can prevent pathogenic invasion while maintaining tolerance to the commensal microbes.

Intraepithelial lymphocytes can be divided into “type a” intestinal T cells, which express TCR $\alpha\beta$ along with CD4 or CD8 $\alpha\beta$ as coreceptors, and “type b”, which express

either TCR $\alpha\beta$ or TCR $\gamma\delta$ and, typically, CD8 $\alpha\alpha$ homodimers. Intestinal T cells such as CD4⁺ T helper cells 1, 2 and 17, CD8⁺ cytotoxic T cells, or FoxP3⁺ regulatory T cells can also be found in the lamina propria. Together with intraepithelial lymphocytes, these lamina propria lymphocytes fight harmful microbes and contribute to mucosal tolerance. Lamina propria lymphocytes are derived from conventional T cells that are developed in the thymus, primed in secondary lymphoid organs, and reach the lamina propria with an effector memory phenotype (Ma et al., 2019). Finally, B cells are also key players in the symbiotic maintenance of the gut system because activated and differentiated B cells (plasma cells) produce secretory immunoglobulin A, which binds commensals, preventing their adhesion to the enterocytes and limiting microbial motility without activating the complement cascade. Thus, B cells prevent inflammatory damage to the epithelial barrier and preserve a healthy microbial ecosystem.

The immune function carried out in the intestine involves a group of organs and structures called the gut-associated lymphoid tissue, which coordinates B cell function and drains mesenteric and caudal lymph nodes. Gut-associated lymphoid tissue includes Peyer's patches, isolated lymphoid follicles, and the appendix. Peyer's patches are located primarily along the small intestine and constitute important sites for the initiation of the immune response in the gut, since they are in close contact with the microbes in the lumen. These structures contain a large number of follicles with small T cell areas and germinal centers in which B cells differentiate and proliferate. They are a major immunoglobulin A induction site and play a key role in the maintenance of commensal bacteria and host protection (Botia-Sanchez et al., 2021). Therefore, similarly to an intact intestinal barrier, the efficient crosstalk between microbes, epithelial cells, and immune system cells guarantees a safe relationship between the microbes and their host. Host health can be compromised when the intestinal barrier is breached, as may occur with pathogen invasion or other factors causing bacterial dysbiosis. Increased permeability of the intestinal barrier, known as "leaky gut", allows the uncontrolled extravasation of microbes and their structural components out of the

gut, where they can reach other tissues and cause an exacerbated immune response and numerous diseases, including mental health disorders (Figure 1).

Bacterial components, such as the lipopolysaccharide from the membrane of Gram-negative bacteria, have been found at increased levels in the serum of depressed people (Caso et al., 2021; Stevens et al., 2018). These bacterial components may leak from the intestinal lumen and activate an innate immune response in the intestine. Thus, on penetrating the lamina propria, antigens are recognized, and trigger the production of pro-inflammatory cytokines, including interleukin-1 β , interleukin-6, tumor necrosis factor alpha (TNF α), interferon- γ , and cytokines involved in the interleukin-23/T helper 17 pathway, in local tissues (Guan, 2019; Sano et al., 2015). Subsequently, the intestinal barrier dysfunction facilitates the passage of local inflammation to the peripheral circulation and other organs, thereby contributing to depression. Research suggests that the T helper 17 pathway plays a key role in depression (Beurel et al., 2013; Beurel et al., 2022; Medina-Rodriguez et al., 2020), as does the abundance of circulating cytokines. Indeed, elevated levels of TNF α , interleukin-6, and interleukin-1 β have been found in blood and cerebrospinal fluid of individuals with depression compared to healthy controls (Medina-Rodriguez et al., 2018).

In summary, maintaining a robust gut barrier and a balanced immune response is important not only for the physical wellbeing of the host but also for their mental health.

4. The gut microbiome and manipulation of the gut microbes and their metabolites as a strategy to treat depression

The gut microbiome is the ecosystem of microorganisms that populate the gastrointestinal tract. The most common bacterial phyla among the microbiome are Bacteroidetes and Firmicutes, although more than 50 phyla have been described in the human gut. Ample research has shown that the effects of the gut microbes go beyond the gastrointestinal system. Apart from participating in human metabolism and nutrition

and being linked to gastrointestinal conditions such as inflammatory bowel disease, irritable bowel syndrome, and obesity, the gut microbiome appears to participate in immune function and cardiovascular, respiratory, liver, kidney, and mental diseases (Hou et al., 2022). Regarding depression, observational studies have shown that the gut microbiome and its metabolites differ between depressed people and healthy controls (Yang et al., 2020). Further, different teams of researchers found that transferring the intestinal microbial community from depressed people to rodents was sufficient to induce depressive-like behaviors in the recipient animals (Kelly et al., 2016). In addition, accumulating data indicate a negative impact of antibiotics on mental health. Overall, antibiotics increase depression prevalence, which is likely linked to a decrease of microbial diversity and intestinal barrier disruption (Dinan and Dinan, 2022). Thus, manipulating or re-establishing the microbiome composition might help to ameliorate depression.

4.1 Probiotic supplementation

Although still controversial, the use of some probiotics has demonstrated a beneficial effect on depression. For instance, one randomized controlled trial (RCT) reported that add-on probiotic treatment (Vivomixx®, containing eight different strains: *Streptococcus thermophilus* NCIMB 30438, *Bifidobacterium breve* NCIMB 30441, *Bifidobacterium longum* NCIMB 30435 (Re-classified as *B. lactis*), *Bifidobacterium infantis* NCIMB 30436 (Re-classified as *B. lactis*), *Lactobacillus acidophilus* NCIMB 30442, *Lactobacillus plantarum* NCIMB 30437, *Lactobacillus paracasei* NCIMB 30439, *Lactobacillus delbrueckii subsp. Bulgaricus* NCIMB 30440 (Re-classified as *L. helveticus*)) led to changes in the gut microbiota and brain, as assessed by functional magnetic resonance imaging (fMRI), and improved depressive symptoms (Schaub et al., 2022). In another clinical trial, NVP-1704 (a mixture of *Lactobacillus reuteri* NK33 and *Bifidobacterium adolescentis* NK98) alleviated subclinical depression and exerted anti-inflammatory effects, with a decrease in serum interleukin-6 concentrations after

eight weeks of intervention (Lee et al., 2021). Another study found that *Bifidobacterium breve* CCFM1025 attenuated major depressive disorder (MDD) by regulating the gut microbiome and tryptophan metabolites, measured in feces by ultra-high performance liquid chromatography-tandem mass spectrometry (UHPLC-MS/MS); however, participant's diet and medication (other than antibiotics) were not strictly controlled, which could have influenced the results (Tian et al., 2022). Another clinical trial with *Lactobacillus plantarum* 299 v showed a decrease in kynurenine concentration determined by high-performance liquid chromatography (HPLC) in blood, together with ameliorated cognitive functions in people with major depression. However, there were no differences in depression score (HAMD-17 scale) observed between probiotic and placebo groups over an eight-week trial (Rudzki et al., 2019). Nor did other probiotics, such as *L. casei* Shirota, decrease depression scores over a nine-week period (Zhang et al., 2021).

Owing to these inconsistencies across different studies, together with the growth of big-data analysis, researchers have begun to synthesize data from different clinical trials. One meta-analysis of five clinical trials (183 cases and 182 controls) showed that probiotics significantly lowered depression scores in people aged under 60 years with MDD (Huang et al., 2016). Another meta-analysis including the same five clinical trials plus five more (1349 participants in total) showed no difference in mood between the probiotic-treated group and the placebo group. However, a separate subgroup analysis of studies conducted in healthy individuals versus those conducted in individuals with MDD or mild to moderate anxiety and/or depression revealed that probiotic supplementation significantly improved the moods of individuals with mild-moderate depressive symptoms (Ng et al., 2018). Interestingly, one recent systematic review of five studies in people with MDD found adjuvant probiotic or synbiotic treatment to be more efficacious for improving symptoms than antidepressant medication alone (Forth et al., 2023).

Thus, probiotic treatment may be effective as monotherapy for depression or as an adjunct to standard antidepressant medication.

In general, most studies agree that common probiotic strains for depression belong to the genus *Lactobacillus*, *Bifidobacterium*, *Akkermansia*, *Clostridium*, *Streptococcus*, and *Enterococcus* (Gao et al., 2023). However, because these genera include a huge number of strains, and because of interpersonal variability, determining a microbial signature and a universal set of probiotics for depression is likely unfeasible. Nevertheless, techniques for individual microbiome screening are advancing fast, and further studies of the gut microbiome composition and potential probiotics might open new avenues for more precise, effective, and personalized ways to improve depression.

4.2 Engineered probiotics

Based on genetic engineering, advances in psychiatric therapies might include the use of genetically modified bacteria. Bacteria can be engineered with immune regulatory proteins, chemotactic response systems, and protein delivery systems (Hou et al., 2022).

Engineered probiotics have been tested in pre-clinical models of colitis, diabetes, cancer, obesity, and pathogenic infections. *Lactococcus lactis* genetically modified to deliver interleukin-10 has been proven safe and effective in people with Crohn's disease (Baat et al., 2006). This may constitute a new approach to continuous expression of beneficial metabolites after intestinal colonization to achieve therapeutic effects, though it has yet to be tested in psychiatric diseases

4.3 Fecal transplants

Another way of manipulating the gut microbiome that has garnered considerable scientific interest is fecal transplantation. The process consists of collecting feces from a healthy donor and introducing them into patient's intestine. Fecal transplantation is

usually performed by colonoscopy, but the medically processed feces can be also be introduced through a nasoduodenal tube, an oral capsule, or an enema. Fecal transplants are safe and well tolerated. They are showing good results in people with a *Clostridium difficile* infection, but their effects in other conditions such as inflammatory bowel disease, obesity, or mental diseases remain unclear. Although evidence from preclinical studies in animal models suggests this approach may improve depression (Knudsen et al., 2021), there are few published studies in humans. One study in people with gastrointestinal disorders found that fecal transplant may improve depressive symptoms (Kurokawa et al., 2018). Furthermore, a case report of a 79-year-old woman with depression who underwent antidepressant and probiotic treatments for six months with no beneficial outcome, showed an improvement in her symptoms following a transplant via gastroscope with a bacterial solution from her six-year-old great-grandson (Cai et al., 2019). Another recent publication reported improved depressive symptoms in two people with MDD after they received fecal microbiota transplantation (oral frozen capsules) as add-on therapy; however, because these people received additional pharmacological and psychotherapy treatments, the authors could not attribute the positive outcomes to the fecal transplant alone (Doll et al., 2022). Further studies are needed to corroborate the efficiency of fecal transplantation in depressed individuals.

4.4 Probiotic food

Diet-mediated approaches to shaping the gut microbiome, such as the ingestion of food containing beneficial bacteria (probiotic foods), might constitute another therapeutic approach to improving mental health. Fermented foods represent a source of live microbes that may be beneficial for depression. These include fermented vegetables (i.e. kimchi) or fermented milk products (i.e. yogurt or kefir) among others (Rao and Samak, 2013). For instance, studies have shown that kefir improves

depression in menopausal women (Ozcan et al., 2019); (Peluzio et al., 2021). Similarly, one cross-sectional study found that consumption of kimchi and fermented milk products reduced the severity and prevalence of depression, though only in men (Kim and Shin, 2019). One recent meta-analysis of eight studies found a significant association between yogurt consumption and decreased depression risk (Luo et al., 2023).

These types of studies are not exempt from limitations, and readers must interpret their results with special caution. Many other dietary components can influence mental health and might be confounding factors in studies of probiotic dietary interventions. Distinguishing the health benefits of fermented food versus other components of the diet is often difficult, and researchers must control for many other confounding factors (e.g. physical activity, socioeconomic status, education, and quality of life) to avoid biased results.

However, observational studies of this nature can still give insights into potential solutions for improving mental health.

4.5 Dietary changes

As well as probiotic food, some other dietary patterns that directly impact gut microbiome composition may be beneficial for the treatment of depression (Dinan et al., 2019). One cross-sectional study of 400 women attending healthcare centers found that diets rich in fruits and vegetables reduced the risk of depression (Baharzadeh et al., 2018). It is known that sulforaphane, present in cruciferous vegetables such as broccoli or sprouts, may prevent the onset of chronic inflammation-related depression (Zhang et al., 2017). In addition, one meta-analysis of 18 studies concluded that total dietary fiber intake is associated with a 10% lower odds of depression in adults and a 57% lower odds in adolescents (Saghafian et al., 2023). Conversely, high-salt diets may lead to greater depressive symptom severity in adolescents (Mrug et al., 2019).

The effects of specific diets on our mental health are mediated by different dietary components, which are related to the gut microbiome:

a) Short-chain fatty acids (SCFAs) and lactate

SCFAs are the main metabolic products of anaerobic bacterial fermentation in the intestine. Dietary fibers are converted to SCFAs after being broken down by bacteria. SCFAs and their metabolites have antidepressant properties (Hamad et al., 2022). Accordingly, compared with healthy controls, people with depression have reduced levels of SCFAs such as acetic acid, propionic acid, and pentanoic acid (Skonieczna-Zydecka et al., 2018), and elevated levels of the saturated fatty acids stearic acid and myristic acid (Frost et al., 2014). Similarly, several studies support an association between depression and other bacterial metabolites abnormalities such as lactate. For example, one study found increased concentrations of urinary lactate in people with severe MDD compared with controls (Chen et al., 2017).

b) Secondary bile acids

The primary bile acids are synthesized from cholesterol in the liver and released into the intestine in response to dietary fat. The action of intestinal bacteria on primary bile acids results in the formation of secondary bile acid species. The secondary bile acid tauroursodeoxycholic acid has been found at increased levels in the blood of people with depression (Pu et al., 2021). One study found that people with MDD had significantly higher levels of 23-nordeoxycholic acid and significantly lower levels of tauroolithocholic acid, glycolithocholic acid, and lithocholic acid 3-sulfate compared with healthy controls (Sun et al., 2022).

c) Tryptophan metabolites

Tryptophan is an essential amino acid necessary to produce proteins. It is found naturally in red meat, poultry, eggs, and dairy products. Thus, diet can also influence the production of several tryptophan metabolites, which have been linked to depression. The intestinal microbiome plays a key role in regulating tryptophan metabolism towards the synthesis of molecules that influence immunity, metabolism,

and neurotransmission (Hou et al., 2023). Kynurenine is one of the main products from tryptophan metabolism. Elevated kynurenine levels are found in social anxiety disorder patients with a history of suicide attempts (Butler et al., 2022; Gimenez-Gomez et al., 2021; Marin et al., 2017). However, several meta-analyses have shown decreased kynurenine levels across MDD patients (Marx et al., 2021; Ogyu et al., 2018). Kynurenine can be metabolized into either kynurenic acid, an N-methyl-D-aspartate (NMDA) receptor antagonist involved in neuroprotection, or quinolinic acid, a potentially neurotoxic NMDA agonist. One meta-analysis showed a reduction in serum kynurenic acid/quinolinic acid in individuals with MDD (Marx et al., 2021), and the decrease in this ratio has been correlated with lower hippocampal and amygdalar volumes (Paul et al., 2022). Levels of quinolinic acid are also elevated in the cerebrospinal fluid of people with a history of suicide attempts (Erhardt et al., 2013). Another recent meta-analysis including 51 studies (7056 participants) found that people with current episodes of MDD had significantly reduced levels of tryptophan, kynurenic acid, kynurenic acid/quinolinic acid, KYNA/3-hydroxykynurenine, and kynurenic acid/kynurenine, and significantly increased levels of kynurenine/tryptophan (Ou et al., 2023). Another intestinal bacteria-derived tryptophan metabolite, known as indole acetic acid, may also be reduced in the cerebrospinal fluid of depressed people (Chen et al., 2022; Coppen, 1972).

Clinical trials on tryptophan supplementation have shown that high tryptophan diets result in fewer depressive symptoms and decreased anxiety (Lindseth et al., 2015).

d) Vitamins

Vitamins from diet can modify the composition and metabolic activity of intestinal bacteria (Pham et al., 2021). Vitamins also appear to affect stress and depression (Bell et al., 1991; Mahdavifar et al., 2021; Young et al., 2019). Studies have linked moderate intake of niacin (vitamin B3) to lower odds of anxiety (Mahdavifar et al., 2021), and moderate intake of folic acid (vitamin B9) to a reduced likelihood of depression

(Mahdavifar et al., 2021). High dietary intake of vitamin B5 and B8 appears to be inversely associated with the prevalence of depression, but only in women (Mahdavifar et al., 2021). Lastly, vitamin A intake has led to significant improvements in depression in people with multiple sclerosis (Bitarafan et al., 2016).

e) Polyphenols

Dietary polyphenols change the gut microbiome composition; at the same time, they are metabolized by the intestinal microbes to produce small active and bioavailable products, which further regulate host functions (Wang et al., 2022). Polyphenols are the most abundant natural compounds in plants, occurring naturally in fruits, vegetables, tea, coffee, cocoa, and wine. In one meta-analysis, 12 of 18 included RCTs showed significant improvements in depression with polyphenol use (Lin et al., 2021). However, many of the data were for post-menopausal women following short-term interventions. In addition, there was a high level of heterogeneity among the trials, possibly because of differences in dosage, forms of supplementation, and duration of the intervention.

f) Polyunsaturated fatty acids (PUFAs)

Modulation of dietary PUFAs by gut microbiome enzymes results in bioactive compounds that could influence human health (Brown et al., 2023). It is known that PUFAs such as omega-3 and omega-9, derived from plant-based products and fish, have beneficial effects on depression (Rao et al., 2008). In this respect, some omega-3 fatty acids, such as eicosapentaenoic acid (EPA), have been found to elicit antidepressant effects in humans, whereas docosahexaenoic acid does not exhibit such benefits (Liao et al., 2019). EPA also alleviated MDD in overweight individuals with elevated inflammatory markers in a randomized dose-finding clinical trial (Lamon-Fava et al., 2023; Mischoulon et al., 2022). Another RCT showed that increases in 18-hydroxyeicosapentaenoic acid (18-HEPE), a metabolite of omega-3 fatty acid, correlates with a reduction in depressive symptoms (Lamon-Fava et al., 2023). One

cross-sectional study found a positive association between serum omega-3 fatty acid (oleic acid) levels and depression (Yin et al., 2023).

Therefore, the above dietary strategies might represent an alternative approach to alleviating or preventing depression by acting on the gut to reach the brain. However, the results of these studies must be interpreted with caution. Observational studies of the effects of one specific compound are usually biased due to the difficulty of eliminating the rest of compounds from the regular diet of participants, which leads to strong confounding effects derived from individual dietary habits. However, diets enriched in several or most of the above components have demonstrated beneficial effects on mental health. For instance, cross-sectional studies and RCTs have found lower risk of depression in people who follow a Mediterranean diet (Oddo et al., 2022; Seethaler et al., 2022).

Consequently, experts in the field recommend that people with depression or susceptibility to depression should supplement a plant-based diet with a high intake of grains/fibers and fish (Dinan et al., 2019; Sanz et al., 2018).

5. Diet-mediated pathways involving the microbiome, the host intestinal barrier, and the immune system as potential targets to treat depression.

This review has presented evidence on the contribution of intestinal microbes and their metabolites, epithelial barrier integrity, and the immune system on depression. However, we have yet to discover how the gut microbiome, the epithelium, and the immune cells interact to control mood. The following section of the review will present potential pathways of microbial-epithelium-immune system communication that might form the basis of future depression therapies (Figure 2). Intestinal epithelial cells collect signals from intestinal microbes to modulate the mucosal barrier and transfer signals to immune cells in the intestine. Indeed, the immune system is intimately linked to the intestinal microbes: early-life colonization of the gut by microbes plays a pivotal role in maturation of the host's immune system (Gensollen et al., 2016), and changes in the

microbiota throughout life affect the intestinal immune function. Likewise, diet has an important direct or indirect effect on intestinal immunity (through the metabolism of the microbiota) and might represent a therapeutic aid to strengthen the intestinal barrier and control inflammation. For instance, some studies have shown that fermented food such as kefir controls inflammation in humans (Bellikci-Koyu et al., 2019). Conversely, high-salt diet induces interleukin-17-dependent gut inflammation (Aguiar et al., 2017).

Diet-linked signals can be triggered by microbial metabolites through different pathways, but the microbes themselves can also control the intestinal barrier and inflammation in different ways.

a) SCFAs and lactate-linked pathways

Bacterial metabolites, such as SCFAs, appear to play important roles in the maintenance of gut epithelial integrity and intestinal immunity (Kayama et al., 2020) (Figure 2A.1, Figure 2B.1). They can be regulated by dietary habits, as mentioned before. SCFAs such as butyrate facilitate the proliferation of epithelial cells as an energy source (Salvi and Cowles, 2021) and modulate their immunological function. Epithelial cells express receptors for SCFAs, such as G protein-coupled receptor 41 (GPR41), GPR43, and GPR109a, and their activation controls the production of cytokines and chemokines by the intestinal epithelium (Kim et al., 2013; Singh et al., 2014). SCFAs also enhance mucus production and release by goblet cells (Burger-van Paassen et al., 2009; Finnie et al., 1995). In the innate immune system, butyrate maintains hypo-responsiveness to the commensal bacteria by repressing the expression of proinflammatory mediators in macrophages and suppressing dendritic cell maturation, inducing therein anti-inflammatory properties to accumulate Foxp3+ regulatory T cells and interleukin-10-producing CD4⁺ T cells (Chang et al., 2014; Singh et al., 2014; Singh et al., 2010). Acetate induces both effector T cells and regulatory T cells by suppression of histone deacetylases and regulation of the mTOR-S6K pathway

(Park et al., 2015) (Figure 2). Lactate, also generated by microbial fermentation, accelerates intestinal stem cell-mediated epithelial development (Lee et al., 2018).

b) Secondary bile acids routes

Bile acids such as deoxycholic acid and ursodeoxycholic acid also regulate epithelial cell activity and protect against intestinal barrier breakdown by promoting enterocyte migration via epidermal growth factor receptor (EGFR)- and cyclooxygenase (COX)-2-dependent mechanisms (Golden et al., 2018; Maran et al., 2009; Mroz et al., 2018) (Figure 2A.2). Bile acids downregulate the production of proinflammatory cytokines by macrophages through recognition by several receptors, such as the farnesoid X receptor (FXR), the sphingosine 1 phosphate receptor 2 (S1PR2), the Vitamin D Receptor or the G-protein-coupled receptor 5 (TGR5), expressed in macrophages (Biagioli et al., 2017; Calzadilla et al., 2022; Guo et al., 2016; Vavassori et al., 2009). In addition, bile acids bind retinoic acid-related orphan receptor (ROR)γt as inverse agonists and decrease interleukin-17 and T helper 17 cells with a resulting improvement in inflammation (Calzadilla et al., 2022) (Figure 2B.2).

c) Tryptophan-linked pathways

Indole generated by bacteria from dietary tryptophan is recognized by the aryl hydrocarbon receptor (AhR) expressed by epithelial cells, and plays an important role in the maintenance of epithelial barrier integrity by participating in mucus secretion, in the expression of cell-junction-associated molecules, and in the appropriate epithelial cell differentiation from crypt stem cells (Metidji et al., 2018; Shimada et al., 2013). Tryptophan-derived phytochemical indole-3-carbinol, found in cruciferous vegetables, act on the AhR, and the absence of AhR ligands affects intraepithelial lymphocytes and results in heightened immune activation (Li et al., 2011). In addition, indole stimulation of ILC3s, which also express the AhR, leads to interleukin-22 increase, which has numerous beneficial effects on epithelial cells and the intestinal barrier (Miljkovic et al., 2021). Microbial-specific indoles can also regulate intestinal barrier function through the

xenobiotic sensor, pregnane X receptor. For instance, indole 3-propionic acid as a ligand for pregnane X receptor up-regulates tight junction protein expression (Venkatesh et al., 2014). (Figure 2A.3, B.3).

d) Vitamin-mediated pathways

Vitamins have known effects on the immune system, since many receptors for vitamins (such as GPR109a, RORs, retinoic acid receptors, retinoid X receptors, liver X receptors, and peroxisome proliferator-activated receptors (PPARs)) are expressed by immune cells such as regulatory T cells and T helper 17 (Veldhoen and Brucklacher-Waldert, 2012). Vitamin B3 (niacin), as a ligand of GPR109a, induces differentiation of regulatory T cells (Singh et al., 2014); vitamin B9 (folic acid) is a survival factor for regulatory T cells (Yoshii et al., 2019); and vitamin A modulates the imbalance of T helper 17 and regulatory T cells (Abdolahi et al., 2015) (Figure 2B.4).

e) Polyphenols-induced routes

Polyphenols have immunomodulatory effects on macrophages; induce B cell proliferation; and decrease T helper 1, T helper 2, T helper 17, and T helper 9 cells, and the inflammatory cytokines interleukin-1 β , interleukin-6, and TNF- α (Shakoor et al., 2021; Tayab et al., 2022) (Figure 2B.5).

f) PUFA-related pathways

PUFAs are potent regulators of inflammation (Figure 2B.6) (Veldhoen and Brucklacher-Waldert, 2012). EPA reduces plasma levels of the inflammatory marker C-reactive protein in MDD patients, which is correlated with depression symptom reduction (Lamon-Fava et al., 2023; Mischoulon et al., 2022). EPA is the precursor of specialized pro-resolving lipid mediators (SPMs) known as resolvins, which are involved in the reduction of inflammation (Lamon-Fava et al., 2021). One very recent RCT of EPA-supplemented MDD patients reported a strong association between changes in 18-HEPE and 15-HEPE, the precursors of resolvins E1-4, with changes in depression symptoms and plasma C-reactive protein concentrations (Lamon-Fava et

al., 2023). The inverse association of 18-HEPE with both systemic inflammation and symptoms of depression highlights the activation of inflammation resolution as a likely mechanism in the treatment of MDD with ω -3 fatty acid supplementation. These findings suggest a mechanistic involvement of EPA-derived SPMs on inflammation and depression improvement. In addition, omega-9 (oleic acid) has demonstrated antidepressant effects through the reduction of stress-induced T helper 17 signaling (Medina-Rodriguez et al., 2020).

g) Pathways induced by microbial components

Regarding bacterial components, epithelial cells express pattern recognition receptors such as toll-like receptors and NOD-like receptors, which sense bacterial components, such as lipopolysaccharide, flagellin, or muramyl dipeptides, and contribute to epithelial cell proliferation and the production of cytokines, antimicrobial molecules, and mucus (Kayama et al., 2020). Toll-like receptor-mediated microbial recognition is required for constitutive anti-inflammatory interleukin-10 production by macrophages in the intestine (Ueda et al., 2010). Toll-like receptor 4 or toll-like receptor 2 activation stimulates regulatory T cells and enhances their immunosuppressive activity (Caramalho et al., 2003; Liu et al., 2006) (Figure 2).

Furthermore, some intestinal bacteria, such as *Citrobacter rodentium* or *Segmented filamentous bacteria* (SFB), attach to the mucosal surface, inducing T helper 17 cell differentiation in the lamina propria through the induction of reactive oxygen species (ROS) or inflammatory molecules in the epithelial cells (Atarashi et al., 2015). Interestingly, SFB can directly induce changes in the host cells by transferring microbial content into epithelial cells through the generation of endocytic vesicles (Ladinsky et al., 2019).

Thus, the abovementioned bacterial signals induce rapid responses, which can later be converted to specific responses by the adaptive immune system. Crosstalk between host cells in direct contact with microbes, their antigens or their metabolites,

and the adaptative immune system, is performed by inflammatory mediators, such as cytokines, and by cells with antigen-presenting properties, such as some epithelial cells, dendritic cells, macrophages, or ILCs (Flannigan et al., 2015; Heuberger et al., 2021; Wosen et al., 2018).

Therefore, although the gut immune system has evolved to live and cooperate with high microbial loads by maintaining a basal level of inflammation and the intestinal barrier function (which together contribute to homeostasis), a failure in the regulatory mechanisms of the intestinal milieu can lead to psychiatric disorders.

6. Conclusion

Our understanding of the complex crosstalk between the brain, the gut microbiome, and the immune system in depression is still very limited despite the large amount of scientific data generated in recent years. To date, most studies on the subject have been observational, meaning they could infer associations but not establish true causal links. In addition, clinical trials have included insufficient numbers of participants and failed to control for many influencing sociodemographic, medical, dietary, and lifestyle variables. As a result, we are unable to draw robust conclusions from the available evidence.

There is a need for large-scale longitudinal and clinical cohort studies including multi-omics data sets, and parallel translational studies in experimental models, to progress from associations to causality, decode the mechanisms underlying microbiota-immune interactions, and enable a more precise identification of biomarkers and therapeutic targets of depression. Microbiome-based biomarkers could be useful for predicting, diagnosing, and establishing the prognosis of depression, as current clinical practice is still based on questionnaires and rating scales. New information on the mechanisms underlying the gut microbiota-mediated effects on the host's immune system could also serve as a basis for designing adequate drug- and lifestyle-based

therapies and identifying new therapeutic targets to improve current management strategies. Furthermore, given the wide variety of causes and phenotypes of the condition, and the considerable individual variability of the gut microbiota, it is reasonable to envisage future progress towards personalized, microbiome-informed management programs to treat or prevent depression.

Figure legends

Figure 1. Intestinal bacteria interplay with the host intestinal epithelium and the immune system to maintain mental wellbeing. Disruption of this well-orchestrated microbe-host ecosystem leads to depression. (Figure created with Biorender.com)

Figure 2 Potential mechanisms controlling mood. Some bacterial metabolites favor intestinal integrity **(A)** and/or reduce inflammation **(B)**, potentially alleviating depression and promoting mental wellbeing. **A.1.** SCFAs bind to receptors in epithelial cells such as GPR41, GPR43, and GPR109a to favor intestinal integrity (e.g. enhance mucus release by goblet cells and facilitate the proliferation of epithelial cells) **B.1.** SCFAs reduce inflammation by increasing Tregs and IL-10-producing Th cells, repressing macrophages and suppressing dendritic cell maturation). **A.2.** SBAs protect against intestinal barrier breakdown by promoting enterocyte migration via EGFR- and COX-2-dependent mechanisms. **B.2.** SBAs downregulate proinflammatory cytokines in macrophages and decrease IL-17 and Th17, which improves inflammation. **A.3.** Indole is recognized by AhR or PXR expressed by epithelial cells to maintain epithelial barrier and participates in mucus secretion, in the expression of tight junction molecules, and in the appropriate epithelial cell differentiation from crypt stem cells. **B.3** Indole stimulates ILC3s increasing IL-22. **B.4.** Vitamins are recognized by many receptors expressed by immune cells such as Tregs and Th17. Vitamins induce differentiation

and survival of Tregs and modulate Th17 cells. **B.5.** PPs have immunomodulatory effects on macrophages and decrease Th1, Th2, Th17, and Th9 cells and inflammatory cytokines (IL-17, IL-1 β , IL-6, and TNF- α). **B.6.** PUFAs decrease immune response by changing plasma CRP levels and Rv concentration and reducing Th17 signaling. AhR: aryl hydrocarbon receptor; COX: cyclooxygenase; CRP: C-reactive protein; D: dendritic cell; EGFR: epidermal growth factor receptor; GPR: G-protein-coupled receptor; IL: interleukin; ILC: innate lymphoid cell; M: macrophage; PP: Polyphenol; PUFA: polyunsaturated fatty acid; PXR: pregnane X receptor; Rv: resolvin; SBA: secondary bile acid; SCFA: short chain fatty acid; Th: T helper cell; TJ: tight junction; TNF: tumor necrosis factor; Tregs: regulatory T cells. (Figure created with Biorender.com)

Data Availability Statement

This article contains no datasets generated or analyzed during the current study.

Authorship contributions

Contributed to conception: Medina-Rodriguez EM, Sanz Y, Martinez-Raga J

Wrote the manuscript: Medina-Rodriguez EM

Revised the manuscript critically: Sanz Y, Martinez-Raga J

References

- Abdolahi M, Yavari P, Honarvar NM, Bitarafan S, Mahmoudi M and Saboor-Yaraghi AA (2015) Molecular Mechanisms of the Action of Vitamin A in Th17/Treg Axis in Multiple Sclerosis. *J Mol Neurosci* **57**:605-613.
- Aguiar SLF, Miranda MCG, Guimaraes MAF, Santiago HC, Queiroz CP, Cunha PDS, Cara DC, Foureaux G, Ferreira AJ, Cardoso VN, Barros PA, Maioli TU and Faria AMC (2017) High-Salt Diet Induces IL-17-Dependent Gut Inflammation and Exacerbates Colitis in Mice. *Front Immunol* **8**:1969.

Alli SR, Gorbovskaia I, Liu JCW, Kolla NJ, Brown L and Muller DJ (2022) The Gut Microbiome in Depression and Potential Benefit of Prebiotics, Probiotics and Synbiotics: A Systematic Review of Clinical Trials and Observational Studies. *Int J Mol Sci* **23**.

Atarashi K, Tanoue T, Ando M, Kamada N, Nagano Y, Narushima S, Suda W, Imaoka A, Setoyama H, Nagamori T, Ishikawa E, Shima T, Hara T, Kado S, Jinnohara T, Ohno H, Kondo T, Toyooka K, Watanabe E, Yokoyama S, Tokoro S, Mori H, Noguchi Y, Morita H, Ivanov II, Sugiyama T, Nunez G, Camp JG, Hattori M, Umesaki Y and Honda K (2015) Th17 Cell Induction by Adhesion of Microbes to Intestinal Epithelial Cells. *Cell* **163**:367-380.

Baharzadeh E, Siassi F, Qorbani M, Koohdani F, Pak N and Sotoudeh G (2018) Fruits and vegetables intake and its subgroups are related to depression: a cross-sectional study from a developing country. *Ann Gen Psychiatry* **17**:46.

Bell IR, Edman JS, Morrow FD, Marby DW, Mirages S, Perrone G, Kayne HL and Cole JO (1991) B complex vitamin patterns in geriatric and young adult inpatients with major depression. *J Am Geriatr Soc* **39**:252-257.

Bellikci-Koyu E, Sarer-Yurekli BP, Akyon Y, Aydin-Kose F, Karagozlu C, Ozgen AG, Brinkmann A, Nitsche A, Ergunay K, Yilmaz E and Buyuktuncer Z (2019) Effects of Regular Kefir Consumption on Gut Microbiota in Patients with Metabolic Syndrome: A Parallel-Group, Randomized, Controlled Study. *Nutrients* **11**.

Beurel E, Harrington LE and Jope RS (2013) Inflammatory T helper 17 cells promote depression-like behavior in mice. *Biol Psychiatry* **73**:622-630.

Beurel E, Medina-Rodriguez EM and Jope RS (2022) Targeting the Adaptive Immune System in Depression: Focus on T Helper 17 Cells. *Pharmacol Rev* **74**:373-386.

Bevins CL and Salzman NH (2011) Paneth cells, antimicrobial peptides and maintenance of intestinal homeostasis. *Nat Rev Microbiol* **9**:356-368.

Biagioli M, Carino A, Cipriani S, Francisci D, Marchiano S, Scarpelli P, Sorcini D, Zampella A and Fiorucci S (2017) The Bile Acid Receptor GPBAR1 Regulates the M1/M2 Phenotype of Intestinal Macrophages and Activation of GPBAR1 Rescues Mice from Murine Colitis. *J Immunol* **199**:718-733.

Bitarafan S, Saboor-Yaraghi A, Sahraian MA, Soltani D, Nafissi S, Togha M, Beladi Moghadam N, Roostaei T, Mohammadzadeh Honarvar N and Harirchian MH (2016) Effect of Vitamin A Supplementation on fatigue and depression in Multiple Sclerosis patients: A Double-Blind Placebo-Controlled Clinical Trial. *Iran J Allergy Asthma Immunol* **15**:13-19.

Botia-Sanchez M, Alarcon-Riquelme ME and Galicia G (2021) B Cells and Microbiota in Autoimmunity. *Int J Mol Sci* **22**.

Braat H, Rottiers P, Hommes DW, Huyghebaert N, Remaut E, Remon JP, van Deventer SJ, Neiryck S, Peppelenbosch MP and Steidler L (2006) A phase I trial with transgenic bacteria expressing interleukin-10 in Crohn's disease. *Clin Gastroenterol Hepatol* **4**:754-759.

Brown EM, Clardy J and Xavier RJ (2023) Gut microbiome lipid metabolism and its impact on host physiology. *Cell Host Microbe* **31**:173-186.

Burger-van Paassen N, Vincent A, Puiman PJ, van der Sluis M, Bouma J, Boehm G, van Goudoever JB, van Seuning I and Renes IB (2009) The regulation of intestinal mucin MUC2 expression by short-chain fatty acids: implications for epithelial protection. *Biochem J* **420**:211-219.

Butler MI, Long-Smith C, Moloney GM, Morkl S, O'Mahony SM, Cryan JF, Clarke G and Dinan TG (2022) The immune-kynurenine pathway in social anxiety disorder. *Brain Behav Immun* **99**:317-326.

Cai T, Shi X, Yuan LZ, Tang D and Wang F (2019) Fecal microbiota transplantation in an elderly patient with mental depression. *Int Psychogeriatr* **31**:1525-1526.

Calzadilla N, Comiskey SM, Dudeja PK, Saksena S, Gill RK and Alrefai WA (2022) Bile acids as inflammatory mediators and modulators of intestinal permeability. *Front Immunol* **13**:1021924.

Caramalho I, Lopes-Carvalho T, Ostler D, Zelenay S, Haury M and Demengeot J (2003) Regulatory T cells selectively express toll-like receptors and are activated by lipopolysaccharide. *J Exp Med* **197**:403-411.

Caso JR, MacDowell KS, Gonzalez-Pinto A, Garcia S, de Diego-Adelino J, Carceller-Sindreu M, Sarramea F, Caballero-Villarraso J, Gracia-Garcia P, De la Camara C, Aguera L, Gomez-Lus ML, Alba C, Rodriguez JM and Leza JC (2021) Gut microbiota, innate immune pathways, and inflammatory control mechanisms in patients with major depressive disorder. *Transl Psychiatry* **11**:645.

Chang PV, Hao L, Offermanns S and Medzhitov R (2014) The microbial metabolite butyrate regulates intestinal macrophage function via histone deacetylase inhibition. *Proc Natl Acad Sci U S A* **111**:2247-2252.

Chen JJ, Zhou CJ, Zheng P, Cheng K, Wang HY, Li J, Zeng L and Xie P (2017) Differential urinary metabolites related with the severity of major depressive disorder. *Behav Brain Res* **332**:280-287.

Chen Y, Tian P, Wang Z, Pan R, Shang K, Wang G, Zhao J and Chen W (2022) Indole Acetic Acid Exerts Anti-Depressive Effects on an Animal Model of Chronic Mild Stress. *Nutrients* **14**.

Coppen A (1972) Indoleamines and affective disorders. *J Psychiatr Res* **9**:163-171.

Dinan K and Dinan T (2022) Antibiotics and mental health: The good, the bad and the ugly. *J Intern Med* **292**:858-869.

Dinan TG, Stanton C and Cryan JF (2013) Psychobiotics: a novel class of psychotropic. *Biol Psychiatry* **74**:720-726.

Dinan TG, Stanton C, Long-Smith C, Kennedy P, Cryan JF, Cowan CSM, Cenit MC, van der Kamp JW and Sanz Y (2019) Feeding melancholic microbes: MyNewGut recommendations on diet and mood. *Clin Nutr* **38**:1995-2001.

Disease GBD, Injury I and Prevalence C (2018) Global, regional, and national incidence, prevalence, and years lived with disability for 354 diseases and injuries for 195 countries and territories, 1990-2017: a systematic analysis for the Global Burden of Disease Study 2017. *Lancet* **392**:1789-1858.

Doll JPK, Vazquez-Castellanos JF, Schaub AC, Schweinfurth N, Kettelhack C, Schneider E, Yamanbaeva G, Mahlmann L, Brand S, Beglinger C, Borgwardt S, Raes J, Schmidt A and Lang UE (2022) Fecal Microbiota Transplantation (FMT) as an Adjunctive Therapy for Depression-Case Report. *Front Psychiatry* **13**:815422.

Eberl G, Colonna M, Di Santo JP and McKenzie AN (2015) Innate lymphoid cells. Innate lymphoid cells: a new paradigm in immunology. *Science* **348**:aaa6566.

Erhardt S, Lim CK, Linderholm KR, Janelidze S, Lindqvist D, Samuelsson M, Lundberg K, Postolache TT, Traskman-Bendz L, Guillemin GJ and Brundin L (2013) Connecting inflammation with glutamate agonism in suicidality. *Neuropsychopharmacology* **38**:743-752.

Finnie IA, Dwarakanath AD, Taylor BA and Rhodes JM (1995) Colonic mucin synthesis is increased by sodium butyrate. *Gut* **36**:93-99.

Flannigan KL, Geem D, Harusato A and Denning TL (2015) Intestinal Antigen-Presenting Cells: Key Regulators of Immune Homeostasis and Inflammation. *Am J Pathol* **185**:1809-1819.

Forth E, Buehner B, Storer A, Sgarbossa C, Milev R and Chinna Meyyappan A (2023) Systematic review of probiotics as an adjuvant treatment for psychiatric disorders. *Front Behav Neurosci* **17**:1111349.

Frost G, Sleeth ML, Sahuri-Arisoylu M, Lizarbe B, Cerdan S, Brody L, Anastasovska J, Ghourab S, Hankir M, Zhang S, Carling D, Swann JR, Gibson G, Viardot A, Morrison D, Louise Thomas E and Bell JD (2014) The short-chain fatty acid acetate reduces appetite via a central homeostatic mechanism. *Nat Commun* **5**:3611.

Gao J, Zhao L, Cheng Y, Lei W, Wang Y, Liu X, Zheng N, Shao L, Chen X, Sun Y, Ling Z and Xu W (2023) Probiotics for the treatment of depression and its comorbidities: A systemic review. *Front Cell Infect Microbiol* **13**:1167116.

Gensollen T, Iyer SS, Kasper DL and Blumberg RS (2016) How colonization by microbiota in early life shapes the immune system. *Science* **352**:539-544.

Gimenez-Gomez P, Ballestin R, Gil de Biedma-Elduayen L, Vidal R, Ferrer-Perez C, Reguilon MD, O'Shea E, Minarro J, Colado MI and Rodriguez-Arias M (2021) Decreased kynurenine pathway potentiate resilience to social defeat effect on cocaine reward. *Neuropharmacology* **197**:108753.

Golden JM, Escobar OH, Nguyen MVL, Mallicote MU, Kavarian P, Frey MR and Gayer CP (2018) Ursodeoxycholic acid protects against intestinal barrier breakdown by promoting enterocyte migration via EGFR- and COX-2-dependent mechanisms. *Am J Physiol Gastrointest Liver Physiol* **315**:G259-G271.

Goto Y (2019) Epithelial Cells as a Transmitter of Signals From Commensal Bacteria and Host Immune Cells. *Front Immunol* **10**:2057.

Guan Q (2019) A Comprehensive Review and Update on the Pathogenesis of Inflammatory Bowel Disease. *J Immunol Res* **2019**:7247238.

Guo C, Xie S, Chi Z, Zhang J, Liu Y, Zhang L, Zheng M, Zhang X, Xia D, Ke Y, Lu L and Wang D (2016) Bile Acids Control Inflammation and Metabolic Disorder through Inhibition of NLRP3 Inflammasome. *Immunity* **45**:802-816.

Gutzeit C, Magri G and Cerutti A (2014) Intestinal IgA production and its role in host-microbe interaction. *Immunol Rev* **260**:76-85.

Hamad I, Cardilli A, Corte-Real BF, Dyczko A, Vangronsveld J and Kleinewietfeld M (2022) High-Salt Diet Induces Depletion of Lactic Acid-Producing Bacteria in Murine Gut. *Nutrients* **14**.

Heuberger C, Pott J and Maloy KJ (2021) Why do intestinal epithelial cells express MHC class II? *Immunology* **162**:357-367.

Hou K, Wu ZX, Chen XY, Wang JQ, Zhang D, Xiao C, Zhu D, Koya JB, Wei L, Li J and Chen ZS (2022) Microbiota in health and diseases. *Signal Transduct Target Ther* **7**:135.

Hou Y, Li J and Ying S (2023) Tryptophan Metabolism and Gut Microbiota: A Novel Regulatory Axis Integrating the Microbiome, Immunity, and Cancer. *Metabolites* **13**.

Huang R, Wang K and Hu J (2016) Effect of Probiotics on Depression: A Systematic Review and Meta-Analysis of Randomized Controlled Trials. *Nutrients* **8**.

Johansson ME, Sjovall H and Hansson GC (2013) The gastrointestinal mucus system in health and disease. *Nat Rev Gastroenterol Hepatol* **10**:352-361.

Kayama H, Okumura R and Takeda K (2020) Interaction Between the Microbiota, Epithelia, and Immune Cells in the Intestine. *Annu Rev Immunol* **38**:23-48.

Kelly JR, Borre Y, C OB, Patterson E, El Aidy S, Deane J, Kennedy PJ, Beers S, Scott K, Moloney G, Hoban AE, Scott L, Fitzgerald P, Ross P, Stanton C, Clarke G, Cryan JF and Dinan TG (2016) Transferring the blues: Depression-associated gut microbiota induces neurobehavioural changes in the rat. *J Psychiatr Res* **82**:109-118.

Kim CS and Shin DM (2019) Probiotic food consumption is associated with lower severity and prevalence of depression: A nationwide cross-sectional study. *Nutrition* **63-64**:169-174.

Kim MH, Kang SG, Park JH, Yanagisawa M and Kim CH (2013) Short-chain fatty acids activate GPR41 and GPR43 on intestinal epithelial cells to promote inflammatory responses in mice. *Gastroenterology* **145**:396-406 e391-310.

Knoop KA and Newberry RD (2018) Goblet cells: multifaceted players in immunity at mucosal surfaces. *Mucosal Immunol* **11**:1551-1557.

Knudsen JK, Michaelsen TY, Bundgaard-Nielsen C, Nielsen RE, Hjerrild S, Leutscher P, Wegener G and Sorensen S (2021) Faecal microbiota transplantation from patients with depression or healthy individuals into rats modulates mood-related behaviour. *Sci Rep* **11**:21869.

Kunugi H (2021) Gut Microbiota and Pathophysiology of Depressive Disorder. *Ann Nutr Metab* **77 Suppl 2**:11-20.

Kurokawa S, Kishimoto T, Mizuno S, Masaoka T, Naganuma M, Liang KC, Kitazawa M, Nakashima M, Shindo C, Suda W, Hattori M, Kanai T and Mimura M (2018) The effect of fecal microbiota transplantation on psychiatric symptoms among patients with irritable bowel syndrome, functional diarrhea and functional constipation: An open-label observational study. *J Affect Disord* **235**:506-512.

Ladinsky MS, Araujo LP, Zhang X, Veltri J, Galan-Diez M, Soualhi S, Lee C, Irie K, Pinker EY, Narushima S, Bandyopadhyay S, Nagayama M, Elhenawy

W, Coombes BK, Ferraris RP, Honda K, Iliev ID, Gao N, Bjorkman PJ and Ivanov, II (2019) Endocytosis of commensal antigens by intestinal epithelial cells regulates mucosal T cell homeostasis. *Science* **363**.

Lamon-Fava S, Liu M, Dunlop BW, Kinkead B, Schettler PJ, Felger JC, Ziegler TR, Fava M, Mischoulon D and Rapaport MH (2023) Clinical response to EPA supplementation in patients with major depressive disorder is associated with higher plasma concentrations of pro-resolving lipid mediators. *Neuropsychopharmacology* **48**:929-935.

Lamon-Fava S, So J, Mischoulon D, Ziegler TR, Dunlop BW, Kinkead B, Schettler PJ, Nierenberg AA, Felger JC, Maddipati KR, Fava M and Rapaport MH (2021) Dose- and time-dependent increase in circulating anti-inflammatory and pro-resolving lipid mediators following eicosapentaenoic acid supplementation in patients with major depressive disorder and chronic inflammation. *Prostaglandins Leukot Essent Fatty Acids* **164**:102219.

Lee HJ, Hong JK, Kim JK, Kim DH, Jang SW, Han SW and Yoon IY (2021) Effects of Probiotic NVP-1704 on Mental Health and Sleep in Healthy Adults: An 8-Week Randomized, Double-Blind, Placebo-Controlled Trial. *Nutrients* **13**.

Lee YS, Kim TY, Kim Y, Lee SH, Kim S, Kang SW, Yang JY, Baek IJ, Sung YH, Park YY, Hwang SW, O E, Kim KS, Liu S, Kamada N, Gao N and Kweon MN (2018) Microbiota-Derived Lactate Accelerates Intestinal Stem-Cell-Mediated Epithelial Development. *Cell Host Microbe* **24**:833-846 e836.

Li Y, Innocentin S, Withers DR, Roberts NA, Gallagher AR, Grigorieva EF, Wilhelm C and Veldhoen M (2011) Exogenous stimuli maintain intraepithelial lymphocytes via aryl hydrocarbon receptor activation. *Cell* **147**:629-640.

Liao Y, Xie B, Zhang H, He Q, Guo L, Subramanieapillai M, Fan B, Lu C and McIntyre RS (2019) Efficacy of omega-3 PUFAs in depression: A meta-analysis. *Transl Psychiatry* **9**:190.

Lin K, Li Y, Toit ED, Wendt L and Sun J (2021) Effects of Polyphenol Supplementations on Improving Depression, Anxiety, and Quality of Life in Patients With Depression. *Front Psychiatry* **12**:765485.

Lindseth G, Helland B and Caspers J (2015) The effects of dietary tryptophan on affective disorders. *Arch Psychiatr Nurs* **29**:102-107.

Liu H, Komai-Koma M, Xu D and Liew FY (2006) Toll-like receptor 2 signaling modulates the functions of CD4+ CD25+ regulatory T cells. *Proc Natl Acad Sci U S A* **103**:7048-7053.

Luo Y, Li Z, Gu L and Zhang K (2023) Fermented dairy foods consumption and depressive symptoms: A meta-analysis of cohort studies. *PLoS One* **18**:e0281346.

Ma H, Tao W and Zhu S (2019) T lymphocytes in the intestinal mucosa: defense and tolerance. *Cell Mol Immunol* **16**:216-224.

Mahdaviifar B, Hosseinzadeh M, Salehi-Abargouei A, Mirzaei M and Vafa M (2021) Dietary intake of B vitamins and their association with depression, anxiety, and stress symptoms: A cross-sectional, population-based survey. *J Affect Disord* **288**:92-98.

Maran RR, Thomas A, Roth M, Sheng Z, Esterly N, Pinson D, Gao X, Zhang Y, Ganapathy V, Gonzalez FJ and Guo GL (2009) Farnesoid X receptor deficiency in mice leads to increased intestinal epithelial cell proliferation and tumor development. *J Pharmacol Exp Ther* **328**:469-477.

Margolis KG, Cryan JF and Mayer EA (2021) The Microbiota-Gut-Brain Axis: From Motility to Mood. *Gastroenterology* **160**:1486-1501.

Marin IA, Goertz JE, Ren T, Rich SS, Onengut-Gumuscu S, Farber E, Wu M, Overall CC, Kipnis J and Gaultier A (2017) Microbiota alteration is associated with the development of stress-induced despair behavior. *Sci Rep* **7**:43859.

Marx W, McGuinness AJ, Rocks T, Ruusunen A, Cleminson J, Walker AJ, Gomes-da-Costa S, Lane M, Sanches M, Diaz AP, Tseng PT, Lin PY, Berk M, Clarke G, O'Neil A, Jacka F, Stubbs B, Carvalho AF, Quevedo J, Soares JC and Fernandes BS (2021) The kynurenine pathway in major depressive disorder, bipolar disorder, and schizophrenia: a meta-analysis of 101 studies. *Mol Psychiatry* **26**:4158-4178.

Medina-Rodriguez EM, Lowell JA, Worthen RJ, Syed SA and Beurel E (2018) Involvement of Innate and Adaptive Immune Systems Alterations in the Pathophysiology and Treatment of Depression. *Front Neurosci* **12**:547.

Medina-Rodriguez EM, Madorma D, O'Connor G, Mason BL, Han D, Deo SK, Oppenheimer M, Nemeroff CB, Trivedi MH, Daunert S and Beurel E (2020) Identification of a Signaling Mechanism by Which the Microbiome Regulates Th17 Cell-Mediated Depressive-Like Behaviors in Mice. *Am J Psychiatry* **177**:974-990.

Metidji A, Omenetti S, Crotta S, Li Y, Nye E, Ross E, Li V, Maradana MR, Schiering C and Stockinger B (2018) The Environmental Sensor AHR Protects from Inflammatory Damage by Maintaining Intestinal Stem Cell Homeostasis and Barrier Integrity. *Immunity* **49**:353-362 e355.

Miljkovic D, Jevtic B, Stojanovic I and Dimitrijevic M (2021) ILC3, a Central Innate Immune Component of the Gut-Brain Axis in Multiple Sclerosis. *Front Immunol* **12**:657622.

Mischoulon D, Dunlop BW, Kinkead B, Schettler PJ, Lamon-Fava S, Rakofsky JJ, Nierenberg AA, Clain AJ, Mletzko Crowe T, Wong A, Felger JC, Sangermano L, Ziegler TR, Cusin C, Fisher LB, Fava M and Rapaport MH (2022) Omega-3 Fatty Acids for Major Depressive Disorder With High Inflammation: A Randomized Dose-Finding Clinical Trial. *J Clin Psychiatry* **83**.

Mroz MS, Lajczak NK, Goggins BJ, Keely S and Keely SJ (2018) The bile acids, deoxycholic acid and ursodeoxycholic acid, regulate colonic epithelial

wound healing. *Am J Physiol Gastrointest Liver Physiol* **314**:G378-G387.

Mrug S, Orihuela C, Mrug M and Sanders PW (2019) Sodium and potassium excretion predict increased depression in urban adolescents. *Physiol Rep* **7**:e14213.

Ng QX, Peters C, Ho CYX, Lim DY and Yeo WS (2018) A meta-analysis of the use of probiotics to alleviate depressive symptoms. *J Affect Disord* **228**:13-19.

Oddo VM, Welke L, McLeod A, Pezley L, Xia Y, Maki P, Koenig MD, Kominiarek MA, Langenecker S and Tussing-Humphreys L (2022) Adherence to a Mediterranean Diet Is Associated with Lower Depressive Symptoms among U.S. Adults. *Nutrients* **14**.

Ogyu K, Kubo K, Noda Y, Iwata Y, Tsugawa S, Omura Y, Wada M, Tarumi R, Plitman E, Moriguchi S, Miyazaki T, Uchida H, Graff-Guerrero A, Mimura M and Nakajima S (2018) Kynurenine pathway in depression: A systematic review and meta-analysis. *Neurosci Biobehav Rev* **90**:16-25.

Ormel J, Kessler RC and Schoevers R (2019) Depression: more treatment but no drop in prevalence: how effective is treatment? And can we do better? *Curr Opin Psychiatry* **32**:348-354.

Ou W, Chen Y, Ju Y, Ma M, Qin Y, Bi Y, Liao M, Liu B, Liu J, Zhang Y and Li L (2023) The kynurenine pathway in major depressive disorder under different disease states: A systematic review and meta-analysis. *J Affect Disord* **339**:624-632.

Ozcan H, Oskay U and Bodur AF (2019) Effects of Kefir on Quality of Life and Sleep Disturbances in Postmenopausal Women. *Holist Nurs Pract* **33**:207-213.

Park J, Kim M, Kang SG, Jannasch AH, Cooper B, Patterson J and Kim CH (2015) Short-chain fatty acids induce both effector and regulatory T cells by

suppression of histone deacetylases and regulation of the mTOR-S6K pathway. *Mucosal Immunol* **8**:80-93.

Paul ER, Schwieler L, Erhardt S, Boda S, Trepici A, Kampe R, Asratian A, Holm L, Yngve A, Dantzer R, Heilig M, Hamilton JP and Samuelsson M (2022) Peripheral and central kynurenine pathway abnormalities in major depression. *Brain Behav Immun* **101**:136-145.

Peluzio M, Dias MME, Martinez JA and Milagro FI (2021) Kefir and Intestinal Microbiota Modulation: Implications in Human Health. *Front Nutr* **8**:638740.

Pham VT, Fehlbaum S, Seifert N, Richard N, Bruins MJ, Sybesma W, Rehman A and Steinert RE (2021) Effects of colon-targeted vitamins on the composition and metabolic activity of the human gut microbiome- a pilot study. *Gut Microbes* **13**:1-20.

Pu J, Liu Y, Zhang H, Tian L, Gui S, Yu Y, Chen X, Chen Y, Yang L, Ran Y, Zhong X, Xu S, Song X, Liu L, Zheng P, Wang H and Xie P (2021) An integrated meta-analysis of peripheral blood metabolites and biological functions in major depressive disorder. *Mol Psychiatry* **26**:4265-4276.

Rao RK and Samak G (2013) Protection and Restitution of Gut Barrier by Probiotics: Nutritional and Clinical Implications. *Curr Nutr Food Sci* **9**:99-107.

Rao TS, Asha MR, Ramesh BN and Rao KS (2008) Understanding nutrition, depression and mental illnesses. *Indian J Psychiatry* **50**:77-82.

Rudzki L, Ostrowska L, Pawlak D, Malus A, Pawlak K, Waszkiewicz N and Szulc A (2019) Probiotic *Lactobacillus Plantarum* 299v decreases kynurenine concentration and improves cognitive functions in patients with major depression: A double-blind, randomized, placebo controlled study. *Psychoneuroendocrinology* **100**:213-222.

- Saghafian F, Hajishafiee M, Rouhani P and Saneei P (2023) Dietary fiber intake, depression, and anxiety: a systematic review and meta-analysis of epidemiologic studies. *Nutr Neurosci* **26**:108-126.
- Salvi PS and Cowles RA (2021) Butyrate and the Intestinal Epithelium: Modulation of Proliferation and Inflammation in Homeostasis and Disease. *Cells* **10**.
- Sano T, Huang W, Hall JA, Yang Y, Chen A, Gavzy SJ, Lee JY, Ziel JW, Miraldi ER, Domingos AI, Bonneau R and Littman DR (2015) An IL-23R/IL-22 Circuit Regulates Epithelial Serum Amyloid A to Promote Local Effector Th17 Responses. *Cell* **163**:381-393.
- Sanz Y, Romani-Perez M, Benitez-Paez A, Portune KJ, Brigidi P, Rampelli S, Dinan T, Stanton C, Delzenne N, Blachier F, Neyrinck AM, Beaumont M, Olivares M, Holzer P, Gunther K, Wolters M, Ahrens W, Claus SP, Campoy C, Murphy R, Sadler C, Fernandez L and Kamp JV (2018) Towards microbiome-informed dietary recommendations for promoting metabolic and mental health: Opinion papers of the MyNewGut project. *Clin Nutr* **37**:2191-2197.
- Sarkar A, Lehto SM, Harty S, Dinan TG, Cryan JF and Burnet PWJ (2016) Psychobiotics and the Manipulation of Bacteria-Gut-Brain Signals. *Trends Neurosci* **39**:763-781.
- Schaub AC, Schneider E, Vazquez-Castellanos JF, Schweinfurth N, Kettelhack C, Doll JPK, Yamanbaeva G, Mahlmann L, Brand S, Beglinger C, Borgwardt S, Raes J, Schmidt A and Lang UE (2022) Clinical, gut microbial and neural effects of a probiotic add-on therapy in depressed patients: a randomized controlled trial. *Transl Psychiatry* **12**:227.
- Schultz I and Keita AV (2020) The Intestinal Barrier and Current Techniques for the Assessment of Gut Permeability. *Cells* **9**.
- Seethaler B, Nguyen NK, Basrai M, Kiechle M, Walter J, Delzenne NM and Bischoff SC (2022) Short-chain fatty acids are key mediators of the favorable effects

of the Mediterranean diet on intestinal barrier integrity: data from the randomized controlled LIBRE trial. *Am J Clin Nutr* **116**:928-942.

Shakoor H, Feehan J, Apostolopoulos V, Platat C, Al Dhaheri AS, Ali HI, Ismail LC, Bosevski M and Stojanovska L (2021) Immunomodulatory Effects of Dietary Polyphenols. *Nutrients* **13**.

Shimada Y, Kinoshita M, Harada K, Mizutani M, Masahata K, Kayama H and Takeda K (2013) Commensal bacteria-dependent indole production enhances epithelial barrier function in the colon. *PLoS One* **8**:e80604.

Singh N, Gurav A, Sivaprakasam S, Brady E, Padia R, Shi H, Thangaraju M, Prasad PD, Manicassamy S, Munn DH, Lee JR, Offermanns S and Ganapathy V (2014) Activation of Gpr109a, receptor for niacin and the commensal metabolite butyrate, suppresses colonic inflammation and carcinogenesis. *Immunity* **40**:128-139.

Singh N, Thangaraju M, Prasad PD, Martin PM, Lambert NA, Boettger T, Offermanns S and Ganapathy V (2010) Blockade of dendritic cell development by bacterial fermentation products butyrate and propionate through a transporter (Slc5a8)-dependent inhibition of histone deacetylases. *J Biol Chem* **285**:27601-27608.

Skonieczna-Zydecka K, Grochans E, Maciejewska D, Szkup M, Schneider-Matyka D, Jurczak A, Loniewski I, Kaczmarczyk M, Marlicz W, Czerwinska-Rogowska M, Pelka-Wysiecka J, Dec K and Stachowska E (2018) Faecal Short Chain Fatty Acids Profile is Changed in Polish Depressive Women. *Nutrients* **10**.

Stagg AJ (2018) Intestinal Dendritic Cells in Health and Gut Inflammation. *Front Immunol* **9**:2883.

Stevens BR, Goel R, Seungbum K, Richards EM, Holbert RC, Pepine CJ and Raizada MK (2018) Increased human intestinal barrier permeability plasma

biomarkers zonulin and FABP2 correlated with plasma LPS and altered gut microbiome in anxiety or depression. *Gut* **67**:1555-1557.

Sun N, Zhang J, Wang J, Liu Z, Wang X, Kang P, Yang C, Liu P and Zhang K (2022) Abnormal gut microbiota and bile acids in patients with first-episode major depressive disorder and correlation analysis. *Psychiatry Clin Neurosci* **76**:321-328.

Tamoutounour S, Henri S, Lelouard H, de Bovis B, de Haar C, van der Woude CJ, Woltman AM, Reyat Y, Bonnet D, Sichien D, Bain CC, Mowat AM, Reiss e Sousa C, Poulin LF, Malissen B and Guilliams M (2012) CD64 distinguishes macrophages from dendritic cells in the gut and reveals the Th1-inducing role of mesenteric lymph node macrophages during colitis. *Eur J Immunol* **42**:3150-3166.

Tayab MA, Islam MN, Chowdhury KAA and Tasnim FM (2022) Targeting neuroinflammation by polyphenols: A promising therapeutic approach against inflammation-associated depression. *Biomed Pharmacother* **147**:112668.

Tian P, Chen Y, Zhu H, Wang L, Qian X, Zou R, Zhao J, Zhang H, Qian L, Wang Q, Wang G and Chen W (2022) Bifidobacterium breve CCFM1025 attenuates major depression disorder via regulating gut microbiome and tryptophan metabolism: A randomized clinical trial. *Brain Behav Immun* **100**:233-241.

Ueda Y, Kayama H, Jeon SG, Kusu T, Isaka Y, Rakugi H, Yamamoto M and Takeda K (2010) Commensal microbiota induce LPS hyporesponsiveness in colonic macrophages via the production of IL-10. *Int Immunol* **22**:953-962.

Vavassori P, Mencarelli A, Renga B, Distrutti E and Fiorucci S (2009) The bile acid receptor FXR is a modulator of intestinal innate immunity. *J Immunol* **183**:6251-6261.

- Veldhoen M and Brucklacher-Waldert V (2012) Dietary influences on intestinal immunity. *Nat Rev Immunol* **12**:696-708.
- Venkatesh M, Mukherjee S, Wang H, Li H, Sun K, Benechet AP, Qiu Z, Maher L, Redinbo MR, Phillips RS, Fleet JC, Kortagere S, Mukherjee P, Fasano A, Le Ven J, Nicholson JK, Dumas ME, Khanna KM and Mani S (2014) Symbiotic bacterial metabolites regulate gastrointestinal barrier function via the xenobiotic sensor PXR and Toll-like receptor 4. *Immunity* **41**:296-310.
- Vitale S, Picascia S and Gianfrani C (2016) The cross-talk between enterocytes and intraepithelial lymphocytes. *Mol Cell Pediatr* **3**:20.
- Vivier E, Artis D, Colonna M, Diefenbach A, Di Santo JP, Eberl G, Koyasu S, Locksley RM, McKenzie ANJ, Mebius RE, Powrie F and Spits H (2018) Innate Lymphoid Cells: 10 Years On. *Cell* **174**:1054-1066.
- Wang X, Qi Y and Zheng H (2022) Dietary Polyphenol, Gut Microbiota, and Health Benefits. *Antioxidants (Basel)* **11**.
- Wosen JE, Mukhopadhyay D, Macaubas C and Mellins ED (2018) Epithelial MHC Class II Expression and Its Role in Antigen Presentation in the Gastrointestinal and Respiratory Tracts. *Front Immunol* **9**:2144.
- Yang J, Zheng P, Li Y, Wu J, Tan X, Zhou J, Sun Z, Chen X, Zhang G, Zhang H, Huang Y, Chai T, Duan J, Liang W, Yin B, Lai J, Huang T, Du Y, Zhang P, Jiang J, Xi C, Wu L, Lu J, Mou T, Xu Y, Perry SW, Wong ML, Licinio J, Hu S, Wang G and Xie P (2020) Landscapes of bacterial and metabolic signatures and their interaction in major depressive disorders. *Sci Adv* **6**.
- Yin J, Li S, Li J, Gong R, Jia Z, Liu J, Jin Z, Yang J and Liu Y (2023) Association of serum oleic acid level with depression in American adults: a cross-sectional study. *BMC Psychiatry* **23**:845.

Yip JLK, Balasuriya GK, Spencer SJ and Hill-Yardin EL (2021) The Role of Intestinal Macrophages in Gastrointestinal Homeostasis: Heterogeneity and Implications in Disease. *Cell Mol Gastroenterol Hepatol* **12**:1701-1718.

Yoshii K, Hosomi K, Sawane K and Kunisawa J (2019) Metabolism of Dietary and Microbial Vitamin B Family in the Regulation of Host Immunity. *Front Nutr* **6**:48.

Young LM, Pipingas A, White DJ, Gauci S and Scholey A (2019) A Systematic Review and Meta-Analysis of B Vitamin Supplementation on Depressive Symptoms, Anxiety, and Stress: Effects on Healthy and 'At-Risk' Individuals. *Nutrients* **11**.

Yuan Q and Walker WA (2004) Innate immunity of the gut: mucosal defense in health and disease. *J Pediatr Gastroenterol Nutr* **38**:463-473.

Zhang JC, Yao W, Dong C, Yang C, Ren Q, Ma M, Han M, Wu J, Ushida Y, Suganuma H and Hashimoto K (2017) Prophylactic effects of sulforaphane on depression-like behavior and dendritic changes in mice after inflammation. *J Nutr Biochem* **39**:134-144.

Zhang X, Chen S, Zhang M, Ren F, Ren Y, Li Y, Liu N, Zhang Y, Zhang Q and Wang R (2021) Effects of Fermented Milk Containing *Lactocaseibacillus paracasei* Strain Shirota on Constipation in Patients with Depression: A Randomized, Double-Blind, Placebo-Controlled Trial. *Nutrients* **13**.

Zheng D, Liwinski T and Elinav E (2020) Interaction between microbiota and immunity in health and disease. *Cell Res* **30**:492-506.

FOOTNOTES

Financial support: This work was funded by the Spanish State Research Agency (AEI; Spain; PID2022-142106OA-I00) and the Miguel Servet program from Carlos III Health Institute (ISCIII; Spain; CP22/00030), co-funded by the European Union, and supported by the EarlyCause project of the European Union's Horizon 2020 research and innovation program (grant n° 848158), the European Commission – NextGenerationEU, through the CSIC Interdisciplinary Thematic Platform (PTI+) NEURO-AGING+ (PTI-NEURO-AGING+).

Conflicts of interest: The authors declare no conflicts of interest, financial or otherwise.

Reprint requests:

Eva M. Medina-Rodríguez, PhD

Psychiatry Service

Doctor Peset University Hospital, FISABIO

Av. Gaspar Aguilar, 90

46017, Valencia, Spain

Tel.: +34 963 900 022

Email: eva.medina@iata.csic.es

Numbered Footnotes:

¹ Psychiatry Service, Doctor Peset University Hospital. FISABIO, Valencia, Spain.

² Institute of Agrochemistry and Food Technology, Spanish National Research Council (IATA-CSIC), Valencia, Spain.

Figure 1




Figure 2

